mathematics

MDPI

Article

Exponential Convergence-(t,s)-Weak Tractability of Approximation in Weighted Hilbert Spaces

Huichao Yan [1] and Jia Chen [2,*]

[1] School of Computer and Network Engineering, Shanxi Datong University, Datong 037009, China; yanhuichao@sxdtdx.edu.cn
[2] School of Mathematics and Statistics, Shanxi Datong University, Datong 037009, China
* Correspondence: jiachen@sxdtdx.edu.cn

Abstract: We study L_2-approximation problems in the weighted Hilbert spaces in the worst case setting. Three interesting weighted Hilbert spaces appear in this paper, whose weights are equipped with two positive parameters γ_j and α_j for $j \in \mathbb{N}$. We consider algorithms using the class of arbitrary linear functionals. We discuss the exponential convergence-(t,s)-weak tractability of these L_2-approximation problems. In particular, we obtain the sufficient and necessary conditions on the weights for exponential convergence-weak tractability and exponential convergence-$(t,1)$-weak tractability with $t < 1$.

Keywords: L_2-approximation; information complexity; tractability; weighted Hilbert spaces

MSC: 41A81; 47A58; 47B02

1. Introduction

We study multivariate approximation problems $\mathrm{APP} = \{\mathrm{APP}_d\}_{d\in\mathbb{N}}$ of functions defined over Hilbert spaces with large or huge d in the worst case setting (approximation error by the worst case error). Such problems appear in quantum physics (see [1]), computational chemistry (see [2]), and economics (see [3]). We consider algorithms using the class of arbitrary linear functionals. The information complexity $n(\varepsilon, \mathrm{APP}_d)$ is the minimal number n of linear functionals for which the approximation error of some algorithm is at most ε. Tractability describes the dependence of the information complexity $n(\varepsilon, \mathrm{APP}_d)$ on the threshold ε and the dimension d. We consider the classical tractability which is polynomially convergent, and the exponential convergence-tractability (EC-tractability) which is exponentially convergent. Recently many authors discuss classical tractability and EC-tractability in weighted Hilbert spaces (see [4] by linear information, ref. [5] by standard information for functionals, and [6] by standard information for operators), especially in analytic Korobov spaces, such as exponential convergence and uniform exponential convergence (see [7]), classical tractability (see [8]) and EC-tractability for L_2-approximation (see [9] for exponential convergence-(t,s)-weak tractability and [10] for other EC-tractability results by algorithms using continuous linear functionals, and see [11] for EC-tractability by algorithms using function values), and EC-tractability for L_p-approximation with $1 \le p \le \infty$ by algorithms using continuous linear functionals (see [12]). Some authors consider tractability in weighted Hilbert spaces, such as classical tractability in weighted Korobov spaces (see [13] for strong polynomial tractability and polynomial tractability, [14] for other classical tractability results by algorithms using continuous linear functionals, and [15] by algorithms using function values), EC-tractability in weighted Korobov spaces (see [16]), and classical tractability in weighted Gaussian ANOVA spaces (see [17,18] with different weights, respectively).

In this paper, we investigate EC-tractability of L_2-approximation problems from the weighted Hilbert spaces with some weights. Let $H(K_{R_{d,\boldsymbol{\alpha},\gamma}})$ be a Hilbert space with

check for updates

Citation: Yan, H.; Chen, J. Exponential Convergence-(t,s)-Weak Tractability of Approximation in Weighted Hilbert Spaces. *Mathematics* **2024**, *12*, 2067. https://doi.org/10.3390/math12132067

Academic Editors: Paola Lamberti and Incoronata Notarangelo

Received: 16 May 2024
Revised: 27 June 2024
Accepted: 30 June 2024
Published: 1 July 2024

Copyright: © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https://creativecommons.org/licenses/by/4.0/).

weight $R_{d,\boldsymbol{\alpha},\gamma}$, where $\gamma = \{\gamma_j\}_{j\in\mathbb{N}}$ and $\boldsymbol{\alpha} = \{\alpha_j\}_{j\in\mathbb{N}}$ are two positive sequences satisfying $1 \geq \gamma_1 \geq \gamma_2 \geq \cdots \geq 0$ and $1 < \alpha_1 \leq \alpha_2 \leq \cdots$. In the worst case setting, we consider the L_2-approximation problem

$$\mathrm{APP}_d : H(K_{R_{d,\boldsymbol{\alpha},\gamma}}) \to L_2([0,1]^d) \ \text{ with } \ \mathrm{APP}_d(f) = f.$$

The classical tractability for L_2-approximation problem $\mathrm{APP} = \{\mathrm{APP}_d\}$ in weighted Korobov spaces $H(K_{R_{d,\boldsymbol{\alpha},\gamma}})$ such as strong polynomial tractability and polynomial tractability were discussed in [13,15,17]; quasi-polynomial tractability, uniform weak tractability, weak tractability and (t,s)-weak tractability were investigated in [14,17]. Additionally, ref. [17] also discussed classical tractability in several weighted Hilbert spaces, including weighted Korobov spaces and weighted Gaussian ANOVA spaces. The EC-tractability of the problem $\mathrm{APP} = \{\mathrm{APP}_d\}$ in weighted Korobov spaces such as EC-$(t,1)$-weak tractability for $0 < t \leq 1$ were studied in [16]. However, the above weighted Hilbert spaces $H(K_{R_{d,\boldsymbol{\alpha},\gamma}})$ with weights $R_{d,\boldsymbol{\alpha},\gamma}$ satisfy $1 \geq \gamma_1 \geq \gamma_2 \geq \cdots \geq 0$ and $1 < \alpha_1 = \alpha_2 = \cdots$.

In this paper we present three cases of weighted Hilbert spaces $H(K_{R_{d,\boldsymbol{\alpha},\gamma}})$ with weights $R_{d,\boldsymbol{\alpha},\gamma}$ for $1 \geq \gamma_1 \geq \gamma_2 \geq \cdots \geq 0$ and $1 < \alpha_1 \leq \alpha_2 \leq \cdots$ that appear in the reference [18]. These weighted Hilbert spaces are similar but also different. The authors in [18] studied the polynomial tractability, strong polynomial tractability, weak tractability, and (t,s)-weak tractability for $t > 1$ and $s > 0$ of the problems $\mathrm{APP} = \{\mathrm{APP}_d\}$ in these three weighted Hilbert spaces. However, there are no results about EC-tractability of the approximation problems $\mathrm{APP} = \{\mathrm{APP}_d\}$ in the above three weighted Hilbert spaces. We will study exponential convergence-(t,s)-weak tractability (EC-(t,s)-WT) for some $t > 0$, $s > 0$ and obtain the complete sufficient and necessary conditions for $t = s = 1$ and $t < 1$, $s = 1$, respectively.

The paper is structured in the following ways. We present three cases of weighted Hilbert spaces in Section 2. Section 3 gives preliminaries about the L_2-approximation problem in the weighted Hilbert space. Section 4.1 is devoted to recall some notions about the tractability, such as classical tractability and exponential convergence-tractability and state the main results. In Section 4.2 we give the proof of Theorem 1. In Section 5 we present a summary.

2. Weighted Reproducing Kernel Hilbert Spaces

In this section we consider weighted reproducing kernel Hilbert spaces with different weights.

Let $H(K_d)$ be a Hilbert space defined in $[0,1]^d$. The function $K_d(\boldsymbol{x},\boldsymbol{y})$ of $\boldsymbol{x},\boldsymbol{y} \in [0,1]^d$ is called a reproducing kernel of $H(K_d)$ if for every $\boldsymbol{y} \in [0,1]^d$ and every $f \in H(K_d)$,

$$f(\boldsymbol{y}) = \langle f(\boldsymbol{x}), K_d(\boldsymbol{x},\boldsymbol{y})\rangle_{H(K_d)}.$$

The Hilbert space is a so-called reproducing kernel Hilbert space. We can study more details on reproducing kernel Hilbert spaces in the reference [19].

In this paper, let $\gamma = \{\gamma_j\}_{j\in\mathbb{N}}$ and $\boldsymbol{\alpha} = \{\alpha_j\}_{j\in\mathbb{N}}$ be two positive sequences of the Hilbert space $H(K_{R_{d,\boldsymbol{\alpha},\gamma}})$ with $R_{d,\boldsymbol{\alpha},\gamma}$ satisfying

$$1 \geq \gamma_1 \geq \gamma_2 \geq \cdots \geq 0, \ \text{ and } \ 1 < \alpha_1 \leq \alpha_2 \leq \cdots. \tag{1}$$

Assume that the function $K_{R_{d,\boldsymbol{\alpha},\gamma}}$ of the space $H(K_{R_{d,\boldsymbol{\alpha},\gamma}})$ with $K_{R_{d,\boldsymbol{\alpha},\gamma}} : [0,1]^d \times [0,1]^d \to \mathbb{C}$ is of product form

$$K_{R_{d,\boldsymbol{\alpha},\gamma}}(\boldsymbol{x},\boldsymbol{y}) := \prod_{k=1}^{d} K_{R_{1,\alpha_k,\gamma_k}}(x_k, y_k),$$

where $K_{R_{1,\alpha,\gamma}} : [0,1] \times [0,1] \to \mathbb{C}$ is a universal weighted function

$$K_{R_{1,\alpha,\gamma}}(x,y) := \sum_{h\in\mathbb{N}_0} R_{\alpha,\gamma}(h)\exp(2\pi i h(x-y)), \ \ x, \ y \in [0,1].$$

Here, let weight $R_{\alpha,\gamma} : \mathbb{N}_0 \to \mathbb{R}^+$ be a summable function, i.e., $\sum_{k\in\mathbb{N}_0} R_{\alpha,\gamma}(k) < \infty$. Then we have

$$K_{R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}}(\boldsymbol{x},\boldsymbol{y}) = \sum_{\boldsymbol{h}\in\mathbb{N}_0^d} R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}(\boldsymbol{h}) \exp(2\pi i \boldsymbol{h}\cdot(\boldsymbol{x}-\boldsymbol{y})),\ \boldsymbol{x},\ \boldsymbol{y} \in [0,1]^d, \tag{2}$$

the inner product

$$\langle f, g\rangle_{H(K_{R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}})} = \sum_{\boldsymbol{h}\in\in\mathbb{N}_0^d} \frac{1}{R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}(\boldsymbol{h})} \widehat{f}(\boldsymbol{h})\overline{\widehat{g}}(\boldsymbol{h}), \tag{3}$$

and

$$||f||_{H(K_{R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}})} = \sqrt{\langle f, f\rangle_{H(K_{R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}})}},$$

where

$$R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}(\boldsymbol{h}) := \prod_{j=1}^{d} R_{\alpha_j,\gamma_j}(h_j),\ \boldsymbol{h} = (h_1, h_2, \ldots, h_d) \in \mathbb{N}_0^d,$$

$$\boldsymbol{x}\cdot\boldsymbol{y} := \sum_{h=1}^{d} x_h \cdot y_h,\ \ \boldsymbol{x} = (x_1, x_2, \ldots, x_d),\ \boldsymbol{y} = (y_1, y_2, \ldots, y_d) \in [0,1]^d,$$

and

$$\widehat{f}(\boldsymbol{h}) = \int_{[0,1]^d} f(\boldsymbol{x}) \exp(-2\pi i \boldsymbol{h}\cdot\boldsymbol{x}) d\boldsymbol{x}.$$

We can ascertain that $K_{R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}}(\boldsymbol{x},\boldsymbol{y})$ is well defined for $1 < \alpha_1 \le \alpha_2 \le \cdots$ and for all $\boldsymbol{x},\ \boldsymbol{y} \in [0,1]^d$, since

$$|K_{R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}}(\boldsymbol{x},\boldsymbol{y})| \le \sum_{\boldsymbol{h}\in\mathbb{N}_0^d} R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}(\boldsymbol{h}) = \prod_{j=1}^{d} \Big(\sum_{h\in\mathbb{N}_0} R_{\alpha_j,\gamma_j}(h)\Big) < \infty.$$

Note that the Hilbert space $H(K_{R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}})$ is a reproducing kernel Hilbert space with the reproducing kernel $K_{R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}}$. Indeed, for every $f \in H(K_{R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}})$ we have

$$f(\boldsymbol{y}) = \langle f(\boldsymbol{x}), K_{R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}}(\boldsymbol{x},\boldsymbol{y})\rangle_{H(K_{R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}})}.$$

The kernel $K_{R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}}$ with weight $R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}$ is called a weighted reproducing kernel and the space $H(K_{R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}})$ is called a weighted reproducing kernel Hilbert space. If $\gamma_1 = \gamma_2 = \cdots = 1$ and $1 < \alpha_1 = \alpha_2 = \cdots$, then the space $H(K_{R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}})$ is called unweighted space. Here, $\mathbb{N}_0 = \{0, 1, \ldots\}$ and $\mathbb{N} = \{1, 2, \ldots\}$.

There are many ways to introduce weighted reproducing kernel Hilbert spaces with weights $R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}$. In this paper we consider three weights like the cases in the reference [18].

2.1. A Weighted Korobov Space

Let $\boldsymbol{\alpha} = \{\alpha_j\}_{j\in\mathbb{N}}$ and $\boldsymbol{\gamma} = \{\gamma_j\}_{j\in\mathbb{N}}$ be two sequences satisfying (1). We consider a weighted Korobov space $H(K_{R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}})$ with weight

$$R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}(\boldsymbol{h}) = r_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}(\boldsymbol{h}) := \prod_{j=1}^{d} r_{\alpha_j,\gamma_j}(h_j),$$

where

$$r_{\alpha,\gamma}(h) := \begin{cases} 1, & \text{for } h = 0, \\ \frac{\gamma}{h^{|\alpha|}}, & \text{for } h \ge 1 \end{cases}$$

for $\alpha > 1$ and $\gamma \in (0,1]$. We can see the case in the references [18,20]. Then we have the kernel function (2) with

$$K_{R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}}(\boldsymbol{x},\boldsymbol{y}) = K_{r_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}}(\boldsymbol{x},\boldsymbol{y}) = \sum_{\boldsymbol{h}\in\mathbb{N}_0^d} r_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}(\boldsymbol{h}) \exp(2\pi i \boldsymbol{h}\cdot(\boldsymbol{x}-\boldsymbol{y}))$$

for $\boldsymbol{x},\ \boldsymbol{y} \in [0,1]^d$, and the inner product (3) with

$$\langle f,g\rangle_{H(K_{R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}})} = \langle f,g\rangle_{H(K_{r_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}})} = \sum_{\boldsymbol{h}\in\in\mathbb{N}_0^d} \frac{1}{r_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}(\boldsymbol{h})} \widehat{f}(\boldsymbol{h})\overline{\widehat{g}}(\boldsymbol{h}).$$

Remark 1. *Obviously, the kernel $K_{r_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}}(\boldsymbol{x},\boldsymbol{y})$ is well defined for $\boldsymbol{\alpha}$ and $\boldsymbol{\gamma}$ satisfying (1), due to*

$$|K_{r_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}}(\boldsymbol{x},\boldsymbol{y})| \le \sum_{\boldsymbol{k}\in\mathbb{N}_0^d} r_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}(\boldsymbol{k}) = \prod_{j=1}^{d}(1+\zeta(\lceil\alpha_j\rceil)\gamma_j) < \infty,$$

where $\zeta(\cdot)$ is the Riemann zeta function.

2.2. A First Variant of the Weighted Korobov Space

Let $\boldsymbol{\alpha} = \{\alpha_j\}_{j\in\mathbb{N}}$ and $\boldsymbol{\gamma} = \{\gamma_j\}_{j\in\mathbb{N}}$ be two sequences satisfying (1). We discuss a first variant of the weighted Korobov space with weight

$$R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}(\boldsymbol{h}) = \psi_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}(\boldsymbol{h}) := \prod_{j=1}^{d} \psi_{\alpha_j,\gamma_j}(h_j),$$

where

$$\psi_{\alpha,\gamma}(h) := \begin{cases} 1, & \text{for } h = 0, \\ \frac{\gamma}{h!}, & \text{for } 1 \le h < \lceil\alpha\rceil, \\ \frac{\gamma(h-\lceil\alpha\rceil)!}{h!}, & \text{for } h \ge \lceil\alpha\rceil \end{cases}$$

for $\alpha > 1$ and $\gamma \in (0,1]$.

Then we have the kernel function (2) with

$$K_{R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}}(\boldsymbol{x},\boldsymbol{y}) = K_{\psi_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}}(\boldsymbol{x},\boldsymbol{y}) = \sum_{\boldsymbol{h}\in\mathbb{N}_0^d} \psi_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}(\boldsymbol{h}) \exp(2\pi i \boldsymbol{h}\cdot(\boldsymbol{x}-\boldsymbol{y}))$$

for $\boldsymbol{x},\ \boldsymbol{y} \in [0,1]^d$ and the inner product (3) with

$$\langle f,g\rangle_{H(K_{R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}})} = \langle f,g\rangle_{H(K_{\psi_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}})} = \sum_{\boldsymbol{h}\in\in\mathbb{N}_0^d} \frac{1}{\psi_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}(\boldsymbol{h})} \widehat{f}(\boldsymbol{h})\overline{\widehat{g}}(\boldsymbol{h}).$$

Lemma 1 ([18] Lemma 2)**.** *For all $j,\ k \in \mathbb{N}$ we have*

$$\psi_{\alpha_j,\gamma_j}(k) \le \lceil\alpha_j\rceil^{\lceil\alpha_j\rceil} r_{\alpha_j,\gamma_j}(k).$$

Remark 2. *From Lemma 1 and $1 < \alpha_1 \le \alpha_2 \le \cdots$ we get*

$$\begin{aligned} |K_{\psi_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}}(\boldsymbol{x},\boldsymbol{y})| &\le \sum_{\boldsymbol{k}\in\mathbb{N}_0^d} \psi_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}(\boldsymbol{k}) = \prod_{j=1}^{d}(1+\sum_{k\in\mathbb{N}} \psi_{\alpha_j,\gamma_j}(k)) \\ &\le \prod_{j=1}^{d}(1+\sum_{k\in\mathbb{N}} \lceil\alpha_j\rceil^{\lceil\alpha_j\rceil} r_{\alpha_j,\gamma_j}(k)) \\ &= \prod_{j=1}^{d}(1+\lceil\alpha_j\rceil^{\lceil\alpha_j\rceil}\zeta(\lceil\alpha_j\rceil)\gamma_j) \\ &< \infty. \end{aligned}$$

Hence, the kernel $K_{\psi_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}}(\boldsymbol{x},\boldsymbol{y})$ is well defined.

2.3. A Second Variant of the Weighted Korobov Space

Let $\boldsymbol{\alpha} = \{\alpha_j\}_{j\in\mathbb{N}}$ and $\boldsymbol{\gamma} = \{\gamma_j\}_{j\in\mathbb{N}}$ be two sequences satisfying (1). We study a second variant of the weighted Korobov space $H(K_{R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}})$ (see the references [18,21]) with weight

$$R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}(\boldsymbol{h}) = \omega_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}(\boldsymbol{h}) := \prod_{j=1}^{d} \omega_{\alpha_j,\gamma_j}(h_j),$$

where

$$\omega_{\alpha,\gamma}(h) := \left(1 + \frac{1}{\gamma}\sum_{l=1}^{\lceil\alpha\rceil}\theta_l(h)\right)^{-1}$$

for $\alpha > 1$ and $\gamma \in (0,1]$ and

$$\theta_l(h) := \begin{cases} \frac{h!}{(h-l)!}, & \text{for } h \geq l, \\ 0, & \text{for } 0 \leq h < l. \end{cases}$$

Then we have the kernel function (2) with

$$K_{R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}}(\boldsymbol{x},\boldsymbol{y}) = K_{\omega_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}}(\boldsymbol{x},\boldsymbol{y}) = \sum_{\boldsymbol{h}\in\mathbb{N}_0^d} \omega_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}(\boldsymbol{h}) \exp(2\pi i \boldsymbol{h}\cdot(\boldsymbol{x}-\boldsymbol{y}))$$

for $\boldsymbol{x},\ \boldsymbol{y} \in [0,1]^d$, and the inner product (3) with

$$\langle f,g\rangle_{H(K_{R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}})} = \langle f,g\rangle_{H(K_{\omega_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}})} = \sum_{\boldsymbol{h}\in\in\mathbb{N}_0^d} \frac{1}{\omega_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}(\boldsymbol{h})} \widehat{f}(\boldsymbol{h})\overline{\widehat{g}}(\boldsymbol{h}).$$

Lemma 2 ([18] Lemma 3)**.** *For all $j,\ k \in \mathbb{N}$ we have*

$$\omega_{\alpha_j,\gamma_j}(k) \leq \lceil\alpha_j\rceil^{\lceil\alpha_j\rceil} r_{\alpha_j,\gamma_j}(k).$$

Remark 3. *We note that the kernel $K_{R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}}(\boldsymbol{x},\boldsymbol{y})$ is also well defined. Indeed, it follows from Lemma 2 and $1 < \alpha_1 \leq \alpha_2 \leq \cdots$ that*

$$\begin{aligned} |K_{\omega_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}}(\boldsymbol{x},\boldsymbol{y})| &\leq \sum_{\boldsymbol{k}\in\mathbb{N}_0^d} \omega_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}(\boldsymbol{k}) = \prod_{j=1}^{d}(1 + \sum_{k\in\mathbb{N}} \omega_{\alpha_j,\gamma_j}(k)) \\ &\leq \prod_{j=1}^{d}(1 + \sum_{k\in\mathbb{N}} \lceil\alpha_j\rceil^{\lceil\alpha_j\rceil} r_{\alpha_j,\gamma_j}(k)) \\ &= \prod_{j=1}^{d}(1 + \lceil\alpha_j\rceil^{\lceil\alpha_j\rceil}\zeta(\lceil\alpha_j\rceil)\gamma_j) \\ &< \infty. \end{aligned}$$

Lemma 3. *Let $R_{\alpha_j,\gamma_j} \in \left\{r_{\alpha_j,\gamma_j}, \psi_{\alpha_j,\gamma_j}, \omega_{\alpha_j,\gamma_j}\right\}$ for all $j \in \mathbb{N}$. Then we have for all $j \in \mathbb{N}$, $k \in \mathbb{N}_0$*

$$R_{\alpha_j,\gamma_j}(k) \leq \lceil\alpha_j\rceil^{\lceil\alpha_j\rceil} r_{\alpha_j,\gamma_j}(k).$$

Particularly, we have for all $j \in \mathbb{N}$, $k \in \mathbb{N}_0$

$$R_{\alpha_j,\gamma_j}(k) \leq \lceil\alpha_1\rceil^{\lceil\alpha_1\rceil} r_{\alpha_1,\gamma_j}(k).$$

Proof. On the one hand, it is obvious from Lemmas 1 and 2 that

$$R_{\alpha_j,\gamma_j}(k) \le \lceil \alpha_j \rceil^{\lceil \alpha_j \rceil} r_{\alpha_j,\gamma_j}(k) \tag{4}$$

for all j, $k \in \mathbb{N}$. Since for all $j \in \mathbb{N}$

$$r_{\alpha_j,\gamma_j}(0) = \psi_{\alpha_j,\gamma_j}(0) = \omega_{\alpha_j,\gamma_j}(0) = 1,$$

we have

$$R_{\alpha_j,\gamma_j}(0) = 1 \le \lceil \alpha_j \rceil^{\lceil \alpha_j \rceil} = \lceil \alpha_j \rceil^{\lceil \alpha_j \rceil} r_{\alpha_j,\gamma_j}(0).$$

Thus we have for all $j \in \mathbb{N}$, $k \in \mathbb{N}_0$ that

$$R_{\alpha_j,\gamma_j}(k) \le \lceil \alpha_j \rceil^{\lceil \alpha_j \rceil} r_{\alpha_j,\gamma_j}(k).$$

On the other hand, noting for all j, $k \in \mathbb{N}$

$$r_{\alpha_j,\gamma_j}(k) \le r_{\alpha_1,\gamma_j}(k), \;\; \psi_{\alpha_j,\gamma_j}(k) \le \psi_{\alpha_1,\gamma_j}(k), \;\; \omega_{\alpha_j,\gamma_j}(k) \le \omega_{\alpha_1,\gamma_j}(k),$$

and for all $j \in \mathbb{N}$

$$r_{\alpha_j,\gamma_j}(0) = r_{\alpha_1,\gamma_j}(0) = 1, \;\; \psi_{\alpha_j,\gamma_j}(0) = \psi_{\alpha_1,\gamma_j}(0) = 1, \;\; \omega_{\alpha_j,\gamma_j}(0) = \omega_{\alpha_1,\gamma_j}(0) = 1,$$

we have for all $j \in \mathbb{N}$, $k \in \mathbb{N}_0$ that

$$R_{\alpha_j,\gamma_j}(k) \le R_{\alpha_1,\gamma_j}(k).$$

Hence, by (4) we further get for all $j \in \mathbb{N}$, $k \in \mathbb{N}_0$ that

$$R_{\alpha_j,\gamma_j}(k) \le R_{\alpha_1,\gamma_j}(k) \le \lceil \alpha_1 \rceil^{\lceil \alpha_1 \rceil} r_{\alpha_1,\gamma_j}(k).$$

□

Remark 4. *Let* $R_{\alpha_j,\gamma_j} \in \left\{ r_{\alpha_j,\gamma_j}, \psi_{\alpha_j,\gamma_j}, \omega_{\alpha_j,\gamma_j} \right\}$ *for all* $j \in \mathbb{N}$. *Then we obtain*

$$R_{\alpha_j,\gamma_j}(0) = 1 \quad \textit{and} \quad R_{\alpha_j,\gamma_j}(1) \ge \frac{\gamma_j}{2} \tag{5}$$

for all $j \in \mathbb{N}$. *Indeed, for all* $j \in \mathbb{N}$ *we have*

$$\psi_{\alpha_j,\gamma_j}(0) = r_{\alpha_j,\gamma_j}(0) = \omega_{\alpha_j,\gamma_j}(0) = 1,$$

which means $R_{\alpha_j,\gamma_j}(0) = 1$. *As a result of all* $j \in \mathbb{N}$, *we get*

$$\psi_{\alpha_j,\gamma_j}(1) = r_{\alpha_j,\gamma_j}(1) = \gamma_j \quad \textit{and} \quad \omega_{\alpha_j,\gamma_j}(1) = \left(1 + \frac{1}{\gamma_j}\right)^{-1} \ge \frac{\gamma_j}{2},$$

which yields $R_{\alpha_j,\gamma_j}(1) \ge \frac{\gamma_j}{2}$.

3. L_2-Approximation in the Weighted Hilbert Spaces

In this paper we investigate the L_2-approximation $\mathrm{APP}_d : H(K_{R_{d,\boldsymbol{\alpha},\gamma}}) \to L_2([0,1]^d)$ given by

$$\mathrm{APP}_d(f) = f \;\; \text{for all} \;\; f \in H(K_{R_{d,\boldsymbol{\alpha},\gamma}})$$

in weighted Hilbert space $H(K_{R_{d,\boldsymbol{\alpha},\gamma}})$ with weight $R_{d,\boldsymbol{\alpha},\gamma} \in \{r_{d,\boldsymbol{\alpha},\gamma}, \psi_{d,\boldsymbol{\alpha},\gamma}, \omega_{d,\boldsymbol{\alpha},\gamma}\}$. We note from Remarks 1–3, and [15] that this L_2-approximation is compact for $1 < \alpha_1 \le \alpha_2 \le \cdots$.

We approximate APP_d by using the algorithm $A_{n,d}$ of the form

$$A_{n,d}(f) = \sum_{i=1}^{n} T_i(f) g_i, \text{ for } f \in H(K_{R_{d,\boldsymbol{\alpha},\gamma}}), \tag{6}$$

where $g_1, g_2, \ldots, g_n$ belong to $L_2([0,1]^d)$ and $T_1, T_2, \ldots, T_n$ are continuous linear functionals on $H(K_{R_{d,\boldsymbol{\alpha},\gamma}})$.

We consider the worst case setting in which the error of the algorithm $A_{n,d}$ of the form (6) is defined as

$$e(A_{n,d}) := \sup_{||f||_{H(K_{R_{d,\boldsymbol{\alpha},\gamma}})} \le 1} ||\mathrm{APP}_d(f) - A_{n,d}(f)||_{L_2}.$$

The error $e(A_{n,d})$ is also called the worst case error. The nth minimal worst case error is defined as

$$e(n, \mathrm{APP}_d) := \inf_{A_{n,d}} e(A_{n,d}) \text{ for } n \ge 1,$$

which is the infimum error among all algorithms (6). For $n = 0$, we set $A_{0,d} = 0$. We call

$$e(0, \mathrm{APP}_d) = \sup_{||f||_{H(K_{R_{d,\boldsymbol{\alpha},\gamma}})} \le 1} ||\mathrm{APP}_d(f)||_{L_2}$$

the initial error of the problem APP_d.

We are interested in how the worst case error for the algorithm $A_{n,d}$ depends on the numbers n and d. We define the information complexity as

$$n(\varepsilon, \mathrm{APP}_d) := \min\{n \in \mathbb{N}_0 : e(n, \mathrm{APP}_d) \le \varepsilon\},$$

where $\varepsilon \in (0,1)$ and $d \in \mathbb{N}$. In this paper, we set $\mathbb{N}_0 = \{0, 1, \ldots\}$ and $\mathbb{N} = \{1, 2, \ldots\}$.

By the references [2,4] we know that the nth minimal worst case errors $e(n, \mathrm{APP}_d)$ and the information complexity $n(\varepsilon, \mathrm{APP}_d)$ are related to the eigenvalues of the continuously linear operator $W_d = \mathrm{APP}_d^* \mathrm{APP}_d : H(K_{R_{d,\boldsymbol{\alpha},\gamma}}) \to H(K_{R_{d,\boldsymbol{\alpha},\gamma}})$, where APP_d^* is the operator dual to APP_d. The eigenvalues of W_d are denoted by $\{\lambda_{d,j}\}_{j\in\mathbb{N}}$ satisfying

$$\lambda_{d,1} \ge \lambda_{d,2} \ge \cdots \ge 0$$

and the corresponding orthogonal eigenvectors of $\{\lambda_{d,j}\}_{j\in\mathbb{N}}$ by $\{\eta_{d,j}\}_{j\in\mathbb{N}}$ satisfying

$$\langle \eta_{d,i}, \eta_{d,j} \rangle_{H(K_{R_{d,\boldsymbol{\alpha},\gamma}})} = \delta_{i,j}, \text{ for all } i, j \in \mathbb{N},$$

where

$$W_d \eta_{d,j} = \lambda_{d,j} \eta_{d,j}, \text{ for all } j \in \mathbb{N}.$$

Here $\delta_{i,j} = 1$ for $i = j$ and $\delta_{i,j} = 0$ for $i \ne j$. Then the nth minimal worst case error is attained for the algorithm

$$A_{n,d}^{\diamond} f = \sum_{i=1}^{n} \langle f, \eta_{d,i} \rangle_{H(K_{R_{d,\boldsymbol{\alpha},\gamma}})} \eta_{d,i}, \text{ for all } n \in \mathbb{N}$$

and

$$e(n, \mathrm{APP}_d) = e(A_{n,d}^{\diamond}) = \sqrt{\lambda_{d,n+1}}, \text{ for all } n \in \mathbb{N}.$$

The initial error $e(0, \mathrm{APP}_d) = \sqrt{\lambda_{d,1}}$. Hence, we have $e(n, \mathrm{APP}_d) = \sqrt{\lambda_{d,n+1}}$ for all $n \in \mathbb{N}_0$. This deduces that the information complexity is equal to

$$n(\varepsilon, \mathrm{APP}_d) = \min\left\{n \in \mathbb{N}_0 : \sqrt{\lambda_{d,n+1}} \le \varepsilon\right\} = \min\left\{n \in \mathbb{N}_0 : \lambda_{d,n+1} \le \varepsilon^2\right\}. \tag{7}$$

Since the eigenvalues $\lambda_{d,j}$ with $j \in \mathbb{N}$ of the operator W_d are $R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}(\boldsymbol{k})$ with $\boldsymbol{k} \in \mathbb{N}_0^d$ (see [4] p. 215), by (7) the information complexity of APP_d from the space $H(K_{R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}})$ is equal to

$$n(\varepsilon, \mathrm{APP}_d) = \min\left\{n \in \mathbb{N}_0 : \lambda_{d,n+1} \leq \varepsilon^2\right\} = \left|\left\{n \in \mathbb{N} : \lambda_{d,n} > \varepsilon^2\right\}\right|$$
$$= \left|\left\{\boldsymbol{h} \in \mathbb{N}_0^d : R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}(\boldsymbol{h}) > \varepsilon^2\right\}\right| = \left|\left\{\boldsymbol{h} \in \mathbb{N}_0^d : \prod_{j=1}^{d} R_{\alpha_j,\gamma_j}(h_j) > \varepsilon^2\right\}\right|, \quad (8)$$

with $\varepsilon \in (0,1)$ and $d \in \mathbb{N}$, where $|A|$ denotes the cardinality of set A.

Note that for the L_2-approximation APP_d from the space $H(K_{R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}})$ the absolute error criterion and the normalized error criterion are the same, since the initial error $e(0, \mathrm{APP}_d) = \sqrt{\lambda_{d,1}} = 1$.

4. Tractability in Weighted Hilbert Spaces and Main Results

In this paper we will study the classical tractability and the exponential convergence-tractability (EC-tractability) for the problem $\mathrm{APP} = \{\mathrm{APP}_d\}_{d\in\mathbb{N}}$ in the weighted Hilbert space $H_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}$.

4.1. Tractability and Main Results

We focus on the behaviours of the information complexity $n(\varepsilon, \mathrm{APP}_d)$ depending on the dimension d and the error threshold ε. Hence, we will study several notions about the classical tractability and the exponential convergence-tractability (EC-tractability) notions (see [4–9,11,12,16,22]).

Definition 1. *Let $APP = \{APP_d\}_{d\in\mathbb{N}}$. We say the following:*

- *Strong polynomial tractability (SPT) if there are positive numbers C and p such that*

$$n(\varepsilon, APP_d) \leq C(\varepsilon^{-1})^p \quad \text{for all} \quad d \in \mathbb{N}, \varepsilon \in (0,1).$$

In this case we define the exponent p^{str} of SPT as

$$p^{str} := \inf\{p : \exists\, C > 0 \text{ such that } n(\varepsilon, APP_d) \leq C(\varepsilon^{-1})^p, \forall\, d \in \mathbb{N}, \varepsilon \in (0,1)\}.$$

- *Polynomial tractability (PT) if there are positive numbers C, p, and q such that*

$$n(\varepsilon, APP_d) \leq C d^q (\varepsilon^{-1})^p \quad \text{for all} \quad d \in \mathbb{N}, \varepsilon \in (0,1).$$

- *Quasi-polynomial tractability (QPT) if there are positive numbers C and t such that*

$$n(\varepsilon, APP_d) \leq C \exp\left(t\left(1 + \ln d\right)\left(1 + \ln \varepsilon^{-1}\right)\right) \quad \text{for all} \quad d \in \mathbb{N}, \varepsilon \in (0,1).$$

- *Uniform weak tractability (UWT) if for all $t, s > 0$,*

$$\lim_{\varepsilon^{-1}+d\to\infty} \frac{\ln n(\varepsilon, APP_d)}{d^t + (\varepsilon^{-1})^s} = 0.$$

- *Weak tractability (WT) if*

$$\lim_{\varepsilon^{-1}+d\to\infty} \frac{\ln n(\varepsilon, APP_d)}{d + \varepsilon^{-1}} = 0.$$

- *(t,s)-weak tractability ((t,s)-WT) for fixed positive t and s if*

$$\lim_{\varepsilon^{-1}+d\to\infty} \frac{\ln n(\varepsilon, APP_d)}{d^t + (\varepsilon^{-1})^s} = 0.$$

We find that (1,1)-WT is the same as WT and

$$\text{SPT} \Longrightarrow \text{PT} \Longrightarrow \text{QPT} \Longrightarrow \text{UWT} \Longrightarrow \text{WT}.$$

In the above definitions regarding classical tractability, replacing ε^{-1} with $(1+\ln(\varepsilon^{-1}))$, we will have the following definitions about exponential convergence-tractability (EC-tractability).

Definition 2. *Let $APP = \{APP_d\}_{d\in\mathbb{N}}$. We say we have the following:*

- *Exponential convergence-strong polynomial tractability (EC-SPT) if there are positive numbers C and p such that*

$$n(\varepsilon, APP_d) \le C\big(1+\ln(\varepsilon^{-1})\big)^p \ \text{ for all } \ d \in \mathbb{N}, \varepsilon \in (0,1).$$

The exponent of EC-SPT is defined as

$$\inf\{p : \exists\, C>0 \text{ such that } n(\varepsilon, APP_d) \le C\big(1+\ln(\varepsilon^{-1})\big)^p, \forall\, d \in \mathbb{N}, \varepsilon \in (0,1)\}.$$

- *Exponential convergence-polynomial tractability (EC-PT) if there are positive numbers C, p, and q such that*

$$n(\varepsilon, APP_d) \le C d^q \big(1+\ln(\varepsilon^{-1})\big)^p \ \text{ for all } \ d \in \mathbb{N}, \varepsilon \in (0,1).$$

- *Exponential convergence-uniform weak tractability (EC-UWT) if for all $t, s > 0$*

$$\lim_{\varepsilon^{-1}+d\to\infty} \frac{\ln n(\varepsilon, APP_d)}{d^t + \big(1+\ln(\varepsilon^{-1})\big)^s} = 0.$$

- *Exponential convergence-weak tractability (EC-WT) if*

$$\lim_{\varepsilon^{-1}+d\to\infty} \frac{\ln n(\varepsilon, APP_d)}{d + \ln(\varepsilon^{-1})} = 0.$$

- *Exponential convergence-(t,s)-weak tractability (EC-(t,s)-WT) for fixed positive t and s if*

$$\lim_{\varepsilon^{-1}+d\to\infty} \frac{\ln n(\varepsilon, APP_d)}{d^t + \big(1+\ln(\varepsilon^{-1})\big)^s} = 0.$$

We note that EC-$(1,1)$-WT is the same as EC-WT, and

$$\text{EC-SPT} \Longrightarrow \text{EC-PT} \Longrightarrow \text{EC-QPT} \Longrightarrow \text{EC-UWT} \Longrightarrow \text{EC-WT}.$$

Obviously, if the problem APP has exponential convergence-tractability, then it has classical tractability and

$$\text{EC-}(t,s)\text{-WT} \Longrightarrow (t,s)\text{-WT}, \quad \text{EC-UWT} \Longrightarrow \text{UWT}, \quad \text{EC-WT} \Longrightarrow \text{WT}.$$

In the worst case setting the classical tractability and EC-tractability of the problem $\text{APP} = \{\text{APP}_d\}_{d\in\mathbb{N}}$ in the weighted Hilbert space $H(K_{R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}})$ with $\boldsymbol{\gamma} = \{\gamma_j\}_{j\in\mathbb{N}}$ and $\boldsymbol{\alpha} = \{\alpha_j\}_{j\in\mathbb{N}}$ satisfying

$$1 \ge \gamma_1 \ge \gamma_2 \ge \cdots \ge 0, \ \text{ and } \ 1 < \alpha^* = \alpha_1 = \alpha_2 = \cdots$$

have been solved by [13,14,16,18] as follows:

- For $R_{d,\alpha^*,\gamma} \in \{r_{d,\alpha^*,\gamma}, \psi_{d,\alpha^*,\gamma}, \omega_{d,\alpha^*,\gamma}\}$, SPT holds iff PT holds iff

$$s_\gamma := \inf\left\{\kappa > 0 : \sum_{j=1}^{\infty} \gamma_j^\kappa < \infty\right\} < \infty$$

and the exponent of SPT is

$$p^{\mathrm{str}} = 2\max\left(s_\gamma, \frac{1}{\alpha}\right).$$

- For $R_{d,\alpha^*,\gamma} = r_{d,\alpha^*,\gamma}$, QPT, UWT, and WT are equivalent and hold iff

$$\gamma_I := \inf_{j\in\mathbb{N}} \gamma_j < 1.$$

For $R_{d,\alpha^*,\gamma} \in \{\psi_{d,\alpha^*,\gamma}, \omega_{d,\alpha^*,\gamma}\}$,

$$\gamma_I < \infty$$

implies QPT.
- For $R_{d,\alpha^*,\gamma} \in \{r_{d,\alpha^*,\gamma}, \psi_{d,\alpha^*,\gamma}, \omega_{d,\alpha^*,\gamma}\}$ and $t > 1$, (t,s)-WT holds for all $1 \geq \gamma_1 \geq \gamma_2 \geq \cdots \geq 0$.
- For $R_{d,\alpha^*,\gamma} = r_{d,\alpha^*,\gamma}$, EC-WT holds iff

$$\lim_{j\to\infty} \gamma_j = 0.$$

- For $R_{d,\alpha^*,\gamma} = r_{d,\alpha^*,\gamma}$ and $t < 1$, EC-$(t,1)$-WT holds iff

$$\lim_{j\to\infty} \frac{\ln j}{\ln(\gamma_j^{-1})} = 0.$$

In the worst case setting the classical tractability such as SPT, PT, and WT of the problem APP $= \{\mathrm{APP}_d\}_{d\in\mathbb{N}}$ in the weighted Hilbert space $H(K_{R_{d,\boldsymbol{\alpha},\gamma}})$ with $\boldsymbol{\gamma} = \{\gamma_j\}_{j\in\mathbb{N}}$ and $\boldsymbol{\alpha} = \{\alpha_j\}_{j\in\mathbb{N}}$ satisfying (1), i.e.,

$$1 \geq \gamma_1 \geq \gamma_2 \geq \cdots \geq 0, \text{ and } 1 < \alpha_1 \leq \alpha_2 \leq \cdots$$

has been solved by [18] as follows:

- For $R_{d,\boldsymbol{\alpha},\gamma} \in \{r_{d,\boldsymbol{\alpha},\gamma}, \psi_{d,\boldsymbol{\alpha},\gamma}, \omega_{d,\boldsymbol{\alpha},\gamma}\}$, SPT holds iff PT holds iff

$$\delta := \liminf_{j\to\infty} \frac{\ln \gamma_j^{-1}}{\ln j} > 0.$$

The exponent of SPT is

$$p^{\mathrm{str}} = 2\max\left\{\frac{1}{\delta}, \frac{1}{\lceil \alpha_1 \rceil}\right\}.$$

- For $R_{d,\boldsymbol{\alpha},\gamma} = r_{d,\boldsymbol{\alpha},\gamma}$, WT holds iff

$$\lim_{j\to\infty} \gamma_j < 1.$$

- For $R_{d,\boldsymbol{\alpha},\gamma} \in \{r_{d,\boldsymbol{\alpha},\gamma}, \psi_{d,\boldsymbol{\alpha},\gamma}, \omega_{d,\boldsymbol{\alpha},\gamma}\}$ and $t > 1$, (t,s)-WT holds.

In this paper, we investigate the EC-tractability of the problem APP $= \{\mathrm{APP}_d\}_{d\in\mathbb{N}}$ in the weighted Hilbert space $H(K_{R_{d,\boldsymbol{\alpha},\gamma}})$ with $\boldsymbol{\gamma} = \{\gamma_j\}_{j\in\mathbb{N}}$ and $\boldsymbol{\alpha} = \{\alpha_j\}_{j\in\mathbb{N}}$ satisfying (1). We obtain sufficient and necessary conditions for EC-$(t,1)$-WT with $0 < t < 1$ and $t = 1$.

Theorem 1. *Let $\boldsymbol{\gamma} = \{\gamma_j\}_{j\in\mathbb{N}}$ and $\boldsymbol{\alpha} = \{\alpha_j\}_{j\in\mathbb{N}}$ satisfy (1). Then the problem APP $= \{APP_d\}_{d\in\mathbb{N}}$ in the weighted Hilbert spaces $H(K_{R_{d,\boldsymbol{\alpha},\gamma}})$ with $R_{d,\boldsymbol{\alpha},\gamma} \in \{r_{d,\boldsymbol{\alpha},\gamma}, \psi_{d,\boldsymbol{\alpha},\gamma}, \omega_{d,\boldsymbol{\alpha},\gamma}\}$*

(1) is EC-WT, if and only if

$$\lim_{j\to\infty}\gamma_j = 0.$$

(2) is EC-$(t,1)$-WT with $t<1$, if and only if

$$\lim_{j\to\infty}\frac{\ln j}{\ln(\gamma_j^{-1})} = 0.$$

4.2. The Proof

In order to prove Theorem 1 we need the following Lemmas.

Lemma 4. *Let $\eta>0$, $\varepsilon\in(0,1)$. We have for any $d\in\mathbb{N}$*

$$n(\varepsilon, APP_d) \le \varepsilon^{-2\eta}\prod_{j=1}^{d}\Big(1+\lceil\alpha_1\rceil^{\lceil\alpha_1\rceil\eta}\zeta(\lceil\alpha_1\rceil\eta)\gamma_j^{\eta}\Big).$$

Proof. By Lemma 3 we have

$$\begin{aligned}
\sum_{k=1}^{\infty}\lambda_{d,k}^{\eta} &= \sum_{\boldsymbol{k}\in\mathbb{N}_0^d}\big(R_{d,\boldsymbol{\alpha},\gamma}(\boldsymbol{k})\big)^{\eta} = \prod_{j=1}^{d}\left(1+\sum_{k=1}^{\infty}\Big(R_{\alpha_j,\gamma_j}(k)\Big)^{\eta}\right)\\
&\le \prod_{j=1}^{d}\left(1+\sum_{k=1}^{\infty}\Big(\lceil\alpha_1\rceil^{\lceil\alpha_1\rceil}r_{\alpha_1,\gamma_j}(k)\Big)^{\eta}\right)\\
&= \prod_{j=1}^{d}\left(1+\lceil\alpha_1\rceil^{\lceil\alpha_1\rceil\eta}\sum_{k=1}^{\infty}\Big(r_{\alpha_1,\gamma_j}(k)\Big)^{\eta}\right)\\
&= \prod_{j=1}^{d}\left(1+\lceil\alpha_1\rceil^{\lceil\alpha_1\rceil\eta}\sum_{k=1}^{\infty}\left(\frac{\gamma_j}{k^{\lceil\alpha_1\rceil}}\right)^{\eta}\right)\\
&= \prod_{j=1}^{d}\Big(1+\lceil\alpha_1\rceil^{\lceil\alpha_1\rceil\eta}\zeta(\lceil\alpha_1\rceil\eta)\gamma_j^{\eta}\Big).
\end{aligned}$$

This yields

$$n\lambda_{d,n}^{\eta} \le \sum_{k=1}^{n}\lambda_{d,k}^{\eta} \le \sum_{k=1}^{\infty}\lambda_{d,k}^{\eta} \le \prod_{j=1}^{d}\Big(1+\lceil\alpha_1\rceil^{\lceil\alpha_1\rceil\eta}\zeta(\lceil\alpha_1\rceil\eta)\gamma_j^{\eta}\Big),$$

which means

$$\lambda_{d,n} \le \frac{\prod_{j=1}^{d}\Big(1+\lceil\alpha_1\rceil^{\lceil\alpha_1\rceil\eta}\zeta(\lceil\alpha_1\rceil\eta)\gamma_j^{\eta}\Big)^{1/\eta}}{n^{1/\eta}}.$$

It follows from the above inequality and (7)

$$n(\varepsilon, \mathrm{APP}_d) = \min\Big\{n\in\mathbb{N}_0 : \lambda_{d,n+1}\le\varepsilon^2\Big\},$$

that

$$n(\varepsilon, \mathrm{APP}_d) \le \varepsilon^{-2\eta}\prod_{j=1}^{d}\Big(1+\lceil\alpha_1\rceil^{\lceil\alpha_1\rceil\eta}\zeta(\lceil\alpha_1\rceil\eta)\gamma_j^{\eta}\Big).$$

This proof is complete. □

Lemma 5. *Let $\varepsilon \in (0,1)$. We have for any $d \geq 2$*

$$n(\varepsilon, APP_d) \geq \left\lceil \left(\varepsilon^{-2}\lceil\alpha_1\rceil^{\lceil\alpha_1\rceil}\gamma_d\right)^{\frac{1}{\lceil\alpha_1\rceil}} \right\rceil.$$

Proof. Set

$$H = H(\varepsilon, d, \alpha) := \left\{ h \in \mathbb{N}_0 : h \leq \left\lceil \left(\varepsilon^{-2}\lceil\alpha_1\rceil^{\lceil\alpha_1\rceil}\gamma_d\right)^{\frac{1}{\lceil\alpha_1\rceil}} \right\rceil - 1 \right\}.$$

If $h > \left\lceil \left(\varepsilon^{-2}\lceil\alpha_1\rceil^{\lceil\alpha_1\rceil}\gamma_d\right)^{\frac{1}{\lceil\alpha_1\rceil}} \right\rceil - 1$ and $d \geq 2$, by Lemma 3 we have

$$\prod_{j=1}^{d-1} R_{\alpha_j,\gamma_j}(h_j) R_{\alpha_d,\gamma_d}(h) \leq R_{\alpha_d,\gamma_d}(h) \leq \lceil\alpha_1\rceil^{\lceil\alpha_1\rceil} r_{\alpha_1,\gamma_d}(h) = \lceil\alpha_1\rceil^{\lceil\alpha_1\rceil} \frac{\gamma_d}{h^{\lceil\alpha_1\rceil}} \leq \varepsilon^2$$

for any $\{h_1, \cdots, h_{d-1}\} \in \mathbb{N}_0^{d-1}$, which means

$$\left\{ \boldsymbol{h} \in \mathbb{N}_0^{d-1} : \prod_{j=1}^{d-1} R_{\alpha_j,\gamma_j}(h_j) R_{\alpha_d,\gamma_d}(h) > \varepsilon^2 \right\} = \varnothing \tag{9}$$

for all $h > \left\lceil \left(\varepsilon^{-2}\lceil\alpha_1\rceil^{\lceil\alpha_1\rceil}\gamma_d\right)^{\frac{1}{\lceil\alpha_1\rceil}} \right\rceil - 1$. It follows from (8) and (9) that

$$\begin{aligned}
n(\varepsilon, \mathrm{APP}_d) &= \left| \left\{ \boldsymbol{h} \in \mathbb{N}_0^d : \prod_{j=1}^{d} R_{\alpha_j,\gamma_j}(h_j) > \varepsilon^2 \right\} \right| \\
&= \left| \left\{ \boldsymbol{h} \in \mathbb{N}_0^d : \prod_{j=1}^{d-1} R_{\alpha_j,\gamma_j}(h_j) R_{\alpha_d,\gamma_d}(h_d) > \varepsilon^2 \right\} \right| \\
&= \sum_{h \in \mathbb{N}_0} \left| \left\{ \boldsymbol{h} \in \mathbb{N}_0^{d-1} : \prod_{j=1}^{d-1} R_{\alpha_j,\gamma_j}(h_j) R_{\alpha_d,\gamma_d}(h) > \varepsilon^2 \right\} \right| \\
&= \sum_{h \in H} \left| \left\{ \boldsymbol{h} \in \mathbb{N}_0^{d-1} : \prod_{j=1}^{d-1} R_{\alpha_j,\gamma_j}(h_j) R_{\alpha_d,\gamma_d}(h) > \varepsilon^2 \right\} \right| \\
&= \sum_{h \in (H \setminus \{0\})} \left| \left\{ \boldsymbol{h} \in \mathbb{N}_0^{d-1} : \prod_{j=1}^{d-1} R_{\alpha_j,\gamma_j}(h_j) > \varepsilon^2 R_{\alpha_d,\gamma_d}^{-1}(h) \right\} \right| \\
&\quad + \left| \left\{ \boldsymbol{h} \in \mathbb{N}_0^{d-1} : \prod_{j=1}^{d-1} R_{\alpha_j,\gamma_j}(h_j) > \varepsilon^2 \right\} \right| \\
&= \sum_{h \in (H \setminus \{0\})} n(\varepsilon R_{\alpha_d,\gamma_d}^{-1/2}(h), \mathrm{APP}_{d-1}) + n(\varepsilon, \mathrm{APP}_{d-1}) \\
&= \sum_{h=1}^{\left\lceil \left(\varepsilon^{-2}\lceil\alpha_1\rceil^{\lceil\alpha_1\rceil}\gamma_d\right)^{\frac{1}{\lceil\alpha_1\rceil}} \right\rceil - 1} n(\varepsilon R_{\alpha_d,\gamma_d}^{-1/2}(h), \mathrm{APP}_{d-1}) + n(\varepsilon, \mathrm{APP}_{d-1}) \\
&\geq \left\lceil \left(\varepsilon^{-2}\lceil\alpha_1\rceil^{\lceil\alpha_1\rceil}\gamma_d\right)^{\frac{1}{\lceil\alpha_1\rceil}} \right\rceil.
\end{aligned}$$

This finishes the proof. □

Lemma 6. *For* $\prod_{j=1}^{d}\left(\frac{\gamma_j}{2}\right) > \varepsilon^2$ *and* $\varepsilon \in (0,1)$ *we have*

$$n(\varepsilon, APP_d) \geq 2^d.$$

Proof. Set

$$\mathcal{A}(\varepsilon, d) = \left\{ \boldsymbol{h} \in \mathbb{N}_0^d : \prod_{j=1}^{d} R_{\alpha_j,\gamma_j}(h_j) > \varepsilon^2 \right\}.$$

If $\boldsymbol{h} = \{h_1, h_2, \ldots, h_d\} \in \{0,1\}^d$, we have from (5) that

$$R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}(\boldsymbol{h}) = \prod_{j=1}^{d} R_{\alpha_j,\gamma_j}(h_j) \geq \prod_{j=1}^{d}\left(\frac{\gamma_j}{2}\right).$$

Thus, we have $\{0,1\}^d \in \mathcal{A}(\varepsilon, d)$ for $\prod_{j=1}^{d}\left(\frac{\gamma_j}{2}\right) > \varepsilon^2$. Hence, it follows from (8) that

$$n(\varepsilon, \mathrm{APP}_d) = |\mathcal{A}(\varepsilon, d)| \geq \left| \left\{ \boldsymbol{h} \in \{0,1\}^d : \prod_{j=1}^{d} R_{\alpha_j,\gamma_j}(h_j) > \varepsilon^2 \right\} \right| = 2^d$$

for $\prod_{j=1}^{d}\left(\frac{\gamma_j}{2}\right) > \varepsilon^2$. This proof is complete. □

Proof of Theorem 1.

If there are infinitely many $\gamma_j = 0$ for $j \in \mathbb{N}$, the results are obviously true. Without loss of generality we discuss only that the γ_j are positive for $j \in \mathbb{N}$.

(1) Let $\delta > 0$ and take $\varepsilon = \prod_{j=1}^{d}\left(\frac{\gamma_j}{2}\right)^{\frac{1+\delta}{2}}$, then we have

$$\prod_{j=1}^{d}\left(\frac{\gamma_j}{2}\right) > \varepsilon^2.$$

It follows from Lemma 6 that

$$\begin{aligned} \frac{\ln n(\varepsilon, \mathrm{APP}_d)}{d + \ln(\varepsilon^{-1})} &\geq \frac{d \ln 2}{d + \frac{1+\delta}{2} \cdot \ln\left(\prod_{j=1}^{d}(2\gamma_j^{-1})\right)} \\ &\geq \frac{d \ln 2}{d + \frac{1+\delta}{2} \cdot d \cdot \ln(2\gamma_d^{-1})} \\ &= \frac{\ln 2}{1 + \frac{1+\delta}{2} \cdot \left(\ln 2 + \ln(\gamma_d^{-1})\right)}. \end{aligned} \tag{10}$$

Assume that App is EC-WT, i.e., for the above fixed ε

$$\lim_{d\to\infty} \frac{\ln n(\varepsilon, \mathrm{APP}_d)}{d + \ln(\varepsilon^{-1})} = 0.$$

Combing (10) and the above equality we have

$$0 = \lim_{d\to\infty} \frac{\ln n(\varepsilon, \mathrm{APP}_d)}{d + \ln(\varepsilon^{-1})} \geq \lim_{d\to\infty} \frac{\ln 2}{1 + \frac{1+\delta}{2} \cdot \left(\ln 2 + \ln(\gamma_d^{-1})\right)}.$$

This implies $\lim_{d\to\infty} \gamma_d = 0$.

On the other hand, assume that we have $\lim_{d\to\infty} \gamma_d = 0$. For $\eta > 0$ we obtain from the upper bound in Lemma 4 that

$$\begin{aligned}
\limsup_{d+\varepsilon^{-1}\to\infty} \frac{\ln n(\varepsilon, \mathrm{APP}_d)}{d+\ln(\varepsilon^{-1})} &\le \limsup_{d+\varepsilon^{-1}\to\infty} \frac{2\eta\ln(\varepsilon^{-1}) + \sum_{j=1}^{d} \ln\left(1 + \lceil\alpha_1\rceil^{\lceil\alpha_1\rceil\eta}\zeta(\lceil\alpha_1\rceil\eta)\gamma_j^{\eta}\right)}{d+\ln(\varepsilon^{-1})} \\
&\le \limsup_{d+\varepsilon^{-1}\to\infty} \frac{2\eta\ln(\varepsilon^{-1}) + \sum_{j=1}^{d} \lceil\alpha_1\rceil^{\lceil\alpha_1\rceil\eta}\zeta(\lceil\alpha_1\rceil\eta)\gamma_j^{\eta}}{d+\ln(\varepsilon^{-1})} \\
&\le 2\eta + \limsup_{d\to\infty} \frac{\lceil\alpha_1\rceil^{\lceil\alpha_1\rceil\eta}\zeta(\lceil\alpha_1\rceil\eta)\sum_{j=1}^{d}\gamma_j^{\eta}}{d} \\
&= 2\eta,
\end{aligned}$$

where we used $\ln(1+x) \le x$ for all $x \ge 0$ and $\limsup_{d\to\infty} \frac{\sum_{j=1}^{d}\gamma_j^{\eta}}{d} = 0$ if $\lim_{d\to\infty}\gamma_d = 0$. Setting $\eta \to 0$, we have

$$\limsup_{d+\varepsilon^{-1}\to\infty} \frac{\ln n(\varepsilon, \mathrm{APP}_d)}{d+\ln(\varepsilon^{-1})} = 0,$$

which yields that ET-WT holds.

(2) Assume that APP is EC-$(t,1)$-WT for $t < 1$. First, we note that $\lim_{d\to\infty}\gamma_d = 0$. Indeed, if $\lim_{d\to\infty}\gamma_d \neq 0$, we deduce from Theorem 1 (1) that EC-WT doesn't hold, i.e.,

$$0 < \limsup_{d+\varepsilon^{-1}\to\infty} \frac{\ln n(\varepsilon, \mathrm{APP}_d)}{d+\ln(\varepsilon^{-1})} \le \limsup_{d+\varepsilon^{-1}\to\infty} \frac{\ln n(\varepsilon, \mathrm{APP}_d)}{d^t+\ln(\varepsilon^{-1})}.$$

This deduces that EC-$(t,1)$-WT for $t < 1$ does not hold.

Next, we will prove $\lim_{j\to\infty} \frac{\ln j}{\ln(\gamma_j^{-1})} = 0$. Let $\varepsilon = \varepsilon_d \in (0,1)$ such that

$$\ln\left(\varepsilon^{-2}\lceil\alpha_1\rceil^{\lceil\alpha_1\rceil}\gamma_d\right)^{\frac{1}{\lceil\alpha_1\rceil}} = d^t$$

for large $d \in \mathbb{N}$. From the lower bound in Lemma 5 we obtain

$$\begin{aligned}
\frac{\ln n(\varepsilon, \mathrm{APP}_d)}{d^t+\ln(\varepsilon^{-1})} &\ge \frac{\ln\left\lceil\left(\varepsilon^{-2}\lceil\alpha_1\rceil^{\lceil\alpha_1\rceil}\gamma_d\right)^{\frac{1}{\lceil\alpha_1\rceil}}\right\rceil}{d^t+\ln(\varepsilon^{-1})} \ge \frac{\ln\left(\varepsilon^{-2}\lceil\alpha_1\rceil^{\lceil\alpha_1\rceil}\gamma_d\right)^{\frac{1}{\lceil\alpha_1\rceil}}}{d^t+\ln(\varepsilon^{-1})} \\
&= \frac{d^t}{d^t+\ln(\varepsilon^{-1})} = \frac{d^t}{d^t+\lceil\alpha_1\rceil d^t/2+\ln(\gamma_d^{-1})/2-\lceil\alpha_1\rceil(\ln\lceil\alpha_1\rceil)/2} \\
&= \frac{1}{1+\lceil\alpha_1\rceil/2+\ln(\gamma_d^{-1})/(2d^t)-\lceil\alpha_1\rceil(\ln\lceil\alpha_1\rceil)/(2d^t)}.
\end{aligned}$$

It follows from the assumption that

$$0 = \limsup_{d+\varepsilon^{-1}\to\infty} \frac{\ln n(\varepsilon, \mathrm{APP}_d)}{d^t + \ln(\varepsilon^{-1})} \geq \limsup_{d+\varepsilon^{-1}\to\infty} \frac{1}{1 + \lceil \alpha_1 \rceil/2 + \ln(\gamma_d^{-1})/(2d^t) - \lceil \alpha_1 \rceil (\ln\lceil \alpha_1 \rceil)/(2d^t)}$$
$$= \limsup_{d+\varepsilon^{-1}\to\infty} \frac{1}{1 + \lceil \alpha_1 \rceil/2 + \ln(\gamma_d^{-1})/(2d^t)},$$

which implies

$$\lim_{d\to\infty} \frac{d^t}{\ln(\gamma_d^{-1})} = 0.$$

Using the fact that $d^t \geq \ln d^t = t \ln d \geq 0$ for large $d \in \mathbb{N}$, we have

$$0 \leq \lim_{d\to\infty} \frac{\ln d}{\ln(\gamma_d^{-1})} \leq \lim_{d\to\infty} \frac{d^t}{t \ln(\gamma_d^{-1})} = 0,$$

i.e.,

$$\lim_{d\to\infty} \frac{\ln d}{\ln(\gamma_d^{-1})} = 0.$$

On the other hand, assume that $\lim_{j\to\infty} \frac{\ln j}{\ln(\gamma_j^{-1})} = 0$. Then we obtain that for all $\delta > 0$ there exists a positive number $N_\delta > 0$ such that

$$\gamma_j \leq j^{-\delta} \text{ for all } j \geq N_\delta. \tag{11}$$

Let $\eta > 0$. We get from Lemma 4 that

$$\ln n(\varepsilon, \mathrm{APP}_d) \leq 2\eta \ln(\varepsilon^{-1}) + \sum_{j=1}^{d} \ln\left(1 + \lceil \alpha_1 \rceil^{\lceil \alpha_1 \rceil \eta} \zeta(\lceil \alpha_1 \rceil \eta) \gamma_j^\eta\right)$$
$$\leq 2\eta \ln(\varepsilon^{-1}) + \sum_{j=1}^{d} \lceil \alpha_1 \rceil^{\lceil \alpha_1 \rceil \eta} \zeta(\lceil \alpha_1 \rceil \eta) \gamma_j^\eta, \tag{12}$$

where we used $\ln(1+x) \leq x$ for all $x > 0$. Choose $\delta = \frac{2}{\eta}$. By (11) and (12) we get

$$\frac{\ln n(\varepsilon, \mathrm{APP}_d)}{d^t + \ln(\varepsilon^{-1})} \leq \frac{2\eta \ln(\varepsilon^{-1}) + \sum_{j=1}^{N_{2/\eta}-1} \lceil \alpha_1 \rceil^{\lceil \alpha_1 \rceil \eta} \zeta(\lceil \alpha_1 \rceil \eta) \gamma_j^\eta + \sum_{j=N_{2/\eta}}^{\max\{d, N_{2/\eta}\}} \lceil \alpha_1 \rceil^{\lceil \alpha_1 \rceil \eta} \zeta(\lceil \alpha_1 \rceil \eta) \gamma_j^\eta}{d^t + \ln(\varepsilon^{-1})}$$
$$\leq 2\eta + \frac{\lceil \alpha_1 \rceil^{\lceil \alpha_1 \rceil \eta} \zeta(\lceil \alpha_1 \rceil \eta) \gamma_1^\eta (N_{2/\eta} - 1) + \lceil \alpha_1 \rceil^{\lceil \alpha_1 \rceil \eta} \zeta(\lceil \alpha_1 \rceil \eta) \sum_{j=N_{2/\eta}}^{\max\{d, N_{2/\eta}\}} j^{-2}}{d^t + \ln(\varepsilon^{-1})}.$$

It follows that

$$\limsup_{d+\varepsilon^{-1}\to\infty} \frac{\ln n(\varepsilon, \mathrm{APP}_d)}{d^t + \ln(\varepsilon^{-1})} \leq 2\eta + \lceil \alpha_1 \rceil^{\lceil \alpha_1 \rceil \eta} \zeta(\lceil \alpha_1 \rceil \eta) \limsup_{d+\varepsilon^{-1}\to\infty} \frac{\sum_{j=N_{2/\eta}}^{\max\{d, N_{2/\eta}\}} j^{-2}}{d^t + \ln(\varepsilon^{-1})}$$
$$\leq 2\eta + \lceil \alpha_1 \rceil^{\lceil \alpha_1 \rceil \eta} \zeta(\lceil \alpha_1 \rceil \eta) \limsup_{d+\varepsilon^{-1}\to\infty} \frac{\sum_{j=1}^{\infty} j^{-2}}{d^t + \ln(\varepsilon^{-1})}$$
$$= 2\eta.$$

Setting $\eta \to 0$, we have

$$\limsup_{d+\varepsilon^{-1}\to\infty} \frac{\ln n(\varepsilon, \mathrm{APP}_d)}{d^t + \ln(\varepsilon^{-1})} = 0.$$

Therefore, Theorem 1 is proved.

□

Example 1. *An example for EC-WT.*

Assume that $\gamma_j = j^{-2}$ and $\alpha_j = j+1$ for all $j \in \mathbb{N}$. Next, we will study EC-WT for the weighted Hilbert spaces $H(K_{R_{d,\boldsymbol{\alpha},\gamma}})$ with weight $R_{d,\boldsymbol{\alpha},\gamma} \in \{r_{d,\boldsymbol{\alpha},\gamma}, \psi_{d,\boldsymbol{\alpha},\gamma}, \omega_{d,\boldsymbol{\alpha},\gamma}\}$.

Obviously, we have $\lim_{j\to\infty} \gamma_j = 0$. By Lemma 4 we get

$$\begin{aligned}
\limsup_{d+\varepsilon^{-1}\to\infty} \frac{\ln n(\varepsilon, APP_d)}{d + \ln(\varepsilon^{-1})} &\le \limsup_{d+\varepsilon^{-1}\to\infty} \frac{2\eta \ln(\varepsilon^{-1}) + \sum_{j=1}^{d} \ln\left(1 + \lceil\alpha_1\rceil^{\lceil\alpha_1\rceil\eta}\zeta(\lceil\alpha_1\rceil\eta)\gamma_j^{\eta}\right)}{d + \ln(\varepsilon^{-1})} \\
&\le \limsup_{d+\varepsilon^{-1}\to\infty} \frac{2\eta \ln(\varepsilon^{-1}) + \sum_{j=1}^{d} \lceil\alpha_1\rceil^{\lceil\alpha_1\rceil\eta}\zeta(\lceil\alpha_1\rceil\eta)\gamma_j^{\eta}}{d + \ln(\varepsilon^{-1})} \\
&\le 2\eta + \limsup_{d\to\infty} \frac{\lceil\alpha_1\rceil^{\lceil\alpha_1\rceil\eta}\zeta(\lceil\alpha_1\rceil\eta) \sum_{j=1}^{d} \gamma_j^{\eta}}{d} \\
&= 2\eta + \limsup_{d\to\infty} \frac{\lceil\alpha_1\rceil^{\lceil\alpha_1\rceil\eta}\zeta(\lceil\alpha_1\rceil\eta) \sum_{j=1}^{d} j^{-2\eta}}{d} \\
&= 2\eta,
\end{aligned}$$

where in the second inequality we used $\ln(1+x) \le x$ for all $x \ge 0$ and $\limsup_{d\to\infty} \frac{\sum_{j=1}^{d} j^{-2\eta}}{d} = 0$. Setting $\eta \to 0$, we have

$$\limsup_{d+\varepsilon^{-1}\to\infty} \frac{\ln n(\varepsilon, APP_d)}{d + \ln(\varepsilon^{-1})} = 0.$$

Hence, APP is EC-WT.

Example 2. *An example for EC-$(t,1)$-WT for $t < 1$.*

Assume that $\gamma_j = 2^{-j}$ and $\alpha_j = 2j$ for all $j \in \mathbb{N}$. Next, we will study EC-$(t,1)$-WT for $t < 1$ for the weighted Hilbert spaces $H(K_{R_{d,\boldsymbol{\alpha},\gamma}})$ with weight $R_{d,\boldsymbol{\alpha},\gamma} \in \{r_{d,\boldsymbol{\alpha},\gamma}, \psi_{d,\boldsymbol{\alpha},\gamma}, \omega_{d,\boldsymbol{\alpha},\gamma}\}$.

Note that $\lim_{j\to\infty} \frac{\ln j}{\ln(\gamma_j^{-1})} = \lim_{j\to\infty} \frac{\ln j}{j \ln 2} = 0$. It follows from Lemma 4 that

$$\begin{aligned}
\ln n(\varepsilon, APP_d) &\le 2\eta \ln(\varepsilon^{-1}) + \sum_{j=1}^{d} \ln\left(1 + \lceil\alpha_1\rceil^{\lceil\alpha_1\rceil\eta}\zeta(\lceil\alpha_1\rceil\eta)\gamma_j^{\eta}\right) \\
&= 2\eta \ln(\varepsilon^{-1}) + \sum_{j=1}^{d} \ln\left(1 + \lceil\alpha_1\rceil^{\lceil\alpha_1\rceil\eta}\zeta(\lceil\alpha_1\rceil\eta)2^{-\eta j}\right) \\
&\le 2\eta \ln(\varepsilon^{-1}) + \sum_{j=1}^{d} \lceil\alpha_1\rceil^{\lceil\alpha_1\rceil\eta}\zeta(\lceil\alpha_1\rceil\eta)2^{-\eta j},
\end{aligned}$$

where in the last inequality we used $\ln(1+x) \le x$ *for all* $x > 0$. *It yields that*

$$\begin{aligned}\limsup_{d+\varepsilon^{-1}\to\infty} \frac{\ln n(\varepsilon, APP_d)}{d^t + \ln(\varepsilon^{-1})} &\le \limsup_{d+\varepsilon^{-1}\to\infty} \frac{2\eta \ln(\varepsilon^{-1}) + \sum_{j=1}^{d} \lceil\alpha_1\rceil^{\lceil\alpha_1\rceil\eta}\zeta(\lceil\alpha_1\rceil\eta)2^{-\eta j}}{d^t + \ln(\varepsilon^{-1})} \\ &\le 2\eta + \limsup_{d+\varepsilon^{-1}\to\infty} \frac{\lceil\alpha_1\rceil^{\lceil\alpha_1\rceil\eta}\zeta(\lceil\alpha_1\rceil\eta)\sum_{j=1}^{d} 2^{-\eta j}}{d^t + \ln(\varepsilon^{-1})} \\ &\le 2\eta + \limsup_{d+\varepsilon^{-1}\to\infty} \frac{\lceil\alpha_1\rceil^{\lceil\alpha_1\rceil\eta}\zeta(\lceil\alpha_1\rceil\eta)\sum_{j=1}^{\infty} 2^{-\eta j}}{d^t + \ln(\varepsilon^{-1})} \\ &= 2\eta.\end{aligned}$$

Setting $\eta \to 0$, *we have*

$$\limsup_{d+\varepsilon^{-1}\to\infty} \frac{\ln n(\varepsilon, APP_d)}{d^t + \ln(\varepsilon^{-1})} = 0.$$

Hence, APP is EC-$(t,1)$*-WT for* $t < 1$.

Remark 5. *We note that for Example 1 with* $\gamma_j = j^{-2}$ *and* $\alpha_j = j+1$ *for all* $j \in \mathbb{N}$, *APP is EC-WT, but not EC-*$(t,1)$*-WT for* $t < 1$. *Indeed, let* $\varepsilon = \varepsilon_d \in (0,1)$ *such that*

$$\ln\left(\varepsilon^{-2}\lceil\alpha_1\rceil^{\lceil\alpha_1\rceil}\gamma_d\right)^{\frac{1}{\lceil\alpha_1\rceil}} = d, \quad i.e., \quad \varepsilon^{-1} = \frac{de^{d\lceil\alpha_1\rceil/2}}{\lceil\alpha_1\rceil^{\lceil\alpha_1\rceil/2}}$$

for large $d \in \mathbb{N}$. *From Lemma 5 we have*

$$\begin{aligned}\frac{\ln n(\varepsilon, APP_d)}{d^t + \ln(\varepsilon^{-1})} &\ge \frac{\ln\left\lceil\left(\varepsilon^{-2}\lceil\alpha_1\rceil^{\lceil\alpha_1\rceil}\gamma_d\right)^{\frac{1}{\lceil\alpha_1\rceil}}\right\rceil}{d^t + \ln(\varepsilon^{-1})} \ge \frac{\ln\left(\varepsilon^{-2}\lceil\alpha_1\rceil^{\lceil\alpha_1\rceil}\gamma_d\right)^{\frac{1}{\lceil\alpha_1\rceil}}}{d^t + \ln(\varepsilon^{-1})} \\ &= \frac{d}{d^t + \ln(\varepsilon^{-1})} = \frac{d}{d^t + \lceil\alpha_1\rceil d/2 + \ln d - \lceil\alpha_1\rceil(\ln\lceil\alpha_1\rceil)/2} \\ &= \frac{1}{d^{t-1} + \lceil\alpha_1\rceil/2 + \ln d/d - \lceil\alpha_1\rceil(\ln\lceil\alpha_1\rceil)/(2d)}.\end{aligned}$$

For the above fixed ε *and* $t < 1$ *we obtain*

$$\lim_{d\to\infty} \frac{\ln n(\varepsilon, APP_d)}{d + \ln(\varepsilon^{-1})} \ge \lim_{d\to\infty} \frac{1}{d^{t-1} + \lceil\alpha_1\rceil/2 + \ln d/d - \lceil\alpha_1\rceil(\ln\lceil\alpha_1\rceil)/(2d)} = \frac{2}{\lceil\alpha_1\rceil}.$$

This means that APP is not EC-$(t,1)$*-WT for* $t < 1$.

Remark 6. *Obviously, for Example 2 with* $\gamma_j = 2^{-j}$ *and* $\alpha_j = 2j$ *for all* $j \in \mathbb{N}$, *APP is also EC-WT. Indeed, if APP is EC-*$(t,1)$*-WT for* $t < 1$, *then it is EC-WT. Assume that APP is EC-*$(t,1)$*-WT for* $t < 1$, *then we have*

$$\lim_{d+\varepsilon^{-1}\to\infty} \frac{\ln n(\varepsilon, APP_d)}{d^t + \ln(\varepsilon^{-1})} = 0.$$

Since

$$0 \le \lim_{d+\varepsilon^{-1}\to\infty} \frac{\ln n(\varepsilon, APP_d)}{d + \ln(\varepsilon^{-1})} \le \lim_{d+\varepsilon^{-1}\to\infty} \frac{\ln n(\varepsilon, APP_d)}{d^t + \ln(\varepsilon^{-1})},$$

we further get

$$\lim_{d+\varepsilon^{-1}\to\infty} \frac{\ln n(\varepsilon, APP_d)}{d + \ln(\varepsilon^{-1})} = 0,$$

which means that APP is EC-WT.

5. Conclusions

In this paper we discuss the EC-WT and EC-$(t,1)$-WT with $t < 1$ for the approximation problem APP in weighted Hilbert spaces $H_{R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}}$ for $R_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}} \in \{r_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}, \psi_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}, \omega_{d,\boldsymbol{\alpha},\boldsymbol{\gamma}}\}$ with parameters $1 \geq \gamma_1 \geq \gamma_2 \geq \cdots \geq 0$ and $1 < \alpha_1 \leq \alpha_2 \leq \cdots$. We obtain the matching necessary and sufficient condition

$$\lim_{j\to\infty} \gamma_j = 0$$

on EC-WT, and the matching necessary and sufficient condition

$$\lim_{j\to\infty} \frac{\ln j}{\ln(\gamma_j^{-1})} = 0$$

on EC-$(t,1)$-WT with $t < 1$. The weights are used to model the importance of the functions from the weighted Hilbert spaces, so we will further research the other EC-tractability notions such as EC-SPT, EC-PT, EC-QWT, and EC-UWT.

Author Contributions: Conceptualization, J.C. and H.Y.; methodology, J.C. and H.Y.; validation, J.C.; formal analysis, J.C.; investigation, H.Y.; resources, H.Y.; data curation, H.Y.; writing—original draft preparation, J.C.; writing—review and editing, H.Y.; visualization, J.C.; supervision, J.C. and H.Y.; project administration, J.C. and H.Y. All authors have read and agreed to the published version of the manuscript.

Funding: Jia Chen is supported by the National Natural Science Foundation of China (Project 12001342), and the Doctoral Foundation Project of Shanxi Datong University (Project 2019-B-10). Huichao Yan is supported by the Scientific and Technological Innovation Project of Colleges and Universities in Shanxi Province (Project 2022L438), the Basic Youth Research Found Project of Shanxi Datong University (Project 2022Q10), and the Doctoral Foundation Project of Shanxi Datong University (Project 2021-B-17).

Data Availability Statement: The original contributions presented in the study are included in the article, further inquiries can be directed to the corresponding author.

Acknowledgments: The authors would like to thank all referees and the editor for suggestions on this paper.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Glimm, J.; Jaffe, A. *Quantum Physics*; Springer: New York, NY, USA, 1987.
2. Le Bris, C. Computational chemistry from the perspective of numerical analysis. *Acta Numer.* **2005**, *14*, 363–444. [CrossRef]
3. Rust, J.; Traub, J.F.; Woźniakowski, H. Is there a curse of dimensionality for contraction fixed points in the worst case. *Econometrica* **2002**, *70*, 285–329. [CrossRef]
4. Novak, E.; Woźniakowski, H. *Tractability of Multivariate Problems, Volume I: Linear Information*; EMS Zürich: Zurich, Swtizerland, 2008.
5. Novak, E.; Woźniakowski, H. *Tractability of Multivariate Problems, Volume II: Standard Information for Functionals*; EMS Zürich: Zurich, Swtizerland, 2010.
6. Novak, E.; Woźniakowski, H. *Tractability of Multivariate Problems, Volume III: Standard Information for Operators*; EMS Zürich: Zurich, Swtizerland, 2012.
7. Dick, J.; Kritzer, P.; Pillichshammer, F.; Woźniakowski, H. Approximation of analytic functions in Korobov spaces. *J. Complex.* **2014**, *30*, 2–28. [CrossRef]
8. Irrgeher, C.; Kritzer, P.; Pillichshammer, F.; Woźniakowski, H. Tractability of multivariate approximation defined over Hilbert spaces with exponential weights. *J. Approx. Theory* **2016**, *207*, 301–338. [CrossRef]
9. Wang, H. A note about EC-(s,t)-weak tractability of multivariate approximation with analytic Korobov kernels. *J. Complex.* **2019**, *55*, 101412. [CrossRef]

10. Liu, Y.; Xu, G. Average case tractability of a multivariate approximation problem. *J. Complex.* **2017**, *43*, 76–102. [CrossRef]
11. Zhang, J. A note on EC-tractability of multivariate approximation in weighted Korobov spaces for the standard information class. *J. Complex.* **2021**, *67*, 101573. [CrossRef]
12. Xu, G. EC-tractability of L_p-approximation in Korobov spaces with exponential weights. *J. Approx. Theory* **2020**, *249*, 1–20. [CrossRef]
13. Wasilkowski, G.W.; Woźniakowski, H. Weighted tensor product algorithms for linear multivariate problems. *J. Complex.* **1999**, *15*, 402–447. [CrossRef]
14. Eberta, A.; Pillichshammer, F. Tractability of approximation in the weighted Korobov space in the worst-case setting—A complete picture. *J. Complex.* **2021**, *67*, 101571. [CrossRef]
15. Novak, E.; Sloan, I.H.; Woźniakowski, H. Tractability of approximation for weighted Korobov spaces on classical and quantum computers, *Found. Comput. Math.* **2004**, *4*, 121–156.
16. Chen, J. EC-(t_1, t_2)-tractability of approximation in weighted Korobov spaces in the worst case setting. *J. Complex.* **2022**, *73*, 101680. [CrossRef]
17. Leobacher, G.; Pillichshammer, F.; Ebert, A. Tractability of L_2-approximation and integration in weighted Hermite spaces of finite smoothness. *J. Complex.* **2023** *78*, 101768.
18. Yan, H.; Chen, J. Tractability of approximation of functions defined over weighted Hilbert spaces. *Axioms* **2024**, *13*, 108. [CrossRef]
19. Aronszajn, N. Theory of reproducing kernels. *Trans. Am. Math. Soc.* **1950**, *68*, 337–404. [CrossRef]
20. Irrgeher, C.; Leobacher, G. High-dimensional integration on the $\mathbb{R}^d$, weighted Hermite spaces, and orthogonal transforms. *J. Complex.* **2015**, *31*, 174–205.
21. Dick, J.; Irrgeher, C.; Leobacher, G; Pillichshammer, F. On the optimal order of integration in Hermite spaces with finite smoothness. *SIAM J. Numer. Anal.* **2018**, *56*, 684–707. [CrossRef]
22. Dick, J.; Larcher, G.; Pillichshammer, F.; Woźniakowski, H. Exponential convergence and tractability of multivariate integration for korobov spaces. *Math. Comp.* **2011**, *80*, 905–930. [CrossRef]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.