



Numerical Solution of the Cauchy-Type Singular Integral Equation with a Highly Oscillatory Kernel Function

SAIRA ^{1,†}, Shuhuang Xiang ^{1,*,†} and Guidong Liu ²

- ¹ School of Mathematics and Statistics, Central South University, Changsha 410083, China; sairahameed@csu.edu.cn
- ² School of Statistics and Mathematics, Nanjing Audit University, Nanjing 211815, China; liugd@nau.edu.cn
- * Correspondence: xiangsh@csu.edu.cn.; Tel.: +86-1397-314-3907
- † Current address: School of Mathematics and Statistics, Central South University, Changsha 410083, China.

Received: 16 August 2019; Accepted: 18 September 2019; Published: 20 September 2019



Abstract: This paper aims to present a Clenshaw–Curtis–Filon quadrature to approximate the solution of various cases of Cauchy-type singular integral equations (CSIEs) of the second kind with a highly oscillatory kernel function. We adduce that the zero case oscillation (k = 0) proposed method gives more accurate results than the scheme introduced in Dezhbord at el. (2016) and Eshkuvatov at el. (2009) for small values of *N*. Finally, this paper illustrates some error analyses and numerical results for CSIEs.

Keywords: Clenshaw-Curtis-Filon; high oscillation; singular integral equations; boundary singularities

1. Introduction

Integral equations have broad roots in branches of science and engineering [1–6]. Cauchy-type singular integral equations (CSIEs) of the second kind occur in electromagnetic scattering and quantum mechanics [7] and are defined as:

$$au(x) + \frac{b}{\pi} \int_{-1}^{1} \frac{u(y)K(x,y)}{y-x} dy = f(x), \quad x \in (-1,1).$$
(1)

A singular integral equation with a Cauchy principal value is a generalized form of an airfoil equation [8]. Here *a* and *b* are constants such that $a^2 + b^2 = 1$, $b \neq 0$ and $K(x, y) = e^{ik(y-x)}$ are the highly oscillatory kernel function. The function f(x) is the Hölder continuous function, whereas u(x) is an unknown function. The solution to the above-mentioned Equation (1) contains boundary singularities $w(x) = (x + 1)^{\alpha}(1 - x)^{\beta}$, i.e., u(x) = w(x)g(x) and g(x) is a smooth function [9,10]. Then the above Equation (1) transforms into:

$$aw(x)g(x) + \frac{b}{\pi} \int_{-1}^{1} \frac{w(y)g(y)e^{ik(y-x)}}{y-x} dy = f(x), \quad x \in (-1,1),$$
(2)

where $\alpha, \beta \in (-1, 1)$ depend on *a* and *b*, such that:

$$\alpha = \frac{1}{2\pi i} \log\left(\frac{a-ib}{a+ib}\right) - N \quad , \quad \beta = \frac{-1}{2\pi i} \log\left(\frac{a-ib}{a+ib}\right) - M, \tag{3}$$
$$\kappa = -(\alpha + \beta) = M + N.$$

Here *M* and *N* are integers in [-1, 1], whereas the index of the integral equation is called κ , analogous to a class of functions, wherein the solution is to be sought. It is pertinent to mention that to



produce integrable singularities in the solution, the index κ is restricted to three cases, [-1, 0, 1], but the addressed paper considers only two cases for κ , i.e., $\kappa \leq 0$. The value of the index κ depends on different values for M and N [11–13]. A great number of real life practical problems, e.g., for $\kappa = -1$, the so-called natched half-plane problem and another problem of a crack parallel to the free boundary of an isotropic semi-infinite plane, that can be reduced to Cauchy singular integral equations are addressed in [14–17]. Writing Equation (2) in operator form, we get [18]:

$$Hg = f, \tag{4}$$

where:

$$Hg = aw(x)g(x) + \frac{b}{\pi} \int_{-1}^{1} \frac{w(y)g(y)e^{ik(y-x)}}{y-x} dy.$$

Let us define another operator:

$$H'f = aw^{*}(x)f(x) - \frac{b}{\pi} \int_{-1}^{1} \frac{w^{*}(y)f(y)e^{ik(y-x)}}{y-x} dy,$$
(5)

further:

$$HH' = I \quad if\kappa > 0$$

$$HH' = H'H = I \quad if\kappa = 0$$

$$H'H = I \quad if\kappa < 0$$
(6)

where $w^*(x) = (1+x)^{-\alpha}(1-x)^{-\beta}$.

It is worthy mentioning the fact that the solution for CSIE exists but unfortunately it is not unique, as CSIE has three solution cases for different values of κ . The aforementioned theorem appertains to the existence of the solution of CSIE for case $\kappa = 0$.

Theorem 1. [13,15] (Existence of CSIEs) Let the singular integral Equation (2) be equivalent to a Fredholm integral equation, which implies that every solution of a Fredholm integral equation is the solution of a singular integral equation and vice versa.

Proof. Based on Equations (4)–(6) the SIE (2) can be transforms into:

$$g = H'f.$$

Furthermore, it can be written as a Fredholm integral equation:

$$u(y) + \int_{-1}^{1} N(y,\tau) y(\tau) d\tau = F(y).$$
⁽⁷⁾

where:

$$F(y) = \frac{b}{\pi}w(y)\int_{-1}^{1}\frac{w^{*}(x)f(x)}{y-x}dx$$

and:

$$N(y,\tau) = aK(x,\tau)w^{-1} - \frac{b}{\pi}w(y)\int_{-1}^{1}\frac{w^{*}(x)K(x,\tau)}{y-x}dx.$$

Thus the claimed theorem is proven. \Box

Moreover, for Equation (1) we have three cases for κ :

$$\kappa = \begin{cases} 1, & \alpha < 0, -1 < \beta, & \alpha \neq \beta, \\ -1, & 0 < \beta, \alpha < 1, & \alpha \neq \beta, \\ 0, & \alpha = -\beta, & |\beta| \neq \frac{1}{2}. \end{cases}$$
(8)

Similarly, solution cases of the CSIE of the second type depending on values of κ are:

• 1: The solution u(x) for $\kappa = 1$ is unbounded at both end points $x = \pm 1$:

$$u(x) = af(x) - \frac{bw(x)}{\pi} e^{-ikx} \int_{-1}^{1} \frac{w^*(y)f(y)e^{iky}}{y-x} dy + Cw(x),$$
(9)

where C is an arbitrary constant such that:

$$\int_{-1}^{1} u(y)e^{iky}dy = C.$$
 (10)

Equation (2) gets infinitely many solutions but is unique for the above condition.

• 2: The solution u(x) is bounded for $\kappa = 0$ at $x = \pm 1$ and unbounded at $x = \pm 1$:

$$u(x) = af(x) - \frac{bw(x)}{\pi} e^{-ikx} \int_{-1}^{1} \frac{w^*(y)f(y)e^{iky}}{y-x} dy,$$
(11)

Equation (2) gets a unique solution.

• 3: The solution u(x) is bounded at both end points $x = \pm 1$ for $\kappa = -1$:

$$u(x) = af(x) - \frac{bw(x)}{\pi} e^{-ikx} \int_{-1}^{1} \frac{w^*(y)f(y)e^{iky}}{y-x} dy.$$
 (12)

Equation (2) has no solution unless it satisfies the following condition:

$$\int_{-1}^{1} \frac{f(y)e^{iky}}{w(y)} dy = 0.$$
 (13)

For many decades researchers have been struggling to find an efficient method to get these solutions. The Galerkin method, polynomial collocation method, Clenshaw–Curtis–Filon method and the steepest descent method are some of the eminent methods among many others for the solution of SIEs [19–24]. Moreover, Chakarbarti and Berge [25] for a linear function f(x) gave an approximated method based on polynomial approximation and Chebyshev points. Z.K. Eshkuvatov [10] introduced the method taking Chebyshev polynomials of all four kinds for all four different solution cases of the CSIE. Reproducing the kernel Hilbert space (RKHS) method has been proposed by A. Dezhbord at el. [26]. The representation of solution u(x) is in the form of a series in reproducing kernel spaces.

This research work introduces the Clenshaw–Curtis–Filon quadrature to approximate the solution for various cases of a Cauchy singular integral equation of the second kind, Equation (1.1), at equally spaced points x_i . So the integral equation takes the form:

$$u_N(x_i) = af(x_i) - \frac{bw(x_i)}{\pi} e^{-ikx_i} \int_{-1}^1 \frac{w^*(y) f_N(y) e^{iky}}{y - x_i} dy,$$
(14)

depending on the κ . Furthermore, the results of the numerical example are compared with [10,26] for k = 0. Comparison reveals that the addressed method gives a more accurate approximation than these

methods, Section 4 provides this phenomena. The rest of the paper is organised as follows; Section 2 defines the numerical evaluation of the Cauchy integral in CSIE and approximates the solution at equally spaced points x_i . Section 3 represents some error analyses for CSIE. Section 4 concludes this paper by giving numerical results.

2. Description of the Method

The presented Clenshaw–Curtis–Filon quadrature to approximate the integral term $I(\alpha, \beta, k, x) = f_{-1}^1 \frac{w(y)f(y)e^{iky}}{y-x} dy$ consists of replacing function f(y) by its interpolation polynomial $P_N(y)$ at Clenshaw–Curtis point set, $y_j = cos \frac{j\pi}{N}$, $j = 0, 1, \dots N$. Rewriting the interpolation in terms of the Chebyshev series:

$$f(y) \approx P_N(y) = \sum_{n=0}^{N} {}^{"}c_n T_n(y).$$
 (15)

Here $T_n(y)$ is the Chebyshev polynomial of the first kind of degree *n*. Double prime denotes a summation, wherein the first and last terms are divided by 2. The FFT is used for proficient calculation of the coefficient c_n [27,28], defined as:

$$c_n = \frac{2}{N} \sum_{j=0}^N "f(y_j) T_n(y_j).$$

Let it be that for any fixed x we can elect N s.t $x \notin \{y_j\}$; then the interpolation polynomial is rewritten in the form of a Chebyshev series as:

$$\bar{P}_{N+1}(y) = \sum_{n=0}^{N+1} a_n T_n(y)$$

where a_n can be computed in O(N) operations once c_n are calculated [27,29]. The Clenshaw–Curtis–Filon quadrature rule for integral $I(\alpha, \beta, k, x)$ is defined as:

$$I(\alpha,\beta,k,x) = \int_{-1}^{1} \frac{w(y)f(y)e^{iky}}{y-x} dy = \int_{-1}^{1} \frac{w(y)\bar{P}_{N+1}(y)e^{iky}}{y-x} dy = \sum_{n=0}^{N+1} a_n M_n(\alpha,\beta,k,x),$$
(16)

where $M_n(\alpha, \beta, k, x) = \int_{-1}^1 \frac{w(y)T_n(y)e^{iky}}{y-x} dy$ are the modified moments. The forthcoming subsection defines the method to compute the moments $M_n(\alpha, \beta, k, x)$ efficiently.

Computation of Moments

A well known property for $T_n(y)$ is defined as [30]:

$$\frac{T_n(y) - T_n(x)}{y - x} = 2 \sum_{j=0}^{n-1} U_{n-1-j}(y) T_j(x) = 2 \sum_{j=0}^{n-1} U_{n-1-j}(x) T_j(y),$$
(17)

where the prime indicates the summation whose first term is divided by 2 and $U_n(y)$ is the Chebyshev polynomial of the second kind.

$$M_{n}(\alpha,\beta,k,x) = \int_{-1}^{1} \frac{w(y)T_{n}(y)e^{iky}}{y-x}dy$$

$$= \int_{-1}^{1} \frac{w(y)(T_{n}(y) - T_{n}(x) + T_{n}(x))e^{iky}}{y-x}dy$$

$$= \int_{-1}^{1} \frac{w(y)(T_{n}(y) - T_{n}(x))e^{iky}}{y-x}dy + T_{n}(x)\int_{-1}^{1} \frac{w(y)e^{iky}}{y-x}dy$$

$$= \int_{-1}^{1} w(y) \left(2\sum_{j=0}^{n-1} U_{n-1-j}(x)T_{j}(y)\right)e^{iky}dy + T_{n}(x)\int_{-1}^{1} \frac{w(y)e^{iky}}{y-x}dy$$

$$= 2\sum_{j=0}^{n-1} U_{n-1-j}(x)\int_{-1}^{1} w(y)T_{j}(y)e^{iky}dy + T_{n}(x)\int_{-1}^{1} \frac{w(y)e^{iky}}{y-x}dy$$
(18)

Piessens and Branders [31] have addressed the fourth homogenous recurrence relation for the integral without singularity $\overline{M}_n(\alpha, \beta, k) = \int_{-1}^1 w(y) T_j(y) e^{iky} dy$.

$$ik\overline{M}_{n+2} + 2(n+\alpha+\beta+2)\overline{M}_{n+1} - 2(2\alpha-2\beta+ik)\overline{M}_n - 2(n-\alpha-\beta-2)\overline{M}_{n-1} + ik\overline{M}_{n-2} = 0, \quad n \ge 2, \quad (19)$$

along with four initial values:

$$\overline{M}_{0}^{0} = 2^{\alpha+\beta+1} e^{-ik} \frac{\Gamma(\alpha+1)\Gamma(\beta+1)}{\Gamma(\alpha+\beta+2)} F_{1}(\alpha+1;\alpha+\beta+2;2ik),$$

$$\overline{M}_{1}^{0} = M_{0}(x,\alpha+1,\beta,k) - M_{0}(x,\alpha,\beta,k),$$

$$\overline{M}_{2}^{0} = \frac{i}{k} [2(\alpha+\beta+2)M_{1} - (2\alpha-2\beta+ik)M_{0}],$$

$$\overline{M}_{3}^{0} = \frac{i}{k} [2(\alpha+\beta+3)M_{2} - (4\alpha-4\beta+ik)M_{1} + 2(\alpha+\beta+1)M_{0}],$$
(20)

where $F_1(\alpha + 1; \alpha + \beta + 2; 2ik)$ stands for confluent hypergeometric function of the first kind. Unfortunately the discussed recurrence relation for moments $\overline{M}_n(\alpha, \beta, k)$ is numerically unstable in the forward direction for n > k; in this sense by applying Oliver's algorithm these modified moments can be computed efficiently [31,32].

The integral $\int_{-1}^{1} \frac{w(y)e^{iky}}{y-x} dy$ is computed by the steepest descent method; the original idea was given by Huybrenchs and Vandewalle [33] for sufficiently high oscillatory integrals.

Proposition 1. The Cauchy singular integral $\int_{-1}^{1} \frac{w(y)e^{iky}}{y-x} dy$ can be transformed into:

$$\int_{-1}^{1} \frac{w(y)e^{iky}}{y-x} dy = S_{-1} - S_1 + i\pi w(x)e^{ikx}$$
(21)

where:

$$S_{-1} = i^{\alpha+1} e^{-ik} \int_0^\infty \frac{y^{\alpha} (2-iy)^{\beta}}{-1+iy-x} e^{-ky} dy$$

$$S_1 = (-i)^{\beta+1} e^{ik} \int_0^\infty \frac{y^{\beta} (2+iy)^{\alpha}}{1+iy-x} e^{-ky} dy.$$
(22)

Proof. Readers are referred to [34] for more details. \Box

The generalized Gauss Laguerre quadrature rule can be used to evaluate the integrals S_{-1} and S_1 in the above equation by using the command lagpts in chebfun [35]. Let $\left\{y_j^{\alpha}, w_j^{\alpha}\right\}_{j=1}^k$ be the nodes and weights of the weight functions $y^{\alpha}e^{-y}$ and $\left\{y_j^{\beta}, w_j^{\beta}\right\}_{j=1}^k$ be the nodes and weights of the weight functions $y^{\beta}e^{-y}$ in accordance with the generalized Gauss Laguerre quadrature rule. Moreover, these integrals can be approximated by:

$$S_{-1} \approx Q_{k} = \left(\frac{i}{k}\right)^{\alpha+1} e^{-ik} \sum_{j=1}^{k} w_{j}^{\alpha} \frac{(2 - (i/k)y_{j}^{\alpha})^{\beta}}{(-1 + (i/k)y_{j}^{\alpha} - x)}$$

$$S_{1} \approx Q_{k} = \left(\frac{i}{k}\right)^{\beta+1} e^{ik} \sum_{j=1}^{k} w_{j}^{\beta} \frac{(2 + (i/k)y_{j}^{\beta})^{\alpha}}{(-1 + (i/k)y_{j}^{\beta} - x)}.$$
(23)

 $M_n(\alpha, \beta, k, x)$ is obtained by substituting Equations (19) and (21) into the last equality of Equation (18). Finally, together with Equations (16) and (14), the approximate solution:

$$u_N(x_i) = af(x_i) - \frac{bw(x_i)}{\pi} e^{-ikx_i} \sum_{n=0}^{N+1} a_n M_n(\alpha, \beta, k, x),$$
(24)

for CSIE (1) is derived for different solution cases at equally spaced points.

3. Error Analysis

Lemma 1. [36,37] Let f(x) be a Lipschitz continuous function on [-1, 1] and $P_N[f]$ be the interpolation polynomial of f(x) at N + 1 Clenshaw–Curtis points. Then it follows that:

$$\lim_{N \to \infty} \|f - P_N[f]\|_{\infty} = 0.$$
⁽²⁵⁾

In particular,

• (i) if f(x) is analytic with $|f(x)| \le M$ in an ellipse ε_{ρ} (Bernstein ellipse) with foci ± 1 and major and minor semiaxis lengths summing to $\rho > 1$, then:

$$||f - P_N[f]||_{\infty} \le \frac{4M}{\rho^N(\rho - 1)}.$$
 (26)

• (ii) if f(x) has an absolutely continuous $(\kappa_0 - 1)$ st derivative and a κ_0 th derivative $f^{(\kappa_0)}$ of bounded variation V_{κ_0} on [-1,1] for some $\kappa_0 \ge 1$, then for $N \ge \kappa_0 + 1$:

$$\|f - P_N[f]_{\infty}\| \le \frac{4V_{\kappa_0}}{\kappa_0 \pi N(N-1)\cdots(N-\kappa_0+1)}.$$
(27)

Proposition 2. [29] Suppose that $f(y) \in C^{R+2}[-1,1]$ with $R = [min\{\alpha,\beta\}]$, then the error of the Clenshaw–Curtis–Filon quadrature rule for integral I[f] satisfies:

$$E_N = |I(\alpha, \beta, k, x) - I_N(\alpha, \beta, k, x)| = O(k^{-2 - \min\{\alpha, \beta\}}), \qquad k \to \infty.$$
⁽²⁸⁾

Theorem 2. Suppose that $u_N(x)$ is the approximate solution of u(x) of CSIE for case $\kappa \le 0$, then for error $|u(x) - u_N(x)|, x \in (-1, 1)$, the Clenshaw–Curtis–Filon quadrature is convergent, i.e.:

$$\lim_{N \to \infty} |u(x) - u_N(x)| = 0.$$
 (29)

Proof. Suppose that $x \notin Y_{N+1}$, $f \in C^2[-1, 1]$ and let

$$Q(y) = \begin{cases} \frac{f(y)-f(x)}{y-x}, & y \neq x\\ f'(x), & y = x. \end{cases}$$

It is stated that $Q(y) \in C^1[-1,1]$ and $||Q'||_{\infty} \leq \frac{3}{2}||f''||_{\infty}$, in addition $R(y) = \frac{P_{N+1}(y)-f(x)}{y-x}$ is a polynomial of degree at most *N*. Then error for solutions u(x) and $u_N(x)$ to CSIE for cases $\kappa \leq 0$ is defined as:

$$u(x) = af(x) - \frac{b}{\pi}e^{-ikx}w(x)\int_{-1}^{1}\frac{w^{*}(y)f(y)e^{iky}}{y-x}dy,$$
$$u_{N}(x) = af(x) - \frac{b}{\pi}e^{-ikx}w(x)\int_{-1}^{1}\frac{w^{*}(y)\bar{P}_{N+1}(y)e^{iky}}{y-x}dy.$$

Then:

$$\begin{split} |u(x) - u_N(x)| &= \left| a(f(x) - f(x)) - \frac{b}{\pi} e^{-ikx} w(x) \int_{-1}^1 w^*(y) (Q(y) - R(y)) e^{iky} dy \right| \\ &\leq \frac{b}{\pi} w(x) \int_{-1}^1 w^*(y) dy ||Q(y) - R(y)||_{\infty} \\ &= D ||Q(y) - R(y)||_{\infty}. \end{split}$$

where $D = \frac{bw(x)2^{\alpha+\beta+1}\Gamma(\alpha+1)\Gamma(\beta+1)}{\pi\Gamma(\alpha+\beta+2)}$.

4. Numerical Examples

Example 1. Let us consider the CSIE of the second kind:

$$\frac{u(x)}{\sqrt{2}} + \frac{1}{\sqrt{2}\pi} e^{-ikx} \int_{-1}^{1} \frac{u(y)e^{iky}}{y-x} dy = \frac{f(x)}{\sqrt{2}}$$
(30)

where f(x) = cos(x). For x = 0.5 and $a = b = \frac{1}{\sqrt{2}}$, we get values of $\alpha = 0.25$ and $\beta = 0.25$ from Equation (3) for $\kappa = 0$. The absolute error for u(x) is presented in Tables 1 and 2 below.

N = 5**N** = 10 **N** = 20 k 3.9207×10^{-14} 1.1102×10^{-16} 4.6387×10^{-9} 50 1.0881×10^{-9} $4.9564 \times {10}^{-15}$ 100 0 3.8093×10^{-11} 4.0030×10^{-16} 2.4825×10^{-16} 1000 5.1593×10^{-13} 2.2204×10^{-16} 1.1102×10^{-16} 10000

Table 1. Absolute error for $\kappa = 0$, bounded at x = 1.

Table 2. Absolute error for $\kappa = 0$, bounded at x = -1.

k	N = 5	N = 10	N = 20
50	1.1156×10^{-9}	9.1854×10^{-15}	1.1102×10^{-16}
100	3.2791×10^{-10}	5.6610×10^{-16}	1.1102×10^{-16}
1000	1.7225×10^{-12}	2.2204×10^{-16}	2.2204×10^{-16}
10000	7.3056×10^{-15}	3.3307×10^{-16}	3.3307×10^{-16}





Figure 1. The mixed boundary value problem.

Taken from [18], it has the analytic solution $\phi(x,t) = \frac{2}{\pi} \arctan \frac{2y}{1-x^2-t^2}$. It can further be reduced to the following integral equation for $\kappa = -1$ and for $\alpha = \beta = \frac{1}{2}$.

$$\frac{-1}{\pi} \int_{-1}^{1} \frac{u(y)}{y-x} dy = C_1 + \frac{1}{\pi} \left[\frac{1-x}{2} \log(1-x) + \frac{1+x}{2} \log(1+x) - \log(2+x) - 1 \right]$$
(31)

Here C_1 *is a constant defined as* $C_1 = 0.4192007182789807$. *Furthermore if* u(x) *is known, the solution of the above boundary value can be derived as:*

$$\phi(\mu,\nu) = \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{\nu u(y,0)}{(y-\mu)^2 + \nu^2} dy$$

where:

$$u(y,0) = \begin{cases} u(y) + (1-y)/2, & |y| \le 1, \\ 1 & t \in [-2, -1], \\ 0, & otherwise. \end{cases}$$
(32)

So here we just solve u(x) for simplicity. Figure 2 illustrates the absolute error for u(x).



Figure 2. The absolute error for u(x), for x = 0.6.

Figure 2 shows that absolute error for u(x) decreases for greater values of N.

Example 3. [10,26] For CSIE with *k* = 0:

$$\int_{-1}^{1} \frac{u(y)}{y-x} dy = x^4 + 5x^3 + 2x^2 + x - \frac{11}{8}$$
(33)

in the case a = 0 *and b* = 1, *where* α *and* β *are derived from Equation* (3) *and the exact values of* u(y) *for cases* $\kappa \leq 0$ *for the solution bounded at* x = -1, x = 1, $x = \pm 1$ *are given as:*

$$u(y) = \frac{1}{\pi} \sqrt{\frac{1+y}{1-y}} \left[y^4 + 4y^3 - 5/2y^2 + y - 7/2 \right]$$

$$u(y) = \frac{-1}{\pi} \sqrt{\frac{1-y}{1+y}} \left[y^4 + 6y^3 + 15/2y^2 + 6y + 7/2 \right]$$

$$u(y) = \frac{-1}{\pi} \sqrt{1-y^2} \left[y^3 + 5y^2 + 5/2y + 7/2 \right].$$

(34)

Table 3 presents the absolute error for the above three cases.

Table 3. Absolute error for case $\kappa \le 0, k = 0$.

x .	Error			
	$\kappa = -1$	$\kappa = 0$, boundned at $x = -1$	$\kappa = 0$, boundned at $x = 1$	
-0.6	0	1.1102×10^{-16}	4.4409×10^{-16}	
-0.2	3.3307×10^{-16}	2.2204×10^{-16}	4.4409×10^{-16}	
0.2	2.2204×10^{-16}	4.4409×10^{-16}	0	
0.6	0	2.2204×10^{-16}	4.4409×10^{-16}	

Clearly, Table 3 shows that obtained absolute errors are significantly good for really small values of N, N = 5, that can never be achieved in [10,26]. The exact value for u(x) in the above examples is obtained through Mathematica 11, while the approximated results are calculated using Matlab R2018a on a 4 GHz personal laptop with 8 GB of RAM.

5. Conclusions

In the presented research work, the Clenshaw–Curtise–Filon quadrature is used to get higher order accuracy. Absolute errors are presented in Tables 1 and 2 for solutions of highly oscillatory CSIEs for $\kappa = 0$. For larger values of *N*, Figure 2 shows the absolute error for u(x) for mixed the boundary value problem, whereas for frequency k = 0, the proposed quadrature posseses higher accuracy than the schemes claimed in [10,26]; Table 3 addresses this very well. This shows that the quadrature rule is quite accurate with the exact solution.

Author Contributions: Conceptualization, SAIRA, S.X. and G.L.; Methodology, SAIRA; Supervision, S.X.; Writing (original draft), SAIRA; Writing (review and editing), SAIRA, S.X. and G.L.

Funding: This research received no external funding.

Conflicts of Interest: The authors declare no conflict of interest.

References

- 1. Polyanin, A.D.; Manzhirov, A.V. Handbook of Integral Equations; CRC Press: Boca Raton, FL, USA, 1998.
- 2. Li, J.; Wang, X.; Xiao, S.; Wang, T. A rapid solution of a kind of 1D Fredholm oscillatory integral equation. *J. Comput. Appl. Math.* **2012**, *236*, 2696–2705.
- 3. Ursell, F. Integral equations with a rapidly oscillating kernel. J. Lond. Math. Soc. 1969, 1, 449–459.
- Yalcinbas, S.; Aynigul, M. Hermite series solutions of linear Fredholm integral equations. *Math. Comput. Appl.* 2011, 16, 497–506.
- 5. Fang, C.; He, G.; Xiang, S. Hermite-Type Collocation Methods to Solve Volterra Integral Equations with Highly Oscillatory Bessel Kernels. *Symmetry* **2019**, *11*, 168.
- 6. Babolian, E.; Hajikandi, A.A. The approximate solution of a class of Fredholm integral equations with a weakly singular kernel. *J. Comput. Appl. Math.* **2011**, *235*, 1148–1159.
- 7. Aimi, A.; Diligenti, M.; Monegato, G. Numerical integration schemes for the BEM solution of hypersingular integral equations. *Int. J. Numer. Method Eng.* **1999**, *45*, 1807–1830.
- 8. Beyrami, H.; Lotfi, T.; Mahdiani, K. A new efficient method with error analysis for solving the second kind Fredholm integral equation with Cauchy kernel. *J. Comput. Appl. Math.* **2016**, *300*, 385–399.
- 9. Setia, A. Numerical solution of various cases of Cauchy type singular integral equation. *Appl. Math. Comput.* **2014**, *230*, 200–207.
- 10. Eshkuvatov, Z.K.; Long, N.N.; Abdulkawi, M. Approximate solution of singular integral equations of the first kind with Cauchy kernel. *Appl. Math. Lett.* **2009**, *22*, 651–657.
- 11. Cuminato, J.A. Uniform convergence of a collocation method for the numerical solution of Cauchy-type singular integral equations: A generalization. *IMA J. Numer. Anal.* **1992**, *12*, 31–45.
- 12. Cuminato, J.A. On the uniform convergence of a perturbed collocation method for a class of Cauchy integral equations. *Appl. Numer. Math.* **1995**, *16*, 439–455.
- 13. Karczmarek, P.; Pylak, D.; Sheshko, M.A. Application of Jacobi polynomials to approximate solution of a singular integral equation with Cauchy kernel. *Appl. Math. Comput.* **2006**, *181*, 694–707.
- 14. Lifanov, I.K. *Singular Integral Equations and Discrete Vortices*; Walter de Gruyter GmbH : Berlin, Germany, 1996.
- 15. Ladopoulos, E.G. *Singular Integral Equations: Linear and Non-Linear Theory and Its Applications in Science and Engineering;* Springer Science and Business Media: Berlin, Germany, 2013.
- 16. Muskhelishvili, N.I. *Some Basic Problems of the Mathematical Theory of Elasticity;* Springer Science and Business Media: Berlin, Germany, 2013.
- 17. Martin, P.A.; Rizzo, F.J. On boundary integral equations for crack problems. *Proc. R. Soc. Lond. A Math. Phys. Sci.* **1989**, 421, 341–355.

- 18. Cuminato, J.A. Numerical solution of Cauchy-type integral equations of index- 1 by collocation methods. *Adv. Comput. Math.* **1996**, *6*, 47–64.
- 19. Asheim, A.; Huybrechs, D. Complex Gaussian quadrature for oscillatory integral transforms. *IMA J. Num. Anal.* **2013**, *33*, 1322–1341.
- 20. Chen, R.; An, C. On evaluation of Bessel transforms with oscillatory and algebraic singular integrands. *J. Comput. Appl. Math.* **2014**, *264*, 71–81.
- 21. Erdelyi, A. Asymptotic representations of Fourier integrals and the method of stationary phase. *SIAM* **1955**, *3*, 17–27.
- 22. Olver, S. Numerical Approximation of Highly Oscillatory Integrals. Ph.D. Thesis, University of Cambridge, Cambridge, UK, 2008.
- 23. Milovanovic, G.V. Numerical calculation of integrals involving oscillatory and singular kernels and some applications of quadratures. *Comput. Math. Appl.* **1998**, *36*, 19–39.
- 24. Dzhishkariani, A.V. The solution of singular integral equations by approximate projection methods. *USSR Comput. Math. Math. Phys.* **1979**, *19*, 61–74.
- 25. Chakrabarti, A.; Berghe, G.V. Approximate solution of singular integral equations. *Appl. Math. Lett.* **2004**, 17, 553–559.
- 26. Dezhbord, A.; Lotfi, T.; Mahdiani, K. A new efficient method for cases of the singular integral equation of the first kind. *J. Comput. Appl. Math.* **2016**, *296*, 156–169.
- 27. He, G.; Xiang, S. An improved algorithm for the evaluation of Cauchy principal value integrals of oscillatory functions and its application. *J. Comput. Appl. Math.* **2015**, *280*, 1–13.
- 28. Trefethen, L.N. C hebyshev Polynomials and Series, Approximation theorey and approximation practice. *Soc. Ind. Appl. Math.* **2013**, *128*, 17-19.
- 29. Liu, G.; Xiang, S. Clenshaw Curtis type quadrature rule for hypersingular integrals with highly oscillatory kernels. *Appl. Math. Comput.* **2019**, *340*, 251–267.
- 30. Wang, H.; Xiang, S. Uniform approximations to Cauchy principal value integrals of oscillatory functions. *Appl. Math. Comput.* **2009**, *215*, 1886–1894.
- 31. Piessens, R.; Branders, M. On the computation of Fourier transforms of singular functions. *J. Comput. Appl. Math.* **1992**, *43*, 159–169.
- 32. Oliver, J. The numerical solution of linear recurrence relations. *Numer. Math.* 1968, 114, 349–360.
- 33. Huybrechs, D.; Vandewalle, S. On the evaluation of highly oscillatory integrals by analytic continuation. *Siam J. Numer. Anal.* **2006**, *44*, 1026–1048.
- 34. Wang, H.; Xiang, S. On the evaluation of Cauchy principal value integrals of oscillatory functions. *J. Comput. Appl. Math.* **2010**, *234*, 95–100.
- 35. Dominguez, V.; Graham, I.G.; Smyshlyaev, V.P. Stability and error estimates for Filon Clenshaw Curtis rules for highly oscillatory integrals. *IMA J. Numer. Anal.* **2011**, *31*, 1253–1280.
- 36. Xiang, S.; Chen, X.; Wang, H. Error bounds for approximation in Chebyshev points. *Numer. Math.* **2010**, *116*, 463–491.
- 37. Xiang, S. Approximation to Logarithmic-Cauchy Type Singular Integrals with Highly Oscillatory Kernels. *Symmetry* **2019**, *11*, 728.



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses/by/4.0/).