



Article L_p Radial Blaschke-Minkowski Homomorphisms and L_p Dual Affine Surface Areas

Zhonghuan Shen ^{1,2} and Weidong Wang ^{1,2,*}

- ¹ Department of Mathematics, China Three Gorges University, Yichang 443002, China; szh@ctgu.edu.cn
- ² Three Gorges Mathematical Research Center, China Three Gorges University, Yichang 443002, China
- * Correspondence: wdwxh722@163.com

Received: 17 February 2019; Accepted: 3 April 2019; Published: 10 April 2019



Abstract: Schuster introduced the notion of radial Blaschke-Minkowski homomorphism and considered the Busemann-Petty problem for volume forms. Whereafter, Wang, Liu and He presented the L_p radial Blaschke-Minkowski homomorphisms and extended Schuster's results. In this paper, associated with L_p dual affine surface areas, we give an affirmative and a negative form of the Busemann-Petty problem and establish two Brunn-Minkowski inequalities for the L_p radial Blaschke-Minkowski homomorphisms.

Keywords: Busemann-Petty problem; L_p radial Blaschke-Minkowski homomorphism; L_p dual affine surface area; Brunn-Minkowski inequality

MSC: 52A20; 52A39; 52A40

1. Introduction

If *K* is a compact star shaped (about the origin) in *n*-dimensional Euclidean space \mathbb{R}^n , then its radial function, $\rho_K = \rho(K, \cdot) : \mathbb{R}^n \setminus \{0\} \to [0, \infty)$, is defined by (see [1])

$$\rho(K, x) = \max\{\lambda \ge 0 : \lambda x \in K\}, \ x \in \mathbb{R}^n \setminus \{0\}.$$

If $\rho(K, \cdot)$ is positive and continuous, K will be called a star body (about the origin). The set of all star bodies in \mathbb{R}^n denotes by S_o^n . For the set of all origin-symmetric star bodies, we write S_{os}^n . Let S^{n-1} denote the unit sphere in \mathbb{R}^n . Two star bodies K and L are said to be dilated (of one another) if $\rho_K(u)/\rho_L(u)$ is independent of $u \in S^{n-1}$.

Intersection bodies were explicitly defined and named by Lutwak (see [2]). For $K \in S_o^n$, the intersection body, *IK*, of *K* is a star body whose radial function is defined by

$$\rho(IK, u) = V_{n-1}(K \cap u^{\perp})$$

for all $u \in S^{n-1}$. Here u^{\perp} is the (n-1)-dimensional hyperplane orthogonal to u and V_{n-1} denotes the (n-1)-dimensional volume.

During past three decades, the investigations of intersection bodies have received great attention from many articles (see [1–12]). In particular, intersection bodies led to the following Busemann-Petty problem (see [2]).

Problem 1 (Busemann-Petty problem). *For* $K, L \in S_0^n$, *is there the implication*

$$IK \subseteq IL \Rightarrow V(K) \leq V(L)$$
?

Here V(K) *denotes the n-dimensional volume of body* K*.*

Remark 1. Problem 1 was stated by Lutwak (see [2]). If K, $L \in S_{os}^n$, then Problem 1 is called the symmetric Busemann-Petty problem. Gardner [13], Zhang [14] showed that the symmetric Busemann-Petty problem has an affirmative answer for $n \leq 4$ and a negative answer for $n \geq 5$.

For Problem 1, Lutwak [2] gave its an affirmative answer if K is restricted to the class of intersection bodies and two negative answers if K is not origin-symmetric star body or L is not intersection body. For more research on solutions to the Busemann-Petty problem, see e.g., [1,13–20].

In 2006, based on the properties of intersection bodies, Schuster [21] introduced the radial Blaschke-Minkowski homomorphism, which is the more general intersection operator as follows:

Definition 1. A map $\Psi : S_o^n \to S_o^n$ is called a radial Blaschke-Minkowski homomorphism if it satisfies the following conditions:

- (1) Ψ is continuous;
- For all $M, N \in \mathcal{S}_{o}^{n}, \Psi(M \widehat{+} N) = \Psi M \widehat{+} \Psi N;$ (2)
- $\Psi(\varphi M) = \varphi \Psi M$ for all $M \in S_0^n$ and all $\varphi \in SO(n)$. (3)

Here " $\hat{+}$ " and " $\hat{+}$ " denote radial Blaschke addition and radial Minkowski addition, respectively; SO(n) denotes the group of rotation transformations.

Whereafter, Schuster ([22]) considered the following Busemann-Petty problem for radial Blaschke-Minkowski homomorphisms.

Problem 2. Let $\Psi : S_0^n \to S_0^n$ be a radial Blaschke-Minkowski homomorphism. For $K, L \in S_0^n$, is there the implication

$$\Psi K \subseteq \Psi L \Rightarrow V(K) \le V(L)?$$

Obviously, Problem 2 is a more general Busemann-Petty problem compared with Problem 1. For Problem 2, Schuster [22] gave an affirmative answer if K belongs to ΨS_0^{n} (the range of Ψ) and two negative forms.

In 2011, Wang, Liu and He [23] extended Schuster's radial Blaschke-Minkowski homomorphisms to L_p analogies, and gave the notion of L_p radial Blaschke-Minkowski homomorphisms as follows:

Definition 2. For p > 0, a map $\Psi_p : S_o^n \to S_o^n$ is called a L_p radial Blaschke-Minkowski homomorphism if it satisfies the following conditions:

- (1) Ψ_p is continuous;
- (2) For all $M, N \in \mathcal{S}_{o}^{n}, \Psi_{p}(M \widehat{+}_{p}N) = \Psi_{p}M \widetilde{+}_{p}\Psi_{p}N;$ (3) $\Psi_{p}(\varphi M) = \varphi \Psi_{p}M$ for all $M \in \mathcal{S}_{o}^{n}$ and all $\varphi \in SO(n).$

Here " $\hat{+}_p$ " and " $\hat{+}_p$ " denote L_p radial Blaschke addition and L_p radial Minkowski addition, respectively.

Meanwhile, associated with L_p radial Blaschke-Minkowski homomorphisms, Wang et al. [23] extended Schuster's results including the Busemann-Petty Problem 2. In recent years, a lot of important conclusions for the radial Blaschke-Minkowski homomorphisms and their L_p analogies were obtained (see e.g., [22–33]).

The L_p dual affine surface areas firstly were introduced by Wang, Yuan and He (see [34]). Here, we improve Wang, Yuan and He's definition as follows: For $K \in S_0^n$ and p > 0, the L_p dual affine surface area, $\tilde{\Omega}_p(K)$, of *K* is defined by

$$n^{-\frac{p}{n}}\widetilde{\Omega}_p(K)^{\frac{n+p}{n}} = \sup\{n\widetilde{V}_p(K,Q^*)V(Q)^{\frac{p}{n}}: Q \in \mathcal{S}_o^n\}.$$
(1)

Here $\tilde{V}_p(M, N)$ denotes the L_p dual mixed volume of $M, N \in S_o^n$, and Q^* denotes the polar of Q which is defined by (see [1])

$$Q^* = \{x : x \cdot y \le 1, y \in Q\}, \ x \in \mathbb{R}^n.$$

If *Q* belongs to the set of convex bodies (or star bodies) whose centroid at the origin, then Equation (1) is just Wang, Yuan and He's (or Wang and Wang's) definition (see [34] or [35]). For the studies of L_p dual affine surface areas, also see [36–41].

Remark 2. Recall that Lutwak's L_p affine surface area was defined as follows (see [42]): For $K \in \mathcal{K}_o^n$ and $p \ge 1$, the L_p affine surface area, $\Omega_p(K)$, of K is defined by

$$n^{-\frac{p}{n}}\Omega_p(K)^{\frac{n+p}{n}} = \inf\{nV_p(K,Q^*)V(Q)^{\frac{p}{n}} : Q \in S_o^n\}.$$
(2)

Here, \mathcal{K}_{o}^{n} denotes the set of convex bodies (compact, convex subsets with nonempty interiors) containing the origin in their interiors in \mathbb{R}^{n} and $V_{p}(M, N)$ is the L_{p} mixed volume of M and N (see [42]). Compare to Equation (1) and Equation (2), we see that Equation (1) is really the duality of Equation (2).

In this paper, associated with L_p dual affine surface areas, we research the L_p radial Blaschke-Minkowski homomorphisms. We firstly consider the following Busemann-Petty problem of L_p radial Blaschke-Minkowski homomorphisms.

Problem 3. Let p > 0 and $\Psi_p : S_o^n \to S_o^n$ be a L_p radial Blaschke-Minkowski homomorphism. For $K, L \in S_o^n$, is there the implication

$$\Psi_p K \subseteq \Psi_p L \Rightarrow \widetilde{\Omega}_p(K) \le \widetilde{\Omega}_p(L)?$$

Let $\Psi_p S_o^n$ denote the range of Ψ_p and $\Psi_p^* S_o^n$ denote the set of polars of all elements in $\Psi_p S_o^n$, then $\Psi_p^* S_o^n \subseteq S_o^n$. From this, we write that

$$n^{-\frac{p}{n}}\widetilde{\Omega}_{p}^{\circ}(K)^{\frac{n+p}{n}} = \sup\{n\widetilde{V}_{p}(K,Q^{*})V(Q)^{\frac{p}{n}}: Q \in \Psi_{p}^{*}\mathcal{S}_{o}^{n}\}.$$
(3)

For Problem 3, according to Equation (3), we obtain an affirmative form as follows:

Theorem 1. For p > 0, let $\Psi_p : S_o^n \to S_o^n$ be a L_p radial Blaschke-Minkowski homomorphism. If $K, L \in S_o^n$, then

$$\Psi_p K \subseteq \Psi_p L \Rightarrow \widetilde{\Omega}_p^{\circ}(K) \le \widetilde{\Omega}_p^{\circ}(L).$$

In addition, $\widetilde{\Omega}_p^{\circ}(K) = \widetilde{\Omega}_p^{\circ}(L)$ when $\Psi_p K = \Psi_p L$.

Furthermore, when $K \notin S_{os}^n$, by Equation (1) we give the following a negative form of Problem 3.

Theorem 2. For $K, L \in S_o^n$ and p > 0, let $\Psi_p : S_o^n \to S_o^n$ be a L_p radial Blaschke-Minkowski homomorphism. If $K \notin S_{os'}^n$ then there exists $L \in S_o^n$ such that

$$\Psi_p K \subset \Psi_p L.$$

However,

$$\widetilde{\Omega}_p(K) > \widetilde{\Omega}_p(L).$$

Next, associated with L_q radial Minkowski sum and L_q harmonic Blaschke sum of star bodies, we establish the following L_p dual affine surface area forms of Brunn-Minkowski inequalities for the L_p radial Blaschke-Minkowski homomorphisms, respectively.

Theorem 3. *If* $K, L \in S_o^n$, $0 and <math>0 < q \le n - p$, then

$$\widetilde{\Omega}_p(\Psi_p(K\widetilde{+}_qL))^{\frac{pq(n+p)}{n(n-p)^2}} \le \widetilde{\Omega}_p(\Psi_pK)^{\frac{pq(n+p)}{n(n-p)^2}} + \widetilde{\Omega}_p(\Psi_pL)^{\frac{pq(n+p)}{n(n-p)^2}},\tag{4}$$

with equality if and only if K and L are dilated.

Let q = n - p in Theorem 3 and notice that $K +_{n-p}L = K +_pL$ (see Equation (7)), we obtain a Brunn-Minkowski inequality for the L_p radial Blaschke sum $K +_pL$.

Corollary 1. If $K, L \in S_0^n$, 0 , then

$$\widetilde{\Omega}_p(\Psi_p(K\widehat{+}_pL))^{\frac{p(n+p)}{n(n-p)}} \leq \widetilde{\Omega}_p(\Psi_pK)^{\frac{p(n+p)}{n(n-p)}} + \widetilde{\Omega}_p(\Psi_pL)^{\frac{p(n+p)}{n(n-p)}},$$

with equality if and only if K and L are dilated.

Theorem 4. *If* $K, L \in S_0^n$, 0 and <math>0 , then

$$\frac{\widetilde{\Omega}_{p}(\Psi_{p}(K \mp_{q} L))^{\frac{p(n+p)(n+q)}{n(n-p)^{2}}}}{V(K \mp_{q} L)} \leq \frac{\widetilde{\Omega}_{p}(\Psi_{p} K)^{\frac{p(n+p)(n+q)}{n(n-p)^{2}}}}{V(K)} + \frac{\widetilde{\Omega}_{p}(\Psi_{p} L)^{\frac{p(n+p)(n+q)}{n(n-p)^{2}}}}{V(L)},$$
(5)

with equality if and only if K and L are dilated. Here $K \mp_q L$ denotes the L_q harmonic Blaschke sum of K and L.

The proofs of Theorem 1 and Theorem 2 are completed in Section 3. In Section 4, we will give the proofs of Theorem 3 and Theorem 4.

2. Background Materials

2.1. General L_p Radial Blaschke Bodies

For $K, L \in S_o^n$, real $p \neq 0$ and $\lambda, \mu \geq 0$ (not both 0), the L_p radial Minkowski combination, $\lambda \cdot K + \mu \mu \cdot L \in S_o^n$, of *K* and *L* is defined by (see [43,44])

$$\rho(\lambda \cdot K + p\mu \cdot L, \cdot)^p = \lambda \rho(K, \cdot)^p + \mu \rho(L, \cdot)^p.$$

Here " $\tilde{+}_p$ " denotes the L_p radial Minkowski sum and $\lambda \cdot K = \lambda^{1/p} K$. The case p = 1 yields the radial Minkowski combination $\lambda \cdot K + \mu \cdot L$.

In 2015, Wang and Wang [35] defined the L_p radial Blaschke combinations of star bodies as follows: For $K, L \in S_o^n$, n > p > 0 and $\lambda, \mu \ge 0$ (not both 0), the L_p radial Blaschke combination, $\lambda \circ K + p\mu \circ L \in S_o^n$, of K and L is defined by

$$\rho(\lambda \circ K \widehat{+}_p \mu \circ L, \cdot)^{n-p} = \lambda \rho(K, \cdot)^{n-p} + \mu \rho(L, \cdot)^{n-p}.$$
(6)

Here " $\hat{+}_p$ " denotes the L_p radial Blaschke sum and $\lambda \circ K = \lambda^{1/(n-p)}K$. If p = 1, then $\lambda \circ K \hat{+}_p \mu \circ L$ is the radial Blaschke combination $\lambda \circ K \hat{+} \mu \circ L$ (see [1]).

From the definitions of above two combinations, we easily see

$$\lambda \cdot K \widetilde{+}_{n-p} \mu \cdot L = \lambda \circ K \widehat{+}_p \mu \circ L. \tag{7}$$

In Equation (6), let

$$\lambda = f_1(\tau) = \frac{(1+\tau)^2}{2(1+\tau^2)}, \quad \mu = f_2(\tau) = \frac{(1-\tau)^2}{2(1+\tau^2)}$$
(8)

with $\tau \in [-1, 1]$ and L = -K, and write

$$\widehat{\nabla}_{p}^{\tau}K = f_{1}(\tau) \circ K \widehat{+}_{p}f_{2}(\tau) \circ (-K).$$
(9)

We call $\widehat{\nabla}_p^{\tau} K$ the general L_p radial Blaschke body of K. From Equations (8) and (9), we easily see that $\widehat{\nabla}_p^1 K = K$, $\widehat{\nabla}_p^{-1} K = -K$ and

$$\widehat{\nabla}_{p}^{0}K = \frac{1}{2} \circ K \widehat{+}_{p} \frac{1}{2} \circ (-K).$$
(10)

For the general L_p radial Blaschke bodies, by Equation (8) we know that $f_1(\tau) + f_2(\tau) = 1$. This and Equation (9) give that if $K \in S_{os}^n$ then $\widehat{\nabla}_p^{\tau} K \in S_{os}^n$. If $K \notin S_{os}^n$, we have the following conclusion.

Theorem 5. For $K, L \in S_o^n$ and p > 0. If $K \notin S_{os}^n$, then for $\tau \in [-1, 1]$,

$$\widehat{\nabla}_{p}^{\tau}K \in \mathcal{S}_{os}^{n} \Leftrightarrow \tau = 0.$$
(11)

Proof. If $\tau = 0$, by Equation (10) we immediately get $\widehat{\nabla}_{p}^{\tau} K \in \mathcal{S}_{os}^{n}$.

Conversely, since $\rho_M(-u) = \rho_{-M}(u)$ for each $M \in S_o^n$ and any $u \in S^{n-1}$, thus if $\widehat{\nabla}_p^{\tau} K \in S_{os}^n$, then for all $u \in S^{n-1}$,

$$\rho_{\widehat{\nabla}_p^{\tau}K}^{n-p}(u) = \rho_{-\widehat{\nabla}_p^{\tau}K}^{n-p}(u) = \rho_{\widehat{\nabla}_p^{\tau}K}^{n-p}(-u).$$

By Equation (9) we have

$$\rho_{f_1(\tau)\circ K\hat{+}_p f_2(\tau)\circ (-K)}^{n-p}(u) = \rho_{f_1(\tau)\circ K\hat{+}_p f_2(\tau)\circ (-K)}^{n-p}(-u).$$

This together with Equation (6) yields

$$f_1(\tau)\rho_K^{n-p}(u) + f_2(\tau)\rho_{-K}^{n-p}(u) = f_1(\tau)\rho_K^{n-p}(-u) + f_2(\tau)\rho_{-K}^{n-p}(-u),$$

i.e.,

$$f_1(\tau)\rho_K^{n-p}(u) + f_2(\tau)\rho_{-K}^{n-p}(u) = f_1(\tau)\rho_{-K}^{n-p}(u) + f_2(\tau)\rho_K^{n-p}(u),$$

hence

$$[f_1(\tau) - f_2(\tau)][\rho_K^{n-p}(u) - \rho_{-K}^{n-p}(u)] = 0.$$

Since $K \notin S_{os}^n$ implies $\rho_K^{n-p}(u) - \rho_{-K}^{n-p}(u) \neq 0$ for all $u \in S^{n-1}$, thus we obtain

 $f_1(\tau) - f_2(\tau) = 0.$

This and Equation (8) give $\tau = 0$. \Box

2.2. L_p Dual Mixed Volumes

Based on the L_p radial Minkowski combinations of star bodies, a class of L_p dual mixed volumes were introduced as follows (see [45,46]): For $M, N \in S_o^n$, $p \neq 0$ and $\varepsilon > 0$, the L_p dual mixed volume, $\tilde{V}_p(M, N)$, of M and N is defined by

$$\frac{n}{p}\widetilde{V}_p(M,N) = \lim_{\varepsilon \to 0^+} \frac{V(M + p\varepsilon \cdot N) - V(M)}{\varepsilon}.$$

From the above definition, L_p dual mixed volume has the following integral representation (see [45,46]):

$$\widetilde{V}_p(M,N) = \frac{1}{n} \int_{S^{n-1}} \rho_M^{n-p}(u) \rho_N^p(u) du.$$
(12)

2.3. L_q Harmonic Blaschke Sums

The harmonic Blaschke sums of star bodies were introduced by Lutwak (see [47]). For $M, N \in S_o^n$, the harmonic Blaschke sum, $M \mp N \in S_o^n$, of M and N is defined by

$$\frac{\rho(M \mp N, \cdot)^{n+1}}{V(M \mp N)} = \frac{\rho(M, \cdot)^{n+1}}{V(M)} + \frac{\rho(N, \cdot)^{n+1}}{V(N)}.$$

Based on above definition, Feng and Wang ([48]) defined the L_q harmonic Blaschke sums as follows: For $M, N \in S_o^n$, real q > -n, the L_q harmonic Blaschke sum, $M \mp_q N \in S_o^n$, of M and N is given by

$$\frac{\rho(M \mp_q N, \cdot)^{n+q}}{V(M \mp_q N)} = \frac{\rho(M, \cdot)^{n+q}}{V(M)} + \frac{\rho(N, \cdot)^{n+q}}{V(N)}.$$
(13)

3. A Type of Busemann-Petty Problem

Theorems 1 and 2 show a type of Busemann-Petty Problem of the L_p radial Blaschke-Minkowski homomorphisms for the L_p dual affine surface areas. In this section, we will prove them. In order to prove Theorem 1, the following lemma is essential.

Lemma 1 ([23]). *If* $M, N \in S_o^n$ *and* p > 0*, then*

$$\widetilde{V}_p(M, \Psi_p N) = \widetilde{V}_p(N, \Psi_p M).$$
(14)

Proof of Theorem 1. Since $\Psi_p K \subseteq \Psi_p L$, thus using Equation (12) we know that for p > 0 and any $M \in S_o^n$,

$$\widetilde{V}_p(M, \Psi_p K) \leq \widetilde{V}_p(M, \Psi_p L).$$

This together with Equation (14) yields

$$\widetilde{V}_p(K, \Psi_p M) \le \widetilde{V}_p(L, \Psi_p M).$$
(15)

Hence, by Equation (3) we have

$$n^{-\frac{p}{n}}\widetilde{\Omega}_{p}^{\circ}(K)^{\frac{n+p}{n}} = \sup\{n\widetilde{V}_{p}(K,Q^{*})V(Q)^{\frac{p}{n}}: Q \in \Psi_{p}^{*}\mathcal{S}_{o}^{n}\}$$
$$= \sup\{n\widetilde{V}_{p}(K,\Psi_{p}M)V(\Psi_{p}^{*}M)^{\frac{p}{n}}:\Psi_{p}^{*}M \in \Psi_{p}^{*}\mathcal{S}_{o}^{n}\}$$
$$\leq \sup\{n\widetilde{V}_{p}(L,\Psi_{p}M)V(\Psi_{p}^{*}M)^{\frac{p}{n}}:\Psi_{p}^{*}M \in \Psi_{p}^{*}\mathcal{S}_{o}^{n}\}$$
$$= n^{-\frac{p}{n}}\widetilde{\Omega}_{p}^{\circ}(L)^{\frac{n+p}{n}},$$

this gives

$$\widetilde{\Omega}_p^{\circ}(K) \leq \widetilde{\Omega}_p^{\circ}(L).$$

Obviously, we see that $\widetilde{\Omega}_p^{\circ}(K) = \widetilde{\Omega}_p^{\circ}(L)$ when $\Psi_p K = \Psi_p L$. \Box

The proof of Theorem 2 needs the following lemmas.

Lemma 2. If $K, L \in S_o^n$, $\lambda, \mu \ge 0$ (not both zero) and 0 , then

$$\widetilde{\Omega}_p(\lambda \circ K \widehat{+}_p \mu \circ L)^{\frac{n+p}{n}} \le \lambda \widetilde{\Omega}_p(K)^{\frac{n+p}{n}} + \mu \widetilde{\Omega}_p(L)^{\frac{n+p}{n}}.$$
(16)

with equality in Equation (16) for λ , $\mu > 0$ if and only if K and L are dilated. For $\lambda = 0$ or $\mu = 0$, Equation (16) becomes an equality.

Proof. From Equations (2) and (6), we have

$$\begin{split} n^{-\frac{p}{n}}\widetilde{\Omega}_{p}(\lambda\circ K\widehat{+}_{p}\mu\circ L)^{\frac{n+p}{n}} \\ &= \sup\left\{n\widetilde{V}_{p}(\lambda\circ K\widehat{+}_{p}\mu\circ L,Q^{*})V(Q)^{\frac{p}{n}}:Q\in\mathcal{S}_{o}^{n}\right\} \\ &= \sup\left\{\left[\int_{S^{n-1}}\rho_{\lambda\circ K\widehat{+}_{p}\mu\circ L}^{n-p}(u)\rho_{Q^{*}}^{p}(u)du\right]V(Q)^{\frac{p}{n}}:Q\in\mathcal{S}_{o}^{n}\right\} \\ &= \sup\left\{\left[\int_{S^{n-1}}[\lambda\rho_{K}^{n-p}(u)+\mu\rho_{L}^{n-p}(u)]\rho(Q^{*},u)^{p}du\right]V(Q)^{\frac{p}{n}}:Q\in\mathcal{S}_{o}^{n}\right\} \\ &= \sup\left\{\lambda\left[\int_{S^{n-1}}\rho_{K}^{n-p}(u)\rho_{Q^{*}}^{p}(u)du\right]V(Q)^{\frac{p}{n}} \\ &+\mu\left[\int_{S^{n-1}}\rho_{L}^{n-p}(u)\rho_{Q^{*}}^{p}(u)du\right]V(Q)^{\frac{p}{n}}:Q\in\mathcal{S}_{o}^{n}\right\} \\ &= \sup\left\{n\lambda\widetilde{V}_{p}(K,Q^{*})V(Q)^{\frac{p}{n}}+n\mu\widetilde{V}_{p}(L,Q^{*})V(Q)^{\frac{p}{n}}:Q\in\mathcal{S}_{o}^{n}\right\} \\ &\leq \sup\left\{n\lambda\widetilde{V}_{p}(K,Q^{*})V(Q)^{\frac{p}{n}}:Q\in\mathcal{S}_{o}^{n}\right\} \\ &+\sup\left\{n\mu\widetilde{V}_{p}(L,Q^{*})V(Q)^{\frac{p}{n}}:Q\in\mathcal{S}_{o}^{n}\right\} \\ &= \lambda n^{-\frac{p}{n}}\widetilde{\Omega}_{p}(K)^{\frac{n+p}{n}}+\mu n^{-\frac{p}{n}}\widetilde{\Omega}_{p}(L)^{\frac{n+p}{n}}. \end{split}$$

This gives inequality Equation (16).

We easily know that equality holds in Equation (16) for $\lambda, \mu > 0$ if and only if *K* and *L* are dilated. For $\lambda = 0$ or $\mu = 0$, Equation (16) becomes an equality. \Box

Lemma 3. *If* $K \in S_o^n$, $0 and <math>\tau \in [-1, 1]$, then

$$\widetilde{\Omega}_p(\widehat{\nabla}_p^{\tau}K) \le \widetilde{\Omega}_p(K). \tag{17}$$

Equality holds in Equation (17) for $\tau \in (-1, 1)$ if and only if K is origin-symmetric. For $\tau = \pm 1$, Equation (17) becomes an equality.

Proof. Taking $\lambda = f_1(\tau)$, $\mu = f_2(\tau)$ and L = -K in Equation (16), these and Equation (9) yield

$$\widetilde{\Omega}_{p}(\overline{\nabla}_{p}^{\tau}K) \leq f_{1}(\tau)\widetilde{\Omega}_{p}(K) + f_{2}(\tau)\widetilde{\Omega}_{p}(-K).$$
(18)

However, by Equations (2) and (12) we have

$$\begin{split} n^{-\frac{p}{n}} \widetilde{\Omega}_{p}(-K)^{\frac{n+p}{n}} &= \sup\{n\widetilde{V}_{p}(-K,Q^{*})V(Q)^{\frac{p}{n}}:Q\in\mathcal{S}_{o}^{n}\}\\ &= \sup\left\{\left[\int_{S^{n-1}} \rho_{-K}^{n-p}(u)\rho_{Q^{*}}^{p}(u)du\right]V(Q)^{\frac{p}{n}}:Q\in\mathcal{S}_{o}^{n}\right\}\\ &= \sup\left\{\left[\int_{S^{n-1}} \rho_{K}^{n-p}(-u)\rho_{-Q^{*}}^{p}(-u)du\right]V(Q)^{\frac{p}{n}}:Q\in\mathcal{S}_{o}^{n}\right\}\\ &= \sup\{n\widetilde{V}_{p}(K,-Q^{*})V(Q)^{\frac{p}{n}}:Q\in\mathcal{S}_{o}^{n}\}\\ &= \sup\{n\widetilde{V}_{p}(K,(-Q)^{*})V(-Q)^{\frac{p}{n}}:-Q\in\mathcal{S}_{o}^{n}\}\\ &= n^{-\frac{p}{n}}\widetilde{\Omega}_{p}(K)^{\frac{n+p}{n}}, \end{split}$$

i.e.,

$$\widetilde{\Omega}_p(-K) = \widetilde{\Omega}_p(K).$$

This together with Equation (18), and notice that $f_1(\tau) + f_2(\tau) = 1$, we obtain Equation (17).

According to the equality conditions of Equation (16), we see that equality holds in Equation (17) for $\tau \in (-1, 1)$ if and only if *K* and -K are dilated, i.e., *K* is origin-symmetric. For $\tau = \pm 1$, Equation (17) becomes an equality. \Box

Lemma 4 ([23]). For p > 0, a map $\Psi_p : S_o^n \to S_o^n$ is a L_p radial Blaschke-Minkowski homomorphism if and only if there is a non-negative measure $\mu \in \mathcal{M}(S^{n-1}, \hat{e})$ such that for $K \in S_o^n$, $\rho(\Psi_p K, \cdot)^p$ is the convolution of $\rho(K, \cdot)^{n-p}$ and μ , namely

$$\rho(\Psi_p K, \cdot)^p = \rho(K, \cdot)^{n-p} * \mu.$$
⁽¹⁹⁾

Here \hat{e} *denotes the pole point of* S^{n-1} *and* $\mathcal{M}(S^{n-1}, \hat{e})$ *denotes the signed finite Borel measure space on* S^{n-1} *(see* [21]).

Obviously, Equation (19) gives that for c > 0,

$$\Psi_p(cK) = c^{\frac{n-p}{p}} \Psi_p K.$$
(20)

Lemma 5. For $0 , let <math>\Psi_p$ be an even L_p radial Blaschke-Minkowski homomorphism. If $K \in S_o^n$ and $\tau \in [-1,1]$, then

$$\Psi_p(\widehat{\nabla}_p^\tau K) = \Psi_p K. \tag{21}$$

Proof. Since Ψ_p is an even L_p radial Blaschke-Minkowski homomorphism, thus for any $K \in S_o^n$, $\Psi_p(-K) = \Psi_p K$. From this, according to Equation (19), Equation (6) and Equation (9), we have for $0 and <math>\mu \in \mathcal{M}(S^{n-1}, \hat{e})$,

$$\begin{split} \rho(\Psi_{p}(\widehat{\nabla}_{p}^{\tau}K),\cdot) &= \rho(\widehat{\nabla}_{p}^{\tau}K,\cdot)^{n-p} * \mu \\ &= [f_{1}(\tau)\rho(K,\cdot)^{n-p} + f_{2}(\tau)\rho(-K,\cdot)^{n-p}] * \mu \\ &= f_{1}(\tau)\rho(K,\cdot)^{n-p} * \mu + f_{2}(\tau)\rho(-K,\cdot)^{n-p} * \mu \\ &= f_{1}(\tau)\rho(\Psi_{p}K,\cdot) + f_{2}(\tau)\rho(\Psi_{p}(-K),\cdot) \\ &= f_{1}(\tau)\rho(\Psi_{p}K,\cdot) + f_{2}(\tau)\rho(\Psi_{p}K,\cdot) = \rho(\Psi_{p}K,\cdot). \end{split}$$

This gives Equation (21). \Box

Proof of Theorem 2. Since $K \notin S_{os}^n$, thus for $0 , by Equation (17) we know that for <math>\tau \in (-1, 1)$,

$$\widetilde{\Omega}_p(\widehat{\nabla}_p^{\tau}K) < \widetilde{\Omega}_p(K).$$

Choose $\varepsilon > 0$ such that

$$\widetilde{\Omega}_p((1+\varepsilon)\widehat{\nabla}_p^{\tau}K) < \widetilde{\Omega}_p(K).$$

From this, let $L = (1 + \varepsilon) \widehat{\nabla}_p^{\tau} K$, then $L \in S_o^n$ (Theorem 5 gives that for $\tau = 0, L \in S_{os}^n$; for $\tau \in (-1, 1)$ and $\tau \neq 0, L \in S_o^n \setminus S_{os}^n$) and satisfies $\widetilde{\Omega}_p(L) < \widetilde{\Omega}_p(K)$.

However, by Equations (20) and (21) we obtain for 0 ,

$$\Psi_p L = \Psi_p((1+\varepsilon)\widehat{\nabla}_p^{\tau}K) = (1+\varepsilon)^{\frac{n-p}{p}}\Psi_p(\widehat{\nabla}_p^{\tau}K) = (1+\varepsilon)^{\frac{n-p}{p}}\Psi_pK \supset \Psi_pK.$$

4. Brunn-Minkowski Inequalities for the L_p Radial Blaschke-Minkowski Homomorphisms

Associated with L_q radial Minkowski sum and L_q harmonic Blaschke sum of star bodies, Theorems 3 and 4 respectively give the L_p dual affine surface area forms of Brunn-Minkowski inequalities for the L_p radial Blaschke-Minkowski homomorphisms. In this section, we will complete their proofs. For the proof of Theorem 3, the following lemmas are essential.

Lemma 6. If $K, L \in S_o^n$, p > 0 and $0 < q \le n - p$, then for any $u \in S^{n-1}$,

$$\rho_{\Psi_{p}(K\tilde{+}_{qL})}^{\frac{pq}{n-p}}(u) \le \rho_{\Psi_{pK}}^{\frac{pq}{n-p}}(u) + \rho_{\Psi_{pL}}^{\frac{pq}{n-p}}(u),$$
(22)

with equality for 0 < q < n - p if and only if K and L are dilated. For q = n - p, Equation (22) becomes an equality.

Proof. Because of 0 < q < n - p implies $\frac{n-p}{q} > 1$, thus by Equation (19) and the Minkowski integral inequality we have for $\mu \in \mathcal{M}(S^{n-1}, \hat{e})$ and any $u \in S^{n-1}$,

$$\begin{split} \rho_{\Psi_{p}(K\tilde{+}_{q}L)}^{\frac{pq}{n-p}}(u) &= \left(\rho_{\Psi_{p}(K\tilde{+}_{q}L)}^{p}(u)\right)^{\frac{q}{n-p}} = \left(\rho_{K\tilde{+}_{q}L}^{n-p}(u)*\mu\right)^{\frac{q}{n-p}} \\ &= \left[\left(\rho_{K\tilde{+}_{q}L}^{q}(u)\right)^{\frac{n-p}{q}}*\mu\right]^{\frac{q}{n-p}} \\ &= \left[\left(\rho_{K}^{q}(u)+\rho_{L}^{q}(u)\right)^{\frac{n-p}{q}}*\mu\right]^{\frac{q}{n-p}} \\ &\leq \left(\rho_{K}^{n-p}(u)*\mu\right)^{\frac{q}{n-p}} + \left(\rho_{L}^{n-p}(u)*\mu\right)^{\frac{q}{n-p}} \\ &= \rho_{\Psi_{p}K}^{\frac{pq}{n-p}}(u) + \rho_{\Psi_{p}L}^{\frac{qq}{n-p}}(u). \end{split}$$

This yields inequality Equation (22).

From the equality condition of Minkowski integral inequality, we know that equality holds in Equation (22) for 0 < q < n - p if and only if *K* and *L* are dilated. Clearly, if q = n - p, Equation (22) becomes an equality. \Box

Lemma 7. If $K, L \in S_0^n$, $0 and <math>0 < q \le n - p$, then for any $M \in S_0^n$,

$$\widetilde{V}_p(\Psi_p(K\widetilde{+}_qL), M)^{\frac{pq}{(n-p)^2}} \le \widetilde{V}_p(\Psi_pK, M)^{\frac{pq}{(n-p)^2}} + \widetilde{V}_p(\Psi_pL, M)^{\frac{pq}{(n-p)^2}},$$
(23)

with equality if and only if K and L are dilated.

Proof. Since $0 and <math>0 < q \le n - p$, thus $\frac{(n-p)^2}{pq} > 1$. Using Equation (12), Equation (22) and the Minkowski integral inequality, we have for any $M \in S_o^n$,

$$\begin{split} \widetilde{V}_{p}(\Psi_{p}(K\widetilde{+}_{q}L), M)^{\frac{pq}{(n-p)^{2}}} &= \left[\frac{1}{n} \int_{S^{n-1}} \rho_{\Psi_{p}(K\widetilde{+}_{q}L)}^{n-p}(u) \rho_{M}^{p}(u) du\right]^{\frac{pq}{(n-p)^{2}}} \\ &= \left[\frac{1}{n} \int_{S^{n-1}} \left(\rho_{\Psi_{p}(K\widetilde{+}_{q}L)}^{\frac{pq}{n-p}}(u)\right)^{\frac{(n-p)^{2}}{pq}} \rho_{M}^{p}(u) du\right]^{\frac{pq}{(n-p)^{2}}} \end{split}$$

$$\leq \left[\frac{1}{n} \int_{S^{n-1}} \left(\rho_{\Psi_{p}K}^{\frac{pq}{n-p}}(u) + \rho_{\Psi_{p}L}^{\frac{pq}{n-p}}(u)\right)^{\frac{(n-p)^{2}}{pq}} \rho_{M}^{p}(u) du\right]^{\frac{pq}{(n-p)^{2}}} \\ \leq \left[\frac{1}{n} \int_{S^{n-1}} \rho_{\Psi_{p}K}^{n-p}(u) \rho_{M}^{p}(u) du\right]^{\frac{pq}{(n-p)^{2}}} \\ + \left[\frac{1}{n} \int_{S^{n-1}} \rho_{\Psi_{p}L}^{n-p}(u) \rho_{M}^{p}(u) du\right]^{\frac{pq}{(n-p)^{2}}} \\ = \widetilde{V}_{p}(\Psi_{p}K, M)^{\frac{pq}{(n-p)^{2}}} + \widetilde{V}_{p}(\Psi_{p}L, M)^{\frac{pq}{(n-p)^{2}}}.$$

This yields inequality Equation (23).

For 0 < q < n - p, according to the equality conditions of Equation (22) and Minkowski integral inequality, we see that equality holds in Equation (23) if and only if *K* and *L* are dilated, and $\Psi_p K$ and $\Psi_p L$ are dilated. However, by Equation (19) we know that *K* and *L* are dilated is equivalent to $\Psi_p K$ and $\Psi_p L$ are dilated. Therefore, equality holds in Equation (23) if and only if *K* and *L* are dilated.

For q = n - p, from equality condition of Minkowski integral inequality, we are aware of equality holds in Equation (23) if and only if $\Psi_p K$ and $\Psi_p L$ are dilated, this is equivalent to K and L are dilated. \Box

Proof of Theorem 3. Because of $\frac{pq}{(n-p)^2} > 0$, thus by Equations (1) and (23), we obtain that

$$\begin{split} & \left[n^{-\frac{p}{n}} \widetilde{\Omega}_{p} (\Psi_{p}(K\widetilde{+}_{q}L))^{\frac{n+p}{n}} \right]^{\frac{pq}{(n-p)^{2}}} \\ &= \left[\sup\{n\widetilde{V}_{p}(\Psi_{p}(K\widetilde{+}_{q}L),Q^{*})V(Q)^{\frac{p}{n}} : Q \in \mathcal{S}_{o}^{n} \} \right]^{\frac{pq}{(n-p)^{2}}} \\ &= \sup\left\{ n^{\frac{pq}{(n-p)^{2}}} \widetilde{V}_{p}(\Psi_{p}(K\widetilde{+}_{q}L),Q^{*})^{\frac{pq}{(n-p)^{2}}}V(Q)^{\frac{p^{2}q}{n(n-p)^{2}}} : Q \in \mathcal{S}_{o}^{n} \right\} \\ &\leq \sup\left\{ n^{\frac{pq}{(n-p)^{2}}} \left[\widetilde{V}_{p}(\Psi_{p}K,Q^{*})^{\frac{pq}{(n-p)^{2}}} + \widetilde{V}_{p}(\Psi_{p}L,Q^{*})^{\frac{pq}{(n-p)^{2}}} \right] V(Q)^{\frac{p^{2}q}{n(n-p)^{2}}} : Q \in \mathcal{S}_{o}^{n} \right\} \\ &\leq \sup\left\{ n^{\frac{pq}{(n-p)^{2}}} \widetilde{V}_{p}(\Psi_{p}K,Q^{*})^{\frac{pq}{(n-p)^{2}}} V(Q)^{\frac{p^{2}q}{n(n-p)^{2}}} : Q \in \mathcal{S}_{o}^{n} \right\} \\ &+ \sup\left\{ n^{\frac{pq}{(n-p)^{2}}} \widetilde{V}_{p}(\Psi_{p}L,Q^{*})^{\frac{pq}{(n-p)^{2}}} V(Q)^{\frac{p^{2}q}{n(n-p)^{2}}} : Q \in \mathcal{S}_{o}^{n} \right\} \\ &= \left[\sup\{n\widetilde{V}_{p}(\Psi_{p}K,Q^{*})V(Q)^{\frac{p}{n}} : Q \in \mathcal{S}_{o}^{n} \} \right]^{\frac{pq}{(n-p)^{2}}} \\ &+ \left[\sup\{n\widetilde{V}_{p}(\Psi_{p}K,Q^{*})V(Q)^{\frac{p}{n}} : Q \in \mathcal{S}_{o}^{n} \} \right]^{\frac{pq}{(n-p)^{2}}} \\ &= \left[n^{-\frac{p}{n}} \widetilde{\Omega}_{p}(\Psi_{p}K)^{\frac{n+p}{n}} \right]^{\frac{pq}{(n-p)^{2}}} + \left[n^{-\frac{p}{n}} \widetilde{\Omega}_{p}(\Psi_{p}L)^{\frac{n+p}{n}} \right]^{\frac{pq}{(n-p)^{2}}}. \end{split}$$

From this, we get inequality Equation (4), and equality holds in Equation (4) if and only if *K* and *L* are dilated. \Box

The proof of Theorem 1 requires the following lemmas.

Lemma 8. If $K, L \in S_0^n$, $0 , then for any <math>u \in S^{n-1}$,

$$\frac{\rho_{\Psi_p(K=qL)}^{(n+q)}(u)}{V(K=qL)} \le \frac{\rho_{\Psi_pK}^{(n+q)}(u)}{V(K)} + \frac{\rho_{\Psi_pL}^{(n+q)}(u)}{V(L)},$$
(24)

with equality if and only if K and L are dilated.

Proof. Since $0 , thus <math>\frac{n-p}{n+q} > 1$. Hence, by Equation (13), Equation (19) and the Minkowski integral inequality we have for $\mu \in \mathcal{M}(S^{n-1}, \hat{e})$ and any $u \in S^{n-1}$,

$$\begin{split} \frac{\rho_{\Psi_{p}(K\mp qL)}^{n(n+q)}(u)}{V(K\mp qL)} &= \frac{\left[\rho_{\Psi_{p}(K\mp qL)}^{p}(u)\right]^{\frac{n+q}{n-p}}}{V(K\mp qL)} = \frac{\left[\rho_{K\mp qL}^{n-p}(u)*\mu\right]^{\frac{n+q}{n-p}}}{V(K\mp qL)} \\ &= \left[\left(\frac{\rho_{K\mp qL}^{n+q}(u)}{V(K\mp qL)}\right)^{\frac{n-p}{n+q}}*\mu\right]^{\frac{n+q}{n-p}} \\ &= \left[\left(\frac{\rho_{K}^{n+q}(u)}{V(K)} + \frac{\rho_{L}^{n+q}(u)}{V(L)}\right)^{\frac{n-p}{n+q}}*\mu\right]^{\frac{n+q}{n-p}} \\ &\leq \left[\frac{\rho_{K}^{n-p}(u)*\mu}{V(K)^{\frac{n-p}{n+q}}}\right]^{\frac{n+q}{n-p}} + \left[\frac{\rho_{L}^{n-p}(u)*\mu}{V(L)^{\frac{n-p}{n+q}}}\right]^{\frac{n+q}{n-p}} \\ &= \frac{\rho_{\Psi_{p}K}^{\frac{p(n+q)}{n-p}}(u)}{V(K)} + \frac{\rho_{\Psi_{p}L}^{\frac{p(n+q)}{n-p}}(u)}{V(L)}. \end{split}$$

This deduces Equation (24).

According to the equality condition of Minkowski integral inequality, we know that equality holds in Equation (24) if and only if K and L are dilated. \Box

Lemma 9. If $K, L \in S_o^n$, $0 and <math>0 , then for any <math>M \in S_o^n$,

$$\frac{\widetilde{V}_{p}(\Psi_{p}(K \mp_{q} L), M)^{\frac{p(n+q)}{(n-p)^{2}}}}{V(K \mp_{q} L)} \leq \frac{\widetilde{V}_{p}(\Psi_{p}K, M)^{\frac{p(n+q)}{(n-p)^{2}}}}{V(K)} + \frac{\widetilde{V}_{p}(\Psi_{p}L, M)^{\frac{p(n+q)}{(n-p)^{2}}}}{V(L)},$$
(25)

with equality if and only if K and L are dilated.

Proof. Since $0 and <math>0 , thus <math>\frac{(n-p)^2}{p(n+q)} > 1$. Using Equation (12), Equation (24) and the Minkowski integral inequality, we have for any $M \in S_o^n$,

$$\begin{split} & \frac{\widetilde{V}_{p}(\Psi_{p}(K \mp_{q} L), M)^{\frac{p(n+q)}{(n-p)^{2}}}}{V(K \mp_{q} L)} \\ &= \frac{\left[\frac{1}{n} \int_{S^{n-1}} \rho_{\Psi_{p}(K \mp_{q} L)}^{n-p}(u) \rho_{M}^{p}(u) du\right]^{\frac{p(n+q)}{(n-p)^{2}}}}{V(K \mp_{q} L)} \\ &= \left[\frac{1}{n} \int_{S^{n-1}} \left(\frac{\rho_{\Psi_{p}(K \mp_{q} L)}^{\frac{p(n+q)}{n-p}}(u)}{V(K \mp_{q} L)}\right)^{\frac{(n-p)^{2}}{p(n+q)}} \rho_{M}^{p}(u) du\right]^{\frac{p(n+q)}{(n-p)^{2}}} \end{split}$$

$$\leq \left[\frac{1}{n} \int_{S^{n-1}} \left(\frac{\rho_{\Psi_p K}^{p(n+q)}(u)}{V(K)} + \frac{\rho_{\Psi_p L}^{p(n+q)}(u)}{V(L)}\right)^{\frac{(n-p)^2}{p(n+q)}} \rho_M^p(u) du\right]^{\frac{p(n+q)}{(n-p)^2}} \\ \leq \left[\frac{1}{n} \int_{S^{n-1}} \frac{\rho_{\Psi_p K}^{n-p}(u) \rho_M^p(u)}{V(K)^{\frac{(n-p)^2}{p(n+q)}}} du\right]^{\frac{p(n+q)}{(n-p)^2}} + \left[\frac{1}{n} \int_{S^{n-1}} \frac{\rho_{\Psi_p L}^{n-p}(u) \rho_M^p(u)}{V(L)^{\frac{(n-p)^2}{p(n+q)}}} du\right]^{\frac{p(n+q)}{(n-p)^2}} \\ = \frac{\widetilde{V}_p(\Psi_p K, M)^{\frac{p(n+q)}{(n-p)^2}}}{V(K)} + \frac{\widetilde{V}_p(\Psi_p L, M)^{\frac{p(n+q)}{(n-p)^2}}}{V(L)}.$$

From this, inequality Equation (25) is obtained.

By the equality conditions of Equation (24) and the Minkowski integral inequality, we see that equality holds in Equation (25) if and only if *K* and *L* are dilated. \Box

Proof of Theorem 1. From $0 and <math>0 , we know that <math>\frac{p(n+q)}{(n-p)^2} > 0$. Thus by Equations (1) and (25), we obtain that

$$\begin{split} & \frac{\left[n^{-\frac{p}{n}}\widetilde{\Omega}_{p}(\Psi_{p}(K\mp_{q}L))^{\frac{n+p}{n}}\right]^{\frac{p(n+q)}{(n-p)^{2}}}}{V(K\mp_{q}L)} \\ & = \frac{\left[\sup\{n\widetilde{V}_{p}(\Psi_{p}(K\mp_{q}L),Q^{*})V(Q)^{\frac{p}{n}}:Q\in\mathcal{S}_{0}^{n}\}\right]^{\frac{p(n+q)}{(n-p)^{2}}}}{V(K\mp_{q}L)} \\ & = \sup\left\{\frac{\left[n\widetilde{V}_{p}(\Psi_{p}(K,Q^{*})(X\mp_{q}L),Q^{*})\right]^{\frac{p(n+q)}{(n-p)^{2}}}V(Q)^{\frac{p^{2}(n+q)}{n(n-p)^{2}}}:Q\in\mathcal{S}_{0}^{n}\right\}}{V(K\mp_{q}L)} \\ & \leq \sup\left\{\left[\frac{\left[n\widetilde{V}_{p}(\Psi_{p}K,Q^{*})\right]^{\frac{p(n+q)}{(n-p)^{2}}}+\left[n\widetilde{V}_{p}(\Psi_{p}L,Q^{*})\right]^{\frac{p(n+q)}{(n-p)^{2}}}\right]V(Q)^{\frac{p^{2}(n+q)}{n(n-p)^{2}}}:Q\in\mathcal{S}_{0}^{n}\right\}}{V(L)} \\ & \leq \sup\left\{\frac{\left[n\widetilde{V}_{p}(\Psi_{p}K,Q^{*})\right]^{\frac{p(n+q)}{(n-p)^{2}}}V(Q)^{\frac{p^{2}(n+q)}{n(n-p)^{2}}}:Q\in\mathcal{S}_{0}^{n}\right\}}{V(L)} \\ & +\sup\left\{\frac{\left[n\widetilde{V}_{p}(\Psi_{p}K,Q^{*})V(Q)^{\frac{p}{n}}:Q\in\mathcal{S}_{0}^{n}\right]\right]^{\frac{p(n+q)}{(n-p)^{2}}}}{V(L)} \\ & = \frac{\left[\sup\{\widetilde{V}_{p}(\Psi_{p}K,Q^{*})V(Q)^{\frac{p}{n}}:Q\in\mathcal{S}_{0}^{n}\}\right]^{\frac{p(n+q)}{(n-p)^{2}}}}{V(L)} \\ & = \frac{\left[n^{-\frac{p}{n}}\widetilde{\Omega}_{p}(\Psi_{p}K)^{\frac{n+p}{n}}\right]^{\frac{p(n+q)}{(n-p)^{2}}}}{V(K)} + \frac{\left[n^{-\frac{p}{n}}\widetilde{\Omega}_{p}(\Psi_{p}L)^{\frac{n+p}{n}}\right]^{\frac{p(n+q)}{(n-p)^{2}}}}}{V(L)}. \end{split}$$

This gives inequality Equation (5). In addition, equality holds in Equation (5) if and only if *K* and *L* are dilated. \Box

Author Contributions: Conceptualization, Z.S. and W.W.; methodology, Z.S. and W.W.; software, W.W.; validation, Z.S. and WW.; formal analysis, W.W.; investigation, Z.S. and W.W.; resources, W.W.; writingoriginal draft preparation, Z.S.; writingreview and editing, W.W.; visualization, Z.S. and W.W.; supervision, Z.S.; project administration, W.W.; funding acquisition, W.W.

Funding: Research is supported in part by the Natural Science Foundation of China (Grant No.11371224).

Acknowledgments: The authors like to sincerely thank the referees for very valuable and helpful comments and suggestions which made the paper more accurate and readable.

Conflicts of Interest: The authors declare no conflict of interest.

References

- 1. Gardner, R.J. Geometric Tomography, 2nd ed.; Cambridge University Press: Cambridge, UK, 2006.
- 2. Lutwak, E. Intersection bodies and dual mixed volumes. Adv. Math. 1988, 71, 531–538. [CrossRef]
- 3. Goodey, P.; Weil, W. Intersection bodies and ellipsoids. *Mathematika* 1995, 42, 295–304. [CrossRef]
- 4. Kalton, N.J.; Koldobsky, A. Intersection bodies and L_p spaces. Adv. Math. 2005, 196, 257–275. [CrossRef]
- 5. Koldobsky, A. Intersection bodies in \mathbb{R}^4 . *Adv. Math.* **1998**, 136, 1–14. [CrossRef]
- 6. Koldobsky, A. Second derivative test for intersection bodies. Adv. Math. 1998, 136, 15–25. [CrossRef]
- Koldobsky, A. A functional analytic approach to intersection bodies. *Geom. Funct. Anal.* 2000, 10, 1507–1526. [CrossRef]
- 8. Leng, G.; Zhao, C. Brunn-Minkowski inequality for mixed intersection bodies. *J. Math. Anal. Appl.* **2005**, 301, 115–123.
- 9. Lu, F.H.; Mao, W.H.; Leng, G.S. On star duality of mixed intersection bodies. J. Inequal. Appl. 2007, 2007, 39345.
- 10. Ludwig, M. Intersection bodies and valuations. Am. J. Math. 2006, 128, 1409–1428. [CrossRef]
- 11. Zhang, G.Y. Centered bodies and dual mixed volumes. Trans. Am. Math. Soc. 1994, 345, 777-801. [CrossRef]
- 12. Zhang, G.Y. Intersection bodies and polytopes. *Mathematika* 1999, 46, 29–34. [CrossRef]
- 13. Gardner, R.J. Intersection bodies and the Busemann-Petty problem. *Trans. Am. Math. Soc.* **1994**, 342, 435–445. [CrossRef]
- 14. Zhang, G.Y. A positive solution to the Busemann-Petty problem in \mathbb{R}^4 . *Ann. Math.* **1999**, *149*, 535–543. [CrossRef]
- 15. Gardner, R.J. A positive answer to the Busemann-Petty problem in three dimensions. *Ann. Math.* **1994**, 140, 435–447. [CrossRef]
- 16. Gardner, R.J.; Koldobsky, A.; Schlumprecht, T. An analytic solution to the Busemann-Petty problem on sections of convex bodies. *Ann. Math.* **1999**, *149*, 691–703. [CrossRef]
- 17. Koldobsky, A. Intersection bodies and the Busemann-Petty problem. C. R. Acad. Sci. Paris Ser. I Math. 1997, 325, 1181–1186. [CrossRef]
- 18. Koldobsky, A. Intersection bodies, positive definite distributions, and the Busemann-Petty problem. *Am. J. Math.* **1998**, *120*, 827–840. [CrossRef]
- 19. Zhang, G.Y. Intersection bodies and the four-dimensional Busemann-Petty problem. *Duke Math. J.* **1993**, *71*, 233–240.
- 20. Zhang, G.Y. Intersection bodies and the Busemann-Petty inequalities in \mathbb{R}^4 . Ann. Math. **1994**, 140, 331–346. [CrossRef]
- 21. Schuster, F.E. Volume inequalities and additive maps of convex bodies. *Mathematika* **2006**, *53*, 211–234. [CrossRef]
- 22. Schuster, F.E. Valuations and Busemann-Petty type problems. Adv. Math. 2008, 219, 344–368. [CrossRef]
- 23. Wang, W.; Liu, L.J.; He, B.W. *L_p* radial Blaschke-Minkowski homomorphisms. *Taiwan J. Math.* **2011**, *15*, 1183–1199. [CrossRef]
- 24. Alesker, S.; Bernig, A.; Schuster, F.E. Harmonic analysis of translation invariant valuations. *Geom. Funct. Anal.* **2011**, *10*, 751–773. [CrossRef]
- 25. Feng, Y.B.; Wang, W.D.; Yuan, J. Differences of quermass- and dual quermassintegrals of Blaschke-Minkowski and radial Blaschke-Minkowski homomorphisms. *Bull. Belg. Math. Soc.-Sim* **2014**, *21*, 577–592.
- Liu, L. Mixed radial Blaschke-Minkowski homomorphisms and comparison of volumes. *Math. Inequal. Appl.* 2013, 16, 401–412. [CrossRef]

- 27. Wang, W. L_p Blaschke-Minkowski homomorphisms. J. Inequal. Appl. 2013, 2013, 140. [CrossRef]
- Wei, B.; Wang, W.D.; Lu, F.H. Inequalities for radial Blaschke-Minkowski homomorphisms. *Ann. Pol. Math.* 2015, 113, 243–253. [CrossRef]
- 29. Zhao, C.J. Radial Blaschke-Minkowski homomorphisms and volume differences. *Geom. Dedic.* 2011, 154, 81–91. [CrossRef]
- 30. Zhao, C.J. On radial Blaschke-Minkowski homomorphisms. Geom. Dedic. 2013, 167, 1–10. [CrossRef]
- 31. Zhao, C.J. On raidal and polar Blaschke-Minkowski homomorphisms. *Proc. Am. Math. Soc.* 2013, 141, 667–676. [CrossRef]
- 32. Zhao, C.J. On Blaschke-Minkowski homomorphisms and radial Blaschke-Minkowski homomorphisms. *J. Geom. Anal.* **2015**, *25*, 1523–1538. [CrossRef]
- 33. Zhou, Y.; Wang, W.D. Some Brunn-Minkowski type inequalities for *L_p* radial Blaschke-Minkowski homomorphisms. *J. Inequal. Appl.* **2016**, 2016, 183. [CrossRef]
- 34. Wang, W.; Yuan, J.; He, B.W. Inequalities for L_p-dual affine surface area. *Math. Inequal. Appl.* **2010**, *7*, 34–45.
- 35. Wang, J.Y.; Wang, W.D. General *L*_{*p*}-dual Blaschke bodies and the applications. *J. Inequal. Appl.* **2015**, 2015, 233. [CrossRef]
- 36. Feng, Y.B.; Wang, W.D. Some inequalities for *L*_{*p*}-dual affine surface area. *Math. Inequal. Appl.* **2014**, 17, 431–441.
- Pei, Y.N.; Wang, W.D. A type of Busemann-Petty problems for general L_p-intersection bodies. Wuhan Univ. J. Natural Sci. 2015, 20, 471–475. [CrossRef]
- 38. Wan, X.Y.; Wang, W.D. L_p-dual mixed affine surface area. Ukra. Math. J. 2016, 68, 679–688. [CrossRef]
- 39. Wang, J.Y.; Wang, W.D. *L_p*-dual affine surface area forms of Busemann-Petty type problems. *Proc. Indian Acad. Sci.* (*Math. Sci.*) **2015**, 125, 71–77. [CrossRef]
- 40. Wei, B.; Wang, W.D. Some inequalities for general *L*_{*p*}-harmonic Blaschke bodies. *J. Math. Inequal.* **2016**, *10*, 63–73. [CrossRef]
- 41. Zhang, T.; Wang, W.D.; Si, L. The mixed *L*_p-dual affine surface area for multiple star bodies. *J. Nonlinear Sci. Appl.* **2016**, *9*, 2813–2822.
- 42. Lutwak, E. The Brunn-Minkowski-Firey theory II: Affine and Geominimal Surface Areas. *Adv. Math.* **1996**, 118, 244–294. [CrossRef]
- 43. Gardner, R.J. The Brunn-Minkowski inequality. Bull. Am. Math. Soc. 2002, 39, 355-405. [CrossRef]
- 44. Schneider, R. *Convex Bodies: The Brunn-Minkowski Theory*, 2nd ed.; Cambridge University Press: Cambridge, UK, 2014.
- 45. Grinberg, E.; Zhang, G.Y. Convolutions, transforms and convex bodies. *Proc. Lond. Math. Soc.* **1999**, *78*, 77–115. [CrossRef]
- 46. Haberl, C. L_p intersection bodies. Adv. Math. 2008, 217, 2599–2624. [CrossRef]
- 47. Lutwak, E. Centroid bodies and dual mixed volumes. Proc. Lond. Math. Soc. 1999, 60, 365–391. [CrossRef]
- 48. Feng, Y.B.; Wang, W.D. Shephard type problems for *L*_p-centroid bodies. *Math. Inequal. Appl.* **2014**, 17, 865–877.



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses/by/4.0/).