



Article One-Point Optimal Family of Multiple Root Solvers of Second-Order

Deepak Kumar^{1,*}, Janak Raj Sharma¹ and Clemente Cesarano^{2,*}

- ¹ Department of Mathematics, Sant Longowal Institute of Engineering and Technology, Longowal, Sangrur 148106, India
- ² Section of Mathematics, International Telematic University UNINETTUNO, Corso Vittorio Emanuele II, 39, 00186 Roma, Italy
- * Correspondence: deepak.babbi@gmail.com (D.K.); c.cesarano@uninettunouniversity.net (C.C.)

Received: 6 June 2019; Accepted: 17 July 2019; Published: 21 July 2019



Abstract: This manuscript contains the development of a one-point family of iterative functions. The family has optimal convergence of a second-order according to the Kung-Traub conjecture. This family is used to approximate the multiple zeros of nonlinear equations, and is based on the procedure of weight functions. The convergence behavior is discussed by showing some essential conditions of the weight function. The well-known modified Newton method is a member of the proposed family for particular choices of the weight function. The dynamical nature of different members is presented by using a technique called the "basin of attraction". Several practical problems are given to compare different methods of the presented family.

Keywords: nonlinear equations; multiple roots; one-point methods; optimal convergence

MSC: 65H05; 49M15; 41A25

1. Introduction

Solving nonlinear equations h(x) = 0, where function h(x) is defined as $h : \Omega \subseteq R \to R$ in an open interval Ω , is a delightful and demanding task in many applied scientific branches, such as Mathematical Biology, Physics, Chemistry, Economics, and also Engineering, to name a few [1–4]. This is mainly because problems from these areas usually include needing to find the root of a nonlinear equation. The huge value of this subject has led to the development of many numerical methods, with most of them having an iterative nature (see [5–7]). With the advanced technology of computer hardware and the latest software, the topic of solving nonlinear equations by using numerical methods has gained additional significance. Researchers are utilizing iterative methods for approximating the solution, since closed-form solutions cannot be obtained in general. In particular, here we consider iterative methods to compute a multiple root (say, α) with multiplicity $m \geq 1$, i.e., $h^{(k)}(\alpha) = 0$, k = 0, 1, 2, ..., m - 1 and $h^{(m)}(\alpha) \neq 0$, of the equation h(x) = 0.

There are a plethora of methods of an iterative nature with a different order of convergence, constructed to approximate the zeros of Equation h(x) = 0 (see [8–18]). The computational efficiency index is a very effective mechanism, defined by Ostrowski in [19] which categorizes the iterative algorithms in the form of their convergence order p_c and the function evaluations d required per iteration. It is formulated as $I = p_c^{1/d}$. The higher the computational efficiency index of an iterative scheme, the better the scheme is.

This idea becomes more rigid with Kung-Traub's conjecture [20], which imposes an upper bound for the convergence order to be limited with fixed functional evaluations. According to this conjecture, an iterative scheme which requires a *d* number of function evaluations can attain the convergence order $p_c = 2^{d-1}$. The iterative methods which obey Kung-Traub's conjecture are optimal in nature. The most basic and widely used method is the well-known modified Newton's method:

$$x_{n+1} = x_n - m \frac{h(x_n)}{h'(x_n)} \quad \forall \ n = 0, 1, 2, \dots$$
 (1)

This method can efficiently find the required zero of multiplicity, *m* with a quadratic order of convergence, provided that the initial approximate x_0 is sufficiently nearer to zero [8]. In Traub's terminology (see [2]), Newton's method (1) is called the one-point method. This classical method attracts many researchers because of its huge applications in several kinds of problems, which are formulated as non-linear equations, differential equations, integral equations, systems of non-linear algebraic equations, and even to random operator equations. However, a common issue and main obstacle in the use of Newton's method is its sensitivity to initial guesses, which must be sufficiently nearer to the exact solution for assured convergence. Developing a criterion for selecting these initial guesses is quite difficult, and therefore, a more effective iterative technique that is globally-convergent is yet to be discovered. Some other important higher order methods that are based on Newton's method (1) have been developed in [11,13,14,21–28].

Recently, Chicharro et al. [28] used the weight function technique to design a class of optimal second-order one-point iterative methods for simple roots, including Newton's method. In this paper, we have applied this technique for the development of a class of optimal second-order one-point methods for multiple roots. The new proposed family contains the modified Newton's method and many other efficient methods. These methods exist when particular weight functions are selected. Therefore, with a wide range of initial approximations, we can select those methods from the family which are able to converge towards exact zero, when Newton's method does not.

The rest of the paper is organized as follows. In Section 2, the technique of the second-order method is developed and its convergence is studied. In Section 3, the basins of attractors are studied to check the stability of the methods. Numerical experiments on different equations are performed in Section 4 to demonstrate the applicability and efficiency of the presented methods. We finish the manuscript with some valuable conclusions in Section 5.

2. The Method

where $e_n = x_n$

For a known multiplicity $m \ge 1$, we consider the following one–step scheme for multiple roots:

$$x_{n+1} = x_n - G(\nu_n), \tag{2}$$

where the function $G(\nu_n) : \mathbb{C} \to \mathbb{C}$ is differentiable in a neighborhood of "0" with $\nu_n = \frac{h(x_n)}{h'(x_n)}$. In the next result, we prove a theorem for the order of convergence of the scheme (2).

Theorem 1. Let $f : \mathbb{C} \to \mathbb{C}$ be a differentiable function in a region in which a multiple zero (say, α) with multiplicity *m* lies. Suppose that the initial approximate x_0 is sufficiently close to α —then, the iteration scheme defined by (2) has a second-order of convergence, provided that G(0) = 0, G'(0) = m, and $|G''(0)| < \infty$, and the error is

$$e_{n+1} = \left(\frac{2mC_1 - G''(0)}{2m^2}\right)e_n^2 + O(e_n^3),$$

$$-\alpha \text{ and } C_k = \frac{m!}{(m+k)!} \frac{f^{(m+k)}(\alpha)}{f^{(m)}(\alpha)} \text{ for } k \in \mathbb{N}.$$
(3)

Proof. Let the error at the *n*-th iteration be $e_n = x_n - \alpha$. Using the Taylor's expansion of $f(x_n)$ and $f'(x_n)$ about α , we have that

$$f(x_n) = \frac{f^{(m)}(\alpha)}{m!} e_n^m \Big(1 + C_1 e_n + C_2 e_n^2 + C_3 e_n^3 + O(e_n^4) \Big).$$
(4)

and

$$f'(x_n) = \frac{f^{(m)}(\alpha)}{m!} e_n^{m-1} \Big(m + (m+1)C_1 e_n + (m+2)C_2 e_n^2 + (m+3)C_3 e_n^3 + O(e_n^4) \Big).$$
(5)

By using (4) and (5), we obtained

$$\nu_n = \frac{e_n}{m} - \frac{C_1}{m}e_n^2 + \frac{(1+m) - 2mC_1}{m^3}e_n^3 + O(e_n^4).$$

If we write the expansion-of-weight function $G(\nu_n)$ about the origin by using the Taylor series, then we have that

$$G(\nu_n) \approx G(0) + \nu G'(0) + \frac{1}{2}\nu^2 G''(0).$$
 (6)

By employing the expression (6) in the scheme (2), we were able to obtain the error equation

$$e_{n+1} = -G(0) + \left(1 - \frac{G'(0)}{m}\right)e_n - \left(\frac{-2C_1G'(0) + G''(0)}{2m^2}\right)e_n^2 + O(e_n^3).$$
(7)

To obtain the second-order of convergence, the constant term and coefficient of e_n in (7) should simultaneously be equal to zero. This is possible when G(0) and G'(0) have the following values:

$$G(0) = 0, \quad G'(0) = m.$$
 (8)

By using the above values in (7), the error equation becomes

$$e_{n+1} = \left(\frac{2mC_1 - G''(0)}{2m^2}\right)e_n^2 + O(e_n^3).$$
(9)

Hence, the second-order convergence is established. \Box

Some Particular Forms of $G(\nu_n)$

We were able to obtain numerous methods of the family (2) based on the form of function $G(\nu)$ that satisfies the conditions of Theorem 1. However, we limited the choices to consider only some simple functions. Accordingly, the following simple forms were chosen:

(1)
$$G(\nu_n) = m\nu_n(1+a_1\nu_n)$$
 (2) $G(\nu_n) = \frac{m\nu_n}{1+a_2\nu_n}$ (3) $G(\nu_n) = \frac{m\nu_n}{1+a_3m\nu_n}$, (4) $G(\nu_n) = m(e^{\nu_n}-1)$
(5) $G(\nu_n) = m\log[\nu_n+1]$ (6) $G(\nu_n) = m\sin\nu_n$ (7) $G(\nu_n) = \frac{\nu_n}{(\frac{1}{\sqrt{m}}+a_4\nu_n)^2}$ (8) $G(\nu_n) = \frac{\nu_n^2+\nu_n}{\frac{1}{m}+a_5\nu_n}$,

where a_1 , a_2 , a_3 , a_4 and a_5 are arbitrary constants.

The corresponding methods to each of the above forms are defined as follows: *Method 1* (M1):

$$x_{n+1} = x_n - m\nu_n(1 + a_1\nu_n).$$

Method 2 (M2):

$$x_{n+1} = x_n - \frac{m\nu_n}{1 + a_2\nu_n}.$$

Method 3 (M3):

$$x_{n+1} = x_n - \frac{m\nu_n}{1 + a_3 m\nu_n}$$

Method 4 (M4):

$$x_{n+1} = x_n - m(e^{\nu_n} - 1)$$

Method 5 (M5):

$$x_{n+1} = x_n - m\log(\nu_n + 1).$$

Method 6 (M6):

 $x_{n+1} = x_n - m \sin \nu_n.$

Method 7 (M7):

$$x_{n+1} = x_n - rac{
u_n}{(rac{1}{\sqrt{m}} + a_4
u_n)^2}$$

Method 8 (M8):

$$x_{n+1} = x_n - \frac{\nu_n^2 + \nu_n}{\frac{1}{m} + a_5 \nu_n}.$$

Remark 1. The scheme (2) shows a one-point family of second-order methods which needs only two function evaluations—namely, $h(x_n)$ and $h'(x_n)$.

Remark 2. Note that the modified Newton's method (1) is the special case of the above methods—M1, M2, M3, and M7—if the corresponding constants a_1 , a_2 , a_3 , and a_4 become zero.

3. Complex Dynamics of Methods

Our goal here is to check the complex dynamics of new methods with the help of a graphical tool, *the basins of attraction*, of the zeros for a polynomial P(z) in the Argand plane. The nature of the basins of attraction provides important ideas about the stability and convergence of iterative methods. This idea was initially introduced by Vrscay and Gilbert [29]. In recent times, most researchers have been using this concept in their work—see, for example [30,31]. We consider the special cases corresponding to $G(v_n)$ of (2) to analyze the basins of attraction.

The starting approximate z_0 is taken in a region of rectangular shape $R \in \mathbb{C}$ that contains all the zeros of P(z). A method, when it starts from point z_0 in a rectangle, either converges to zero, P(z), or eventually diverges. Therefore, the stopping criterion is 10^{-3} up to a number of 25 iterations.

To show complex geometry, we checked the basins of attraction of the methods M1–M8 on the following four polynomials:

Problem 1. In this problem, we took the polynomial $P_1(z) = (z^2 + 4)^2$, which has zeros $\{\pm 2i\}$ with a multiplicity of 2. We used a mesh of 400×400 points in a rectangular frame $D \in \mathbb{C}$ of area $[-2, 2] \times [-2, 2]$, and gave the color green for "2i" and red for "-2i". Each initial point from the green region converges towards "2i", and from the red region it converges to "-2i". Basins obtained for the methods M1–M8 are shown in Figure 1. Analyzing the behavior of the methods, we see that the methods M5 and M6 possess lesser numbers of divergent points, followed by M1, M4, M8, and M2. On the contrary, the method M3 has a higher number of divergent points, followed by M7.

Problem 2. Let us consider the polynomial $P_2(z) = (z^3 - z)^3$ having zeros $\{0, \pm 1\}$ with a multiplicity of 3. To see the dynamical structure, we considered a rectangular frame $D = [-2, 2] \times [-2, 2] \in \mathbb{C}$ with 400 × 400 mesh points, and gave the colors red, green, and blue to each point in the basins of attraction of -1, 0, and 1, respectively. Basins obtained for the methods M1–M8 are shown in Figure 2. Analyzing the behavior of the methods, we observe that the methods M5 and M6 have wider convergence regions, followed by M1, M4, M8, M2, M3, and M7.

Figure 1. Basins of attraction of M1–M8 for polynomial $P_1(z)$.



Figure 2. Basins of attraction of M1–M8 for polynomial $P_2(z)$.

Problem 3. Next, consider the polynomial $P_3(z) = (z^4 - 6z^2 + 8)^2$ that has four zeros $\{\pm 2, \pm 1.414...\}$ with a multiplicity of 2. To view attraction basins, we considered a rectangular frame $D = [-2,2] \times [-2,2] \in \mathbb{C}$ with 400 × 400 mesh points and assigned the colors red, blue, green, and yellow to each point in the basin of -2, -1.414..., 1.414..., and 2, respectively. Basins obtained for the methods M1–M8 are shown in Figure 3. Observing the behavior of the methods, we see that the methods M5, M8, M2, M3, M4, M1, and M6 possess a lesser number of divergent points, and therefore, they show good convergence. On the contrary, the method M7 has a higher number of divergent points.



Figure 3. Basins of attraction of M1–M8 for polynomial $P_3(z)$.

Problem 4. Lastly, we consider the polynomial $P_4(z) = z^6 - \frac{1}{2}z^5 + \frac{11}{4}(1+i)z^4 - \frac{1}{4}(19+3i)z^3 + \frac{1}{4}(11+5i)z^2 - \frac{1}{4}(11+i)z + \frac{3}{2} - 3i$ that has six simple zeros $\{1, -1+2i, -\frac{1}{2} - \frac{i}{2}, i, -\frac{3i}{2}, 1-i\}$. To view the attraction basins, we considered a rectangle $D = [-2, 2] \times [-2, 2] \in \mathbb{C}$ with 300×300 grid points, and assigned the colors red, green, yellow, blue, cyan, and purple to each point in the basin of $1, -1+2i, -\frac{1}{2} - \frac{1}{2}i, i, -\frac{3}{2}i$, and 1 - i, respectively. Basins obtained for the methods M1–M8 are shown in Figure 4. Analyzing the basins of the methods, we observe that the methods M5, M8, M2, M3, M4, and M6 possess a lesser number of divergent points. On the contrary, the methods M1 and M7 have a higher number of divergent points.



Figure 4. Basins of attraction of M1–M8 for polynomial $P_4(z)$.

From the graphics of the basins, we can give the judgment on the behavior and suitability of any method in the applications. In the event that we pick an initial point z_0 in a zone where various basins of attraction contact one another, it is difficult to anticipate which root will be achieved by the iterative

technique that begins in z_0 . Subsequently, the choice of z_0 in such a zone is anything but a decent one. Both the dark zones and the zones with various colors are not appropriate for choosing z_0 when we need to obtain a specific root. The most appealing pictures showed up when we had extremely intricate borders between basins of attraction. These borders correspond to the cases where the method is more demanding with respect to the initial point.

4. Numerical Results

In this section, we demonstrate the efficiency, effectiveness, and convergence behavior of the family of new methods by applying them to some practical problems. In this view, we take the special cases M1–M8 of the proposed class and choose $(a_1 = 1/10)$, $(a_2 = 1/4)$, $(a_3 = 1/10)$, $(a_4 = 1/10)$, and $(a_5 = 1/5)$ in the numerical work.

As we know that the constants a_1 , a_2 , a_3 , a_4 and a_5 are arbitrary, there is no particular reason for choosing these values for the constants, and we chose the values randomly. The proposed methods are compared with the existing modified Newton Method (1), also known as MNM.

To verify the theoretical results, we calculate the computational order of convergence (COC) by using the formula (see [32])

$$COC = \frac{\log |(x_{n+1} - \alpha)/(x_n - \alpha)|}{\log |(x_n - \alpha)/(x_{n-1} - \alpha)|}.$$

The computational work was performed in the programming software, Mathematica, by using multiple-precision arithmetic. Numerical results displayed in Tables 1–4 include: (i) the number of approximations (*n*) required to converge to the solution such that $|x_{n+1} - x_n| + |f(x_n)| < 10^{-100}$, (ii) values of the last three consecutive errors $|e_n| = |x_{n+1} - x_n|$, (iii) residual error $|f(x_n)|$, and (iv) computational order of convergence (COC).

For testing, we chose four test problems as follows:

Example 1 (Eigenvalue problem). *Eigen value problem is a difficult problem when characteristic polynomial involves a huge square matrix. Finding the zeros of characteristic equation of a square matrix with order more than 4 can even be a challenging task. So, we think about accompanying 9 \times 9 matrix*

	-12	0	0	19	-19	76	-19	18	437]
	-64	24	0	-24	24	64	-8	32	376
	-16	0	24	4	-4	16	-4	8	92
1	-40	0	0	-10	50	40	2	20	242
$M = \frac{1}{6}$	-4	0	0	-1	41	4	1	0	25
0	-40	0	0	18	-18	104	-18	20	462
	-84	0	0	-29	29	84	21	42	501
	16	0	0	-4	4	-16	4	16	-92
	0	0	0	0	0	0	0	0	24

The characteristic polynomial of the matrix (M) is given as

$$h_1(x) = x^9 - 29x^8 + 349x^7 - 2261x^6 + 8455x^5 - 17663x^4 + 15927x^3 + 6993x^2 - 24732x + 12960.$$

This function has one multiple zero $\alpha = 3$ with a multiplicity of 4. We chose initial approximations $x_0 = 2.75$ and obtained the numerical results as shown in Table 1.

Methods	n	$ e_{n-2} $	$ e_{n-1} $	$ e_n $	$f(x_n)$	СОС
MNM	7	$1.70 imes 10^{-21}$	$6.84 imes 10^{-43}$	1.11×10^{-85}	$5.90 imes10^{-681}$	2.000
M1	7	$2.79 imes10^{-21}$	$1.90 imes 10^{-42}$	$8.77 imes10^{-85}$	$9.90 imes 10^{-674}$	2.000
M2	7	$2.99 imes10^{-24}$	$1.56 imes10^{-48}$	$4.27 imes10^{-97}$	$8.37 imes10^{-773}$	2.000
M3	6	$2.17 imes10^{-13}$	$6.50 imes10^{-27}$	$5.80 imes10^{-54}$	$3.68 imes10^{-428}$	2.000
M4	7	$3.39 imes10^{-18}$	$4.18 imes10^{-36}$	$6.32 imes 10^{-72}$	$3.53 imes10^{-570}$	2.000
M5	6	$5.79 imes10^{-15}$	$3.77 imes 10^{-30}$	$1.60 imes10^{-60}$	$5.51 imes 10^{-481}$	2.000
M6	7	$1.93 imes 10^{-21}$	$8.86 imes10^{-43}$	$1.86 imes10^{-85}$	$3.69 imes 10^{-679}$	2.000
M7	6	$2.28 imes10^{-13}$	$7.15 imes10^{-27}$	$7.03 imes10^{-54}$	$1.71 imes 10^{-427}$	2.000
M8	7	$6.63 imes10^{-20}$	$1.26 imes 10^{-39}$	$4.60 imes10^{-79}$	1.09×10^{-627}	2.000

Table 1. Comparison of performance of methods for Example 1.

Example 2 (Beam Designing Model). *Here, we consider a beam situating problem (see* [4]) *where a beam of length r unit is inclining toward the edge of a cubical box with the length of the sides being 1 unit each, to such an extent that one end of the bar touches the wall and the opposite end touches the floor, as shown in Figure 5.*



Figure 5. Beam situating problem.

The problem is: What should be the distance be along the bottom of the beam to the floor from the base of the wall? Suppose y is the distance along the edge of the box to the beam from the floor, and let x be the distance from the bottom of the box and of the beam. Then, for a particular value of r, we have

$$x^4 + 4x^3 - 24x^2 + 16x + 16 = 0.$$

The non-negative solution of the equation is a root x = 2 with a multiplicity of 2. We consider the initial approximate $x_0 = 3$ to find the root. Numerical results of various methods are shown in Table 2.

Table 2. Comparison of performance of methods for Example 2.

Methods	n	$ e_{n-2} $	$ e_{n-1} $	$ e_n $	$f(x_n)$	COC
MNM	7	$1.61 imes 10^{-20}$	$6.50 imes10^{-41}$	$1.06 imes 10^{-81}$	$1.86 imes 10^{-324}$	2.000
M1	7	$3.37 imes10^{-21}$	$2.55 imes10^{-42}$	$1.47 imes 10^{-84}$	$5.64 imes10^{-336}$	2.000
M2	7	$7.19 imes10^{-18}$	$1.94 imes10^{-35}$	$1.41 imes10^{-70}$	1.34×10^{-279}	2.000
M3	7	$2.52 imes 10^{-18}$	$2.22 imes 10^{-36}$	$1.73 imes 10^{-72}$	2.63×10^{-287}	2.000
M4	6	$1.85 imes 10^{-22}$	$4.10 imes10^{-47}$	$2.02 imes 10^{-96}$	$5.76 imes 10^{-388}$	2.000
M5	7	$6.46 imes10^{-16}$	$2.09 imes10^{-31}$	$2.17 imes10^{-62}$	$1.34 imes10^{-246}$	2.000
M6	7	$1.23 imes10^{-20}$	$3.75 imes10^{-41}$	$3.52 imes 10^{-82}$	2.31×10^{-326}	2.000
M7	7	$1.35 imes 10^{-17}$	$7.17 imes10^{-35}$	$2.01 imes10^{-69}$	$6.02 imes 10^{-275}$	2.000
M8	6	$1.34 imes10^{-17}$	$9.03 imes 10^{-36}$	$4.08 imes 10^{-72}$	$1.66 imes 10^{-287}$	2.000

Example 3. Van der Waals Equation of State, which can be expressed as

$$\left(P + \frac{a_1 n^2}{V^2}\right)(V - na_2) = nRT,$$

explains the conduct of a real gas by taking in the perfect gas conditions two additional parameters, a_1 and a_2 , explicit for every gas. So as to decide the volume V of the gas as far as the rest of the parameters, we are required to explain the nonlinear condition in V.

$$PV^{3} - (na_{2}P + nRT)V^{2} + a_{1}n^{2}V - a_{1}a_{2}n^{3} = 0.$$

Given the constants a_1 and a_2 of a specific gas, one can find values for n, P, and T, with the end goal that this condition has three real roots. By utilizing the specific values, we get the accompanying nonlinear function

$$f1(x) = x^3 - 5.22x^2 + 9.0825x - 5.2675,$$

having three real roots. One root among three roots is a multiple zero $\alpha = 1.75$ with a multiplicity of order two, and other one being a simple zero $\xi = 1.72$. However, our desired zero is $\alpha = 1.75$. We considered the initial guess $x_0 = 1.8$ for this problem. Numerical results of various methods are shown in Table 3.

Methods	n	$ e_{n-2} $	$ e_{n-1} $	$ e_n $	$f(x_n)$	СОС
MNM	10	$2.22 imes 10^{-22}$	8.24×10^{-43}	$1.13 imes 10^{-83}$	1.36×10^{-331}	2.000
M1	10	$1.49 imes10^{-22}$	$3.70 imes10^{-43}$	$2.27 imes10^{-84}$	2.22×10^{-334}	2.000
M2	10	$1.52 imes 10^{-21}$	$3.88 imes 10^{-41}$	$2.52 imes 10^{-80}$	$3.43 imes10^{-318}$	2.000
M3	10	$1.05 imes 10^{-21}$	$1.83 imes10^{-41}$	$5.64 imes10^{-81}$	$8.54 imes10^{-321}$	2.000
M4	10	$3.11 imes10^{-24}$	$1.59 imes10^{-46}$	$4.15 imes10^{-91}$	$2.39 imes10^{-361}$	2.000
M5	10	$8.82 imes 10^{-21}$	$1.32 imes 10^{-39}$	$2.93 imes10^{-77}$	6.32×10^{-306}	2.000
M6	10	$2.42 imes 10^{-22}$	$9.73 imes10^{-43}$	$1.58 imes10^{-83}$	$5.16 imes 10^{-331}$	2.000
M7	10	$1.95 imes 10^{-21}$	$6.42 imes 10^{-41}$	$6.92 imes 10^{-80}$	1.95×10^{-316}	2.000
M8	9	$9.79 imes10^{-17}$	$1.57 imes10^{-31}$	$4.02 imes 10^{-61}$	2.10×10^{-241}	2.000

Table 3. Comparison of performance of methods for Example 3.

Example 4. Lastly, we consider the test function

$$h_4(x) = (x^2 + 1)(2xe^{x^2+1} + x^3 - x)\cosh^2\left(\frac{\pi x}{2}\right).$$

The function has a multiple zero $\alpha = i$ *of multiplicity 4. We chose the initial approximation* $x_0 = 1.25i$ *to obtain the zero of the function.*

Methods	n	$ e_{n-2} $	$ e_{n-1} $	$ e_n $	$f(x_n)$	СОС
MNM	7	$2.87 imes 10^{-15}$	$2.75 imes 10^{-30}$	$2.51 imes 10^{-60}$	5.82×10^{-478}	2.000
M1	7	$2.88 imes 10^{-15}$	$2.76 imes10^{-30}$	$2.53 imes10^{-60}$	$6.21 imes 10^{-478}$	2.000
M2	7	$3.41 imes 10^{-15}$	$3.95 imes10^{-30}$	$5.30 imes10^{-60}$	2.45×10^{-475}	2.000
M3	7	$4.43 imes10^{-15}$	$6.84 imes10^{-30}$	$1.63 imes10^{-59}$	$2.14 imes10^{-471}$	2.000
M4	7	$5.70 imes10^{-15}$	$1.16 imes10^{-29}$	$4.75 imes10^{-59}$	$1.24 imes10^{-467}$	2.000
M5	7	$5.48 imes10^{-15}$	$1.07 imes10^{-29}$	$4.06 imes10^{-59}$	$3.53 imes10^{-468}$	2.000
M6	7	$2.99 imes10^{-15}$	$2.98 imes10^{-30}$	$2.95 imes10^{-60}$	$2.10 imes10^{-477}$	2.000
M7	7	$4.48 imes10^{-15}$	$6.97 imes10^{-30}$	$1.69 imes10^{-59}$	$2.90 imes 10^{-471}$	2.000
M8	7	$3.37 imes10^{-15}$	$3.82 imes 10^{-30}$	$4.91 imes10^{-60}$	1.29×10^{-475}	2.000

Table 4. Comparison of performance of methods for Example 4.

5. Conclusions

In this paper, we presented a new one-point family of iterative methods with quadratic convergence for computing multiple roots with known multiplicity, based on the weight function technique. Analysis of the convergence showed the second order of convergence under some suppositions regarding the nonlinear function whose zeros are to be obtained. Some efficient and simple cases of the class were presented, and their stability was tested by analyzing complex geometry using a graphical tool—namely, the basin of attraction. The methods were employed to solve some real-world problems, such as the Eigenvalue problem, beam positioning problem, and the Van dar

Waal equation of state, and were also compared with existing methods. Numerical comparison of the results revealed that the presented methods had good convergence behavior, similar to the well-known modified Newton's method.

Author Contributions: The contribution of all the authors have been similar. All of them worked together to develop the present manuscript.

Funding: This research received no external funding.

Acknowledgments: We would like to express our gratitude to the anonymous reviewers for their help with the publication of this paper.

Conflicts of Interest: The authors declare no conflict of interest.

References

- 1. Argyros, I.K. Convergence and Applications of Newton-Type Iterations; Springer: New York, NY, USA, 2008.
- 2. Traub, J.F. *Iterative Methods for the Solution of Equations;* Chelsea Publishing Company: New York, NY, USA, 1982.
- 3. Hoffman, J.D. *Numerical Methods for Engineers and Scientists;* McGraw-Hill Book Company: New York, NY, USA, 1992.
- 4. Zachary, J.L. Introduction to Scientific Programming: Computational Problem Solving Using Maple and C; Springer: New York, NY, USA, 2012.
- 5. Cesarano, C. Generalized special functions in the description of fractional diffusive equations. *Commun. Appl. Ind. Math.* **2019**, *10*, 31–40.
- 6. Cesarano, C. Multi-dimensional Chebyshev polynomials: A non-conventional approach. *Commun. Appl. Ind. Math.* **2019**, *10*, 1–19.
- 7. Cesarano, C.; Ricci, P.E. Orthogonality Properties of the Pseudo-Chebyshev Functions (Variations on a Chebyshev's Theme). *Mathematics* **2019**, *7*, 180.
- 8. Schröder, E. Über unendlich viele Algorithmen zur Auflösung der Gleichungen. *Math. Ann.* 1870, 2, 317–365.
- 9. Dong, C. A family of multipoint iterative functions for finding multiple roots of equations. *Int. J. Comput. Math.* **1987**, 21, 363–367.
- 10. Geum, Y.H.; Kim, Y.I.; Neta, B. A class of two-point sixth-order multiple-zero finders of modified double-Newton type and their dynamics. *Appl. Math. Comput.* **2015**, *270*, 387–400.
- 11. Hansen, E.; Patrick, M. A family of root finding methods. *Numer. Math.* **1977**, *27*, 257–269.
- 12. Li, S.; Liao, X.; Cheng, L. A new fourth-order iterative method for finding multiple roots of nonlinear equations. *Appl. Math. Comput.* **2009**, *215*, 1288–1292.
- 13. Neta, B. New third order nonlinear solvers for multiple roots. Appl. Math. Comput. 2008, 202, 162–170.
- 14. Osada, N. An optimal multiple root-finding method of order three. J. Comput. Appl. Math. 1994, 51, 131–133.
- 15. Sharifi, M.; Babajee, D.K.R.; Soleymani, F. Finding the solution of nonlinear equations by a class of optimal methods. *Comput. Math. Appl.* **2012**, *63*, 764–774.
- 16. Sharma, J.R.; Sharma, R. Modified Jarratt method for computing multiple roots. *Appl. Math. Comput.* **2010**, 217, 878–881.
- 17. Victory, H.D.; Neta, B. A higher order method for multiple zeros of nonlinear functions. *Int. J. Comput. Math.* **1983**, *12*, 329–335.
- 18. Zhou, X.; Chen, X.; Song, Y. Constructing higher-order methods for obtaining the multiple roots of nonlinear equations. *J. Comput. Appl. Math.* **2011**, 235, 4199–4206.
- 19. Ostrowski, A.M. Solution of Equations and Systems of Equations; Academic Press: New York, NY, USA, 1966.
- 20. Kung, H.T.; Traub, J.F. Optimal order of one-point and multipoint iteration. J. Assoc. Comput. Mach. 1974, 21, 643–651.
- 21. Geum, Y.H.; Kim, Y.I.; Neta, B. A sixth–order family of three–point modified Newton–like multiple–root finders and the dynamics behind their extraneous fixed points. *Appl. Math. Comput.* **2016**, *283*, 120–140.
- 22. Behl, R.; Cordero, A.; Motsa, S.S.; Torregrosa, J.R. An eighth-order family of optimal multiple root finders and its dynamics. *Numer. Algor.* **2018**, *77*, 1249–1272.
- 23. Zafar, F.; Cordero, A.; Rana, Q.; Torregrosa, J.R. Optimal iterative methods for finding multiple roots of nonlinear equations using free parameters. *J. Math. Chem.* **2017**, doi:10.1007/s10910-017-0813-1.

- 24. Geum, Y.H.; Kim, Y.I.; Neta, B. Constructing a family of optimal eighth-order modified Newton-type multiple-zero finders along with the dynamics behind their purely imaginary extraneous fixed points. *J. Comput. Appl. Math.* **2018**, *333*, 131–156.
- 25. Behl, R.; Zafar, F.; Alshomrani, A.S.; Junjua, M.; Yasmin, N. An optimal eighth-order scheme for multiple zeros of univariate function. *Int. J. Comput. Math.* **2019**, *16*, 1843002.
- 26. Behl, R.; Alshomrani, A.S.; Motsa, S.S. An optimal scheme for multiple roots of nonlinear equations with eighth-order convergence. *J. Math. Chem.* **2018**, doi:10.1007/s10910-018-0857-x.
- 27. Zafar, F.; Cordero, A.; Torregrosa, J.R. An efficient family of optimal eighth-order multiple root finder. *Mathematics* **2018**, *6*, 310.
- 28. Chicharro, F.I.; Cordero, A.; Garrido, N.; Torregrosa, J.R. Genertaing root-finder iterative methods of second-order: Convergence and stability. *Axioms* **2019**, *8*, 55.
- 29. Vrscay, E.R.; Gilbert, W.J. Extraneous fixed points, basin boundaries and chaotic dynamics for Schröder and König rational iteration functions. *Numer. Math.* **1988**, *52*, 1–16.
- 30. Scott, M.; Neta, B.; Chun, C. Basin attractors for various methods. Appl. Math. Comput. 2011, 218, 2584–2599.
- 31. Varona, J.L. Graphic and numerical comparison between iterative methods. Math. Intell. 2002, 24, 37–46.
- 32. Weerakoon, S.; Fernando, T.G.I. A variant of Newton's method with accelerated third-order convergence. *Appl. Math. Lett.* **2000**, *13*, 87–93.



 \odot 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses/by/4.0/).