

Article

Comultiplications on the Localized Spheres and Moore Spaces

Dae-Woong Lee 

Department of Mathematics, and Institute of Pure and Applied Mathematics, Jeonbuk National University, 567 Baekje-daero, Deokjin-gu, Jeonju-si, Jeollabuk-do 54896, Korea; dwlee@jbnu.ac.kr

Received: 2 December 2019; Accepted: 30 December 2019; Published: 5 January 2020



Abstract: Any nilpotent CW-space can be localized at primes in a similar way to the localization of a ring at a prime number. For a collection \mathcal{P} of prime numbers which may be empty and a localization $X_{\mathcal{P}}$ of a nilpotent CW-space X at \mathcal{P} , we let $|C(X)|$ and $|C(X_{\mathcal{P}})|$ be the cardinalities of the sets of all homotopy comultiplications on X and $X_{\mathcal{P}}$, respectively. In this paper, we show that if $|C(X)|$ is finite, then $|C(X)| \geq |C(X_{\mathcal{P}})|$, and if $|C(X)|$ is infinite, then $|C(X)| = |C(X_{\mathcal{P}})|$, where X is the k -fold wedge sum $\bigvee_{i=1}^k S^{n_i}$ or Moore spaces $M(G, n)$. Moreover, we provide examples to concretely determine the cardinality of homotopy comultiplications on the k -fold wedge sum of spheres, Moore spaces, and their localizations.

Keywords: comultiplications; localized spheres; basic Whitehead products; Hilton formula; Moore space

MSC: Primary 55Q20; Secondary 55P60, 55Q40, 55Q15, 20F18

1. Introduction

Homotopy comultiplications, one of the Eckmann–Hilton dual notions of homotopy multiplications, play a fundamental role in classical and rational homotopy theories. One reason for this is that the set of pointed homotopy classes of base point preserving continuous maps from a co-H-space X to a space Y has a canonical binary operation with identity induced by the homotopy comultiplication on a co-H-space. The calculation of homotopy comultiplications is very complicated in that there are usually many homotopy comultiplications on a given co-H-space with many different homotopy properties.

Localization theories in the pointed homotopy category of simply connected (or nilpotent) CW-complexes have been developed in algebraic or topological forms. Moreover, the study of homotopy comultiplications, binary operations for homotopy groups with coefficients and the same n -type structures of co-H-spaces has been carried out by several authors; see [1–8]. For example, homotopy comultiplications on the wedge sum of circles were developed by using methods of group theory in [9]. The techniques of rational homotopy were applied to obtain some rational results for finite 1-connected co-H-spaces in [10,11]. The homotopy comultiplications on the wedge sum of two Moore spaces were investigated using homological algebra in [12]. The cardinality of homotopy comultiplications on a suspension has been determined in [13]. A topological transversality theorem for multivalued maps with continuous, compact selections was developed in [14]. From the equivariant homotopy theoretic point of view, an explicit expression of the behavior of the local cohomology spectral sequence graded on the representation ring was presented with many pictures in [15].

In this paper, we study the cardinality of the set of all homotopy comultiplications on a wedge of (localized) spheres or (localized) Moore spaces, and develop the properties of the homotopy

comultiplications of those CW-spaces. Our methods can be used to study homotopy comultiplications on a wedge of any number of spheres.

The paper is organized as follows: In Section 2, we describe the fundamental notions of homotopy comultiplications and introduce the localized version of Hilton's formula. In Section 3, we establish basic facts regarding the types of homotopy comultiplications on the k -fold wedge sum of localized spheres and define certain homotopy comultiplications on the k -fold wedge sum of localized spheres and Moore spaces. In Section 4, we describe the main result of this paper and give an example to illustrate it.

Convention. In this paper, we work in the pointed homotopy category \mathcal{C} of connected nilpotent CW-complexes and homotopy classes of base point preserving continuous maps. We mostly use '=' for the homotopy ' \simeq ' unless we emphasize the homotopy. We also do not distinguish notationally between a continuous map and its homotopy class.

2. Preliminaries

Let $\{X_\gamma | \gamma \in \Gamma\}$ be a family of pointed topological spaces whose base points are x_γ for each $\gamma \in \Gamma$. The wedge sum $\bigvee_{\gamma \in \Gamma} X_\gamma$ of $\{X_\gamma | \gamma \in \Gamma\}$ is defined as the quotient space

$$\bigvee_{\gamma \in \Gamma} X_\gamma = \coprod_{\gamma \in \Gamma} X_\gamma / \{x_\gamma | \gamma \in \Gamma\},$$

where $\coprod_{\gamma \in \Gamma} X_\gamma$ is the topological sum of $\{X_\gamma | \gamma \in \Gamma\}$.

For any abelian group G and a positive integer $n \geq 2$, there exists a 1-connected CW-complex X such that

$$H_i(X) = \begin{cases} G & \text{if } i = n; \\ 0 & \text{if } i \neq n. \end{cases}$$

The homotopy type of X is uniquely determined up to homotopy and it is said to be a *Moore space* of type (G, n) . We denote the Moore space of type (G, n) (or any CW-space which has the same homotopy type to it) by $M(G, n)$.

Localization plays a pivotal role in both algebra and algebraic topology, especially in homotopy theory. For a collection \mathcal{P} of prime numbers which may be empty, a group G is called a \mathcal{P} -local group if the map

$$q : G \rightarrow G$$

given by

$$q(g) = \underbrace{g + g + \dots + g}_{(q\text{-times})}$$

is bijective for all $q \in \mathcal{P}^c$. A nilpotent CW-complex X is said to be \mathcal{P} -local if $\pi_n(X)$ is \mathcal{P} -local for all $n \geq 1$. Any nilpotent CW-space can be localized at primes in the sense of Sullivan [16] and Bousfield and Kan [17] in a similar way to the localization of a ring at a prime number. As usual, the localization of a nilpotent CW-space X at \mathcal{P} is denoted as $X_{\mathcal{P}}$ which is unique up to homotopy.

A pair (X, ψ) consisting of a space X and a base point preserving continuous map $\psi : X \rightarrow X \vee X$ is said to be a *co-Hopf space*, or *co-H-space* for short, if $\pi^1 \psi \simeq 1_X$ and $\pi^2 \psi \simeq 1_X$, where 1_X is the identity map of X and $\pi^1, \pi^2 : X \vee X \rightarrow X$ are the first and second projections, respectively. In this case, the map $\psi : X \rightarrow X \vee X$ is said to be a *homotopy comultiplication*, or *comultiplication* for short. Homotopy comultiplication is an Eckmann–Hilton dual notion of homotopy multiplication; see [18–20]. It can be seen that (X, ψ) is a co-H-space if

$$j\psi \simeq \Delta : X \rightarrow X \times X,$$

where $j : X \vee X \rightarrow X \times X$ is the inclusion map and Δ is the diagonal map.

We now consider the k -fold wedge sum of spheres $X := \mathbb{S}^{n_1} \vee \mathbb{S}^{n_2} \vee \dots \vee \mathbb{S}^{n_k}$. Let $\xi_i : \mathbb{S}^{n_i} \rightarrow X$, $i = 1, 2, \dots, k$ be the canonical inclusion maps. We now construct the so-called *basic Whitehead products* (see also [21–24]) by using the homotopy classes of those canonical inclusion maps by induction on n . We denote the basic Whitehead products of weight 1 by $\xi_1, \xi_2, \dots, \xi_k$ that are ordered by the canonical inclusion maps in the way so that $\xi_1 < \xi_2 < \dots < \xi_k$. Assume that the basic Whitehead products of weight less than or equal to $n - 1$ have been already defined and ordered so that if $r < s < n$, all basic Whitehead products of weight r are less than all of the basic Whitehead products of weight s . Then a basic Whitehead product of weight n is a basic Whitehead product $[a, b]$, where a and b are basic Whitehead products of weights p and q , respectively, with $p + q = n$ and $a < b$. If b is a basic Whitehead product $[c, d]$ consisting of basic Whitehead products c and d , then the condition $c \leq a$ is necessary. Then the basic Whitehead products of weight n are greater than any basic Whitehead product of weight less than n and are ordered among themselves arbitrarily.

For a basic Whitehead product of weight n , we associate to it a string of canonical inclusion maps $\xi_1, \xi_2, \dots, \xi_k$ that appear in the basic Whitehead products. The *height* h_s of the basic Whitehead product w_s is defined as

$$h_s = \sum_i t_i(n_i - 1),$$

where t_i is the number of the canonical inclusion map ξ_i that appears in the basic Whitehead product w_s for $s = 1, 2, 3, \dots$

We are interested in the cardinality of the set of homotopy comultiplications on a co-H-space. In the case of a wedge of two spheres, we obtain the following.

Lemma 1. *The cardinality of homotopy comultiplications of $\mathbb{S}^n \vee \mathbb{S}^m$ is*

$$\prod_{s=3}^{\infty} |\pi_m(\mathbb{S}^{h_s+1})|,$$

where $2 \leq n < m$ and $|X|$ denotes the cardinality of a set X .

Proof. See ([22], Corollary 2.9) and ([24], Corollary 2.15) for the more general case. \square

One of the most important theorems in algebraic topology is the Hilton's formula [25], which is described as follows.

Theorem 1. *Let $w_1, w_2, \dots, w_s, \dots$ be the basic Whitehead products of the k -fold wedge sum of spheres*

$$X = \mathbb{S}^{n_1} \vee \mathbb{S}^{n_2} \vee \dots \vee \mathbb{S}^{n_k}.$$

Then for all m , we have

$$\pi_m(X) \cong \bigoplus_{s=1}^{\infty} \pi_m(\mathbb{S}^{h_s+1}).$$

Here,

- h_s is the height of w_s ; and
- the isomorphism $\theta : \bigoplus_{s=1}^{\infty} \pi_m(\mathbb{S}^{h_s+1}) \rightarrow \pi_m(X)$ is given by

$$\theta|_{\pi_m(\mathbb{S}^{h_s+1})} = w_{s*} : \pi_m(\mathbb{S}^{h_s+1}) \rightarrow \pi_m(X).$$

The direct sum is finite for each positive integer m since the height h_s goes to infinity as $s \rightarrow \infty$.

3. Comultiplications on a Wedge of (Localized) Spheres and Moore Spaces

In this section, we investigate the cardinality of the set of all homotopy comultiplications on the k -fold wedge sum of localized spheres and Moore spaces. In particular, we are interested in determining inequalities and relationships between the cardinalities of the sets of homotopy comultiplications on the wedge sum of spheres, Moore spaces and their localizations.

3.1. The Localized Version of Hilton's Formula

Let \mathcal{P} be a collection of prime numbers and let $G_{\mathcal{P}}$ denote the \mathcal{P} -localization of a nilpotent group G . It is well known that if $h : G \rightarrow H$ is any homomorphism in the category of nilpotent groups, then we have a unique map $h_{\mathcal{P}} : G_{\mathcal{P}} \rightarrow H_{\mathcal{P}}$ such that the diagram

$$\begin{array}{ccc} G & \xrightarrow{h} & H \\ \downarrow e & & \downarrow e \\ G_{\mathcal{P}} & \xrightarrow{h_{\mathcal{P}}} & H_{\mathcal{P}} \end{array}$$

is strictly commutative, where $e : G \rightarrow G_{\mathcal{P}}$ is a \mathcal{P} -localizing map; similarly, for the nilpotent CW-spaces.

Lemma 2. Let $\psi : X \rightarrow X \vee X$ be a comultiplication on a nilpotent CW-space X . Then the localization $\psi_{\mathcal{P}} : X_{\mathcal{P}} \rightarrow X_{\mathcal{P}} \vee X_{\mathcal{P}}$ at a set \mathcal{P} of primes is a comultiplication.

Proof. Since the localization of a nilpotent CW-space has the functorial property, all the faces including the bottom triangle colored in blue of the following triangular prism are strictly commutative.

Here (in Figure 1),

- $\pi^1 : X \vee X \rightarrow X$ is the first projection;
- $\pi_{\mathcal{P}}^1 : X_{\mathcal{P}} \vee X_{\mathcal{P}} \rightarrow X_{\mathcal{P}}$ is the first projection (obtained by π^1 by taking the \mathcal{P} -localization);
- 1_X is the identity map of X ; and
- $e : X \rightarrow X_{\mathcal{P}}$ is a \mathcal{P} -localizing map.

Similarly, we obtain the same result for the second projection $\pi^2 : X \vee X \rightarrow X$. This implies that if $\psi : X \rightarrow X \vee X$ is a comultiplication on X , then its localization $\psi_{\mathcal{P}} : X_{\mathcal{P}} \rightarrow X_{\mathcal{P}} \vee X_{\mathcal{P}}$ also has the comultiplication structure on the localization $X_{\mathcal{P}}$ of X . \square




Figure 1. The triangular prism.

We now consider the localized version of the Hilton's formula described in Theorem 1 of Section 2 as follows.

Theorem 2. Let $X_{\mathcal{P}}$ be the localization of the k -fold wedge sum of spheres $X := \mathbb{S}^{n_1} \vee \mathbb{S}^{n_2} \vee \dots \vee \mathbb{S}^{n_k}$. Then we have

$$\pi_m(X_{\mathcal{P}}) \cong \bigoplus_{s=1}^{\infty} \pi_m(\mathbb{S}_{\mathcal{P}}^{h_s+1})$$

for all m .

Proof. We note that localization preserves the Cartesian product, the smash product and the wedge sum in the pointed homotopy category of pointed nilpotent CW-complexes; that is,

- $(X \times Y)_{\mathcal{P}} \simeq X_{\mathcal{P}} \times Y_{\mathcal{P}}$;
- $(X \wedge Y)_{\mathcal{P}} \simeq X_{\mathcal{P}} \wedge Y_{\mathcal{P}}$; and
- $(X \vee Y)_{\mathcal{P}} \simeq X_{\mathcal{P}} \vee Y_{\mathcal{P}}$.

Since the localizations of nilpotent CW-complexes are unique up to homotopy equivalence and the localizations of nilpotent groups commute with finite direct sums, we have

$$\begin{aligned} \pi_m(X_{\mathcal{P}}) &\cong \pi_m(X) \otimes \mathbb{Z}_{\mathcal{P}} \\ &\cong \left(\bigoplus_{s=1}^{\infty} \pi_m(\mathbb{S}^{h_s+1}) \right) \otimes \mathbb{Z}_{\mathcal{P}} \\ &\cong \bigoplus_{s=1}^{\infty} \left(\pi_m(\mathbb{S}^{h_s+1}) \otimes \mathbb{Z}_{\mathcal{P}} \right) \\ &\cong \bigoplus_{s=1}^{\infty} \pi_m(\mathbb{S}_{\mathcal{P}}^{h_s+1}), \end{aligned}$$

as required. \square

3.2. The 2-Fold Wedge of (Localized) Spheres

We begin with the 2-fold wedge sum of (localized) spheres to illustrate our methods. The results will be applied to the k -fold wedge sum of (localized) spheres for $k \geq 3$.

Notation. Throughout this (sub)section, we will make use of the following notations.

- $X := \mathbb{S}^n \vee \mathbb{S}^m$ is the wedge sum of spheres, where $2 \leq n < m$.
- $k^1 : \mathbb{S}^n \rightarrow X$ and $k^2 : \mathbb{S}^m \rightarrow X$ are the inclusion maps.
- $i^1, i^2 : \mathbb{S}^n \rightarrow \mathbb{S}^n \vee \mathbb{S}^n$ are the first and second inclusion maps, respectively.
- $p^1, p^2 : \mathbb{S}^n \vee \mathbb{S}^n \rightarrow \mathbb{S}^n$ are the first and second projections, respectively.
- $\iota^1, \iota^2 : X \rightarrow X \vee X$ are the first and second inclusion maps, respectively.
- $\pi^1, \pi^2 : X \vee X \rightarrow X$ are the first and second projections, respectively.
- \mathcal{P} is a collection of prime numbers.
- $f_{\mathcal{P}} : A_{\mathcal{P}} \rightarrow B_{\mathcal{P}}$ is the \mathcal{P} -localization of a map $f : A \rightarrow B$, where A and B are object classes in the pointed homotopy category \mathcal{C} of connected nilpotent CW-spaces.
- $C(X) \subseteq [X; X \vee X]$ is the set of all homotopy classes of comultiplications on a nilpotent CW-space X , e.g., $X := \bigvee_{i=1}^k \mathbb{S}^{n_i}$, Moore spaces, or their localizations.

Lemma 3. Let $X_{\mathcal{P}} = \mathbb{S}_{\mathcal{P}}^n \vee \mathbb{S}_{\mathcal{P}}^m$ be a wedge sum of localized spheres with $2 \leq n < m$. Then any comultiplication $\psi : X_{\mathcal{P}} \rightarrow X_{\mathcal{P}} \vee X_{\mathcal{P}}$ has the following type

$$\begin{cases} \psi k_{\mathcal{P}}^1 &= \iota_{\mathcal{P}}^1 k_{\mathcal{P}}^1 + \iota_{\mathcal{P}}^2 k_{\mathcal{P}}^1 \\ \psi k_{\mathcal{P}}^2 &= \iota_{\mathcal{P}}^1 k_{\mathcal{P}}^2 + \iota_{\mathcal{P}}^2 k_{\mathcal{P}}^2 + Q \end{cases} \quad (1)$$

for some homotopy class $Q : \mathbb{S}_{\mathcal{P}}^m \rightarrow X_{\mathcal{P}} \vee X_{\mathcal{P}}$ satisfying $\pi_{\mathcal{P}}^1 Q = 0 = \pi_{\mathcal{P}}^2 Q$, where the additions are the homotopy additions in the homotopy groups.

We note that if Y is a CW-complex which is filtered by its n -skeletons Y^n for $n \geq 0$, then its localization $Y_{\mathcal{P}}$ of Y at \mathcal{P} is filtered by $Y_{\mathcal{P}}^n$. Moreover, the two CW-spaces $Y_{\mathcal{P}}^{n+1}/Y_{\mathcal{P}}^n$ and $(Y^{n+1}/Y^n)_{\mathcal{P}}$ have the same homotopy type, and each is a wedge sum of the localized spheres $\mathbb{S}_{\mathcal{P}}^{n+1}$.

Proof. We observe that $X_{\mathcal{P}}$ is a localized CW-complex and thus there is a cellular map ([19] p. 77)

$$\psi' : X_{\mathcal{P}} \rightarrow X_{\mathcal{P}} \vee X_{\mathcal{P}}$$

such that

$$\psi \simeq \psi' : X_{\mathcal{P}} \rightarrow X_{\mathcal{P}} \vee X_{\mathcal{P}}.$$

Therefore, we have

$$\psi'(X_{\mathcal{P}}^n) \subset (X_{\mathcal{P}} \vee X_{\mathcal{P}})^n.$$

We note that $X_{\mathcal{P}}^n = \mathbb{S}_{\mathcal{P}}^n$ and that the localized sphere $\mathbb{S}_{\mathcal{P}}^n$ has a CW-decomposition that consists of one localized zero cell and one localized n -cell. Therefore, we obtain

$$(X_{\mathcal{P}} \vee X_{\mathcal{P}})^n = \mathbb{S}_{\mathcal{P}}^n \vee \mathbb{S}_{\mathcal{P}}^n$$

and observe that ψ' gives a base point preserving continuous map $\eta : \mathbb{S}_{\mathcal{P}}^n \rightarrow \mathbb{S}_{\mathcal{P}}^n \vee \mathbb{S}_{\mathcal{P}}^n$ such that the following diagram

$$\begin{array}{ccccc} \mathbb{S}_{\mathcal{P}}^n & \xrightarrow{\eta} & \mathbb{S}_{\mathcal{P}}^n \vee \mathbb{S}_{\mathcal{P}}^n & \xrightarrow{I} & \mathbb{S}_{\mathcal{P}}^n \times \mathbb{S}_{\mathcal{P}}^n \\ \downarrow k_{\mathcal{P}}^1 & & \downarrow k_{\mathcal{P}}^1 \vee k_{\mathcal{P}}^1 & & \downarrow k_{\mathcal{P}}^1 \times k_{\mathcal{P}}^1 \\ X_{\mathcal{P}} & \xrightarrow{\psi \simeq \psi'} & X_{\mathcal{P}} \vee X_{\mathcal{P}} & \xrightarrow{J} & X_{\mathcal{P}} \times X_{\mathcal{P}} \end{array} \quad (2)$$

is commutative up to homotopy, where I and J are the inclusion maps. We also note that the following triangle

$$\begin{array}{ccc} X_{\mathcal{P}} & \xrightarrow{\psi} & X_{\mathcal{P}} \vee X_{\mathcal{P}} \\ & \searrow \Delta & \downarrow J \\ & & X_{\mathcal{P}} \times X_{\mathcal{P}} \end{array} \quad (3)$$

is homotopy commutative, where $\Delta : X_{\mathcal{P}} \rightarrow X_{\mathcal{P}} \times X_{\mathcal{P}}$ is the diagonal map. From the homotopy commutative diagrams (2) and (3), we obtain

$$(k_{\mathcal{P}}^1 \times k_{\mathcal{P}}^1)I\eta \simeq J\psi k_{\mathcal{P}}^1 \simeq \Delta k_{\mathcal{P}}^1 = (k_{\mathcal{P}}^1 \times k_{\mathcal{P}}^1)\bar{\Delta}, \quad (4)$$

where $\bar{\Delta} : \mathbb{S}_{\mathcal{P}}^n \rightarrow \mathbb{S}_{\mathcal{P}}^n \times \mathbb{S}_{\mathcal{P}}^n$ is the diagonal map. Since

$$(k_{\mathcal{P}}^1 \times k_{\mathcal{P}}^1)_* : [\mathbb{S}_{\mathcal{P}}^n; \mathbb{S}_{\mathcal{P}}^n \times \mathbb{S}_{\mathcal{P}}^n] \rightarrow [\mathbb{S}_{\mathcal{P}}^n; X_{\mathcal{P}} \times X_{\mathcal{P}}]$$

is a monomorphism of homotopy groups, from (4) we have

$$I\eta \simeq \bar{\Delta};$$

that is, the following triangle

$$\begin{array}{ccc} \mathbb{S}_{\mathcal{P}}^n & \xrightarrow{\eta} & \mathbb{S}_{\mathcal{P}}^n \vee \mathbb{S}_{\mathcal{P}}^n \\ & \searrow \bar{\Delta} & \downarrow I \\ & & \mathbb{S}_{\mathcal{P}}^n \times \mathbb{S}_{\mathcal{P}}^n \end{array} \quad (5)$$

is also commutative up to homotopy, where $\bar{\Delta} : \mathbb{S}_{\mathcal{P}}^n \rightarrow \mathbb{S}_{\mathcal{P}}^n \times \mathbb{S}_{\mathcal{P}}^n$ is the diagonal map. This implies that the map $\eta : \mathbb{S}_{\mathcal{P}}^n \rightarrow \mathbb{S}_{\mathcal{P}}^n \vee \mathbb{S}_{\mathcal{P}}^n$ is also a comultiplication. Since the localized sphere $\mathbb{S}_{\mathcal{P}}^n$ is 1-connected for

$n \geq 2$, we see that the comultiplication η is unique up to homotopy (see ([1] Proposition 3.1)) and has the form

$$\eta = i_{\mathcal{P}}^1 + {}_{\eta}i_{\mathcal{P}}^2,$$

where $+_{\eta}$ is the addition induced by the comultiplication η . By the uniqueness of the comultiplication

$$\eta : \mathbb{S}_{\mathcal{P}}^n \rightarrow \mathbb{S}_{\mathcal{P}}^n \vee \mathbb{S}_{\mathcal{P}}^n$$

as the standard (or suspension) comultiplication, we have

$$i_{\mathcal{P}}^1 + {}_{\eta}i_{\mathcal{P}}^2 = i_{\mathcal{P}}^1 + i_{\mathcal{P}}^2,$$

where $+$ is the homotopy addition in the homotopy group

$$[\mathbb{S}_{\mathcal{P}}^n; X_{\mathcal{P}} \vee X_{\mathcal{P}}] \cong [\mathbb{S}^n; X_{\mathcal{P}} \vee X_{\mathcal{P}}] \cong \pi_n(X_{\mathcal{P}} \vee X_{\mathcal{P}}).$$

Hence

$$\psi k_{\mathcal{P}}^1 = (k_{\mathcal{P}}^1 \vee k_{\mathcal{P}}^1)\eta = (k_{\mathcal{P}}^1 \vee k_{\mathcal{P}}^1)(i_{\mathcal{P}}^1 + {}_{\eta}i_{\mathcal{P}}^2) = i_{\mathcal{P}}^1 k_{\mathcal{P}}^1 + i_{\mathcal{P}}^2 k_{\mathcal{P}}^1.$$

We now consider the homotopy class Q in the homotopy group $[\mathbb{S}_{\mathcal{P}}^n; X_{\mathcal{P}} \vee X_{\mathcal{P}}]$ as an abelian group given by

$$Q = \psi k_{\mathcal{P}}^2 - i_{\mathcal{P}}^1 k_{\mathcal{P}}^2 - i_{\mathcal{P}}^2 k_{\mathcal{P}}^2;$$

that is,

$$\psi k_{\mathcal{P}}^2 = i_{\mathcal{P}}^1 k_{\mathcal{P}}^2 + i_{\mathcal{P}}^2 k_{\mathcal{P}}^2 + Q.$$

Then we obtain

$$\pi_{\mathcal{P}}^1 Q = \pi_{\mathcal{P}}^1 \psi k_{\mathcal{P}}^2 - \pi_{\mathcal{P}}^1 i_{\mathcal{P}}^1 k_{\mathcal{P}}^2 - \pi_{\mathcal{P}}^1 i_{\mathcal{P}}^2 k_{\mathcal{P}}^2 = k_{\mathcal{P}}^2 - k_{\mathcal{P}}^2 - 0 = 0$$

and

$$\pi_{\mathcal{P}}^2 Q = \pi_{\mathcal{P}}^2 \psi k_{\mathcal{P}}^2 - \pi_{\mathcal{P}}^2 i_{\mathcal{P}}^1 k_{\mathcal{P}}^2 - \pi_{\mathcal{P}}^2 i_{\mathcal{P}}^2 k_{\mathcal{P}}^2 = k_{\mathcal{P}}^2 - 0 - k_{\mathcal{P}}^2 = 0,$$

as required. \square

Lemma 4. If $Q : \mathbb{S}_{\mathcal{P}}^m \rightarrow X_{\mathcal{P}} \vee X_{\mathcal{P}}$ is any homotopy class in $[\mathbb{S}_{\mathcal{P}}^m; X_{\mathcal{P}} \vee X_{\mathcal{P}}]$ such that

$$\pi_{\mathcal{P}}^1 Q = 0 = \pi_{\mathcal{P}}^2 Q,$$

then the map $\psi : X_{\mathcal{P}} \rightarrow X_{\mathcal{P}} \vee X_{\mathcal{P}}$ defined in Equation (1) is a comultiplication.

Proof. We have

- $\pi_{\mathcal{P}}^1 \psi k_{\mathcal{P}}^1 = \pi_{\mathcal{P}}^1 (i_{\mathcal{P}}^1 k_{\mathcal{P}}^1 + i_{\mathcal{P}}^2 k_{\mathcal{P}}^1) = k_{\mathcal{P}}^1 + 0 = k_{\mathcal{P}}^1;$
- $\pi_{\mathcal{P}}^1 \psi k_{\mathcal{P}}^2 = \pi_{\mathcal{P}}^1 (i_{\mathcal{P}}^1 k_{\mathcal{P}}^2 + i_{\mathcal{P}}^2 k_{\mathcal{P}}^2 + Q) = k_{\mathcal{P}}^2 + 0 + \pi_{\mathcal{P}}^1 Q = k_{\mathcal{P}}^2 + 0 + 0 = k_{\mathcal{P}}^2;$
- $\pi_{\mathcal{P}}^2 \psi k_{\mathcal{P}}^1 = \pi_{\mathcal{P}}^2 (i_{\mathcal{P}}^1 k_{\mathcal{P}}^1 + i_{\mathcal{P}}^2 k_{\mathcal{P}}^1) = 0 + k_{\mathcal{P}}^1 = k_{\mathcal{P}}^1;$ and
- $\pi_{\mathcal{P}}^2 \psi k_{\mathcal{P}}^2 = \pi_{\mathcal{P}}^2 (i_{\mathcal{P}}^1 k_{\mathcal{P}}^2 + i_{\mathcal{P}}^2 k_{\mathcal{P}}^2 + Q) = 0 + k_{\mathcal{P}}^2 + \pi_{\mathcal{P}}^2 Q = 0 + k_{\mathcal{P}}^2 + 0 = k_{\mathcal{P}}^2;$

that is, $\pi_{\mathcal{P}}^1 \psi = 1_{X_{\mathcal{P}}}$ and $\pi_{\mathcal{P}}^2 \psi = 1_{X_{\mathcal{P}}}$. This implies that ψ is a comultiplication on $X_{\mathcal{P}}$. \square

Using the results above, we can define a comultiplication on a wedge of localized spheres as follows.

Definition 1. Let $Q : \mathbb{S}_{\mathcal{P}}^m \rightarrow X_{\mathcal{P}} \vee X_{\mathcal{P}}$ be any element of $[\mathbb{S}_{\mathcal{P}}^m; X_{\mathcal{P}} \vee X_{\mathcal{P}}]$ satisfying

$$\pi_{\mathcal{P}}^1 Q = 0 = \pi_{\mathcal{P}}^2 Q.$$

We define a comultiplication $\psi_Q : X_{\mathcal{P}} \rightarrow X_{\mathcal{P}} \vee X_{\mathcal{P}}$ by

$$\begin{cases} \psi_Q k_{\mathcal{P}}^1 &= i_{\mathcal{P}}^1 k_{\mathcal{P}}^1 + i_{\mathcal{P}}^2 k_{\mathcal{P}}^1 \\ \psi_Q k_{\mathcal{P}}^2 &= i_{\mathcal{P}}^1 k_{\mathcal{P}}^2 + i_{\mathcal{P}}^2 k_{\mathcal{P}}^2 + Q. \end{cases}$$

The homotopy class Q is called a perturbation of the comultiplication ψ_Q .

We note that if $e : \mathbb{S}^{2n-1} \rightarrow \mathbb{S}_{\mathcal{P}}^{2n-1}$ is the \mathcal{P} -localizing map, then the homomorphism

$$e^* : [\mathbb{S}_{\mathcal{P}}^{2n-1}; \mathbb{S}_{\mathcal{P}}^n \vee \mathbb{S}_{\mathcal{P}}^n] \xrightarrow{\cong} [\mathbb{S}^{2n-1}; \mathbb{S}_{\mathcal{P}}^n \vee \mathbb{S}_{\mathcal{P}}^n] \quad (6)$$

induced by e is an isomorphism of homotopy groups.

Convention. For our convenience in notation, using the isomorphism of homotopy groups in (6), we make use of the same notations of the homotopy classes on both sides of the isomorphism; that is,

$$\gamma \mapsto e^*(\gamma) = \gamma \circ e \longleftrightarrow \gamma \in [\mathbb{S}_{\mathcal{P}}^{2n-1}; \mathbb{S}_{\mathcal{P}}^n \vee \mathbb{S}_{\mathcal{P}}^n]$$

for any homotopy class γ in the homotopy group $[\mathbb{S}_{\mathcal{P}}^{2n-1}; \mathbb{S}_{\mathcal{P}}^n \vee \mathbb{S}_{\mathcal{P}}^n]$. In particular,

$$\gamma = [i_{\mathcal{P}}^1, i_{\mathcal{P}}^2] \in [\mathbb{S}_{\mathcal{P}}^{2n-1}; \mathbb{S}_{\mathcal{P}}^n \vee \mathbb{S}_{\mathcal{P}}^n]$$

as the generalized Whitehead product in the sense of Arkowitz [26]; see below.

Lemma 5. Let $[i^1, i^2] : \mathbb{S}^{2n-1} \rightarrow \mathbb{S}^n \vee \mathbb{S}^n$ be the Whitehead product. Then we have

$$[i^1, i^2]_{\mathcal{P}} = [i_{\mathcal{P}}^1, i_{\mathcal{P}}^2].$$

Proof. The functorial property of the localization shows that the following diagram

$$\begin{array}{ccccccc} \mathbb{S}^{2n-1} & \xrightarrow{a} & \mathbb{S}^n \vee \mathbb{S}^n & \xrightarrow{i^1 \vee i^2} & \mathbb{S}^n \vee \mathbb{S}^n \vee \mathbb{S}^n \vee \mathbb{S}^n & \xrightarrow{\nabla} & \mathbb{S}^n \vee \mathbb{S}^n \\ \downarrow e & & \downarrow e & & \downarrow e & & \downarrow e \\ \mathbb{S}_{\mathcal{P}}^{2n-1} & \xrightarrow{a_{\mathcal{P}}} & \mathbb{S}_{\mathcal{P}}^n \vee \mathbb{S}_{\mathcal{P}}^n & \xrightarrow{i_{\mathcal{P}}^1 \vee i_{\mathcal{P}}^2} & \mathbb{S}_{\mathcal{P}}^n \vee \mathbb{S}_{\mathcal{P}}^n \vee \mathbb{S}_{\mathcal{P}}^n \vee \mathbb{S}_{\mathcal{P}}^n & \xrightarrow{\nabla_{\mathcal{P}}} & \mathbb{S}_{\mathcal{P}}^n \vee \mathbb{S}_{\mathcal{P}}^n \end{array} \quad (7)$$

is commutative up to homotopy, where $a : \mathbb{S}^{2n-1} \rightarrow \mathbb{S}^n \vee \mathbb{S}^n$ is the attaching map and $e : Y \rightarrow Y_{\mathcal{P}}$ is a \mathcal{P} -localizing map. Since

$$[\mathbb{S}^{2n-1}; \mathbb{S}^n \vee \mathbb{S}^n] \cong [\mathbb{S}_{\mathcal{P}}^{2n-1}; \mathbb{S}_{\mathcal{P}}^n \vee \mathbb{S}_{\mathcal{P}}^n],$$

the commutative diagram in (7) shows the proof. \square

Example 1. Let $\alpha : \mathbb{S}_{\mathcal{P}}^m \rightarrow \mathbb{S}_{\mathcal{P}}^{2n-1}$ be any homotopy class. We define a homotopy class Q in the homotopy group

$$\pi_m(X_{\mathcal{P}} \vee X_{\mathcal{P}}) = [\mathbb{S}^m; X_{\mathcal{P}} \vee X_{\mathcal{P}}] \cong [\mathbb{S}_{\mathcal{P}}^m; X_{\mathcal{P}} \vee X_{\mathcal{P}}]$$

as the composition

$$\mathbb{S}_{\mathcal{P}}^m \xrightarrow{\alpha} \mathbb{S}_{\mathcal{P}}^{2n-1} \xrightarrow{[i^1, i^2]_{\mathcal{P}}} \mathbb{S}_{\mathcal{P}}^n \vee \mathbb{S}_{\mathcal{P}}^n \xrightarrow{k_{\mathcal{P}}^1 \vee k_{\mathcal{P}}^1} X_{\mathcal{P}} \vee X_{\mathcal{P}}$$

of maps; that is,

$$Q = (k_{\mathcal{P}}^1 \vee k_{\mathcal{P}}^1)[i^1, i^2]_{\mathcal{P}}(\alpha) = [i_{\mathcal{P}}^1 k_{\mathcal{P}}^1, i_{\mathcal{P}}^2 k_{\mathcal{P}}^1]_{\alpha},$$

where $[i^1, i^2]_{\mathcal{P}}$ is the localization of the Whitehead product $[i^1, i^2]$. Then the map $\psi_Q : X_{\mathcal{P}} \rightarrow X_{\mathcal{P}} \vee X_{\mathcal{P}}$ defined by Definition 1 is a comultiplication on $X_{\mathcal{P}} \vee X_{\mathcal{P}}$ whose perturbation is Q .

Let $q_{\mathcal{P}}^2 : X_{\mathcal{P}} \rightarrow \mathbb{S}_{\mathcal{P}}^m$ be the projection. Then the cofibration

$$\mathbb{S}_{\mathcal{P}}^n \vee \mathbb{S}_{\mathcal{P}}^n \xrightarrow{k_{\mathcal{P}}^1 \vee k_{\mathcal{P}}^1} X_{\mathcal{P}} \vee X_{\mathcal{P}} \xrightarrow{q_{\mathcal{P}}^2 \vee q_{\mathcal{P}}^2} \mathbb{S}_{\mathcal{P}}^m \vee \mathbb{S}_{\mathcal{P}}^m,$$

asserts that if $Q \in \pi_m(X_{\mathcal{P}} \vee X_{\mathcal{P}})$ is any homotopy class satisfying

$$\pi_{\mathcal{P}}^1 Q = 0 = \pi_{\mathcal{P}}^2 Q,$$

then there exists a unique homotopy class $W \in \pi_m(\mathbb{S}_{\mathcal{P}}^n \vee \mathbb{S}_{\mathcal{P}}^n)$ such that $Q = (k_{\mathcal{P}}^1 \vee k_{\mathcal{P}}^1)W$ and

$$p_{\mathcal{P}}^1 W = 0 = p_{\mathcal{P}}^2 W,$$

where $p_{\mathcal{P}}^1, p_{\mathcal{P}}^2 : \mathbb{S}_{\mathcal{P}}^n \vee \mathbb{S}_{\mathcal{P}}^n \rightarrow \mathbb{S}_{\mathcal{P}}^n$ are the first and second projections induced by p^1 and p^2 , respectively.

To investigate the fundamental property of homotopy comultiplications on a wedge of localized spheres, we need to study the perturbations of all homotopy comultiplications. This naturally raises the following question: How can we construct the perturbation of a comultiplication on the wedge of localized spheres? The following lemma gives an answer to this query.

We denote the (generalized) basic Whitehead products of $i_{\mathcal{P}}^1$ and $i_{\mathcal{P}}^2$ by $w_1, w_2, \dots, w_s, \dots$ and the height of w_s is denoted by h_s for each $s = 1, 2, 3, \dots$. Then we have the following.

Lemma 6. A perturbation $Q : \mathbb{S}_{\mathcal{P}}^m \rightarrow X_{\mathcal{P}} \vee X_{\mathcal{P}}$ of any comultiplication $\psi_Q : X_{\mathcal{P}} \rightarrow X_{\mathcal{P}} \vee X_{\mathcal{P}}$ can be expressed uniquely as in Definition 1 as follows:

$$Q = (k_{\mathcal{P}}^1 \vee k_{\mathcal{P}}^1)_* \left(\sum_{s=3}^{\infty} w_s a_s \right),$$

where w_s is the s th (generalized) basic Whitehead product localized at \mathcal{P} and a_s is any homotopy class in the homotopy group

$$[\mathbb{S}_{\mathcal{P}}^m; \mathbb{S}_{\mathcal{P}}^{h_s+1}] \cong [\mathbb{S}^m; \mathbb{S}_{\mathcal{P}}^{h_s+1}]$$

for $s = 3, 4, 5, \dots$

Proof. By using Lemmas 3–5, we see that every comultiplication $\psi_Q : X_{\mathcal{P}} \rightarrow X_{\mathcal{P}} \vee X_{\mathcal{P}}$ has the type of Definition 1 satisfying

$$\pi_{\mathcal{P}}^1 Q = 0 = \pi_{\mathcal{P}}^2 Q.$$

Using Theorem 2, we obtain

$$\begin{aligned} \pi_m(\mathbb{S}_{\mathcal{P}}^n \vee \mathbb{S}_{\mathcal{P}}^n) &\cong \bigoplus_{s=1}^{\infty} \pi_m(\mathbb{S}_{\mathcal{P}}^{h_s+1}) \\ &= \pi_m(\mathbb{S}_{\mathcal{P}}^n) \oplus \pi_m(\mathbb{S}_{\mathcal{P}}^n) \oplus \bigoplus_{s=3}^{\infty} \pi_m(\mathbb{S}_{\mathcal{P}}^{h_s+1}). \end{aligned}$$

Moreover, every homotopy class Q in the homotopy group $[\mathbb{S}_{\mathcal{P}}^m; X_{\mathcal{P}} \vee X_{\mathcal{P}}]$ can be uniquely expressed as the following type:

$$(k_{\mathcal{P}}^1 \vee k_{\mathcal{P}}^1)_* (i_{\mathcal{P}}^1 a_1 + i_{\mathcal{P}}^2 a_2 + \sum_{s=3}^{\infty} w_s a_s) \in [\mathbb{S}_{\mathcal{P}}^m; X_{\mathcal{P}} \vee X_{\mathcal{P}}] \cong \pi_m(X_{\mathcal{P}} \vee X_{\mathcal{P}}).$$

Here,

- a_1 and a_2 are any homotopy classes in $[\mathbb{S}_{\mathcal{P}}^m; \mathbb{S}_{\mathcal{P}}^n] \cong \pi_m(\mathbb{S}_{\mathcal{P}}^n)$;
- a_s is any homotopy class in $[\mathbb{S}_{\mathcal{P}}^m; \mathbb{S}_{\mathcal{P}}^{h_s+1}] \cong \pi_m(\mathbb{S}_{\mathcal{P}}^{h_s+1})$ for $s = 3, 4, 5, \dots$; and
- w_s is the (generalized) basic Whitehead product for $s = 3, 4, 5, \dots$

We recall that all of the (generalized) basic Whitehead products w_s consists of at least one $i_{\mathcal{P}}^1$ and at least one $i_{\mathcal{P}}^2$, for example, $w_3 = [i_{\mathcal{P}}^1, i_{\mathcal{P}}^2]$, $w_4 = [i_{\mathcal{P}}^1, [i_{\mathcal{P}}^1, i_{\mathcal{P}}^2]]$, $w_5 = [i_{\mathcal{P}}^2, [i_{\mathcal{P}}^1, i_{\mathcal{P}}^2]]$ and so on. We now have

$$\begin{aligned}\pi_{\mathcal{P}}^1 \circ Q &= \pi_{\mathcal{P}}^1(k_{\mathcal{P}}^1 \vee k_{\mathcal{P}}^1)_* \left(\sum_{s=1}^{\infty} w_s a_s \right) \\ &= \pi_{\mathcal{P}}^1(k_{\mathcal{P}}^1 \vee k_{\mathcal{P}}^1)_* (i_{\mathcal{P}}^1 a_1 + i_{\mathcal{P}}^2 a_2 + \sum_{s=3}^{\infty} w_s a_s) \\ &= a_1 + 0_*(a_2) + 0 \\ &= a_1\end{aligned}\tag{8}$$

and

$$\begin{aligned}\pi_{\mathcal{P}}^2 \circ Q &= \pi_{\mathcal{P}}^2(k_{\mathcal{P}}^1 \vee k_{\mathcal{P}}^1)_* \left(\sum_{s=3}^{\infty} w_s a_s \right) \\ &= \pi_{\mathcal{P}}^2(k_{\mathcal{P}}^1 \vee k_{\mathcal{P}}^1)_* (i_{\mathcal{P}}^1 a_1 + i_{\mathcal{P}}^2 a_2 + \sum_{s=3}^{\infty} w_s a_s) \\ &= 0_*(a_1) + a_2 + 0 \\ &= a_2,\end{aligned}\tag{9}$$

where 0_* is the trivial homomorphism between homotopy groups and 0 is the trivial homotopy class. From the fact that $\pi_{\mathcal{P}}^1 Q = 0 = \pi_{\mathcal{P}}^2 Q$, the homotopy classes a_1 and a_2 in (8) and (9) should be zero, as required. \square

For the 2-fold wedge sum of localized spheres, we have

Theorem 3. The cardinality of the set of comultiplications on $\mathbb{S}_{\mathcal{P}}^n \vee \mathbb{S}_{\mathcal{P}}^m$ is

$$\prod_{s=3}^{\infty} |\pi_m(\mathbb{S}_{\mathcal{P}}^{h_s+1})|,$$

where $2 \leq n < m$ and \mathcal{P} is a collection of prime numbers.

Proof. By Lemma 6, we have the proof. \square

Let $|C(X_{\mathcal{P}})|$ be the cardinality of the set $C(X_{\mathcal{P}})$ of comultiplications on $X_{\mathcal{P}}$.

Example 2. If $\mathcal{P} = \{2, 3\}$, then we have

- (1) $|C(\mathbb{S}_{\mathcal{P}}^4 \vee \mathbb{S}_{\mathcal{P}}^7)| = \infty$;
- (2) $|C(\mathbb{S}_{\mathcal{P}}^4 \vee \mathbb{S}_{\mathcal{P}}^8)| = 2$; and
- (3) $|C(\mathbb{S}_{\mathcal{P}}^4 \vee \mathbb{S}_{\mathcal{P}}^9)| = 2$.

Indeed, by Theorem 3, we have

$$\begin{aligned}|C(\mathbb{S}_{\mathcal{P}}^4 \vee \mathbb{S}_{\mathcal{P}}^8)| &= |\pi_8(\mathbb{S}_{\mathcal{P}}^7)| \times |\pi_8(\mathbb{S}_{\mathcal{P}}^{10})| \times \cdots \\ &= |\mathbb{Z}_2 \otimes \mathbb{Z}_{\mathcal{P}}| \times |\{e\} \otimes \mathbb{Z}_{\mathcal{P}}| \times \cdots \\ &= 2 \times 1 \times 1 \times \cdots \times 1 \times \cdots \\ &= 2\end{aligned},$$

where $\{e\}$ is the trivial group, and similarly for the other cases.

3.3. The k -Fold Wedge Sum of (Localized) Spheres

We now consider the k -fold wedge sum of 1-connected spheres $\{\mathbb{S}^{n_i} | i = 1, 2, \dots, k\}$ and their localized spheres $\{\mathbb{S}_{\mathcal{P}}^{n_i} | i = 1, 2, \dots, k\}$ as a generalization of the statements above; that is,

$$X = \mathbb{S}^{n_1} \vee \mathbb{S}^{n_2} \vee \dots \vee \mathbb{S}^{n_k}$$

and

$$X_{\mathcal{P}} = \mathbb{S}_{\mathcal{P}}^{n_1} \vee \mathbb{S}_{\mathcal{P}}^{n_2} \vee \dots \vee \mathbb{S}_{\mathcal{P}}^{n_k}$$

for $2 \leq n_1 < n_2 < \dots < n_k$.

Notation. We make use of the following notations in the rest of this paper.

- $X := \bigvee_{i=1}^k \mathbb{S}^{n_i}$ for $2 \leq n_1 < n_2 < \dots < n_k$.
- $\alpha_i^1, \alpha_i^2 : \mathbb{S}^{n_i} \rightarrow X \vee X$ are the i th and $(k+i)$ th inclusions, respectively, for $i = 1, 2, \dots, k-1$.
- $\zeta^i : \mathbb{S}^{n_i} \rightarrow X$ is the i th inclusion for $i = 1, 2, \dots, k$.
- $\iota^1, \iota^2 : X \rightarrow X \vee X$ are the first and second inclusions, respectively.
- $\pi^1, \pi^2 : X \vee X \rightarrow X$ are the first and second projections, respectively.
- W_j^+ is the set of all (generalized) basic Whitehead products localized at \mathcal{P} containing the homotopy classes from α_i^1 for $i = 1, 2, \dots, j-1$ and $j = 2, 3, \dots, k$.
- $W_j^\#$ is the set of all (generalized) basic Whitehead products localized at \mathcal{P} containing the homotopy classes from α_i^2 for $i = 1, 2, \dots, j-1$ and $j = 2, 3, \dots, k$.
- W_j^* is the set of all (generalized) basic Whitehead products localized at \mathcal{P} containing as a factor at least one of the homotopy classes from α_i^1 and at least one of the homotopy classes from α_i^2 for $i = 1, 2, \dots, j-1$ and $j = 2, 3, \dots, k$.

We note that if $k = 2$, $n_1 = n$ and $n_2 = m$; that is, $X = \mathbb{S}^n \vee \mathbb{S}^m$, then we have

1. $\zeta^1 = k^1 : \mathbb{S}^n \hookrightarrow \mathbb{S}^n \vee \mathbb{S}^m$;
2. $\zeta^2 = k^2 : \mathbb{S}^m \hookrightarrow \mathbb{S}^n \vee \mathbb{S}^m$;
3. $\alpha_1^1 = (k^1 \vee k^1) \circ i^1 : \mathbb{S}^n \xrightarrow{i^1} \mathbb{S}^n \vee \mathbb{S}^n \xrightarrow{k^1 \vee k^1} X \vee X$; and
4. $\alpha_1^2 = (k^1 \vee k^1) \circ i^2 : \mathbb{S}^n \xrightarrow{i^2} \mathbb{S}^n \vee \mathbb{S}^n \xrightarrow{k^1 \vee k^1} X \vee X$

in the notations above.

To develop the basic Whitehead products in the k -fold wedge sum of spheres localized at \mathcal{P} , we order the basic Whitehead products of weight 1 as follows:

- $\alpha_i^1 < \alpha_j^2$ for $i, j = 1, 2, \dots, k-1$; and
- $\iota^1 \zeta^i < \iota^2 \zeta^j$ for $i, j = 1, 2, \dots, k$.

Definition 2. Let $Q_j : \mathbb{S}_{\mathcal{P}}^{n_j} \rightarrow X_{\mathcal{P}} \vee X_{\mathcal{P}}$ be any element of $[\mathbb{S}_{\mathcal{P}}^{n_j}; X_{\mathcal{P}} \vee X_{\mathcal{P}}]$ satisfying

$$\pi_{\mathcal{P}}^1 Q_j = 0 = \pi_{\mathcal{P}}^2 Q_j$$

for each $j = 2, 3, \dots, k$, where $\pi_{\mathcal{P}}^1, \pi_{\mathcal{P}}^2 : X_{\mathcal{P}} \vee X_{\mathcal{P}} \rightarrow X_{\mathcal{P}}$ are the first and second projections, respectively. We define a comultiplication

$$\psi = \psi_{(Q_2, Q_3, \dots, Q_k)} : X_{\mathcal{P}} \rightarrow X_{\mathcal{P}} \vee X_{\mathcal{P}}$$

by

$$\begin{cases} \psi_Q \zeta_P^1 &= \iota_P^1 \zeta_P^1 + \iota_P^2 \zeta_P^1 \\ \psi_Q \zeta_P^2 &= \iota_P^1 \zeta_P^2 + \iota_P^2 \zeta_P^2 + Q_2 \\ \vdots & \\ \psi_Q \zeta_P^j &= \iota_P^1 \zeta_P^j + \iota_P^2 \zeta_P^j + Q_j \\ \vdots & \\ \psi_Q \zeta_P^k &= \iota_P^1 \zeta_P^k + \iota_P^2 \zeta_P^k + Q_k. \end{cases} \quad (10)$$

Here, we call $Q_j \in [\mathbb{S}_{\mathcal{P}}^{n_j}; X_{\mathcal{P}} \vee X_{\mathcal{P}}]$ the j th perturbation of $\psi = \psi_{(Q_2, Q_3, \dots, Q_k)}$ for $j = 2, 3, \dots, k$.

We need to investigate the j th perturbation $Q_j \in [\mathbb{S}_{\mathcal{P}}^{n_j}; X_{\mathcal{P}} \vee X_{\mathcal{P}}]$ of $\psi_{(Q_2, Q_3, \dots, Q_k)} : X_{\mathcal{P}} \rightarrow X_{\mathcal{P}} \vee X_{\mathcal{P}}$ as a generalization of Lemma 6 more concretely.

Lemma 7. The j th perturbation $Q_j \in [\mathbb{S}_{\mathcal{P}}^{n_j}; X_{\mathcal{P}} \vee X_{\mathcal{P}}]$ of any comultiplication $\psi : X_{\mathcal{P}} \rightarrow X_{\mathcal{P}} \vee X_{\mathcal{P}}$ can be uniquely expressed as in Definition 2 by

$$Q_j = \sum_{w_j \in W_j^*} w_j \circ c_j. \quad (11)$$

for $j = 2, 3, \dots, k$. Here,

- $w_j : \mathbb{S}_{\mathcal{P}}^{h_{w_j}+1} \rightarrow X_{\mathcal{P}} \vee X_{\mathcal{P}}$ is the j th (generalized) basic Whitehead product localized at \mathcal{P} ; and
- $c_j : \mathbb{S}_{\mathcal{P}}^{n_j} \rightarrow \mathbb{S}_{\mathcal{P}}^{h_{w_j}+1}$ is any homotopy class in the homotopy group $[\mathbb{S}_{\mathcal{P}}^{n_j}; \mathbb{S}_{\mathcal{P}}^{h_{w_j}+1}] \cong [\mathbb{S}_{\mathcal{P}}^{n_j}; \mathbb{S}_{\mathcal{P}}^{h_{w_j}+1}]$, where h_{w_j} is the height of the (generalized) basic Whitehead product $w_j \in W_j^*$ for each $j = 2, 3, \dots, k$.

Proof. The localized version of the Hilton's formula says that the j th perturbation Q_j of $\psi : X_{\mathcal{P}} \rightarrow X_{\mathcal{P}} \vee X_{\mathcal{P}}$ can be uniquely written by

$$Q_j = \sum_{i=1}^{j-1} (\alpha_i^1)_{\mathcal{P}} \circ \beta_j^i + \sum_{i=1}^{j-1} (\alpha_i^2)_{\mathcal{P}} \circ \gamma_j^i + \sum_{u_j \in W_j^+} u_j \circ a_j + \sum_{v_j \in W_j^\#} v_j \circ b_j + \sum_{w_j \in W_j^*} w_j \circ c_j \quad (12)$$

Here,

- β_j^i and γ_j^i are any homotopy classes in $[\mathbb{S}_{\mathcal{P}}^{n_j}; \mathbb{S}_{\mathcal{P}}^{n_i}] \cong \pi_{n_j}(\mathbb{S}_{\mathcal{P}}^{n_i})$ for $i = 1, 2, \dots, j-1$;
- a_j is any homotopy class in $[\mathbb{S}_{\mathcal{P}}^{n_j}; \mathbb{S}_{\mathcal{P}}^{h_{u_j}+1}] \cong \pi_{n_j}(\mathbb{S}_{\mathcal{P}}^{h_{u_j}+1})$ for $j = 2, 3, \dots, k$;
- b_j is any homotopy class in $[\mathbb{S}_{\mathcal{P}}^{n_j}; \mathbb{S}_{\mathcal{P}}^{h_{v_j}+1}] \cong \pi_{n_j}(\mathbb{S}_{\mathcal{P}}^{h_{v_j}+1})$ for $j = 2, 3, \dots, k$;
- c_j is any homotopy class in $[\mathbb{S}_{\mathcal{P}}^{n_j}; \mathbb{S}_{\mathcal{P}}^{h_{w_j}+1}] \cong \pi_{n_j}(\mathbb{S}_{\mathcal{P}}^{h_{w_j}+1})$ for $j = 2, 3, \dots, k$; and
- u_j , v_j and w_j are the (generalized) basic Whitehead product localized at \mathcal{P} for $j = 2, 3, \dots, k$.

We show that the homotopy classes β_j^i , γ_j^i , a_j and b_j are all zero. By taking the first and second projections $\pi_{\mathcal{P}}^1, \pi_{\mathcal{P}}^2 : X_{\mathcal{P}} \vee X_{\mathcal{P}} \rightarrow X_{\mathcal{P}}$ to the perturbation Q_j of any comultiplication $\psi : X_{\mathcal{P}} \rightarrow X_{\mathcal{P}} \vee X_{\mathcal{P}}$, we have

$$\pi_{\mathcal{P}}^1 \circ Q_j = 0 = \pi_{\mathcal{P}}^2 \circ Q_j \quad (13)$$

for each $j = 2, 3, \dots, k$; that is,

$$\begin{aligned} 0 &= \pi_{\mathcal{P}}^1 \circ Q_j \\ &= \sum_{i=1}^{j-1} \pi_{\mathcal{P}}^1 \circ (\alpha_i^1)_{\mathcal{P}} \circ \beta_j^i + \sum_{i=1}^{j-1} \pi_{\mathcal{P}}^1 \circ (\alpha_i^2)_{\mathcal{P}} \circ \gamma_j^i + \sum_{u_j \in W_j^{\dagger}} \pi_{\mathcal{P}}^1 \circ u_j \circ a_j + \sum_{v_j \in W_j^{\dagger}} \pi_{\mathcal{P}}^1 \circ v_j \circ b_j \\ &\quad + \sum_{w_j \in W_j^*} \pi_{\mathcal{P}}^1 \circ w_j \circ c_j \end{aligned} \quad (14)$$

and similarly, for the second projection $\pi_{\mathcal{P}}^2$; see below. Since the map

$$\pi_{\mathcal{P}}^1 \circ (\alpha_i^1)_{\mathcal{P}} : \mathbb{S}_{\mathcal{P}}^{n_j} \xrightarrow{(\alpha_i^1)_{\mathcal{P}}} X_{\mathcal{P}} \vee X_{\mathcal{P}} \xrightarrow{\pi_{\mathcal{P}}^1} X_{\mathcal{P}} \quad (15)$$

in (14) is the inclusion map, the localized version of the Hilton's formula asserts that the homotopy class β_j^i should be zero for each $i = 1, 2, \dots, j-1$ and $j = 2, 3, \dots, k$. The choice of the (generalized) basic Whitehead products localized at \mathcal{P} in W_j^{\dagger} says that all the homotopy classes a_j should be zero. We note that the maps

$$\pi_{\mathcal{P}}^1 \circ (\alpha_i^2)_{\mathcal{P}} : \mathbb{S}_{\mathcal{P}}^{n_j} \xrightarrow{(\alpha_i^2)_{\mathcal{P}}} X_{\mathcal{P}} \vee X_{\mathcal{P}} \xrightarrow{\pi_{\mathcal{P}}^1} X_{\mathcal{P}}, \quad (16)$$

$$\pi_{\mathcal{P}}^1 \circ v_j : \mathbb{S}_{\mathcal{P}}^{h_{v_j}+1} \xrightarrow{v_j} X_{\mathcal{P}} \vee X_{\mathcal{P}} \xrightarrow{\pi_{\mathcal{P}}^1} X_{\mathcal{P}} \quad (17)$$

and

$$\pi_{\mathcal{P}}^1 \circ w_j : \mathbb{S}_{\mathcal{P}}^{h_{w_j}+1} \xrightarrow{w_j} X_{\mathcal{P}} \vee X_{\mathcal{P}} \xrightarrow{\pi_{\mathcal{P}}^1} X_{\mathcal{P}} \quad (18)$$

are all zero by the choice of $(\alpha_i^2)_{\mathcal{P}}$, v_j and w_j for each $i = 1, 2, \dots, j-1$ and $j = 2, 3, \dots, k$.

Similarly, by taking the second projection $\pi_{\mathcal{P}}^2 : X_{\mathcal{P}} \vee X_{\mathcal{P}} \rightarrow X_{\mathcal{P}}$ to the perturbation Q_j again, we see that the map

$$\pi_{\mathcal{P}}^2 \circ (\alpha_i^2)_{\mathcal{P}} : \mathbb{S}_{\mathcal{P}}^{n_j} \xrightarrow{(\alpha_i^2)_{\mathcal{P}}} X_{\mathcal{P}} \vee X_{\mathcal{P}} \xrightarrow{\pi_{\mathcal{P}}^2} X_{\mathcal{P}} \quad (19)$$

from (12) is the inclusion map so that $\gamma_j^i = 0$ for each $i = 1, 2, \dots, j-1$ and $j = 2, 3, \dots, k$. The choice of the (generalized) basic Whitehead products localized at \mathcal{P} in W_j^{\dagger} says that all the homotopy classes b_j should be zero. Moreover, the localized version of the Hilton's formula asserts that the homotopy class $\pi_{\mathcal{P}}^2 \circ (\alpha_i^1)_{\mathcal{P}}$, $\pi_{\mathcal{P}}^2 \circ u_j$, and $\pi_{\mathcal{P}}^2 \circ w_j$ are all zero by the choice of $(\alpha_i^1)_{\mathcal{P}}$, u_j and w_j for each $i = 1, 2, \dots, j-1$ and $j = 2, 3, \dots, k$, as required. \square

For the k -fold wedge sum of localized spheres, we have the following.

Theorem 4. The cardinality of the set of comultiplications on $X_{\mathcal{P}} := \mathbb{S}_{\mathcal{P}}^{n_1} \vee \mathbb{S}_{\mathcal{P}}^{n_2} \vee \dots \vee \mathbb{S}_{\mathcal{P}}^{n_k}$ is

$$\prod_{w_2 \in W_2^*} |\pi_{n_2}(\mathbb{S}_{\mathcal{P}}^{h_{w_2}+1})| \times \prod_{w_3 \in W_3^*} |\pi_{n_3}(\mathbb{S}_{\mathcal{P}}^{h_{w_3}+1})| \times \dots \times \prod_{w_k \in W_k^*} |\pi_{n_k}(\mathbb{S}_{\mathcal{P}}^{h_{w_k}+1})|,$$

where $2 \leq n_1 < n_2 < \dots < n_k$ and \mathcal{P} is a collection of prime numbers.

Proof. By Lemma 7, we complete the proof. \square

Example 3. If $\mathcal{P} = \{2, 3, 5\}$, then we have

- (1) $|C(\mathbb{S}_{\mathcal{P}}^8 \vee \mathbb{S}_{\mathcal{P}}^{12} \vee \mathbb{S}_{\mathcal{P}}^{26})| = \infty$; and
- (2) $|C(\mathbb{S}_{\mathcal{P}}^8 \vee \mathbb{S}_{\mathcal{P}}^{12} \vee \mathbb{S}_{\mathcal{P}}^{27})| = 512$.

3.4. The (Localized) Moore Spaces

A comultiplication $\varphi : Y \rightarrow Y \vee Y$ is said to be *associative* if

$$(\varphi \vee 1) \circ \varphi \simeq (1 \vee \varphi) \circ \varphi : Y \rightarrow Y \vee Y \vee Y,$$

where $1 : Y \rightarrow Y$ is the identity map. If φ is associative, we call (Y, φ) an *associative co-H-space* or *co-H-group*. It is well known that all Moore spaces $M(G, n)$, $n \geq 2$, are co-H-groups, where G is an abelian group.

In general, the cardinality of the set $C(Y) \subseteq [Y; Y \vee Y]$ of all homotopy classes of comultiplications on Y is extremely difficult to detect. As a particular case, it was shown in [27] that the set of comultiplications $C(M(G, 2))$ is in one-one correspondence with the group $\text{Ext}(G, G \otimes G)$, and that if $n \geq 3$, then $C(M(G, n))$ has one element as the standard comultiplication, where G is an abelian group. We note that this result is also valid for localizations.

4. Main Result and Its Illustration

Let \mathcal{P} be a collection of prime numbers which may be empty and let $X_{\mathcal{P}}$ be the localization of the k -fold wedge sum of spheres

$$X := \bigvee_{i=1}^k \mathbb{S}^{n_i} = \mathbb{S}^{n_1} \vee \mathbb{S}^{n_2} \vee \dots \vee \mathbb{S}^{n_k}$$

for $2 \leq n_1 < n_2 < \dots < n_k$ or a Moore space $M(G, n)$, where G is an abelian group and $n \geq 2$. Then, by using the results in Sections 2 and 3, we have the following theorem.

Theorem 5. Let $|C(X)|$ and $|C(X_{\mathcal{P}})|$ be the cardinalities of the sets of all the homotopy comultiplications on X and $X_{\mathcal{P}}$, respectively.

- (1) If $|C(X)|$ is finite, then $|C(X)| \geq |C(X_{\mathcal{P}})|$.
- (2) If $|C(X)|$ is infinite, then $|C(X)| = |C(X_{\mathcal{P}})|$.

Proof. We prove the theorem in the case of the k -fold wedge sum of spheres and its localization. We first note that if the homotopy group $\pi_{n_j}(\mathbb{S}^{h_{w_j}+1})$ contains a q -torsion subgroup T_q with $(p, q) = 1$ for $w_j \in W_j^*$, $j = 2, 3, \dots, k$, then all of the q -torsion subgroups of $\pi_{n_j}(\mathbb{S}^{h_{w_j}+1})$ could entirely vanish in $\pi_{n_j}(\mathbb{S}_{\mathcal{P}}^{h_{w_j}+1})$ for $w_j \in W_j^*$, $j = 2, 3, \dots, k$, where q is a prime number with $(p, q) = 1$ for all $p \in \mathcal{P}$.

(1) If $|C(X)|$ is finite, then all of the homotopy groups $\pi_{n_j}(\mathbb{S}^{h_{w_j}+1})$, $j = 2, 3, \dots, k$ must be the torsion abelian groups T_{w_j} , $j = 2, 3, \dots, k$, where T_{w_j} is the torsion subgroup of

$$\prod_{j=2}^k \prod_{w_j \in W_j^*} \pi_{n_j}(\mathbb{S}^{h_{w_j}+1}) = \prod_{j=2}^k \prod_{w_j \in W_j^*} T_{w_j}. \quad (20)$$

We also note that

$$T_{w_j} \cong \mathbb{Z}_{p_1^{i_1}} \oplus \mathbb{Z}_{p_2^{i_2}} \oplus \dots \oplus \mathbb{Z}_{p_l^{i_l}}, \quad (21)$$

where p_1, p_2, \dots, p_l are (not necessarily distinct) primes and j_1, j_2, \dots, j_l are (not necessarily distinct) positive integers. We thus have

$$\begin{aligned} |C(X)| &= \prod_{j=2}^k \prod_{w_j \in W_j^*} |\pi_{n_j}(\mathbb{S}^{h_{w_j}+1})| \quad (\text{by Lemma 1}) \\ &= \prod_{j=2}^k \prod_{w_j \in W_j^*} |T_{w_j}| \\ &\geq \prod_{j=2}^k \prod_{w_j \in W_j^*} |T_{w_j} \otimes \mathbb{Z}_{\mathcal{P}}| \\ &= \prod_{j=2}^k \prod_{w_j \in W_j^*} |\pi_{n_j}(\mathbb{S}_{\mathcal{P}}^{h_{w_j}+1})| \quad (\text{by Theorem 4}) \\ &= |C(X_{\mathcal{P}})|. \end{aligned} \tag{22}$$

We note that equality of the inequality in (22) holds only in the case that the collection \mathcal{P} of primes contains all of the prime numbers expressed in the torsion subgroups T_{w_j} , $j = 2, 3, \dots, k$.

(2) If $|C(X)|$ is infinite, then at least one of the homotopy groups $\pi_{n_j}(\mathbb{S}^{h_{w_j}+1})$, $j = 2, 3, \dots, k$ contains an infinite cyclic group, say in dimension $n_s = h_{w_s} + 1$ or $n_s = 2(h_{w_s} + 1) - 1$, where $h_{w_s} + 1$ is an even integer for some $s = 2, 3, \dots, k$. Moreover, it can be seen that the numbers of nonzero homotopy groups $\pi_{n_j}(\mathbb{S}^{h_{w_j}+1})$ or $\pi_{n_j}(\mathbb{S}_{\mathcal{P}}^{h_{w_j}+1})$, $j = 2, 3, \dots, k$ are always finite because the height h_{w_j} goes to infinity.

Let F_{w_s} be a free abelian group of rank 1 and let T_{w_s} be a torsion subgroup of $\pi_{n_s}(\mathbb{S}^{h_{w_s}+1})$ for some $s = 2, 3, \dots, k$. Then we have

$$\begin{aligned} |C(X)| &= \prod_{j=2}^k \prod_{w_j \in W_j^*} |\pi_{n_j}(\mathbb{S}^{h_{w_j}+1})| \\ &= |F_{w_s} \oplus T_{w_s}| \times \prod_{j \neq s} \prod_{w_j \in W_j^*} |\pi_{n_j}(\mathbb{S}^{h_{w_j}+1})| \\ &\approx |\mathbb{Z}|. \end{aligned} \tag{23}$$

Here, \approx means the same cardinality as sets and

$$F_{w_s} \oplus T_{w_s} \cong \begin{cases} \mathbb{Z} & \text{for } n_s = h_{w_s} + 1 \\ \mathbb{Z} \oplus T_{w_s} & \text{for } n_s = 2(h_{w_s} + 1) - 1 \text{ and } h_{w_s} + 1 \text{ even.} \end{cases}$$

By Theorem 4 or by tensoring $\mathbb{Z}_{\mathcal{P}}$ on the homotopy groups described in (23), we see that if $n_s = h_{w_s} + 1$ for some s , then

$$\begin{aligned} |C(X_{\mathcal{P}})| &= \prod_{j=2}^k \prod_{w_j \in W_j^*} |\pi_{n_j}(\mathbb{S}_{\mathcal{P}}^{h_{w_j}+1})| \\ &= |(F_{w_s} \oplus T_{w_s}) \otimes \mathbb{Z}_{\mathcal{P}}| \times \prod_{j \neq s} \prod_{w_j \in W_j^*} |\pi_{n_j}(\mathbb{S}_{\mathcal{P}}^{h_{w_j}+1})| \\ &= |\mathbb{Z} \otimes \mathbb{Z}_{\mathcal{P}}| \times \prod_{j \neq s} \prod_{w_j \in W_j^*} |\pi_{n_j}(\mathbb{S}_{\mathcal{P}}^{h_{w_j}+1})| \\ &\approx |\mathbb{Z}_{\mathcal{P}}| \end{aligned} \tag{24}$$

and if $h_{w_s} + 1$ is even and $n_s = 2(h_{w_s} + 1) - 1$ for some s , then

$$\begin{aligned}
 |C(X_{\mathcal{P}})| &= \prod_{j=2}^k \prod_{w_j \in W_j^*} |\pi_{n_j}(\mathbb{S}_{\mathcal{P}}^{h_{w_j}+1})| \\
 &= |(F_{w_s} \oplus T_{w_s}) \otimes \mathbb{Z}_{\mathcal{P}}| \times \prod_{j \neq s} \prod_{w_j \in W_j^*} |\pi_{n_j}(\mathbb{S}_{\mathcal{P}}^{h_{w_j}+1})| \\
 &= |(\mathbb{Z} \oplus F_{w_s}) \otimes \mathbb{Z}_{\mathcal{P}}| \times \prod_{j \neq s} \prod_{w_j \in W_j^*} |\pi_{n_j}(\mathbb{S}_{\mathcal{P}}^{h_{w_j}+1})| \\
 &\approx |\mathbb{Z}_{\mathcal{P}}|
 \end{aligned} \tag{25}$$

We note that some finite groups in (24) and (25) will vanish for some torsion groups, each of whose orders do not belong to the collection \mathcal{P} of prime numbers. Moreover, because \mathcal{P} is a collection of primes and the free abelian group \mathbb{Z} is a subgroup of $\mathbb{Z}_{\mathcal{P}}$ as a subring of the ring of rational numbers \mathbb{Q} ; that is,

$$\mathbb{Z} \subseteq \mathbb{Z}_{\mathcal{P}} \subseteq \mathbb{Q},$$

their cardinalities are the same as countably infinite sets.

The proof in the case of a Moore space $M(G, n)$ and its localization runs through the similar way to the case of the k -fold wedge sum of spheres and its localization. \square

Remark 1. We note that if the collection \mathcal{P} of prime numbers is the empty collection, then the above statement is also true as a rationalization $X_{\mathbb{Q}}$ of a 1-connected CW-complex $X := \bigvee_{i=1}^k \mathbb{S}^{n_i}$ for $2 \leq n_1 < n_2 < \dots < n_k$.

We finally give an example to illustrate our results.

Example 4. The finite (resp. infinite) cases are described in Tables 1–3 (resp. Tables 4 and 5). In the tables, ϕ denotes the empty collection of primes and \aleph_0 is aleph-zero.

Table 1. The finite case.

$X = \mathbb{S}^n \vee \mathbb{S}^m$	\mathcal{P}	$ C(X) $	$ C(X_{\mathcal{P}}) $
$\mathbb{S}^3 \vee \mathbb{S}^6$	$\{2, 5\}$	2	2
$\mathbb{S}^3 \vee \mathbb{S}^6$	$\{3, 5\}$	2	1
$\mathbb{S}^8 \vee \mathbb{S}^{26}$	$\{11, 13\}$	504	1
$\mathbb{S}^3 \vee \mathbb{S}^6$	ϕ	2	1

Table 2. The finite case for $k \geq 3$.

$X = \bigvee_{i=1}^k \mathbb{S}^{n_i}$	\mathcal{P}	$ C(X) $	$ C(X_{\mathcal{P}}) $
$\mathbb{S}^4 \vee \mathbb{S}^6 \vee \mathbb{S}^8$	$\{3, 5\}$	2	1
$\mathbb{S}^5 \vee \mathbb{S}^6 \vee \mathbb{S}^7$	$\{3, 5\}$	1	1
$\mathbb{S}^5 \vee \mathbb{S}^7 \vee \mathbb{S}^{10}$	$\{3, 5\}$	2	1
$\mathbb{S}^7 \vee \mathbb{S}^9 \vee \mathbb{S}^{11} \vee \mathbb{S}^{22}$	$\{3, 5\}$	19,568,944,742,400	1

Table 3. The finite case: Moore spaces.

$Y = M(G, n)$	\mathcal{P}	$ C(Y) $	$ C(Y_{\mathcal{P}}) $
$M(\mathbb{Z}, 2)$	$\{2, 5\}$	1	1
$M(\mathbb{Z}_3, 2)$	$\{2, 5\}$	3	1
$M(\mathbb{Q}, 2)$	$\{3, 5\}$	1	1
$M(G, n), n \geq 3$	ϕ	1	1

Table 4. The infinite case.

$X = \mathbb{S}^n \vee \mathbb{S}^m$	\mathcal{P}	$ C(X) $	$ C(X_{\mathcal{P}}) $
$\mathbb{S}^4 \vee \mathbb{S}^7$	$\{2, 5\}$	$ \mathbb{Z} = \aleph_0$	$ \mathbb{Z}_{\mathcal{P}} = \aleph_0$
$\mathbb{S}^5 \vee \mathbb{S}^9$	$\{2, 5\}$	$ \mathbb{Z} = \aleph_0$	$ \mathbb{Z}_{\mathcal{P}} = \aleph_0$
$\mathbb{S}^{10} \vee \mathbb{S}^{19}$	$\{2, 5\}$	$ \mathbb{Z} = \aleph_0$	$ \mathbb{Z}_{\mathcal{P}} = \aleph_0$
$\mathbb{S}^4 \vee \mathbb{S}^7$	\emptyset	$ \mathbb{Z} = \aleph_0$	$ \mathbb{Q} = \aleph_0$

Table 5. The infinite case for $k \geq 3$.

$X = \bigvee_{i=1}^k \mathbb{S}^{n_i}$	\mathcal{P}	$ C(X) $	$ C(X_{\mathcal{P}}) $
$\mathbb{S}^3 \vee \mathbb{S}^4 \vee \mathbb{S}^5$	$\{2, 5\}$	$ \mathbb{Z} = \aleph_0$	$ \mathbb{Z}_{\mathcal{P}} = \aleph_0$
$\mathbb{S}^4 \vee \mathbb{S}^6 \vee \mathbb{S}^{12}$	$\{2, 5\}$	$ \mathbb{Z} = \aleph_0$	$ \mathbb{Z}_{\mathcal{P}} = \aleph_0$
$\mathbb{S}^7 \vee \mathbb{S}^{12} \vee \mathbb{S}^{24}$	$\{2, 5\}$	$ \mathbb{Z} = \aleph_0$	$ \mathbb{Z}_{\mathcal{P}} = \aleph_0$
$\mathbb{S}^6 \vee \mathbb{S}^8 \vee \mathbb{S}^{20}$	\emptyset	$ \mathbb{Z} = \aleph_0$	$ \mathbb{Q} = \aleph_0$

Proof. To prove the example, we use the previous results in Sections 2 and 3, and prove the first two cases to illustrate the method. We note that $\pi_6(\mathbb{S}^5)$ is isomorphic to the cyclic group of order 2; that is, $\pi_6(\mathbb{S}^5) \cong \mathbb{Z}_2$; see [28]. If $X = \mathbb{S}^3 \vee \mathbb{S}^6$ and $\mathcal{P} = \{2, 5\}$, then we have

$$\begin{aligned}
 |C(X)| &= |\pi_6(\mathbb{S}^5)| \times |\pi_6(\mathbb{S}^7)| \times |\pi_6(\mathbb{S}^7)| \\
 &\quad \times |\pi_6(\mathbb{S}^9)| \times |\pi_6(\mathbb{S}^9)| \times |\pi_6(\mathbb{S}^9)| \times |\pi_6(\mathbb{S}^{11})| \times \dots \\
 &= 2 \times 1 \times 1 \times 1 \times 1 \times 1 \times 1 \times \dots \\
 &= 2
 \end{aligned}$$

and

$$\begin{aligned}
 |C(X_{\mathcal{P}})| &= |\pi_6(\mathbb{S}^5) \otimes \mathbb{Z}_{\mathcal{P}}| \times |\pi_6(\mathbb{S}^7) \otimes \mathbb{Z}_{\mathcal{P}}| \times |\pi_6(\mathbb{S}^7) \otimes \mathbb{Z}_{\mathcal{P}}| \\
 &\quad \times |\pi_6(\mathbb{S}^9) \otimes \mathbb{Z}_{\mathcal{P}}| \times |\pi_6(\mathbb{S}^9) \otimes \mathbb{Z}_{\mathcal{P}}| \times |\pi_6(\mathbb{S}^9) \otimes \mathbb{Z}_{\mathcal{P}}| \times |\pi_6(\mathbb{S}^{11}) \otimes \mathbb{Z}_{\mathcal{P}}| \times \dots \\
 &= |\mathbb{Z}_2 \otimes \mathbb{Z}_{\mathcal{P}}| \times |\{e\} \otimes \mathbb{Z}_{\mathcal{P}}| \times |\{e\} \otimes \mathbb{Z}_{\mathcal{P}}| \\
 &\quad \times |\{e\} \otimes \mathbb{Z}_{\mathcal{P}}| \times |\{e\} \otimes \mathbb{Z}_{\mathcal{P}}| \times |\{e\} \otimes \mathbb{Z}_{\mathcal{P}}| \times |\{e\} \otimes \mathbb{Z}_{\mathcal{P}}| \times \dots \\
 &= 2 \times 1 \times 1 \times 1 \times 1 \times 1 \times 1 \times \dots \\
 &= 2
 \end{aligned}$$

where $\{e\}$ is the trivial group. If $X = \mathbb{S}^3 \vee \mathbb{S}^6$ and $\mathcal{P} = \{3, 5\}$, then we get

$$\begin{aligned}
 |C(X_{\mathcal{P}})| &= |\pi_6(\mathbb{S}^5) \otimes \mathbb{Z}_{\mathcal{P}}| \times |\pi_6(\mathbb{S}^7) \otimes \mathbb{Z}_{\mathcal{P}}| \times |\pi_6(\mathbb{S}^7) \otimes \mathbb{Z}_{\mathcal{P}}| \\
 &\quad \times |\pi_6(\mathbb{S}^9) \otimes \mathbb{Z}_{\mathcal{P}}| \times |\pi_6(\mathbb{S}^9) \otimes \mathbb{Z}_{\mathcal{P}}| \times |\pi_6(\mathbb{S}^9) \otimes \mathbb{Z}_{\mathcal{P}}| \times |\pi_6(\mathbb{S}^{11}) \otimes \mathbb{Z}_{\mathcal{P}}| \times \dots \\
 &= |\mathbb{Z}_2 \otimes \mathbb{Z}_{\mathcal{P}}| \times |\{e\} \otimes \mathbb{Z}_{\mathcal{P}}| \times |\{e\} \otimes \mathbb{Z}_{\mathcal{P}}| \\
 &\quad \times |\{e\} \otimes \mathbb{Z}_{\mathcal{P}}| \times |\{e\} \otimes \mathbb{Z}_{\mathcal{P}}| \times |\{e\} \otimes \mathbb{Z}_{\mathcal{P}}| \times |\{e\} \otimes \mathbb{Z}_{\mathcal{P}}| \times \dots \\
 &= 1 \times 1 \times 1 \times 1 \times 1 \times 1 \times 1 \times \dots \\
 &= 1
 \end{aligned}$$

Therefore, if $X = \mathbb{S}^3 \vee \mathbb{S}^6$, then we have

$$\begin{cases} |C(X)| = 2 = |C(X_{\mathcal{P}})| & \text{for } \mathcal{P} = \{2, 5\} \\ |C(X)| = 2 > 1 = |C(X_{\mathcal{P}})| & \text{for } \mathcal{P} = \{3, 5\} \end{cases} ,$$

as required. \square

5. Conclusions and Further Prospects

Co-H-spaces, also called spaces with a comultiplication, and localization theory play a pivotal role in (equivariant) homotopy theory. In general, it turns out that the computation of the cardinality of the

set of homotopy comultiplications is very difficult. In this paper, we have investigated the inequalities and relationships between the cardinalities of the sets of homotopy comultiplications on the wedge of localized spheres and localized Moore spaces.

We do hope that our methods can be used to study homotopy comultiplications on the wedge sum of any number localized spheres and localized Moore spaces. We also hope that the results in this paper can be applied to the notions of algebra comultiplications on the (localized) algebraic objects such as (free) Lie algebras by considering the coproduct of Lie algebras or cohomology modules over \mathbb{Z}_p .

Funding: This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korean government (MSIT) (No. 2018R1A2B6004407).

Acknowledgments: The author is grateful to the anonymous referees for a careful reading and many helpful suggestions that improved the quality of this paper.

Conflicts of Interest: The author declares no conflict of interest.

References

1. Arkowitz, M. Co-H-spaces. In *Handbook of Algebraic Topology*; North-Holland: New York, NY, USA, 1995; pp. 1143–1173.
2. Arkowitz, M. Binary operations for homotopy groups with coefficients. *Sci. China Math.* **2018**, *61*, 1543–1552. [[CrossRef](#)]
3. Lee, D.-W. Phantom maps and the Gray index. *Topol. Appl.* **2004**, *138*, 265–275. [[CrossRef](#)]
4. Lee, D.-W. On the same n -type conjecture for the suspension of the infinite complex projective space. *Proc. Am. Math. Soc.* **2009**, *137*, 1161–1168. [[CrossRef](#)]
5. Lee, D.-W. On the same n -type structure for the suspension of the Eilenberg-MacLane spaces. *J. Pure Appl. Algebra* **2010**, *214*, 2027–2032. [[CrossRef](#)]
6. Lee, D.-W. On the same N -type of the suspension of the infinite quaternionic projective space. *J. Pure Appl. Algebra* **2013**, *217*, 1325–1334. [[CrossRef](#)]
7. Lee, D.-W. On the generalized same N -type conjecture. *Math. Proc. Camb. Phil. Soc.* **2014**, *157*, 329–344. [[CrossRef](#)]
8. Lee, D.-W. Algebraic loop structures on algebra comultiplications. *Open Math.* **2019**, *17*, 742–757. [[CrossRef](#)]
9. Arkowitz, M.; Gutierrez, M. Comultiplications on free groups and wedge of circles. *Trans. Amer. Math. Soc.* **1998**, *350*, 1663–1680. [[CrossRef](#)]
10. Arkowitz, M.; Lupton, G. Rational co-H-spaces. *Comment. Math. Helv.* **1991**, *66*, 79–108. [[CrossRef](#)]
11. Arkowitz, M.; Lupton, G. Equivalence classes of homotopy-associative comultiplications of finite complexes. *J. Pure Appl. Algebra* **1995**, *102*, 109–136. [[CrossRef](#)]
12. Golasinski, M.; Goncalves, D.L. Comultiplications of the wedge of two Moore spaces. *Colloq. Math.* **1998**, *76*, 229–242. [[CrossRef](#)]
13. Naylor, C.M. On the number of co-multiplications of a suspension. *Ill. J. Math.* **1968**, *12*, 620–622. [[CrossRef](#)]
14. O'Regan, D. The topological transversality theorem for multivalued maps with continuous selections. *Mathematics* **2019**, *7*, 1113. [[CrossRef](#)]
15. Greenlees, J.P.C.; Lee, D.-W. The representation-ring-graded local cohomology spectral sequence for $B\mathbb{P}\mathbb{R}\langle 3 \rangle$. *Comm. Algebra* **2019**, *47*, 2396–2411. [[CrossRef](#)]
16. Sullivan, D. *Geometric Topology. Part I. Localization, Periodicity, and Galois Symmetry*; Revised Version; Massachusetts Institute of Technology: Cambridge, MA, USA, 1971.
17. Bousfield, A.K.; Kan, D.M. *Homotopy Limits, Completions and Localizations*; Lecture Notes in Mathematics; Springer: Berlin, Germany; New York, NY, USA, 1972; Volume 304.
18. Ganea, T. Cogroups and suspensions. *Invent. Math.* **1970**, *9*, 185–197. [[CrossRef](#)]
19. Whitehead, G.W. *Elements of Homotopy Theory*; Springer: Berlin, Germany, 1978.
20. Zabrodsky, A. Homotopy associativity and finite CW-complexes. *Topology* **1970**, *9*, 121–128. [[CrossRef](#)]
21. Arkowitz, M.; Lee, D.-W. Properties of comultiplications on a wedge of spheres. *Topol. Appl.* **2010**, *157*, 1607–1621. [[CrossRef](#)]

22. Arkowitz, M.; Lee, D.-W. Comultiplications on a wedge of two spheres. *Sci. China Math.* **2011**, *54*, 9–22. [[CrossRef](#)]
23. Lee, D.-W. Comultiplication structures for a wedge of spheres. *Filomat* **2016**, *30*, 3525–3546. [[CrossRef](#)]
24. Lee, D.-W.; Lee, S. Homotopy comultiplications on the k -fold wedge of spheres. *Topol. Appl.* **2019**, *254*, 145–170. [[CrossRef](#)]
25. Hilton, P.J. On the homotopy groups of the union of spheres. *J. Lond. Math. Soc.* **1955**, *30*, 154–172. [[CrossRef](#)]
26. Arkowitz, M. The generalized Whitehead product. *Pac. J. Math.* **1962**, *12*, 7–23. [[CrossRef](#)]
27. Arkowitz, M.; Golasiński, M. Co-H-structures on Moore spaces of type $(G, 2)$. *Canad. J. Math.* **1994**, *46*, 673–686. [[CrossRef](#)]
28. Toda, H. *Composition Methods in Homotopy Groups of Spheres*; Princeton University Press: Princeton, NJ, USA, 1962.



© 2020 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).