

Article

Stability Results for Implicit Fractional Pantograph Differential Equations via ϕ -Hilfer Fractional Derivative with a Nonlocal Riemann-Liouville Fractional Integral Condition

Idris Ahmed ^{1,2,3} , Poom Kumam ^{2,4,*} , Kamal Shah ^{5,6} , Piyachat Borisut ^{1,2} ,
Kanokwan Sitthithakerngkiet ⁷  and Musa Ahmed Demba ^{1,2,8} 

- ¹ KMUTTFixed Point Research Laboratory, Room SCL 802 Fixed Point Laboratory, Science Laboratory Building, Department of Mathematics, Faculty of Science, King Mongkut's University of Technology Thonburi (KMUTT), 126 Pracha-Uthit Road, Bang Mod, Thrung Khru, Bangkok 10140, Thailand; idris.ahamed@mail.kmutt.ac.th (I.A.); piyachat.borisut@mail.kmutt.ac.th (P.B.); musa.demba@mail.kmutt.ac.th (M.A.D.)
- ² Center of Excellence in Theoretical and Computational Science (TaCS-CoE), Science Laboratory Building, King Mongkut's University of Technology Thonburi (KMUTT), 126 Pracha-Uthit Road, Bang Mod, Thrung Khru, Bangkok 10140, Thailand
- ³ Department of Mathematics and Computer Science, Sule Lamido University, Kafin-Hausa, Jigawa State P.M.B 048, Nigeria
- ⁴ Department of Medical Research, China Medical University Hospital, China Medical University, Taichung 40402, Taiwan
- ⁵ Department of Mathematics, University of Malakand, Chakadara Dir(L), Khyber Pakhtunkhwa 18800, Pakistan; kamal@uom.edu.pk
- ⁶ Department of Mathematics and Basic Sciences, Prince Sultan University, Riyadh 11586, Saudi Arabia
- ⁷ Intelligent and Nonlinear Dynamic Innovations Research Center, Department of Mathematics, Faculty of Applied Science, King Mongkut's University of Technology North Bangkok (KMUTNB), Wongsawang, Bangsue, Bangkok 10800, Thailand; kanokwan.s@sci.kmutnb.ac.th
- ⁸ Department of Mathematics, Faculty of Computing and Mathematical Sciences, Kano University of Science and Technology, Wudil, Kano State P.M.B 3244, Nigeria
- * Correspondence: poom.kumam@mail.kmutt.ac.th

Received: 27 November 2019; Accepted: 18 December 2019; Published: 7 January 2020

Abstract: This paper presents a class of implicit pantograph fractional differential equation with more general Riemann-Liouville fractional integral condition. A certain class of generalized fractional derivative is used to set the problem. The existence and uniqueness of the problem is obtained using Schaefer's and Banach fixed point theorems. In addition, the Ulam-Hyers and generalized Ulam-Hyers stability of the problem are established. Finally, some examples are given to illustrative the results.

Keywords: Hilfer fractional derivative; Ulam stability; pantograph differential equation; nonlocal integral condition

MSC: 26A33; 34A34; 34D20; 34A12

1. Introduction

The fractional-order differential equation is the oldest theory in the field of science and engineering. This theory has been used over the years, as the outcomes were found to be important in the field of economics, control theory and material sciences see [1–4]. Because of the nonlocal property

of fractional-order differential equation, researchers are allowed to select the most appropriate operator and use it in order to get a better description of the complex phenomena in the real world. The generalization of classical calculus are the fractional calculus. Nevertheless, there are various definitions of fractional integrals and derivatives of arbitrary order with different types of operator. Recently, Furati et al. [5] proposed a Hilfer fractional derivatives which interpolates with Riemann-Liouville and Caputo fractional derivatives. These fractional operator provide an extra degree of freedom when choosing the initial condition. Furthermore, models based on this operator provide an excellent results compared with the integer-order derivatives, for example, we refer the interesting reader to see [6–18].

Qualitative analysis of fractional differential equations plays a vital role in the field of fractional differential equations. However, many researchers studied the existence and uniqueness of solution of differential equation with different types of fractional integral and derivatives. More recently, motivated by classical Riemann-Liouville, Caputo fractional derivative, Hilfer-fractional derivative, ψ -Riemann-Liouville integral and ψ -Caputo fractional derivatives, Sousa and Oliveira [19] initiated an interesting fractional differential operator called ψ -Hilfer fractional derivatives, that is a fractional derivative of a function with respect to another function ψ . These fractional derivatives generalized the aforementioned fractional derivatives and integrals. The main advantages of these operator is the freedom of choice of the function ψ and its merge and acquire the properties of the aforementioned fractional operators. Results based on these setting can be found in [18–34]. The Ulam-Hyers stability point of view, is the vital and special type of stability that attracts many researchers in the field of mathematical analysis. Moreover, the Ulam-Hyers and Ulam-Hyers-Rassias stability of linear, implicit and nonlinear fractional differential equations were examined in [17,35–49].

Pantograph differential equations are a special class of delay differential equation arising in deterministic situations and are of the form:

$$\begin{cases} g'(s) = kg(s) + lg(\lambda s), & s \in [0, b], \quad b > 0, \quad 0 < \lambda < 1, \\ g(0) = g_0. \end{cases} \quad (1)$$

The pantograph is a device used in electric trains to collect electric current from the overhead lines. This equation was modeled by Ockendon and Tayler [50]. Pantograph equation play a vital role in physics, pure and applied mathematics, such as control systems, electrodynamics, probability, number theory, and quantum mechanics. Motivated by their importance, a lot of researchers generalized these equation in to various forms and introduced the solvability aspect of such problems both theoretically and numerically, (for more details see [16,51–57] and references therein). However, very few works have been proposed with respect to pantograph fractional differential equations.

In [48], the authors considered an implicit fractional differential equations with nonlocal condition described by:

$$\begin{cases} D_{0+}^{\alpha, \beta} w(\tau) = f(\tau, w(\tau), D_{0+}^{\alpha, \beta} w(\tau)), & \tau \in I = [0, T], \\ I_{0+}^{1-\gamma} w(0) = \sum_{i=1}^m c_i w(\eta_i), & \alpha \leq \gamma = \alpha + \beta - \alpha\beta, \quad \eta_i \in [0, T], \end{cases} \quad (2)$$

where $D_{0+}^{\alpha, \beta}(\cdot)$ is the Hilfer fractional derivative of order $(0 < \alpha < 1)$ and type $0 \leq \beta \leq 1$. The existence and uniqueness results were obtained by applying Schaefer's fixed point theorem and Banach's contraction principle. Moreover, the authors discussed the stability analysis via Gronwall's lemma. Sousa and Oliveira [47] discussed the existence, uniqueness and Ulam-Hyers-Rassias stability for a class of φ -Hilfer fractional differential equations described by:

$$\begin{cases} {}^H\mathfrak{D}_{a+}^{\alpha, \beta; \varphi} g(t) = f(t, g(t), {}^H\mathfrak{D}_{a+}^{\alpha, \beta; \varphi} g(t)), & t \in \mathcal{J} = [a, T], \\ \mathfrak{I}_{a+}^{1-\gamma; \varphi} g(a) = g_a, & \alpha \leq \gamma = \alpha + \beta - \alpha\beta, \quad T > a, \end{cases} \quad (3)$$

where ${}^H\mathcal{D}_{0+}^{\alpha,\beta;\varphi}(\cdot)$ is the φ -Hilfer fractional derivative of order $(0 < \alpha \leq 1)$ and operator $(0 \leq \beta \leq 1)$, $\mathcal{I}_{0+}^{1-\gamma;\varphi}(\cdot)$, is the Riemann-Liouville fractional integral of order $1 - \gamma$, with respect to the function φ , $f : [a, T] \times \mathbb{R}^2 \rightarrow \mathbb{R}$ is a continuous function. Recently Harikrishnan et. al [58] established existence and uniqueness of nonlocal initial value problem for fractional pantograph differential equation involving ψ -Hilfer fractional derivative of the form:

$$\begin{cases} {}^H\mathcal{D}_{a+}^{\alpha,\beta;\psi}v(s) = f(s, v(s), v(\lambda s)), & s \in (a, b], \quad s > a, \quad 0 < \lambda < 1, \\ \mathcal{I}_{a+}^{1-\gamma;\psi}v(a) = \sum_{j=1}^k c_j v(\tau_j), & \tau_j \in (a, b], \quad \gamma = \alpha + \beta - \alpha\beta, \end{cases} \quad (4)$$

where ${}^H\mathcal{D}_{a+}^{\alpha,\beta;\psi}(\cdot)$ is the ψ -Hilfer fractional derivative of order $0 < \alpha < 1$ and type $0 \leq \beta \leq 1$, $\mathcal{I}_{a+}^{1-\gamma;\psi}(\cdot)$, is the Riemann-Liouville fractional integral of order $1 - \gamma$, with respect to the continuous function ψ such that $\psi'(\cdot) > 0$, $f \in C(t \in (a, b], \mathbb{R}^2, \mathbb{R})$.

Motivated by the papers [21,47,48] and some familiar results on fractional pantograph differential equations [16,52,55,58]. We discuss the existence and uniqueness of the solution of the implicit pantograph fractional differential equations involving ϕ -Hilfer fractional derivatives. Furthermore, the Ulam-Hyers and generalized Ulam-Hyers stability are also discussed. The implicit pantograph fractional differential equations involving ϕ -Hilfer fractional derivatives is of the form

$$\begin{cases} {}^H\mathcal{D}_{0+}^{r,p;\phi}z(t) = f(t, z(t), z(\gamma t), {}^H\mathcal{D}_{0+}^{r,p;\phi}z(\gamma t)), & t \in \mathcal{J} = (0, T], \quad 0 < \gamma < 1, \\ \mathcal{I}_{0+}^{1-q;\phi}z(0+) = \sum_{i=1}^m b_i \mathcal{I}_{0+}^{\rho;\phi}z(\xi_i), & r \leq q = r + p - rp, \end{cases} \quad (5)$$

where ${}^H\mathcal{D}_{0+}^{r,p;\phi}(\cdot)$ is the generalized ϕ -Hilfer fractional derivatives of order $(0 < r < 1)$ and type $(0 \leq p \leq 1)$, $\mathcal{I}_{0+}^{1-q;\phi}(\cdot)$ and $\mathcal{I}_{0+}^{\rho;\phi}(\cdot)$ are ϕ -Riemann-Liouville fractional integral of order $1 - q$ and $\rho > 0$ respectively with respect to the continuous function ϕ such that $\phi'(\cdot) \neq 0$, $f : (0, T] \times \mathbb{R}^3 \rightarrow \mathbb{R}$ is a given continuous function, $T > 0$, $b_i \in \mathbb{R}$ and $\xi_i \in \mathcal{J}$ satisfying $0 < \xi_1 \leq \xi_2 \leq \dots \leq \xi_m < T$ for $i = 1, 2, \dots, m$.

As far as we know, to the best of our understanding, results of Ulam-Hyers and generalized Ulam-Hyers stability with respect to the pantograph differential equation are very few and in fact most authors discuss existence and uniqueness, while we study existence, uniqueness and stability analysis for a class of implicit pantograph fractional differential equations with ϕ -Hilfer derivatives and nonlocal Riemann-Liouville fractional integral condition.

This paper contributes to the growth of qualitative analysis of fractional differential equation in particular pantograph fractional differential equation when ϕ -Hilfer fractional derivatives involved and the nonlocal initial condition proposed in this paper generalized the following initial conditions:

- If $\rho \rightarrow 0$, the initial condition reduces to multi-point nonlocal condition.
- If $\rho \rightarrow 1$, the initial condition coincide with the nonlocal integral condition.
- In physical problems, the nonlocal condition yields an excellent results compared with the initial condition $z(0) = z_0$ [59,60].

In addition, we notice that the function $f(s, v(s), v(\lambda s))$, $s \in (a, b]$, $0 < \lambda < 1$, defined in Equation (4) is not well-define for some choices of λ .

Therefore, the paper is organized as follows: In Section 2, it recalls some basic and fundamental definitions and lemmas. In Section 3, we prove existence and uniqueness of the proposed problem (5). Ulam-Hyers and generalized Ulam-Hyers stability for the proposed problem were discussed in Section 4. While in Section 5, two examples were given to illustrate the applicability of our results. Lastly, the conclusion part of the paper is given in Section 6.

2. Preliminaries

This section will recall some useful prerequisites facts, definitions and some fundamental lemmas with respect to fractional differential equations.

Throughout the paper, we denote $\mathcal{C}[\mathcal{J}, \mathbb{R}]$ the Banach space of all continuous functions from $\mathcal{J}$ into $\mathbb{R}$ with the norm defined by [1]

$$\|f\| = \sup_{t \in \mathcal{J}} \{|f(t)|\}.$$

The weighted space $\mathcal{C}_{q,\phi}[\mathcal{J}, \mathbb{R}]$ of continuous function f on the interval $[a, T]$ is defined by

$$\mathcal{C}_{q,\phi}[\mathcal{J}, \mathbb{R}] = \{f(t) : (a, T] : (\phi(t) - \phi(0))^q f(t) \in \mathcal{C}[\mathcal{J}, \mathbb{R}]\},$$

with the norm

$$\|f\|_{\mathcal{C}_{q,\phi}[\mathcal{J}, \mathbb{R}]} = \|(\phi(t) - \phi(0))^q f(t)\| = \max |(\phi(t) - \phi(0))^q f(t) : t \in \mathcal{J}|.$$

Moreover, for each $n \in \mathbb{N}$ and $0 \leq q < 1$ with $q = r + p - rp$

$$\mathcal{C}_{q,\phi}^n[\mathcal{J}, \mathbb{R}] = \{f^n \in \mathcal{C}_{q,\phi}[\mathcal{J}, \mathbb{R}]\}$$

$$\mathcal{C}_{q,\phi}^{r,p}[\mathcal{J}, \mathbb{R}] = \{f \in \mathcal{C}_{q,\phi}[\mathcal{J}, \mathbb{R}] : \mathcal{D}_{0+}^{r,p;\phi} f \in \mathcal{C}_{q,\phi}[\mathcal{J}, \mathbb{R}]\}.$$

Indeed, for $n = 0$, we have

$$\mathcal{C}_{q,\phi}^0[\mathcal{J}, \mathbb{R}] = \mathcal{C}_{q,\phi}[\mathcal{J}, \mathbb{R}],$$

with the norm

$$\|f\|_{\mathcal{C}_{q,\phi}^n[\mathcal{J}, \mathbb{R}]} = \sum_{k=0}^{n-1} \|f^k\|_{\mathcal{C}[\mathcal{J}, \mathbb{R}]} + \|f^n\|_{\mathcal{C}_{q,\phi}^n[\mathcal{J}, \mathbb{R}]}.$$

Furthermore, we present the following space $\mathcal{C}_{1-q,\phi}^{r,p}[\mathcal{J}, \mathbb{R}]$ and $\mathcal{C}_{1-q,\phi}^q[\mathcal{J}, \mathbb{R}]$ defined as:

$$\mathcal{C}_{1-q,\phi}^{r,p}[\mathcal{J}, \mathbb{R}] = \{f \in \mathcal{C}_{1-q,\phi}[\mathcal{J}, \mathbb{R}], \mathcal{D}_{0+}^{r,p;\phi} f \in \mathcal{C}_{1-q,\phi}[\mathcal{J}, \mathbb{R}]\}$$

and

$$\mathcal{C}_{1-q,\phi}^q[\mathcal{J}, \mathbb{R}] = \{f \in \mathcal{C}_{1-q,\phi}[\mathcal{J}, \mathbb{R}], \mathcal{D}_{0+}^{q;\phi} f \in \mathcal{C}_{1-q,\phi}[\mathcal{J}, \mathbb{R}]\}.$$

Clearly, $\mathcal{C}_{1-q,\phi}^q[\mathcal{J}, \mathbb{R}] \subset \mathcal{C}_{1-q,\phi}^{r,p}[\mathcal{J}, \mathbb{R}]$.

Definition 1 ([1]). Let $(0, b]$ be a finite or infinite interval on the half-axis $\mathbb{R}^+$, and $\phi(\xi) \geq 0$ be monotone function on $(a, b]$ whose $\phi'(\xi)$ is continuous on $(0, b)$. The ϕ -Hilfer Riemann-Liouville fractional integral of order $r \in \mathbb{R}^+$ of function w is defined by

$$(\mathcal{I}_{0+}^{r,\phi} w)(\xi) = \frac{1}{\Gamma(r)} \int_{0+}^{\xi} \phi'(s) (\phi(\xi) - \phi(s)) w(s) ds, \quad \xi > 0, \quad (6)$$

where $\Gamma(\cdot)$ represent the Gamma function.

Definition 2 ([5]). Let $n - 1 < r < n$, $0 \leq p \leq 1$. The left-sided Hilfer fractional derivative of order r and parameter p of function w is defined by

$$\mathcal{D}_{0+}^{r,p} w(\xi) = \left(\mathcal{I}_{0+}^{p(n-r)} \mathcal{D}^n \mathcal{I}_{0+}^{(1-p)(n-r)} w \right) (\xi), \quad (7)$$

where $\mathcal{D}^n = \left(\frac{d}{d\xi} \right)^n$.

The following Definition generalized Equation (7).

Definition 3 ([19]). Let $f, \phi \in C^n[\mathcal{J}, \mathbb{R}]$ be two functions such that $\phi(\xi) \geq 0$ and $\phi'(t) \neq 0$ for all $\xi \in [\mathcal{J}, \mathbb{R}]$ and $n - 1 < r < n$ with $n \in \mathbb{N}$. The left-side ϕ -Hilfer fractional derivative of a function w of order r and type $(0 \leq p \leq 1)$ is defined by

$${}^H\mathcal{D}_{0+}^{r,p;\phi}w(\xi) = \mathcal{I}_{0+}^{p(n-r);\phi} \left(\frac{1}{\phi'(\xi)} \frac{d}{d\xi} \right)^n \mathcal{I}_{0+}^{(1-p)(n-r);\phi}w(\xi). \quad (8)$$

The following lemma shows the semigroup properties of ϕ -Hilfer fractional integral and derivative.

Lemma 1 ([5]). Let $r \geq 0, 0 \leq p < 1$ and $w \in L^1[\mathcal{J}, \mathbb{R}]$. Then

$$\mathcal{I}_{0+}^{r;\phi} \mathcal{I}_{0+}^{p;\phi}w(\xi) = \mathcal{I}_{0+}^{r+p;\phi}w(\xi),$$

a.e $\xi \in \mathcal{J}$.

In particular, if $w \in \mathcal{C}_{q,\phi}[\mathcal{J}, \mathbb{R}]$ and $w \in \mathcal{C}[\mathcal{J}, \mathbb{R}]$, then

$$\mathcal{I}_{0+}^{r;\phi} \mathcal{I}_{0+}^{p;\phi}w(\xi) = \mathcal{I}_{0+}^{r+p;\phi}w(\xi),$$

for all $\xi \in (0, T]$ and

$${}^H\mathcal{D}_{0+}^{r;\phi} \mathcal{I}_{0+}^{r;\phi}w(\xi) = w(\xi),$$

for all $\xi \in \mathcal{J}$.

The composition of the ϕ -Hilfer fractional integral and derivative operator is given by the following lemmas.

Lemma 2 ([21]). Let $r \geq 0, 0 \leq p < 1$ and $q = r + p - rp$. If $w(\xi) \in \mathcal{C}_{1-q}^q[\mathcal{J}, \mathbb{R}]$, then

$$\mathcal{I}_{0+}^{q;\phi} {}^H\mathcal{D}_{0+}^{q;\phi}w(\xi) = \mathcal{I}_{0+}^{r;\phi} {}^H\mathcal{D}_{0+}^{r,p;\phi}w(\xi)$$

and

$${}^H\mathcal{D}_{0+}^{q;\phi} \mathcal{I}_{0+}^{r;\phi}w(\xi) = {}^H\mathcal{D}_{0+}^{p(1-r);\phi}w(\xi).$$

Lemma 3 ([6,19]). If $w \in \mathcal{C}^n[\mathcal{J}, \mathbb{R}]$ and let $n - 1 < r < n, 0 \leq p \leq 1$ and $q = r + p - rp$. Then

$$\mathcal{I}_{0+}^{r;\phi} {}^H\mathcal{D}_{0+}^{r,p;\phi}w(\xi) = w(\xi) - \sum_{k=1}^n \frac{(\phi(\xi) - \phi(0))^{q-k}}{\Gamma(q-k+1)} w_{\phi}^{[n-k]} \mathcal{I}_{0+}^{(1-p)(n-r);\phi}w(0),$$

for all $\xi \in \mathcal{J}$. Moreover, if $0 < r < 1$, we have

$$\mathcal{I}_{0+}^{r;\phi} {}^H\mathcal{D}_{0+}^{r,p;\phi}w(\xi) = w(\xi) - \frac{(\phi(\xi) - \phi(0))^{q-1}}{\Gamma(q)} \mathcal{I}_{0+}^{(1-p)(1-r);\phi}w(0).$$

In addition, if $w \in \mathcal{C}_{1-q;\phi}[\mathcal{J}, \mathbb{R}]$ and $\mathcal{I}_{0+}^{1-q;\phi}w \in \mathcal{C}_{1-q;\phi}^1[\mathcal{J}, \mathbb{R}]$, then

$$\mathcal{I}_{0+}^{q;\phi} {}^H\mathcal{D}_{0+}^{q;\phi}w(\xi) = w(\xi) - \frac{(\phi(\xi) - \phi(0))^{q-1}}{\Gamma(q)} \mathcal{I}_{0+}^{(1-q);\phi}w(0),$$

for all $0 < q < 1$ and $t \in \mathcal{J}$.

Lemma 4 ([6]). Let $r > 0$, $0 \leq q < 1$ and $w \in \mathcal{C}_{q;\phi}[\mathcal{J}, \mathbb{R}]$. If $r > q$, then $\mathcal{I}_{0+}^{r;\phi} w \in \mathcal{C}[\mathcal{J}, \mathbb{R}]$ and

$$\mathcal{I}_{0+}^{r;\phi} w(0) = \lim_{\xi \rightarrow 0} \mathcal{I}_{0+}^{r;\phi} w(\xi) = 0.$$

Lemma 5 ([21]). Let $r > 0$, $0 \leq p \leq 1$ and $q = r + p - rp$. If $w \in \mathcal{C}_{1-q;\phi}^q[\mathcal{J}, \mathbb{R}]$, then

$$\mathcal{I}_{0+}^{q;\phi} \mathcal{D}_{0+}^{q;\phi} w(\xi) = \mathcal{I}_{0+}^{r;\phi} \mathcal{D}_{0+}^{r,p;\phi} w(\xi)$$

and

$${}^H \mathcal{D}_{0+}^{q;\phi} \mathcal{I}_{0+}^{r;\phi} w(\xi) = \mathcal{D}_{0+}^{q(1-r);\phi} w(\xi).$$

Lemma 6. Let $f \in L^1(\mathcal{J})$ such that $\mathcal{D}_{0+}^{p(1-r);\phi} w \in L^1(\mathcal{J})$ exists, then

$$\mathcal{D}_{0+}^{r,p;\phi} \mathcal{I}_{0+}^{r;\phi} w(\xi) = \mathcal{I}_{0+}^{p(1-r);\phi} \mathcal{D}_{0+}^{p(1-r);\phi} w(\xi).$$

Next, we take into account some important properties of ϕ -fractional derivative and integral operator as follows:

Proposition 1 ([1]). Let $\xi > 0$, $r \geq 0$ and $s > 0$. Then, ϕ -fractional integral and derivative of a power function are given by

$${}^H \mathcal{D}_{0+}^{r;\phi} (\phi(\xi) - \phi(0))^{s-1} = \frac{\Gamma(s)}{\Gamma(s-r)} (\phi(\xi) - \phi(0))^{r+s-1}$$

and

$$\mathcal{I}_{0+}^{r;\phi} (\phi(\xi) - \phi(0))^{s-1} = \frac{\Gamma(s)}{\Gamma(s+r)} (\phi(\xi) - \phi(0))^{r+s-1}.$$

Furthermore, if $0 < r < 1$, then

$${}^H \mathcal{D}_{0+}^{r;\phi} (\phi(\xi) - \phi(0))^{r-1} = 0.$$

Theorem 1 ([19]). If $w \in \mathcal{C}^1[\mathcal{J}, \mathbb{R}]$, $0 < r < 1$ and $0 \leq p \leq 1$. Then we have the followings:

- (i) ${}^H \mathcal{D}_{0+}^{r,p;\phi} \mathcal{I}_{0+}^{r;\phi} w(\xi) = w(\xi)$.
- (ii) $\mathcal{I}_{0+}^{r;\phi} {}^H \mathcal{D}_{0+}^{r,p;\phi} w(\xi) = w(\xi) - \frac{(\phi(\xi) - \phi(0))^{q-1}}{\Gamma(q)} \mathcal{I}_{0+}^{(1-p)(1-r);\phi} w(\xi)$.

Lemma 7 ([6]). Let $h : \mathcal{J} \times \mathbb{R} \rightarrow \mathbb{R}$ such that for any $z \in \mathcal{C}_{1-q;\phi}[\mathcal{J}, \mathbb{R}]$, $h \in \mathcal{C}_{1-q;\phi}[\mathcal{J}, \mathbb{R}]$. A function $z \in \mathcal{C}_{1-q;\phi}^q[\mathcal{J}, \mathbb{R}]$ is a solution of the fractional initial value problem:

$$\begin{cases} {}^H \mathcal{D}_{0+}^{r,p;\phi} z(t) = h(t), & 0 < r \leq 1, \quad 0 \leq p \leq 1, \\ I_{0+}^{1-q,\phi} z(0^+) = z_0 \in \mathbb{R}, & q = r + p - rp, \end{cases}$$

if and only if z satisfies the following integral equation,

$$z(t) = \frac{z_0}{\Gamma(q)} (\phi(t) - \phi(0))^{q-1} + \frac{1}{\Gamma(r)} \int_{0^+}^t \phi'(s) (\phi(t) - \phi(s))^{r-1} h(s) ds.$$

3. Main Results

In this section, we first adopt some techniques from Lemma 7 in order to establish an important mixed-type integral equation of problem (5). Thus, we need the following auxiliary lemma.

Lemma 8. Let $0 < r < 1$, $0 \leq p \leq 1$ and $q = r + p - rp$. Suppose $f : \mathcal{J} \times \mathbb{R}^3 \rightarrow \mathbb{R}$ is a function such that $f \in \mathcal{C}_{1-q;\phi}[\mathcal{J}, \mathbb{R}]$ for any $z \in \mathcal{C}_{1-q;\phi}[\mathcal{J}, \mathbb{R}]$. If $z \in \mathcal{C}_{1-q;\phi}^q[\mathcal{J}, \mathbb{R}]$ then z satisfies the problem (5) if and only if z satisfies the mixed-type integral equation:

$$z(t) = \frac{\delta \Gamma(\rho + q)}{\Gamma(q) \Gamma(\rho + r)} (\phi(t) - \phi(0))^{q-1} \sum_{i=1}^m b_i \int_{0^+}^{\xi_i} \phi'(s) (\phi(\xi_i) - \phi(s))^{\rho+r-1} T_z(s) ds \\ + \frac{1}{\Gamma(r)} \int_{0^+}^t \phi'(s) (\phi(t) - \phi(s))^{r-1} T_z(s) ds, \quad (9)$$

where

$$\delta = \frac{1}{\Gamma(\rho + q) - \sum_{i=1}^m b_i (\phi(\xi_i) - \phi(0))^{\rho+q-1}}, \quad (10)$$

such that $\Gamma(\rho + q) \neq \sum_{i=1}^m b_i (\phi(\xi_i) - \phi(0))^{\rho+q-1}$.

For simplicity, we take

$$T_z(t) = {}^H D_{0^+}^{r,p;\phi} z(t) = f(t, z(t), z(\gamma t), T_z(t)). \quad (11)$$

Proof. Suppose $z \in \mathcal{C}_{1-q;\phi}^q[\mathcal{J}, \mathbb{R}]$ is a solution to the problem (5), then, we show that z is also a solution of (5). Indeed, from Lemma 7, we have

$$z(t) = \frac{(\phi(t) - \phi(0))^{q-1}}{\Gamma(q)} \mathcal{I}_{0^+}^{1-q;\phi} z(0) + \frac{1}{\Gamma(r)} \int_{0^+}^t \phi'(s) (\phi(t) - \phi(s))^{r-1} T_z(s) ds. \quad (12)$$

Now, if we substitute $t = \xi_i$ and multiply both sides by b_i in Equation (12), we obtain

$$b_i z(\xi_i) = \frac{(\phi(\xi_i) - \phi(0))^{q-1}}{\Gamma(q)} b_i \mathcal{I}_{0^+}^{1-q;\phi} z(0) + \frac{b_i}{\Gamma(r)} \int_{0^+}^{\xi_i} \phi'(s) (\phi(\xi_i) - \phi(s))^{r-1} T_z(s) ds. \quad (13)$$

Next, by applying $\mathcal{I}_{0^+}^{\rho;\phi}$ to both sides of Equation (13) and using Lemma 1 and Proposition 1, we get

$$\mathcal{I}_{0^+}^{\rho;\phi} b_i z(\xi_i) = \frac{(\phi(\xi_i) - \phi(0))^{\rho+q-1}}{\Gamma(\rho + q)} b_i \mathcal{I}_{0^+}^{1-q;\phi} z(0) \\ + \frac{b_i}{\Gamma(\rho + r)} \int_{0^+}^{\xi_i} \phi'(s) (\phi(\xi_i) - \phi(s))^{\rho+r-1} T_z(s) ds. \quad (14)$$

This implies that

$$\sum_{i=1}^m \mathcal{I}_{0^+}^{\rho;\phi} b_i z(\xi_i) = \frac{1}{\Gamma(\rho + q)} \left(\sum_{i=1}^m b_i (\phi(\xi_i) - \phi(0))^{\rho+q-1} \right) \mathcal{I}_{0^+}^{1-q;\phi} z(0) \\ + \frac{1}{\Gamma(\rho + r)} \sum_{i=1}^m b_i \int_{0^+}^{\xi_i} \phi'(s) (\phi(\xi_i) - \phi(s))^{\rho+r-1} T_z(s) ds. \quad (15)$$

Inserting the initial condition: $\mathcal{I}_{0+}^{1-q;\phi} z(0^+) = \sum_{i=1}^m \mathcal{I}_{0+}^{\rho;\phi} b_i z(\xi_i)$ in Equation (15) we have

$$\begin{aligned} \mathcal{I}_{0+}^{1-q;\phi} z(0) &= \frac{1}{\Gamma(\rho+q)} \left(\sum_{i=1}^m b_i (\phi(\xi_i) - \phi(0))^{\rho+q-1} \right) \mathcal{I}_{0+}^{1-q;\phi} z(0) \\ &\quad + \frac{1}{\Gamma(\rho+r)} \sum_{i=1}^m b_i \int_{0+}^{\xi_i} \phi'(s) (\phi(\xi_i) - \phi(s))^{\rho+r-1} T_z(s) ds, \end{aligned} \quad (16)$$

which implies that

$$\begin{aligned} \frac{1}{\Gamma(\rho+r)} \sum_{i=1}^m b_i \int_{0+}^{\xi_i} \phi'(s) (\phi(\xi_i) - \phi(s))^{\rho+r-1} T_z(s) ds \\ = \left(1 - \frac{1}{\Gamma(\rho+q)} \sum_{i=1}^m b_i (\phi(\xi_i) - \phi(0))^{\rho+q-1} \right) \mathcal{I}_{0+}^{1-q;\phi} z(0) \\ = \frac{1}{\delta\Gamma(\rho+q)} \mathcal{I}_{0+}^{1-q;\phi} z(0). \end{aligned} \quad (17)$$

Thus,

$$\mathcal{I}_{0+}^{1-q;\phi} z(0) = \frac{\delta\Gamma(\rho+q)}{\Gamma(\rho+r)} \sum_{i=1}^m b_i \int_{0+}^{\xi_i} \phi'(s) (\phi(\xi_i) - \phi(s))^{\rho+r-1} T_z(s) ds. \quad (18)$$

Hence, the result follows by putting Equation (18) in Equation (12). This implies that $z(t)$ satisfies Equation (9).

Conversely, suppose that $z \in \mathcal{C}_{1-q;\phi}^q$ satisfies the mixed-type integral Equation (9), then, we show that z satisfies Equation (5). Applying $\mathcal{D}_{0+}^{q;\phi}$ to both sides of Equation (9) and using Lemma 2 and Proposition 1, we get

$$\begin{aligned} \mathcal{D}_{0+}^{q;\phi} z(t) &= \mathcal{D}_{0+}^{q;\phi} \left(\frac{\delta\Gamma(\rho+q)}{\Gamma(q)\Gamma(r)} (\phi(t) - \phi(0))^{q-1} \sum_{i=1}^m b_i \int_{0+}^{\xi_i} \phi'(s) (\phi(\xi_i) - \phi(s))^{\rho+r-1} T_z(s) ds \right) \\ &\quad + \mathcal{D}_{0+}^{q;\phi} \left(\frac{1}{\Gamma(r)} \int_{0+}^t \phi'(s) (\phi(t) - \phi(s))^{r-1} T_z(s) ds \right) \\ &= \mathcal{D}_{0+}^{p(1-r);\phi} f(t, z(t), z(\gamma t), \mathcal{D}_{0+}^{r,p;\phi} z(\gamma t)). \end{aligned} \quad (19)$$

Since $\mathcal{D}_{0+}^{r,p;\phi} z \in \mathcal{C}_{1-q;\phi}[\mathcal{J}, \mathbb{R}]$, then by definition of $\mathcal{C}_{1-q;\phi}^q[\mathcal{J}, \mathbb{R}]$ and make use of Equation (19), we have

$$\mathcal{D}_{0+}^{p(1-r);\phi} f = \mathcal{D} \mathcal{I}_{0+}^{1-p(1-r);\phi} f \in \mathcal{C}_{1-q;\phi}[\mathcal{J}, \mathbb{R}].$$

For every $f \in \mathcal{C}_{1-q;\phi}[\mathcal{J}, \mathbb{R}]$ and Lemma 3, we can see that $\mathcal{I}_{0+}^{1-p(1-r);\phi} f \in \mathcal{C}_{1-q;\phi}[\mathcal{J}, \mathbb{R}]$, which implies that $\mathcal{I}_{0+}^{1-p(1-r);\phi} f \in \mathcal{C}_{1-q;\phi}^1[\mathcal{J}, \mathbb{R}]$ from the definition of $\mathcal{C}_{q;\phi}^n[\mathcal{J}, \mathbb{R}]$. Applying $\mathcal{I}_{0+}^{p(1-r);\phi}$ on both sides of Equation (19) and using Lemma 3, we have

$$\begin{aligned} \mathcal{I}_{0+}^{p(1-r);\phi} \mathcal{D}_{0+}^{q;\phi} z(t) &= \mathcal{I}_{0+}^{p(1-r);\phi} \mathcal{D}_{0+}^{p(1-r);\phi} T_z(t) \\ &= T_z(t) - \frac{\left(\mathcal{I}_{0+}^{1-p(1-r);\phi} T_z \right) (0^+)}{\Gamma(p(1-r))} (\phi(t) - \phi(0))^{p(r-1)-1} \\ &= T_z(t) = f(t, z(t), z(\gamma t), \mathcal{D}_{0+}^{r,p;\phi} z(\gamma t)). \end{aligned} \quad (20)$$

Finally, we show that if $z \in \mathcal{C}_{1-q}^q[\mathcal{J}, \mathbb{R}]$ satisfies Equation (9), it also satisfies the initial condition. Thus, by applying $\mathcal{I}_{0+}^{1-q;\phi}$ to both sides of Equation (9) and using Lemma 1 and Proposition 1, we obtain

$$\begin{aligned} \mathcal{I}_{0+}^{1-q;\phi} z(t) &= \mathcal{I}_{0+}^{1-q;\phi} \left(\frac{\delta \Gamma(\rho+q)}{\Gamma(q)\Gamma(\rho+r)} (\phi(t) - \phi(0))^{q-1} \sum_{i=1}^m b_i \int_{0+}^{\xi_i} \phi'(s) (\phi(\xi_i) - \phi(s))^{\rho+r-1} T_z(s) ds \right) \\ &+ \mathcal{I}_{0+}^{1-q;\phi} \left(\frac{1}{\Gamma(r)} \int_{0+}^t \phi'(s) (\phi(t) - \phi(s))^{r-1} T_z(s) ds \right) \\ &= \frac{\delta \Gamma(\rho+q)}{\Gamma(\rho+r)} \sum_{i=1}^m b_i \int_{0+}^{\xi_i} \phi'(s) (\phi(\xi_i) - \phi(s))^{\rho+r-1} T_z(s) ds + \mathcal{I}_{0+}^{1-p(1-r);\phi} T_z(t). \end{aligned} \quad (21)$$

Using Lemma 4 and the fact that $1-q < 1-p(1-r)$, then taking limit as $t \rightarrow 0$ in Equation (21) yields

$$\mathcal{I}_{0+}^{1-q;\phi} z(0+) = \frac{\delta \Gamma(\rho+q)}{\Gamma(\rho+r)} \sum_{i=1}^m b_i \int_{0+}^{\xi_i} \phi'(s) (\phi(\xi_i) - \phi(s))^{\rho+r-1} T_z(s) ds. \quad (22)$$

Now, substituting $t = \xi_i$ and multiplying through by b_i in Equation (9), we get

$$\begin{aligned} b_i z(\xi_i) &= \frac{\delta \Gamma(\rho+q)}{\Gamma(q)\Gamma(\rho+r)} b_i (\phi(\xi_i) - \phi(0))^{q-1} \sum_{i=1}^m b_i \int_{0+}^{\xi_i} \phi'(s) (\phi(\xi_i) - \phi(s))^{\rho+r-1} T_z(s) ds \\ &+ \mathcal{I}_{0+}^{1-q;\phi} \frac{b_i}{\Gamma(r)} \int_{0+}^{\xi_i} \phi'(s) (\phi(\xi_i) - \phi(s))^{r-1} T_z(s) ds. \end{aligned} \quad (23)$$

Applying $\mathcal{I}_{0+}^{\rho;\phi}$ to both sides of Equation (23), we obtain

$$\begin{aligned} \mathcal{I}_{0+}^{\rho;\phi} b_i z(\xi_i) &= \frac{\delta b_i (\phi(\xi_i) - \phi(0))^{\rho+q-1}}{\Gamma(\rho+r)} \sum_{i=1}^m b_i \int_{0+}^{\xi_i} \phi'(s) (\phi(\xi_i) - \phi(s))^{\rho+r-1} T_z(s) ds \\ &+ \frac{b_i}{\Gamma(\rho+r)} \int_{0+}^{\xi_i} \phi'(s) (\phi(\xi_i) - \phi(s))^{\rho+r-1} T_z(s) ds, \end{aligned} \quad (24)$$

which implies

$$\begin{aligned} \sum_{i=1}^m b_i \mathcal{I}_{0+}^{\rho;\phi} z(\xi_i) &= \frac{\delta}{\Gamma(\rho+r)} \sum_{i=1}^m b_i \int_{0+}^{\xi_i} \phi'(s) (\phi(\xi_i) - \phi(s))^{\rho+r-1} T_z(s) ds \sum_{i=1}^m b_i (\phi(\xi_i) - \phi(0))^{\rho+q-1} \\ &+ \frac{1}{\Gamma(\rho+r)} \sum_{i=1}^m b_i \int_{0+}^{\xi_i} \phi'(s) (\phi(\xi_i) - \phi(s))^{\rho+r-1} T_z(s) ds \\ &= \frac{1}{\Gamma(\rho+r)} \sum_{i=1}^m b_i \int_{0+}^{\xi_i} \phi'(s) (\phi(\xi_i) - \phi(s))^{\rho+r-1} T_z(s) ds \times \\ &\quad \left(1 + \delta \sum_{i=1}^m b_i (\phi(\xi_i) - \phi(0))^{\rho+q-1} \right) \end{aligned} \quad (25)$$

and

$$\mathcal{I}_{0+}^{1-q;\phi} z(0+) = \frac{\delta \Gamma(\rho+q)}{\Gamma(\rho+r)} \sum_{i=1}^m b_i \int_{0+}^{\xi_i} \phi'(s) (\phi(\xi_i) - \phi(s))^{\rho+r-1} T_z(s) ds. \quad (26)$$

Therefore, in view of Equation (22) and Equation (26), we have

$$\mathcal{I}_{0+}^{1-q;\phi} z(0^+) = \sum_{i=1}^m b_i \mathcal{I}_{0+}^{\rho;\phi} z(\xi_i). \quad (27)$$

□

3.1. Existence Result Via Schaefer's Fixed Point Theorem

This subsection will provide the proof of the existence results of Equation (5) using Schaefer's fixed point theorem.

Theorem 2 ([61]). Let $\mathcal{A} : \mathcal{X} \rightarrow \mathcal{X}$ be a completely continuous operator. Suppose that the set $\mathcal{E}(\mathcal{A}) = \{p \in \mathcal{X} : p = \varrho \mathcal{A}p, \text{ for some } \varrho \in [0, 1]\}$ is bounded, then $\mathcal{A}$ has a fixed point.

Thus we need the following assumptions:

(A₁) Let $f : \mathcal{J} \times \mathbb{R}^3 \rightarrow \mathbb{R}$ be a function such that $f \in \mathcal{C}_{1-q;\phi}[\mathcal{J}, \mathbb{R}]$ for any $z \in \mathcal{C}_{1-q;\phi}[\mathcal{J}, \mathbb{R}]$.

(A₂) There exist $k, l, m, n \in \mathcal{C}_{1-q;\phi}[\mathcal{J}, \mathbb{R}]$ with $k^* = \sup_{t \in \mathcal{J}} |k(t)| < 1$ such that

$$|f(t, u, v, w)| \leq k(t) + l(t)|x| + m(t)|y| + n(t)|z|, \quad t \in \mathcal{J}, \quad u, v, w \in \mathbb{R}.$$

Theorem 3. Let $0 < r < 1$, $0 \leq p \leq 1$ and $q = r + p - rp$. Suppose that the assumptions (A₁) and (A₂) are satisfied. Then there exist at least one solution of the problem (5) in the space $\mathcal{C}_{1-q;\phi}^{r,p}[\mathcal{J}, \mathbb{R}]$.

Proof. Define the operator $F : \mathcal{C}_{1-q;\phi}[\mathcal{J}, \mathbb{R}] \rightarrow \mathcal{C}_{1-q;\phi}[\mathcal{J}, \mathbb{R}]$ by

$$\begin{aligned} (Fz)(t) &= \frac{\delta \Gamma(\rho + q)}{\Gamma(q) \Gamma(\rho + r)} (\phi(t) - \phi(0))^{q-1} \sum_{i=1}^m b_i \int_{0+}^{\xi_i} \phi'(s) (\phi(\xi_i) - \phi(s))^{\rho+r-1} T_z(s) ds \\ &\quad + \frac{1}{\Gamma(r)} \int_{0+}^t \phi'(s) (\phi(t) - \phi(s))^{r-1} T_z(s) ds, \end{aligned} \quad (28)$$

then, clearly the operator F is well-defined. The proof is given in the following steps: Step 1: the operator F is continuous. Let z_n be a sequence such that $z_n \rightarrow z$ in $\mathcal{C}_{1-q;\phi}[\mathcal{J}, \mathbb{R}]$. Then for each $t \in \mathcal{J}$, we have

$$\begin{aligned} &|((Fz_n)(t) - (Fz)(t))(\phi(t) - \phi(0))^{1-q}| \\ &\leq \frac{|\delta| \Gamma(\rho + q)}{\Gamma(q) \Gamma(\rho + r)} \sum_{i=1}^m b_i \int_{0+}^{\xi_i} \phi'(s) (\phi(\xi_i) - \phi(s))^{\rho+r-1} |T_{z_n}(s) - T_z(s)| ds \\ &\quad + \frac{1}{\Gamma(r)} (\phi(t) - \phi(0))^{1-q} \int_{0+}^t \phi'(s) (\phi(t) - \phi(s))^{r-1} |T_{z_n}(s) - T_z(s)| ds \\ &\leq \frac{|\delta| \Gamma(\rho + q)}{\Gamma(q) \Gamma(\rho + r)} \mathcal{B}(q, \rho + r) \sum_{i=1}^m b_i (\phi(\xi_i) - \phi(s))^{\rho+r+q-1} \|T_{z_n}(\cdot) - T_z(\cdot)\|_{\mathcal{C}_{1-q;\phi}} \\ &\quad + \frac{\mathcal{B}(q, r)}{\Gamma(r)} (\phi(t) - \phi(0))^r \|T_{z_n}(\cdot) - T_z(\cdot)\|_{\mathcal{C}_{1-q;\phi}} \\ &\leq \left[\frac{|\delta| \Gamma(\rho + q)}{\Gamma(q) \Gamma(\rho + r)} \mathcal{B}(q, \rho + r) \sum_{i=1}^m b_i (\phi(\xi_i) - \phi(s))^{\rho+r+q-1} \right. \\ &\quad \left. + \frac{\mathcal{B}(q, r)}{\Gamma(r)} (\phi(t) - \phi(0))^r \right] \|T_{z_n}(\cdot) - T_z(\cdot)\|_{\mathcal{C}_{1-q;\phi}}. \end{aligned} \quad (29)$$

Since f is continuous, this implies that T_z is also continuous. Therefore, we have

$$\|T_{z_n} - T_z\|_{\mathcal{C}_{1-q;\phi}} \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

Step 2: F maps bounded sets into bounded sets in $\mathcal{C}_{1-q;\phi}[\mathcal{J}, \mathbb{R}]$.

Indeed, it suffices to show that for any $\kappa > 0$, there exist a $\mu > 0$ such that for any $z \in \mathbf{B}_\kappa = \{z \in \mathcal{C}_{1-q;\phi}[\mathcal{J}, \mathbb{R}] : \|z\| \leq \kappa\}$, thus we have $\|F(z)\|_{\mathcal{C}_{1-q;\phi}} \leq \mu$.

For simplicity, we put

$$E_1 = \frac{|\delta|\Gamma(\rho+q)}{\Gamma(q)\Gamma(\rho+r)} \sum_{i=1}^m b_i \int_{0^+}^{\xi_i} \phi'(s)(\phi(\xi_i) - \phi(s))^{\rho+r-1} |T_z(s)| ds \quad (30)$$

and

$$E_2 = \frac{1}{\Gamma(r)} (\phi(t) - \phi(0))^{1-q} \int_{0^+}^t \phi'(s)(\phi(t) - \phi(s))^{r-1} |T_z(s)| ds. \quad (31)$$

It follows from assumption (A_2) that

$$\begin{aligned} |T_z(t)| &= |f(t, z(t), z(\gamma t), T_z(t))| \\ &\leq k(t) + l(t)|z| + m(t)|z| + n(t)|T_z(t)| \\ &\leq \frac{k^* + (l^* + m^*)|z(t)|}{1 - n^*}. \end{aligned} \quad (32)$$

Thus, in view of Equations (30)–(32), we get

$$\begin{aligned} E_1 &\leq \frac{|\delta|\Gamma(\rho+q)}{\Gamma(q)(1-n^*)} \sum_{i=1}^m b_i \left(\frac{k^*}{\Gamma(\rho+r+1)} (\phi(\xi_i) - \phi(0))^{\rho+r} \right. \\ &\quad \left. + (l^* + m^*) \frac{(\phi(\xi_i) - \phi(0))^{\rho+r+q-1}}{\Gamma(\rho+r)} \mathcal{B}(q, \rho+r) \|z\|_{\mathcal{C}_{1-q;\phi}} \right) \\ E_2 &\leq \frac{1}{(1-n^*)} \left(\frac{k^*}{\Gamma(r+1)} (\phi(T) - \phi(0))^{\rho+r-q+1} \right. \\ &\quad \left. + \frac{(l^* + m^*)\mathcal{B}(q, r)}{\Gamma(r)} (\phi(T) - \phi(0))^r \|z\|_{\mathcal{C}_{1-q;\phi}} \right). \end{aligned}$$

This implies that,

$$\begin{aligned} |(Fz)(t)((\phi(t) - \phi(0))^{q-1})| &\leq \frac{k^*}{(1-n^*)} \left[\frac{|\delta|\Gamma(\rho+q)}{\Gamma(q)\Gamma(\rho+r+1)} \sum_{i=1}^m b_i (\phi(\xi_i) - \phi(0))^{\rho+r} \right. \\ &\quad \left. + \frac{k^*}{\Gamma(r+1)} (\phi(T) - \phi(0))^{\rho+r-q+1} \right] \\ &\quad + \frac{(l^* + m^*)}{(1-n^*)} \left[\frac{|\delta|\Gamma(\rho+q)}{\Gamma(q)\Gamma(\rho+r)} \mathcal{B}(q, \rho+r) \sum_{i=1}^m b_i (\phi(\xi_i) - \phi(0))^{\rho+r+q-1} \right. \\ &\quad \left. + \frac{\mathcal{B}(q, r)}{\Gamma(r)} (\phi(T) - \phi(0))^r \right] \|z\|_{\mathcal{C}_{1-q;\phi}} \\ &= \mu. \end{aligned} \quad (33)$$

Step 3: F maps bounded sets into equicontinuous set of $\mathcal{C}_{1-q;\phi}[\mathcal{J}, \mathbb{R}]$. Let $t_1, t_2 \in \mathcal{J}$ such that $t_1 \geq t_2$ and B_κ be a bounded set of $\mathcal{C}_{1-q;\phi}[\mathcal{J}, \mathbb{R}]$ as defined in Step 2. Let $z \in B_\kappa$, then

$$\begin{aligned} & |((\phi(t_1) - \phi(0))^{q-1})(Fz)(t_1) - ((\phi(t_2) - \phi(0))^{q-1})(Fz)(t_2)| \\ & \leq \left| \frac{1}{\Gamma(r)} (\phi(t_1) - \phi(0))^{1-q} \int_{0^+}^{t_1} \phi'(s) (\phi(t_1) - \phi(s))^{r-1} T_z(s) ds \right. \\ & \quad \left. - \frac{1}{\Gamma(r)} (\phi(t_2) - \phi(0))^{1-q} \int_{0^+}^{t_2} \phi'(s) (\phi(t_2) - \phi(s))^{r-1} T_z(s) ds \right| \\ & \leq \frac{1}{\Gamma(r)} \left| \int_{0^+}^{t_1} \phi'(s) \left[((\phi(t_1) - \phi(0))^{q-1})(\phi(t_1) - \phi(s))^{r-1} \right. \right. \\ & \quad \left. \left. - ((\phi(t_2) - \phi(0))^{q-1})(\phi(t_2) - \phi(s))^{r-1} T_z(s) ds \right] \right| \\ & \quad + \left| \frac{(\phi(t_2) - \phi(0))^{q-1}}{\Gamma(r)} \int_{t_1}^{t_2} \phi'(s) (\phi(t_1) - \phi(0))^{r-1} T_z(s) ds \right| \\ & \rightarrow 0, \quad \text{as } t_1 \rightarrow t_2. \end{aligned} \quad (34)$$

Thus, steps 1–3, together with the Arzela–Ascoli theorem, show that the operator F is completely continuous.

Step 4: a priori bounds.

It is enough to show that the set $\chi = \{z \in \mathcal{C}_{1-q;\phi}[\mathcal{J}, \mathbb{R}] : z = \sigma(Fz), 0 < \sigma < 1\}$ is bounded.

Now, let $z \in \chi$, $z = \sigma(Fz)$ for some $0 < \sigma < 1$. Thus for each $t \in \mathcal{J}$, we obtain

$$\begin{aligned} z(t) = & \sigma \left[\frac{\delta \Gamma(\rho + q)}{\Gamma(q) \Gamma(\rho + r)} (\phi(t) - \phi(0))^{q-1} \sum_{i=1}^m b_i \int_{0^+}^{\xi_i} \phi'(s) (\phi(\xi_i) - \phi(s))^{\rho+r-1} T_z(s) ds \right. \\ & \left. + \frac{1}{\Gamma(r)} \int_{0^+}^t \phi'(s) (\phi(t) - \phi(s))^{r-1} T_z(s) ds \right]. \end{aligned}$$

It follows from assumption (A_2) , that for every $t \in \mathcal{J}$,

$$\begin{aligned} |z(t)(\phi(t) - \phi(0))^{1-q}| & \leq |(Fz)(t)(\phi(t) - \phi(0))^{1-q}| \\ & \leq \frac{k^*}{(1 - n^*)} \left[\frac{|\delta| \Gamma(\rho + q)}{\Gamma(q) \Gamma(\rho + r + 1)} \sum_{i=1}^m b_i (\phi(\xi_i) - \phi(0))^{\rho+r} \right. \\ & \quad \left. + \frac{k^*}{\Gamma(r + 1)} (\phi(T) - \phi(0))^{\rho+r-q+1} \right] \\ & \quad + \frac{(l^* + m^*)}{(1 - n^*)} \left[\frac{|\delta| \Gamma(\rho + q)}{\Gamma(q) \Gamma(\rho + r)} \mathcal{B}(q, \rho + r) \sum_{i=1}^m b_i (\phi(\xi_i) - \phi(0))^{\rho+r-q-1} \right. \\ & \quad \left. + \frac{\mathcal{B}(q, r)}{\Gamma(r)} (\phi(T) - \phi(0))^r \right] \|z\|_{\mathcal{C}_{1-q;\phi}} \\ & < \infty. \end{aligned} \quad (35)$$

This shows that the set χ is bounded. Hence, by the Schaefer's fixed point theorem, problem (5) has at least one solution. $\square$

3.2. Existence Result Via Banach Contraction Principle

Now, we prove the uniqueness of problem (5) by means of Banach contraction principle. Therefore, the following hypotheses are needed.

(A₃) There exist constants $K, L > 0$ such that

$$|f(t, u, v, w) - f(t, \bar{u}, \bar{v}, \bar{w})| \leq K(|u - \bar{u}| + |v - \bar{v}|) + L|w - \bar{w}|$$

for any $u, v, w, \bar{u}, \bar{v}, \bar{w} \in \mathbb{R}$ and $t \in \mathcal{J}$.

(A₄) Suppose that

$$\left(\frac{2K}{1-L} \right) \Omega < 1,$$

where

$$\Omega = \frac{|\delta|\Gamma(\rho+q)}{\Gamma(q)\Gamma(\rho+r)} \mathcal{B}(q, \rho+r) \sum_{i=1}^m b_i (\phi(\xi_i) - \phi(0))^{\rho+r+q-1} + \frac{\mathcal{B}(q, r)}{\Gamma(r)} (\phi(T) - \phi(0))^r. \quad (36)$$

Theorem 4. Let $0 < r < 1$, $0 \leq p \leq 1$ and $q = r + p - rp$. Suppose that the hypotheses (A₁), (A₃) and (A₄) are satisfied. Then, problem (5) has a unique solution in the space $C_{1-q;\phi}^{r,p}[\mathcal{J}, \mathbb{R}]$.

Proof. Define the operator $F : C_{1-q;\phi}[\mathcal{J}, \mathbb{R}] \rightarrow C_{1-q;\phi}[\mathcal{J}, \mathbb{R}]$ by

$$\begin{aligned} (Fz)(t) &= \frac{\delta\Gamma(\rho+q)}{\Gamma(q)\Gamma(\rho+r)} (\phi(t) - \phi(0))^{q-1} \sum_{i=1}^m b_i \int_{0^+}^{\xi_i} \phi'(s) (\phi(\xi_i) - \phi(s))^{\rho+r-1} T_z(s) ds \\ &\quad + \frac{1}{\Gamma(r)} \int_{0^+}^t \phi'(s) (\phi(t) - \phi(s))^{r-1} T_z(s) ds, \end{aligned} \quad (37)$$

then, clearly the operator F is well-defined. Let $z_1, z_2 \in C_{1-q;\phi}^{r,p}[\mathcal{J}, \mathbb{R}]$ and $t \in \mathcal{J}$, then, we have

$$\begin{aligned} &|((Fz_1)(t) - (Fz_2)(t))(\phi(t) - \phi(0))^{1-q}| \\ &\leq \frac{|\delta|\Gamma(\rho+q)}{\Gamma(q)\Gamma(\rho+r)} \sum_{i=1}^m b_i \int_{0^+}^{\xi_i} \phi'(s) (\phi(\xi_i) - \phi(s))^{\rho+r-1} |T_{z_1}(s) - T_{z_2}(s)| ds \\ &\quad + \frac{1}{\Gamma(r)} (\phi(t) - \phi(0))^{1-q} \int_{0^+}^t \phi'(s) (\phi(t) - \phi(s))^{r-1} |T_{z_1}(s) - T_{z_2}(s)| ds \end{aligned} \quad (38)$$

and

$$\begin{aligned} |T_{z_1}(t) - T_{z_2}(t)| &= |f(t, z_1(t), z_1(\gamma t), T_{z_1}(t) - f(t, z_2(t), z_2(\gamma t), T_{z_2}(t)))| \\ &\leq K(|z_1(t) - z_2(t)| + |z_1(\gamma t) - z_2(\gamma t)|) + L|(T_{z_1})(t) - (T_{z_2})(t)| \\ &\leq \left(\frac{2K}{1-L} \right) |z_1(t) - z_2(t)|. \end{aligned} \quad (39)$$

Thus, by substituting Equation (39) in Equation (38), we obtain

$$\begin{aligned} &|((Fz_1)(t) - (Fz_2)(t))(\phi(t) - \phi(0))^{1-q}| \\ &\leq \frac{|\delta|\Gamma(\rho+q)}{\Gamma(q)\Gamma(\rho+r)} \sum_{i=1}^m b_i \left(\frac{2K}{(1-L)} \int_{0^+}^{\xi_i} \phi'(s) (\phi(\xi_i) - \phi(s))^{\rho+r-1} ds \right) \|z_1(t) - z_2(t)\|_{C_{1-q;\phi}} \\ &\quad + \frac{1}{\Gamma(r)} (\phi(t) - \phi(0))^{1-q} \frac{2K}{(1-L)} \left(\int_{0^+}^t \phi'(s) (\phi(t) - \phi(s))^{r-1} ds \right) \|z_1(t) - z_2(t)\|_{C_{1-q;\phi}} \\ &\leq \frac{2K}{(1-L)} \left(\frac{|\delta|\Gamma(\rho+q)}{\Gamma(q)\Gamma(\rho+r)} \mathcal{B}(q, \rho+r) \sum_{i=1}^m b_i (\phi(\xi_i) - \phi(0))^{\rho+r+q-1} \right. \\ &\quad \left. + \frac{\mathcal{B}(q, r)}{\Gamma(r)} (\phi(T) - \phi(0))^r \right) \|z_1(t) - z_2(t)\|_{C_{1-q;\phi}}. \end{aligned} \quad (40)$$

Also,

$$\begin{aligned} \|(Fz_1) - (Fz_2)\|_{C_{1-q;\phi}} &\leq \frac{2K}{(1-L)} \left(\frac{|\delta|\Gamma(\rho+q)}{\Gamma(q)\Gamma(\rho+r)} \mathcal{B}(q, \rho+r) \sum_{i=1}^m b_i(\phi(\xi_i) - \phi(0))^{\rho+r+q-1} \right. \\ &\quad \left. + \frac{\mathcal{B}(q, r)}{\Gamma(r)} (\phi(T) - \phi(0))^r \right) \|z_1(t) - z_2(t)\|_{C_{1-q;\phi}}. \end{aligned} \quad (41)$$

It follows from hypotheses (A_4) that F is a contraction map. Therefore, by Banach contraction principle, we can conclude that problem (5) has a unique solution. $\square$

4. Ulam-Hyers Stability

Two types of Ulam stability for (5) are discussed in this section, namely Ulam-Hyers and generalized Ulam-Hyers stability.

Definition 4. Problem (5) is said to be Ulam-Hyers stable if there exists $\omega \in \mathbb{R}_+ \setminus \{0\}$, such that for each $\epsilon > 0$ and solution $x \in \mathcal{C}_{1-q;\phi}^1[\mathcal{J}, \mathbb{R}]$ of the inequality

$$|{}^H\mathcal{D}_{0+}^{r,p;\phi}x(t) - f(t, x(t), x(\gamma t), {}^H\mathcal{D}_{0+}^{r,p;\phi}x(\gamma t))| \leq \epsilon, \quad t \in \mathcal{J}, \quad (42)$$

there exists a solution $z \in \mathcal{C}_{1-q;\phi}^1[\mathcal{J}, \mathbb{R}]$ of equation (5), such that

$$|x(t) - z(t)| \leq \omega\epsilon, \quad t \in \mathcal{J}.$$

Definition 5. Problem (5) is said to be generalized Ulam-Hyers stable if there exist $\Phi \in \mathcal{C}(\mathbb{R}_+, \mathbb{R}_+)$, $\Phi_f(0) = 0$, such that for each solution $x \in \mathcal{C}_{1-q;\phi}^1[\mathcal{J}, \mathbb{R}]$ of the (42), there exists a solution $z \in \mathcal{C}_{1-q;\phi}^1[\mathcal{J}, \mathbb{R}]$ of Equation (5), such that

$$|x(t) - z(t)| \leq \Phi_f\epsilon, \quad t \in \mathcal{J}.$$

Remark 1. A function $x \in \mathcal{C}_{1-q;\phi}[\mathcal{J}, \mathbb{R}]$ is a solution of the inequality (42), if and only if there exist a function $g \in \mathcal{C}_{1-q;\phi}[\mathcal{J}, \mathbb{R}]$ such that:

$$(i) \quad |g(t)| \leq \epsilon, \quad t \in \mathcal{J}.$$

$$(ii) \quad {}^H\mathcal{D}_{0+}^{r,p;\phi}x(t) = f(t, x(t), x(\gamma t), {}^H\mathcal{D}_{0+}^{r,p;\phi}x(\gamma t)) + g(t), \quad t \in \mathcal{J}.$$

Lemma 9. Let $0 < r < 1$, $0 \leq p \leq 1$, if a function $x \in \mathcal{C}_{1-q;\phi}[\mathcal{J}, \mathbb{R}]$ is a solution of the inequality (42), then x is a solution of the following integral inequality

$$\left| x(t) - A_x - \frac{1}{\Gamma(r)} \int_{0+}^t \phi'(s)(\phi(t) - \phi(s))^{r-1} T_x ds \right| \leq \Omega\epsilon. \quad (43)$$

Proof. Clearly it follow from Remark 1 that

$$\begin{aligned} {}^H\mathcal{D}_{0+}^{r,p;\phi}x(t) &= f(t, x(t), x(\gamma t), {}^H\mathcal{D}_{0+}^{r,p;\phi}x(\gamma t)) + g(t) \\ &= T_x(t) + g(t), \end{aligned}$$

and

$$\begin{aligned} x(t) &= \frac{\delta\Gamma(\rho+q)}{\Gamma(q)\Gamma(\rho+r)}(\phi(t)-\phi(0))^{q-1} \sum_{i=1}^m b_i \left(\int_{0+}^{\xi_i} \phi'(s)(\phi(\xi_i)-\phi(s))^{\rho+r-1} T_x(s) ds \right. \\ &\quad \left. + \int_{0+}^{\xi_i} \phi'(s)(\phi(\xi_i)-\phi(s))^{\rho+r-1} g(s) ds \right) + \frac{1}{\Gamma(r)} \int_{0+}^t \phi'(s)(\phi(t)-\phi(s))^{r-1} T_x(s) ds \\ &\quad + \frac{1}{\Gamma(r)} \int_{0+}^t \phi'(s)(\phi(t)-\phi(s))^{r-1} g(s) ds. \end{aligned} \quad (44)$$

Hence

$$\begin{aligned} &\left| x(t) - A_x - \frac{1}{\Gamma(r)} \int_{0+}^t \phi'(s)(\phi(t)-\phi(s))^{r-1} T_x(s) ds \right| \\ &= \left| \frac{\delta\Gamma(\rho+q)}{\Gamma(q)\Gamma(\rho+r)}(\phi(t)-\phi(0))^{q-1} \sum_{i=1}^m b_i \int_{0+}^{\xi_i} \phi'(s)(\phi(\xi_i)-\phi(s))^{\rho+r-1} g(s) ds \right. \\ &\quad \left. + \frac{1}{\Gamma(r)} \int_{0+}^t \phi'(s)(\phi(t)-\phi(s))^{r-1} g(s) ds \right| \\ &\leq \frac{|\delta|\Gamma(\rho+q)}{\Gamma(q)\Gamma(\rho+r)}(\phi(t)-\phi(0))^{q-1} \sum_{i=1}^m |b_i| \int_{0+}^{\xi_i} \phi'(s)(\phi(\xi_i)-\phi(s))^{\rho+r-1} |g(s)| ds \\ &\quad + \frac{1}{\Gamma(r)} \int_{0+}^t \phi'(s)(\phi(t)-\phi(s))^{r-1} |g(s)| ds \\ &\leq \Omega\epsilon. \end{aligned} \quad (45)$$

□

Theorem 5. Suppose that the hypotheses (A_1) , (A_3) and (A_4) are satisfied. Then problem (5) is both Ulam-Hyers and generalized Ulam-Hyers stable on $\mathcal{J}$.

Proof. Let $\epsilon > 0$ and $x \in \mathcal{C}_{1-q;\phi}[\mathcal{J}, \mathbb{R}]$ be a function which satisfies the inequality (42) and let $z \in \mathcal{C}_{1-q;\phi}[\mathcal{J}, \mathbb{R}]$ be a unique solution of the following implicit fractional pantograph differential equation

$$\begin{aligned} {}^H\mathcal{D}_{0+}^{r,p;\phi} z(t) &= f(t, z(t), z(\gamma t), {}^H\mathcal{D}_{0+}^{r,p;\phi} z(\gamma t)) \leq \epsilon, \quad t \in \mathcal{J}, \quad 0 < r < 1, 0 \leq p \leq 1, \\ \mathcal{I}_{0+}^{1-q;\phi} z(0+) &= \mathcal{I}_{0+}^{1-q;\phi} z(0+) = \sum_{i=1}^m b_i \mathcal{I}_{0+}^{\rho;\phi} z(\xi_i), \quad \xi_i \in (0, T], \quad q = r + p - rp. \end{aligned}$$

Using Lemma 9, we have

$$z(t) = A_z - \frac{1}{\Gamma(r)} \int_{0+}^t \phi'(s)(\phi(t)-\phi(s))^{r-1} T_z(s) ds,$$

where

$$A_z = \frac{\delta\Gamma(\rho+q)}{\Gamma(q)\Gamma(\rho+r)}(\phi(t)-\phi(a))^{q-1} \sum_{i=1}^m b_i \int_{0+}^{\xi_i} \phi'(s)(\phi(\xi_i)-\phi(s))^{\rho+r-1} T_z(s) ds.$$

Clearly, if $z(\xi_i) = x(\xi_i)$ and $\mathcal{I}_{0+}^{1-q;\phi} z(0^+) = \mathcal{I}_{0+}^{1-q;\phi} z(0^+)$, we get $A_z = A_x$ and that

$$\begin{aligned} & |A_z - A_x| \\ &= \frac{|\delta|\Gamma(\rho+q)}{\Gamma(q)\Gamma(\rho+r)} (\phi(t) - \phi(0))^{q-1} \sum_{i=1}^m b_i \int_{0^+}^{\xi_i} \phi'(s) (\phi(\xi_i) - \phi(s))^{\rho+r-1} |T_z(s) - T_x(s)| ds \\ &\leq \frac{|\delta|\Gamma(\rho+q)}{\Gamma(q)\Gamma(\rho+r)} (\phi(t) - \phi(0))^{q-1} \left(\frac{2K}{1-L} \right) \sum_{i=1}^m b_i \mathcal{I}_{0+}^{\rho+r;\phi} |z(s) - x(s)|(\xi_i) \\ &= 0. \end{aligned}$$

Now for any $t \in \mathcal{J}$ and Lemma 9, we have

$$\begin{aligned} |x(t) - z(t)| &= \left| x(t) - A_x - \frac{1}{\Gamma(r)} \int_{0^+}^t \phi'(s) (\phi(t) - \phi(s))^{r-1} T_x(s) ds \right| \\ &\quad + \frac{1}{\Gamma(r)} \int_{0^+}^t \phi'(s) (\phi(t) - \phi(s))^{r-1} |T_x(s) - T_z(s)| ds \\ &\leq \left| x(t) - A_x - \frac{1}{\Gamma(r)} \int_{0^+}^t \phi'(s) (\phi(t) - \phi(s))^{r-1} T_x(s) ds \right| \\ &\quad + \left(\frac{2K}{1-L} \right) \frac{1}{\Gamma(r)} \int_{0^+}^t \phi'(s) (\phi(t) - \phi(s))^{r-1} |x(s) - z(s)| ds \\ &\leq \Omega\epsilon + \left(\frac{2K}{1-L} \right) \frac{\mathcal{B}(r,q)(\phi(T) - \phi(0))^r}{\Gamma(r)} |x(t) - z(t)| ds. \end{aligned}$$

Thus,

$$|x(t) - z(t)| \leq \omega\epsilon,$$

where

$$\omega = \frac{\Omega(1-L)\Gamma(r)}{(1-L)\Gamma(r) - 2K(\phi(T) - \phi(0))^r \mathcal{B}(r,q)}.$$

Therefore, problem (5) is Ulam-Hyers stable. Moreover, if we set $\Phi_f(\epsilon) = \omega\epsilon$ such that $\Phi_f(0) = 0$, then problem (5) is generalized Ulam-Hyers stable. $\square$

5. Examples

Example 1. Consider the implicit fractional pantograph differential equation which involves ϕ -Hilfer fractional derivative of the following form:

$$\begin{cases} {}^H\mathcal{D}_{0+}^{\frac{2}{3}, \frac{1}{2}; t} z(t) = \frac{1}{3(5^{2t}+5)[1+|z(t)|+|z(\frac{1}{2}t)|+|{}^H\mathcal{D}_{1+}^{\frac{2}{3}, \frac{1}{2}; t} z(\frac{1}{2}t)|]}, & t \in \mathcal{J} = (0, 2], \\ \mathcal{I}_{0+}^{1-\frac{5}{6}; t} z(0) = 3\mathcal{I}_{0+}^{\frac{1}{2}; t} z(\frac{3}{2}), & \frac{2}{3} \leq \frac{5}{6} = \frac{2}{3} + (\frac{1}{2}) - (\frac{2}{3})(\frac{1}{2}). \end{cases} \quad (46)$$

By comparing (5) with (46), we have:

$r = \frac{2}{3}$, $p = \gamma = \rho = \frac{1}{2}$, $q = \frac{5}{6}$, $T = 2$ and $\phi(\cdot) = t$. Also from the initial condition we can easily see that $b_1 = 3$ since $m = 1$, $\xi_1 = \frac{3}{2} \in \mathcal{J}$ and $f : \mathcal{J} \times \mathbb{R}^3 \rightarrow \mathbb{R}$ is a function defined by

$$f(t, u, v, w) = \frac{1}{3(5^{2t}+5)(1+|u|+|v|+|w|)}, \quad t \in \mathcal{J}, \quad u, v, w \in \mathbb{R}_+.$$

Obviously, f is continuous and for all $u, v, w, \bar{u}, \bar{v}, \bar{w} \in \mathbb{R}_+$ and $t \in \mathcal{J}$, we have $|f(t, u, v, w) - f(t, \bar{u}, \bar{v}, \bar{w})| \leq \frac{1}{90} (|u - \bar{u}| + |v - \bar{v}| + |w - \bar{w}|)$. Thus, it follows that conditions (A_1) and (A_3) are true with $K = L = \frac{1}{90}$. Therefore, by simple calculation, we get $|\delta| \approx 0.3935$ and

$$\left(\frac{2K}{1-L} \right) \Omega \approx 0.0642 < 1.$$

Since, all the assumptions of Theorem 4 are satisfied. Then problem (5) has a unique solution on $\mathcal{J}$. However, we can also find out that $\Omega \approx 2.8551 > 0$ and $\omega = 2.9321 > 0$. Hence, by Theorem 5, problem (5) is both Ulam-Hyers and also generalized Ulam-Hyers stable.

Example 2. Consider the implicit fractional pantograph differential equation which involves ϕ -Hilfer fractional derivative of form:

$$\begin{cases} {}^H\mathcal{D}_{0+}^{\frac{1}{3}, \frac{2}{3}; \sqrt{t}} z(t) = \frac{2 + |z(t)| + |z(\frac{3}{2}t)| + |{}^H\mathcal{D}_{0+}^{\frac{1}{3}, \frac{2}{3}; \sqrt{t}} z(\frac{3}{2}t)|}{95e^{2t} \cos 2t \left(1 + |z(t)| + |z(\frac{3}{2}t)| + |{}^H\mathcal{D}_{0+}^{\frac{1}{3}, \frac{2}{3}; \sqrt{t}} z(\frac{3}{2}t)| \right)}, & t \in \mathcal{J} = (0, 1], \\ \mathcal{I}_{0+}^{1-\frac{7}{9}; \sqrt{t}} z(0) = z(\frac{1}{2}) + 3z(\frac{4}{5}), & \frac{2}{3} < \frac{7}{9} = \frac{1}{3} + (\frac{2}{3}) - (\frac{1}{3})(\frac{2}{3}). \end{cases} \quad (47)$$

By comparing Equation (47) with Equation (5), we obtain that:

$r = \frac{1}{3}$, $p = \frac{2}{3}$, $q = \frac{7}{9}$, $\rho = 0$, $\gamma = \frac{3}{2}$, $T = 1$ and $\phi(\cdot) = \sqrt{t}$. Also we can easily see that $b_1 = 1$, $b_2 = 3$ since $m = 2$, $\xi_1 = \frac{1}{2}$, $\xi_2 = \frac{4}{5} \in \mathcal{J}$ and $f: \mathcal{J} \times \mathbb{R}^3 \rightarrow \mathbb{R}$ is a function defined by

$$f(t, u, v, w) = \frac{2 + |u| + |v| + |w|}{95e^{2t} \cos 2t (1 + |u| + |v| + |w|)}, \quad t \in \mathcal{J}, \quad u, v, w \in \mathbb{R}_+.$$

Thus, f is continuous and we can see that, for all $u, v, w, \bar{u}, \bar{v}, \bar{w} \in \mathbb{R}_+$ and $t \in \mathcal{J}$,

$$|f(t, u, v, w) - f(t, \bar{u}, \bar{v}, \bar{w})| \leq \frac{1}{95} (|u - \bar{u}| + |v - \bar{v}| + |w - \bar{w}|).$$

So assumptions (A_1) and (A_3) are fulfilled with $K = L = \frac{1}{95}$. Furthermore,

$$|f(t, u, v, w)| \leq \frac{1}{95e^{2t} \cos 2t} (2 + |u| + |v| + |w|), \quad t \in \mathcal{J}.$$

The above implies that (A_2) is true with $k(t) = \frac{2}{95e^{2t} \cos 2t}$, $l(t) = m(t) = n(t) = \frac{1}{95e^{2t} \cos 2t}$ and $k^* = \frac{2}{95}$, $l^* = m^* = n^* = \frac{1}{95}$. Therefore, all the hypotheses of Theorem 4 are satisfied, which means that problem (5) has at least one solution on $\mathcal{J}$. Moreover, by using the same procedure as in example 5.2, we obtain, that $|\delta| \approx 1.1025$, $\Omega \approx 3.6662 > 0$ and

$$\left(\frac{2K}{1-L} \right) \Omega \approx 0.0782 < 1.$$

Thus, all the hypotheses of Theorem 4 holds. Hence, problem (5) has a unique solution on $\mathcal{J}$.

Example 3. Consider the implicit fractional pantograph differential equation which involves ϕ -Hilfer fractional derivative of the following form:

$$\begin{cases} {}^H\mathcal{D}_{0+}^{\frac{1}{2}, \frac{1}{3}; t} z(t) = \frac{1}{4^{t+3} [1 + |z(t)| + |z(\frac{1}{6}t)| + |{}^H\mathcal{D}_{0+}^{\frac{1}{2}, \frac{1}{3}; t} z(\frac{1}{6}t)|]}, & t \in \mathcal{J} = (0, 3], \\ \mathcal{I}_{0+}^{1-q; t} z(0) = \sqrt{2}\mathcal{I}_{0+}^{\frac{2}{5}; t} z(2) + \sqrt{5}\mathcal{I}_{0+}^{\frac{2}{5}; t} z(\frac{5}{2}), & q = \frac{1}{2} + (\frac{1}{3}) - (\frac{1}{2})(\frac{1}{3}). \end{cases} \quad (48)$$

By comparing Equation (5) with Equation (48), we get the followings values:

$r = \frac{1}{2}$, $p = \frac{1}{3}$, $\gamma = \frac{1}{6}$, $\rho = \frac{2}{5}$, $q = \frac{2}{3}$, $T = 3$ and $\phi(\cdot) = t$. Also from the initial condition we can easily see that $b_1 = \sqrt{2}$, $b_1 = \sqrt{5}$ since $m = 2$, $\xi_1 = 2$, $\xi_2 = \frac{5}{2}$ and $f : \mathcal{J} \times \mathbb{R}^3 \rightarrow \mathbb{R}$ is a function defined by

$$f(t, u, v, w) = \frac{1}{4^{t+3}(1 + |u| + |v| + |w|)}, t \in \mathcal{J}, \quad u, v, w \in \mathbb{R}_+.$$

Thus, f is continuous and for all $u, v, w, \bar{u}, \bar{v}, \bar{w} \in \mathbb{R}_+$ and $t \in \mathcal{J}$, yields

$|f(t, u, v, w) - f(t, \bar{u}, \bar{v}, \bar{w})| \leq \frac{1}{64} (|u - \bar{u}| + |v - \bar{v}| + |w - \bar{w}|)$. Hence, it follows that conditions (A_1) and (A_3) are true with $K = L = \frac{1}{90}$. Therefore, by substitution these values, we get $|\delta| \approx 0.3456$, $\Omega \approx 7.4535 > 0$ and

$$\left(\frac{2K}{1-L} \right) \Omega \approx 0.2366 < 1,$$

which implies that, all the assumptions of Theorem 4 are satisfied. Thus, problem (5) has a unique solution on $\mathcal{J}$.

6. Conclusions

In our study, Firstly, we established the equivalence between problem (5) and the Volterra integral equation. Secondly, Banach and Schaefer's fixed point theorems were used to establish the existence and uniqueness solutions for implicit fractional pantograph differential equation which involves ϕ -Hilfer fractional derivatives. Based on ϕ -Hilfer fractional derivatives, we found that the stability of Ulam-Hyers and generalized Ulam-Hyers allowed on the implicit fractional pantograph differential equation, supplemented with a nonlocal Riemann-Liouville condition. In addition, examples were given to illustrate our main results. Moreover, it worthy to mention the following remarks:

- If $\rho \rightarrow 0$ and $\phi(t) = t$, we obtain the results of [48] and [52]. Furthermore, if $\rho \rightarrow 0$ we obtain the Ulam-Hyers and generalized Ulam-Hyers stability for the implicit fractional pantograph differential equations with ϕ -Hilfer fractional derivatives [52,58] and if $q = 0$ we obtain [51].
- If $\rho \rightarrow 1$, the nonlocal Riemann-Liouville integral condition reduces to a nonlocal integral condition which plays an important role in computational fluid dynamics, ill-posed problems and mathematical models [62].
- If $\rho \rightarrow 0$, the initial condition reduces to multi-point nonlocal condition.
- If $t \in [a, b]$ as defined in paper [58], the function $f(t, x(t), x(\lambda t))$ is not well-defined for some choice of $0 < \lambda < 1$. Thus, our results modify and improve the above cited remarks and can be considered as the development of the qualitative analysis of fractional differential equations. The study of Ulam-Hyers stability in the frame of ϕ -Hilfer fractional derivative with a generalized nonlocal boundary condition proposed in this paper and other coupled system will be presented in the near future.

Author Contributions: The authors contributed equally in writing this article. All authors have read and agreed to the published version of the manuscript.

Funding: Petchra Pra Jom Klao Doctoral Scholarship for Ph.D. program of King Mongkut's University of Technology Thonburi (KMUTT). The Center of Excellence in Theoretical and Computational Science (TaCS-CoE), KMUTT. Center of Excellence in Theoretical and Computational Science (TaCS-CoE), KMUTT. King Mongkut's University of Technology North Bangkok, Contract no. KMUTNB-63-KNOW-033.

Acknowledgments: The authors acknowledge the financial support provided by the Center of Excellence in Theoretical and Computational Science (TaCS-CoE), KMUTT. The first author was supported by the "Petchra Pra Jom Klao Ph.D. Research Scholarship from King Mongkut's University of Technology Thonburi" (Grant No. 13/2561). Moreover, this research work was financially supported by King Mongkut's University of Technology Thonburi through the KMUTT 55th Anniversary Commemorative Fund.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

References

- Kilbas, A.A.; Srivastava, H.M.; Trujillo, J.J. *Theory and Applications of fractional derivatival Equations*; Elsevier Science Limited: Amsterdam, The Netherlands, 2006; Volume 204.
- Samko, S.G.; Kilbas, A.A.; Marichev, O.I.; *Fractional Integrals and Derivatives*; Gordon and Breach Science Publishers: Yverdon Yverdon-les-Bains, Switzerland, 1993; Volume 1993.
- Podlubny, I. *Fractional Differential Equations: An Introduction to Fractional Derivatives, Fractional Differential Equations, to Methods of Their Solution and Some of Their Applications*; Elsevier: Amsterdam, The Netherlands, 1998.
- Mainardi, F. *Fractional Calculus and Waves in Linear Viscoelasticity: An Introduction to Mathematical Models*; World Scientific: Singapore, 2010.
- Furati, K.; Kassim, M.; Tata, N. Existence and uniqueness for a problem involving Hilfer fractional derivative. *Comput. Math. Appl.* **2012**, *64*, 1616–1626.
- Oliveira, D.S.; de Oliveira, E.C. Hilfer–Katugampola fractional derivatives. *Comput. Appl. Math.* **2018**, *37*, 3672–3690.
- Osler, T.J. The fractional derivative of a composite function. *SIAM J. Math. Anal.* **1970**, *1*, 288–293.
- Gambo, Y.Y.; Jarad, F.; Baleanu, D.; Abdeljawad, T. On Caputo modification of the Hadamard fractional derivatives. *Adv. Differ. Equ.* **2014**, *2014*, 10.
- Jarad, F.; Abdeljawad, T.; Baleanu, D. Caputo-type modification of the Hadamard fractional derivatives. *Adv. Differ. Equ.* **2012**, *2012*, 142.
- Jarad, F.; Abdeljawad, T.; Baleanu, D. On the generalized fractional derivatives and their Caputo modification. *J. Nonlinear Sci. Appl.* **2017**, *10*, 2607–2619.
- Marin, M.; Baleanu, D.; Vlase, S. Effect of microtemperatures for micropolar thermoelastic bodies. *Struct. Eng. Mech.* **2017**, *61*, 381–387.
- Gladkov, S.; Bogdanova, S. On the question of the magnetic susceptibility of fractal ferromagnetic wires. *Russ. Phys. J.* **2014**, *57*, 469–473.
- Hilfer, R. Fractional calculus and regular variation in thermodynamics. In *Applications of Fractional Calculus in Physics*; World Scientific: Singapore, 2000; pp. 429–463.
- Hilfer, R. Fractional time evolution. In *Applications of Fractional Calculus in Physics*; World Scientific: Singapore, 2000; pp. 87–130.
- Gerolymatou, E.; Vardoulakis, I.; Hilfer, R. Modelling infiltration by means of a nonlinear fractional diffusion model. *J. Phys. Appl. Phys.* **2006**, *39*, 4104.
- Vivek, D.; Kanagarajan, K.; Sivasundaram, S. Dynamics and stability of pantograph equations via Hilfer fractional derivative. *Nonlinear Stud.* **2016**, *23*, 685–698.
- Vivek, D.; Kanagarajan, K.; Harikrishnan, S. Analytic study on nonlocal initial value problems for pantograph equations with Hilfer–Hadamard fractional derivative. *Int. J. Math. Its Appl.* **2018**, *55*, 7.
- Abdo, M.S.; Panchal, S.K.; Bhairat, S.P. Existence of solution for Hilfer fractional differential equations with boundary value conditions. *arXiv* **2019**, arXiv:1909.13680.
- Sousa, J.V.d.C.; de Oliveira, E.C. On the ψ -Hilfer fractional derivative. *Commun. Nonlinear Sci. Numer. Simul.* **2018**, *60*, 72–91.
- Ravichandran, C.; Logeswari, K.; Jarad, F. New results on existence in the framework of Atangana–Baleanu derivative for fractional integro-differential equations. *Chaos Solitons Fractals* **2019**, *125*, 194–200.
- Abdo, M.S.; Panchal, S.K. Fractional integro-differential equations involving ψ -Hilfer fractional derivative. *Adv. Appl. Math. Mech.* **2019**, *11*, 1–22.
- Almeida, R. A Caputo fractional derivative of a function with respect to another function. *Commun. Nonlinear Sci. Numer. Simul.* **2017**, *44*, 460–481.
- Ahmad, B.; Nieto, J.J. Existence results for nonlinear boundary value problems of fractional integrodifferential equations with integral boundary conditions. *Bound. Value Probl.* **2009**, *2009*, 708576.
- Ahmad, B.; Sivasundaram, S. Existence of solutions for impulsive integral boundary value problems of fractional order. *Nonlinear Anal. Hybrid Syst.* **2010**, *4*, 134–141.
- Wang, G.; Ghanmi, A.; Horrigue, S.; Madian, S. Existence Result and Uniqueness for Some Fractional Problem. *Mathematics* **2019**, *7*, 516.
- Ali, K.B.; Ghanmi, A.; Kefi, K. Existence of solutions for fractional differential equations with Dirichlet boundary conditions. *Electron. J. Differ. Equ.* **2016**, *2016*, 1–11.

27. Nieto, J.; Ouahab, A.; Venkatesh, V. Implicit fractional differential equations via the Liouville–Caputo derivative. *Mathematics* **2015**, *3*, 398–411.
28. Zhang, W.; Liu, W.; Xue, T. Existence and uniqueness results for the coupled systems of implicit fractional differential equations with periodic boundary conditions. *Adv. Differ. Equ.* **2018**, *2018*, 413.
29. Staněk, S. Existence results for implicit fractional differential equations with nonlocal boundary conditions. *Mem. Differ. Equations Math. Phys.* **2017**, *72*, 119–130.
30. Srivastava, H. Some families of Mittag-Leffler type functions and associated operators of fractional calculus (Survey). *Turk. World Math. Soc. J. Pure Appl. Math.* **2016**, *7*, 123–145.
31. Benchohra, M.; Bouriah, S.; Nieto, J.J. Existence of periodic solutions for nonlinear implicit Hadamard’s fractional differential equations. *Revista de la Real Academia de Ciencias Exactas Físicas y Naturales. Serie A Matemáticas* **2018**, *112*, 25–35.
32. Srivastava, H.; El-Sayed, A.; Gaafar, F. A class of nonlinear boundary value problems for an arbitrary fractional-order differential equation with the Riemann-Stieltjes functional integral and infinite-point boundary conditions. *Symmetry* **2018**, *10*, 508.
33. Borisut, P.; Kumam, P.; Ahmed, I.; Sitthithakerngkiet, K. Nonlinear Caputo Fractional Derivative with Nonlocal Riemann-Liouville Fractional Integral Condition Via Fixed Point Theorems. *Symmetry* **2019**, *11*, 829.
34. Živorad Tomovski.; Hilfer, R.; Srivastava, H.M. Fractional and operational calculus with generalized fractional derivative operators and Mittag-Leffler type functions. *Integral Transform. Spec. Funct.* **2010**, *21*, 797–814.
35. Kharade, J.P.; Kucche, K.D. On the Impulsive Implicit ψ -Hilfer Fractional Differential Equations with Delay. *arXiv* **2019**, arXiv:1908.07793.
36. Hyers, D.H. On the stability of the linear functional equation. *Proc. Natl. Acad. Sci. USA* **1941**, *27*, 222.
37. Ulam, S.M. *Problems in Modern Mathematics*; Courier Corporation: North Chelmsford, MA, USA, 2004.
38. Ulam, S.M. *A Collection of Mathematical Problems*; Interscience Publishers: Geneva, Switzerland, 1960; Volume 8.
39. Aoki, T. On the stability of the linear transformation in Banach spaces. *J. Math. Soc. Jpn.* **1950**, *2*, 64–66.
40. Rassias, T.M. On the stability of the linear mapping in Banach spaces. *Proc. Am. Math. Soc.* **1978**, *72*, 297–300.
41. Abbas, S.; Benchohra, M.; Lagreg, J.; Alsaedi, A.; Zhou, Y. Existence and Ulam stability for fractional differential equations of Hilfer-Hadamard type. *Adv. Differ. Equ.* **2017**, *2017*, 180.
42. Wang, J.; Lin, Z. Ulam’s type stability of Hadamard type fractional integral equations. *Filomat* **2014**, *28*, 1323–1331.
43. de Oliveira, E.C.; Sousa, J.V.d.C. Ulam-Hyers-Rassias stability for a class of fractional integro-differential equations. *Results Math.* **2018**, *73*, 111.
44. Sousa, J.V.d.C.; Oliveira, D.d.S.; Capelas de Oliveira, E. On the existence and stability for noninstantaneous impulsive fractional integrodifferential equation. *Math. Methods Appl. Sci.* **2019**, *42*, 1249–1261.
45. Liu, K.; Wang, J.; O’Regan, D. Ulam-Hyers–Mittag-Leffler stability for ψ -Hilfer fractional-order delay differential equations. *Adv. Differ. Equ.* **2019**, *2019*, 50.
46. Liu, K.; Fečkan, M.; O’Regan, D.; Wang, J. Hyers-Ulam Stability and Existence of Solutions for Differential Equations with Caputo-Fabrizio Fractional Derivative. *Mathematics* **2019**, *7*, 333.
47. Sousa, J.V.d.C.; de Oliveira, E.C. On the Ulam-Hyers-Rassias stability for nonlinear fractional differential equations using the ψ -Hilfer operator. *J. Fixed Point Theory Appl.* **2018**, *20*, 96.
48. Vivek, D.; Kanagarajan, K.; Elsayed, E. Some existence and stability results for Hilfer-fractional implicit differential equations with nonlocal conditions. *Mediterr. J. Math.* **2018**, *15*, 15.
49. Ahmed, H.M.; El-Borai, M.M.; El-Owaidey, H.M.; Ghanem, A.S. Impulsive Hilfer fractional differential equations. *Adv. Differ. Equations* **2018**, *2018*, 226.
50. Ockendon, J.R.; Tayler, A.B. The dynamics of a current collection system for an electric locomotive. *Proc. R. Soc. London. Math. Phys. Sci.* **1971**, *322*, 447–468.
51. Balachandran, K.; Kiruthika, S.; Trujillo, J. Existence of solutions of nonlinear fractional pantograph equations. *Acta Math. Sci.* **2013**, *33*, 712–720.
52. Vivek, D.; Kanagarajan, K.; Sivasundaram, S. Theory and analysis of nonlinear neutral pantograph equations via Hilfer fractional derivative. *Nonlinear Stud.* **2017**, *24*, 699–712.
53. Anguraj, A.; Vinodkumar, A.; Malar, K. Existence and stability results for random impulsive fractional pantograph equations. *Filomat* **2016**, *30*, 3839–3854.

54. Bhalekar, S.; Patade, J. Series Solution of the Pantograph Equation and Its Properties. *Fractal Fract.* **2017**, *1*, 16.
55. Shah, K.; Vivek, D.; Kanagarajan, K. Dynamics and Stability of ψ -fractional Pantograph Equations with Boundary Conditions. *Boletim da Sociedade Paranaense de Matemática* **2018**, *22*, 1–13.
56. Elsayed, E.M.; Harikrishnan, S.; Kanagarajan, K. Analysis of nonlinear neutral pantograph differential equations with Hilfer fractional derivative. *MathLAB* **2018**, *1*, 231–240.
57. Harikrishnan, S.; Ibrahim, R.; Kanagarajan, K. Establishing the existence of Hilfer fractional pantograph equations with impulses. *Fundam. J. Math. Appl.* **2018**, *1*, 36–42.
58. Harikrishnan, S.; Elsayed, E.; Kanagarajan, K. Existence and uniqueness results for fractional pantograph equations involving ψ -Hilfer fractional derivative. *Dyn. Contin. Discret. Impuls. Syst.* **2018**, *25*, 319–328.
59. Ahmad, B.; Sivasundaram, S. Some existence results for fractional integro-differential equations with nonlinear conditions. *Commun. Appl. Anal.* **2008**, *12*, 107.
60. Ntouyas, S. Nonlocal initial and boundary value problems: a survey. In *Handbook of Differential Equations: Ordinary Differential Equations*; Elsevier: Amsterdam, The Netherlands, 2006; Volume 2, pp. 461–557.
61. Yong, Z.; Jinrong, W.; Lu, Z. *Basic Theory of Fractional Differential Equations*; World Scientific: Singapore, 2016.
62. Ciegis, R.; Bugajev, A. Numerical approximation of one model of bacterial self-organization. *Nonlinear Anal. Model. Control.* **2012**, *17*, 253–270.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).