




Article

The Stability and Stabilization of Infinite Dimensional Caputo-Time Fractional Differential Linear Systems

Hanaa Zitane ^{1,†} , Ali Boutoulout ^{1,†}  and Delfim F. M. Torres ^{2,*,†} 

¹ MACS Laboratory, Department of Mathematics, Faculty of Sciences, University of Moulay Ismail, Meknes 11201, Morocco; h.zitane@edu.umi.ac.ma (H.Z.); a.boutoulout@fs-umi.ac.ma (A.B.)

² Center for Research and Development in Mathematics and Applications (CIDMA), Department of Mathematics, University of Aveiro, 3810-193 Aveiro, Portugal

* Correspondence: delfim@ua.pt; Tel.: +351-234-370-668

† The authors contributed equally to this work.

Received: 31 December 2019; Accepted: 2 March 2020; Published: 5 March 2020



Abstract: We investigate the stability and stabilization concepts for infinite dimensional time fractional differential linear systems in Hilbert spaces with Caputo derivatives. Firstly, based on a family of operators generated by strongly continuous semigroups and on a probability density function, we provide sufficient and necessary conditions for the exponential stability of the considered class of systems. Then, by assuming that the system dynamics are symmetric and uniformly elliptical and by using the properties of the Mittag–Leffler function, we provide sufficient conditions that ensure strong stability. Finally, we characterize an explicit feedback control that guarantees the strong stabilization of a controlled Caputo time fractional linear system through a decomposition approach. Some examples are presented that illustrate the effectiveness of our results.

Keywords: fractional differential equations; fractional diffusion systems; Caputo derivative; stability and stabilization in Hilbert spaces; decomposition method

MSC: 26A33; 93D15

1. Introduction

Fractional order calculus is a natural generalization of classical integer order calculus. It deals with integrals and derivatives of an arbitrary real or complex order. Fractional order calculus has become very popular in recent years, due to its demonstrated applications in many fields of applied sciences and engineering, such as the spread of contaminants in underground water, charge transport in amorphous semiconductors, and diffusion of pollution in the atmosphere [1–3]. Because it generalizes and includes in the limit the integer order calculus, fractional calculus has the potential to accomplish much more than what integer order calculus achieves [4]. In particular, it has proved to be a powerful tool to describe long-term memory and hereditary properties of various dynamical complex processes [5]; diffusion processes, such as those found in batteries [6]; and electrochemical and control processes [7], to model and control epidemics [8,9] and mechanical properties of viscoelastic systems and damping materials, such as stress and strain [10].

One can find in the literature several different fractional calculuses. Here we use the fractional calculus of Caputo, which was introduced by Michele Caputo in his 1967 paper [11]. Such calculus has appeared, in a natural way, for representing observed phenomena in laboratory experiments and field observations, where the mathematical theory was checked with experimental data. Indeed, the operator introduced by Caputo in 1967, and used by us in the present work, represents an observed

linear dissipative mechanism phenomenon with a time derivative of order 0.15 entering the stress-strain relation [11]. More recently, a variational analysis with Caputo operators has been developed, which provides further mathematical substance to the use of Caputo fractional operators [12,13].

In the analysis and design of control systems, the stability issue always has an important role [14,15]. For a dynamical system, an equilibrium state is said to be stable if said system remains close to this state for small disturbances, and for an unstable system the question is how to stabilize it, especially by a feedback control law [16]. The stabilization concept for integer order systems and related problems has been considered in several works; see, e.g., [17–20] and references cited therein. In [17], the relationship between the asymptotic behavior of a system, the spectrum properties of its dynamics, and the existence of a Lyapunov functional is provided. Several techniques are considered to study different kinds of stabilization; for example, the exponential stabilization was studied via a decomposition method [19], and the strong stabilization was developed using the Riccati approach [20].

Similarly to classical dynamical systems, stability analysis is a central task in the study of fractional dynamical systems, which has attracted the increasing interest of many researchers [9,21]. For finite dimensional systems, the stability concept for fractional differential systems equipped with the Caputo derivative was investigated in many works [22]. In [23], Matignon studies the asymptotic behavior for linear fractional differential systems with the Caputo derivative, where their dynamics A are a constant coefficient matrix. In this case, the stability is guaranteed if the eigenvalues of the dynamics matrix A , $\lambda \in \sigma(A)$, satisfy $|\arg(\lambda)| > \frac{\pi\alpha}{2}$ [23]. Since then, many scholars have carried out further studies on the stability for different classes of fractional linear systems [24,25]. In [24], stability theorems for fractional differential systems, which include linear systems, time-delayed systems, and perturbed systems, are established, while in [25], Ge, Chen, and Kou provide results on the Mittag-Leffler stability and propose a Lyapunov direct method, which covers the power law stability and the exponential stability. See also [26], where the Mittag-Leffler and the class-K function stability of fractional differential equations of order $\alpha \in (1, 2)$ are investigated. In 2018, the notion of regional stability was introduced for fractional systems in [27], where the authors study the Mittag-Leffler stability and the stabilization of systems with Caputo derivatives, but only on a sub-region of its geometrical domain. More recently, fractional output stabilization problems for distributed systems in the Riemann–Liouville sense were studied [28–30], where feedback controls, which ensure exponential, strong, and weak stabilization of the state fractional spatial derivatives, with real and complex orders, are characterized.

An analysis of the literature shows that existing results on stability of fractional systems are essentially limited to finite-dimensional fractional order linear systems, while results on infinite-dimensional spaces are a rarity. In contrast, here we investigate global stability and stabilization of infinite dimensional fractional dynamical linear systems in the Hilbert space $L^2(\Omega)$ with Caputo derivatives of fractional order $0 < \alpha < 1$. In particular, we characterize exponential and strong stability for fractional Caputo systems on infinite-dimensional spaces.

The remainder of this paper is organized as follows. In Section 2, some basic knowledge of fractional calculus and some preliminary results, which will be used throughout the paper, are given. In Section 3, we prove results on the global asymptotic and exponential stability of Caputo-time fractional differential linear systems. In contrast with available results in the literature, which are restricted to systems of integer order or to fractional systems in the finite dimensional state space \mathbb{R}^n , here we study a completely different class of systems: we investigate fractional linear systems where the state space is the Hilbert space $L^2(\Omega)$. We also characterize the stabilization of a controlled Caputo diffusion linear system via a decomposition method. Section 4 presents the main conclusions of the work and some interesting open questions that deserve further investigations.

2. Preliminaries and Notation

In this section, we introduce several definitions and results of fractional calculus that are used in the sequel.

Definition 1 ([2]). Let $0 < \alpha < 1$ and $T > 0$. The Caputo derivative of fractional order α for an absolutely continuous function $y(\cdot)$ on $[0, T]$ can be defined as follows:

$${}^C D_t^\alpha y(t) = \frac{1}{\Gamma(1-\alpha)} \int_0^t (t-s)^{-\alpha} \frac{d}{ds} y(s) ds,$$

where $\Gamma(1-\alpha)$ is the Euler Gamma function.

Lemma 1 ([31]). For any given function $g \in L^2(0, T, L^2(\Omega))$, we say that function $y \in C(0, T, L^2(\Omega))$ is a mild solution of the system

$$\begin{cases} {}^C D_t^\alpha y(t) = Ay(t) + g(t) & t \in]0, +\infty[\\ y(0) = y_0 & y_0 \in L^2(\Omega) \end{cases} \quad (1)$$

if it satisfies

$$y(t) = S_\alpha(t)y_0 + \int_0^t (t-s)^{\alpha-1} K_\alpha(t-s)g(s) ds, \quad (2)$$

where

$$S_\alpha(t) = \int_0^{+\infty} \Psi_\alpha(\theta) S(t^\alpha \theta) d\theta \quad (3)$$

and

$$K_\alpha(t) = \alpha \int_0^{+\infty} \theta \Psi_\alpha(\theta) S(t^\alpha \theta) d\theta \quad (4)$$

with

$$\Psi_\alpha(\theta) = \frac{1}{\alpha} \theta^{-1-\frac{1}{\alpha}} T_\alpha(\theta^{-\frac{1}{\alpha}}), \quad (5)$$

$(S(t))_{t \geq 0}$ the strongly continuous semigroup generated by operator A , and T_α the probability density function defined on $(0, \infty)$ by

$$T_\alpha = \frac{1}{\pi} \sum_{n=1}^{+\infty} (-1)^n \theta^{\alpha n-1} \frac{\Gamma(n\alpha+1)}{n!} \sin(n\pi\alpha).$$

Remark 1 ([32]). The probability density function T_α defined on $(0, \infty)$ satisfies

$$T_\alpha(\theta) \geq 0, \quad \theta \in (0, \infty), \quad \text{and} \quad \int_0^{+\infty} T_\alpha(\theta) d\theta = 1.$$

Definition 2 ([33]). The Mittag-Leffler function of one parameter is defined as

$$E_\eta(z) = \sum_{n=0}^{+\infty} \frac{z^n}{\Gamma(\eta n + 1)} \quad \text{with} \quad \operatorname{Re}(\eta) > 0, \quad z \in \mathbb{C}.$$

Definition 3 ([33]). The Mittag-Leffler function of two parameters is defined as

$$E_{\eta,\beta}(z) = \sum_{n=0}^{+\infty} \frac{z^n}{\Gamma(\eta n + \beta)} \quad \text{with} \quad \operatorname{Re}(\eta) > 0, \quad \beta > 0, \quad z \in \mathbb{C}.$$

Remark 2. The Mittag-Leffler function appears naturally in the solution of fractional differential equations and in various applications: see [33] and references therein. The exponential function is a special case of the Mittag-Leffler function [34]: for $\beta = 1$ one has $E_{\eta,1}(z) = E_\eta(z)$ and $E_{1,1}(z) = e^z$.

Lemma 2 ([35]). The Mittag-Leffler function $E_\alpha(-t^\alpha)$ is completely monotonic: for all $0 < \alpha < 1$, and for all $n \in \mathbb{N}$ and $t > 0$, one has

$$(-1)^n \frac{d^n}{dt^n} E_\alpha(-t^\alpha) \geq 0.$$

Lemma 3 ([36]). The generalized Mittag–Leffler function $E_{\eta,\beta}(-x)$, $x \geq 0$, is completely monotonic for $\eta, \beta > 0$ if and only if $\eta \in (0, 1]$ and $\beta \geq \eta$.

Lemma 4 ([37]). Let $\beta > 0$, $0 < \eta < 2$, and μ be an arbitrary real number such that $\frac{\pi\eta}{2} < \mu < \min\{\pi, \pi\eta\}$. Then, the following asymptotic expressions hold:

- If $|\arg(z)| \leq \mu$ and $|z| > 0$, then

$$|E_{\eta,\beta}(z)| \leq M_1(1 + |z|)^{(1-\beta)/\eta} e^{\operatorname{Re}(z^{\frac{1}{\eta}})} + \frac{M_2}{1 + |z|}; \quad (6)$$

- If $\mu < |\arg(z)| \leq \pi$ and $|z| \geq 0$, then

$$|E_{\eta,\beta}(z)| \leq \frac{M_2}{1 + |z|}, \quad (7)$$

where M_1 and M_2 are positive constants.

3. Main Results

Our main goal is to study the stability and provide stabilization for a class of abstract Caputo-time fractional differential linear systems.

3.1. Stability of Time Fractional Differential Systems

Let Ω be an open bounded subset of \mathbb{R}^n , $n = 1, 2, 3, \dots$, and let us consider the following abstract time fractional order differential system:

$$\begin{cases} {}^C D_t^\alpha z(t) = Az(t), & t \in]0, +\infty[, \\ z(0) = z_0, & z_0 \in L^2(\Omega), \end{cases} \quad (8)$$

where ${}^C D_t^\alpha$ is the left-sided Caputo fractional derivative of order $0 < \alpha < 1$; the second order operator $A : D(A) \subset L^2(\Omega) \rightarrow L^2(\Omega)$ is linear with a dense domain, is such that the coefficients do not depend on time t , and is also the infinitesimal generator of the C_0 -semi-group $(S(t))_{t \geq 0}$ on the Hilbert state space $L^2(\Omega)$ endowed with its usual inner product $\langle \cdot, \cdot \rangle$ and the corresponding norm $\|\cdot\|$. The unique mild solution of system (8) can be written, from Lemma 1, as

$$z(t) = S_\alpha(t)z_0,$$

where $S_\alpha(t)$ is defined by (3).

We begin by proving the following lemma, which will be used thereafter.

Lemma 5. Let A be the infinitesimal generator of a C_0 -semi-group $(S(t))_{t \geq 0}$ on the Hilbert space $L^2(\Omega)$. Assume that there exists a function $h(\cdot) \in L^2(0, +\infty; \mathbb{R}^+)$ satisfying

$$\|S_\alpha(t+s)z\| \leq h(t)\|S_\alpha(s)z\|, \quad \forall t, s \geq 0, \quad \forall z \in L^2(\Omega). \quad (9)$$

Then the operators $(S_\alpha(t))_{t \geq 0}$ are uniformly bounded.

Proof. To prove that $(S_\alpha(t))_{t \geq 0}$ are bounded, we have to show that

$$\forall z \in L^2(\Omega) \sup_{t \geq 0} \|S_\alpha(t)z\| < \infty. \quad (10)$$

By *reductio ad absurdum*, let us suppose that (10) does not hold, which means that there exists a sequence $(t_s + \tau_n)$, $t_s > 0$ and $\tau_n \rightarrow +\infty$, satisfying

$$\|S_\alpha(t_s + \tau_n)z\| \rightarrow +\infty \text{ as } n \rightarrow +\infty. \quad (11)$$

From relation

$$\int_0^{+\infty} \|S_\alpha(s + \tau_n)z\|^2 ds = \int_{\tau_n}^{+\infty} \|S_\alpha(s)z\|^2 ds, \quad 0 \leq s < +\infty,$$

it follows that the right-hand side goes to 0 as $n \rightarrow +\infty$. Using Fatou's Lemma yields

$$\liminf_{n \rightarrow +\infty} \|S_\alpha(s + \tau_n)z\| = 0 \quad \forall s > 0.$$

Therefore, for some $s_0 < t_s$, we may find a subsequence τ_{n_k} such that

$$\lim_{k \rightarrow +\infty} \|S_\alpha(s_0 + \tau_{n_k})z\| = 0.$$

By virtue of condition (9), one obtains

$$\|S_\alpha(t_s + \tau_{n_k})z\| \leq h(t_s - s_0) \|S_\alpha(s_0 + \tau_{n_k})z\| \xrightarrow{k \rightarrow +\infty} 0,$$

which contradicts (11). The intended conclusion follows from the uniform boundedness principle. \square

Definition 4. Let $z_0 \in L^2(\Omega)$. System (8) is said to be exponentially stable if there exist two strictly positive constants, $M > 0$ and $\omega > 0$, such that

$$\|z(t)\| \leq Me^{-\omega t} \|z_0\|, \quad \forall t \geq 0.$$

The next theorem provides necessary and sufficient conditions for exponential stability of the abstract fractional order differential system (8).

Theorem 1. Suppose that the operators $(S_\alpha(t))_{t \geq 0}$ fulfill assumption (9) and

$$\forall z \in L^2(\Omega) \quad \|S_\alpha(t+s)z\| \leq \|S_\alpha(t)z\| \cdot \|S_\alpha(s)z\|, \quad \forall t, s \geq 0. \quad (12)$$

Then, system (8) is exponentially stable if, and only if, for every $z \in L^2(\Omega)$ there exists a positive constant $\delta < \infty$ such that

$$\int_0^{+\infty} \|S_\alpha(t)z\|^2 dt < \delta. \quad (13)$$

Proof. One has

$$\begin{aligned} t \|S_\alpha(t)z\|^2 &= \int_0^t \|S_\alpha(t)z\|^2 ds \\ &= \int_0^t \|S_\alpha(t-s+s)z\|^2 ds. \end{aligned}$$

Combining assumption (9), Lemma 5, and condition (13), one gets

$$\begin{aligned} t \|S_\alpha(t)z\|^2 &\leq \int_0^t \|S_\alpha(s)z\|^2 \|S_\alpha(t-s)z\|^2 ds \\ &\leq N\delta \|z\|^2 \end{aligned}$$

for some $N > 0$. Therefore, for t sufficiently large, it follows that

$$\|S_\alpha(t)\| < 1.$$

Then, there exists $t_1 > 0$ such that

$$\ln \|S_\alpha(t)\| < 0, \quad \forall t \geq t_1.$$

Thus,

$$\omega_0 = \inf_{t \geq 0} \frac{\ln \|S_\alpha(t)\|}{t} < 0.$$

Now, let us show that

$$\omega_0 = \lim_{t \rightarrow +\infty} \frac{\ln \|S_\alpha(t)\|}{t}. \quad (14)$$

Let $t_s > 0$ be a fixed number and $N' = \sup_{t \in [0, t_s]} \|S_\alpha(t)\|$. Thus, for each $t > t_s$, there exists $m \in \mathbb{N}$ such that $mt_s \leq t \leq (m+1)t_s$. From (12), it follows that

$$\begin{aligned} \|S_\alpha(t)\| &= \|S_\alpha(mt_s + (t - mt_s))\| \\ &\leq \|S_\alpha(mt_s)\| \|S_\alpha(t - mt_s)\|, \end{aligned}$$

which yields

$$\frac{\ln \|S_\alpha(t)\|}{t} \leq \frac{\ln \|S_\alpha(mt_s)\|}{t} + \frac{\ln \|S_\alpha(t - mt_s)\|}{t}.$$

Using again (12), it results that

$$\frac{\ln \|S_\alpha(t)\|}{t} \leq \frac{mt_s}{t} \frac{\ln \|S_\alpha(t_s)\|}{t_s} + \frac{\ln \|N'\|}{t}.$$

Since $\frac{mt_s}{t} \leq 1$ and t_s is arbitrary, one obtains

$$\limsup_{t \rightarrow +\infty} \frac{\ln \|S_\alpha(t)\|}{t} \leq \inf_{t > 0} \frac{\ln \|S_\alpha(t)\|}{t} \leq \liminf_{t \rightarrow +\infty} \frac{\ln \|S_\alpha(t)\|}{t}.$$

Consequently, (14) holds. Hence, we conclude that for all $\omega \in]0, -\omega_0[$, there exists $M > 0$ such that

$$\forall z \in L^2(\Omega) \quad \|S_\alpha(t)z\| \leq Me^{-\omega t} \|z\|, \quad \forall t \geq 0,$$

which means that system (8) is exponentially stable. The converse is obvious. \square

Remark 3. When $\alpha = 1$, the conditions (9) and (12) are verified, and we retrieve from our Theorem 1 the results established in [17,18] about the exponential stability of system (8) on Ω , which is equivalent to

$$\int_0^{+\infty} \|S(t)z\|^2 dt < \infty, \quad \forall z \in L^2(\Omega).$$

Definition 5. Let $z_0 \in L^2(\Omega)$. System (8) is said to be strongly stable if its corresponding solution $z(t)$ satisfies

$$\|z(t)\| \longrightarrow 0 \text{ as } t \longrightarrow +\infty.$$

In our next theorem, we provide sufficient conditions that guaranty the strong stability of the fractional order differential system (8). The result generalizes the asymptotic result established by Matignon for finite dimensional state spaces, where the dynamics of the system A are considered to be

a matrix with constant coefficients in \mathbb{R}^n [23]. In contrast, here we tackle the stability for a different class of systems. Precisely, we consider fractional systems where the system dynamics A are a linear operator generating a strongly continuous semigroup in the infinite dimensional state space $L^2(\Omega)$.

Theorem 2. Let $(\lambda_p)_{p \geq 1}$ and $(\phi_p)_{p \geq 1}$ be the eigenvalues and the corresponding eigenfunctions of operator A on $L^2(\Omega)$. If A is a symmetric uniformly elliptical operator, then system (8) is strongly stable on Ω .

Proof. Since A is a symmetric uniformly elliptical operator, it follows that system (8) admits a weak solution defined by

$$z(t) = \sum_{p=1}^{+\infty} E_{\alpha}(\lambda_p t^{\alpha}) \langle z_0, \phi_p \rangle \phi_p \quad \forall z_0 \in L^2(\Omega),$$

where $(\lambda_p)_{p \geq 1}$ satisfy

$$0 > \lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_j \geq \dots, \lim_{j \rightarrow \infty} \lambda_j = -\infty,$$

and $(\phi_p)_{p \geq 1}$ forms an orthonormal basis in $L^2(\Omega)$ [38,39]. Using the fact that function $E_{\alpha}(-t^{\alpha})$ is completely monotonic, for all $\alpha \in (0, 1)$ and $t > 0$ (Lemma 2), yields

$$\begin{aligned} \|z(t)\| &= \left\| \sum_{p=1}^{+\infty} E_{\alpha}(\lambda_p t^{\alpha}) \langle z_0, \phi_p \rangle \phi_p \right\| \\ &\leq |E_{\alpha}(\lambda_1 t^{\alpha})| \|z_0\|. \end{aligned}$$

Moreover, from Lemma 4, it follows that

$$\|z(t)\| \leq \frac{M_2}{1 - \lambda_1 t^{\alpha}} \|z_0\| \rightarrow 0 \quad \text{as } t \rightarrow +\infty$$

for some $M_2 > 0$. Hence, system (8) is strongly stable on Ω . \square

Example 1. Let us consider, on $\Omega =]0, 1[$, the following one-dimensional fractional diffusion system defined by

$$\begin{cases} {}^C D_t^{0.5} z(x, t) = \frac{\partial^2 z}{\partial x^2}(x, t), & x \in \Omega, \quad t \in]0, +\infty[, \\ z(0, t) = z(1, t) = 0, & \forall t > 0, \\ z(x, 0) = z_0, & x \in \Omega, \end{cases} \quad (15)$$

where the second order operator $A = \frac{\partial^2}{\partial x^2}$ has its spectrum given by the eigenvalues $\lambda_p = -(p\pi)^2$, $p \geq 1$, and the corresponding eigenfunctions are $\phi_p(x) = \sqrt{\frac{2}{1+(p\pi)^2}} \sin(p\pi x)$, $p \geq 1$. Operator A generates a C_0 -semi-group $(S(t))_{t \geq 0}$ defined by

$$S(t)z_0 = \sum_{p=1}^{+\infty} e^{\lambda_p t} \langle z_0, \phi_p \rangle \phi_p.$$

Moreover, the solution of system (15) is given by

$$S_{0.5}(t)z_0 = \sum_{p=1}^{+\infty} E_{0.5}(\lambda_p t^{0.5}) \langle z_0, \phi_p \rangle \phi_p.$$

One has that operator A is symmetric and uniformly elliptical. Consequently, from our Theorem 2, we deduce that system (15) is strongly stable on Ω . This is illustrated numerically in Figure 1 for $z(x, 0) = \sin(\pi x)$, $t = 0.1$, $t = 0.15$, $t = 0.2$, and $t = 1.0$.




Figure 1. The state of system (15) for $z(x,0) = \sin(\pi x)$, $t = 0.1$, $t = 0.15$, $t = 0.2$, and $t = 1.0$, illustrating the fact that (15) is strongly stable on $\Omega =]0, 1[$.

3.2. Stabilization of Time Fractional Differential Systems

Let Ω be an open bounded subset of \mathbb{R}^n , $n = 1, 2, 3, \dots$. We consider the following Caputo-time fractional differential linear system:

$$\begin{cases} {}^C D_t^\alpha z(t) = Az(t) + Bu(t), & t \in]0, +\infty[, \quad 0 < \alpha < 1, \\ z(0) = z_0, & z_0 \in L^2(\Omega), \end{cases} \quad (16)$$

with the same assumptions on A as in Section 3.1 and where B is a bounded linear operator from U into $L^2(\Omega)$, where U is the space of controls, assumed to be a Hilbert space. By Lemma 1, the unique mild solution $z(\cdot)$ of system (16) is defined by

$$z(t) = S_\alpha(t)z_0 + \int_0^t (t-s)^{\alpha-1} K_\alpha(t-s) Bu(s) ds, \quad (17)$$

where $S_\alpha(t)$ and $K_\alpha(t)$ are given, respectively, by (3) and (4).

Definition 6. System (16) is said to be exponentially (respectively strongly) stabilizable if there exists a bounded operator $K \in \mathcal{L}(L^2(\Omega), U)$ such that the system

$$\begin{cases} {}^C D_t^\alpha z(t) = (A + BK)z(t), & t \in]0, +\infty[, \\ z(0) = z_0, & z_0 \in L^2(\Omega), \end{cases} \quad (18)$$

is exponentially (respectively strongly) stable on Ω .

Remark 4. It is clear that the exponential stabilization of system (16) implies the strong stabilization of (16). Note that the concept is general: when $\alpha = 1$, we obtain the classical definitions of stability and stabilization.

Let $(S^k(t))_{t \geq 0}$ be the strongly continuous semi-group generated by $A + BK$, where $K \in \mathcal{L}(L^2(\Omega), U)$ is the feedback operator. The unique mild solution of system (16) can be written as

$$z(t) = S_\alpha^k(t)z_0$$

with

$$S_\alpha^k(t) = \int_0^{+\infty} \Psi_\alpha(\theta) S^k(t^\alpha \theta) d\theta,$$

where $\Psi_\alpha(\theta)$ is defined by (5).

Theorem 3. Let $A + BK$ generate a strongly continuous semi-group $(S^k(t))_{t \geq 0}$ on $L^2(\Omega)$. If the operator $(S_\alpha^k(t))_{t \geq 0}$ satisfies conditions (9) and (12) and if

$$\forall z \in L^2(\Omega) \quad \int_0^{+\infty} \|S_\alpha^k(t)z\|^2 dt < \infty$$

holds, then system (16) is exponentially stabilizable on Ω .

Proof. The proof is similar to the proof of Theorem 1. \square

Theorem 4. Let $(\lambda_p^k)_{p \geq 1}$ and $(\phi_p^k)_{p \geq 1}$ be the eigenvalues and the corresponding eigenfunctions of operator $A + BK$ on $L^2(\Omega)$. If $A + BK$ is a symmetric uniformly elliptical operator, then system (16) is strongly stabilizable on Ω .

Proof. The proof is similar to the proof of Theorem 2. \square

Example 2. Let us consider, on $\Omega =]0, 1[$, the following fractional differential system of order $\alpha = 0.2$:

$$\begin{cases} {}^C D_t^{0.2} z(x, t) = \frac{1}{100} \frac{\partial^2 z}{\partial x^2}(x, t) + \frac{1}{2} z(x, t) + BKz(x, t), & (x, t) \in \Omega \times]0, +\infty[, \\ z(0, t) = z(1, t) = 0, & \forall t > 0, \\ z(x, 0) = z_0, & x \in \Omega, \end{cases} \quad (19)$$

with the linear bounded operator $B = I$ and where we take $K = -B^* = -I$. The operator

$$A + BK = \frac{1}{100} \frac{\partial^2}{\partial x^2} - \frac{1}{2},$$

with spectrum given by the eigenvalues $\lambda_p^k = -\frac{1}{2} - \frac{1}{100}(p\pi)^2$, $p \geq 1$, and the corresponding eigenfunctions $\phi_p^k(x) = \sqrt{\frac{2}{1+(p\pi)^2}} \cos(p\pi x)$, $p \geq 1$, generates a C_0 -semi-group $(S^k(t))_{t \geq 0}$ defined by

$$S^k(t)z_0 = \sum_{p=1}^{+\infty} e^{\lambda_p^k t} \langle z_0, \phi_p^k \rangle \phi_p^k.$$

Furthermore, the solution of system (19) can be written as

$$z(t) = S_{0.2}^k(t)z_0 = \sum_{p=1}^{+\infty} E_{0.2}(\lambda_p^k t^{0.2}) \langle z_0, \phi_p^k \rangle \phi_p^k.$$

It is clear that $A + BK$ is a symmetric and uniformly elliptical operator. Hence, from Theorem 4, we deduce that system (19) is strongly stabilizable on Ω , i.e., the system

$$\begin{cases} {}^C D_t^{0.2} z(x, t) = \frac{1}{100} \frac{\partial^2 z}{\partial x^2}(x, t) + \frac{1}{2} z(x, t) + Bu(t), & (x, t) \in \Omega \times]0, +\infty[, \\ z(0, t) = z(1, t) = 0, & \forall t > 0, \\ z(x, 0) = z_0, & x \in \Omega, \end{cases}$$

is strongly stabilizable by the feedback control $u(t) = -B^*z(t)$. Figure 2 shows, for $z(x, 0) = x(x - 1)$, that the state $z(x, t)$ of system (19) is unstable at $t = 0$. Moreover, we see that the state evolves close to 0 at $t = 10$. Numerically, the state is stabilized by $u(t) = -B^*z(t)$ with an error equal to 1.75×10^{-04} .




Figure 2. The state of system (19) for $z(x, 0) = x(x - 1)$, $t = 0$, and $t = 10$, illustrating the fact that (19) is unstable at $t = 0$ but it is stabilized at $t = 10$ on $\Omega =]0, 1[$.

3.3. Decomposition Method

Now, we study the stabilization of system (16) using the decomposition method, which consists of decomposing the state space and the system using the spectral properties of operator A .

Let $\xi > 0$ be fixed and assume that there are at most finitely-many nonnegative eigenvalues of A , each with finite-dimensional eigenspace. In other words, assume there exists $l \in \mathbb{N}$ such that

$$\sigma(A) = \sigma_u(A) \cup \sigma_s(A), \quad (20)$$

where $\sigma_u(A) = \sigma(A) \cap \{\lambda_p, p = 1, 2, \dots, l\}$, $\sigma_s(A) = \sigma(A) \cap \{\lambda_p, p = l + 1, l + 2, \dots\}$ with $\lambda_l \geq 0$ and $\lambda_{l+1} \leq -\xi$. Because the sequence $(\phi_p)_{p \geq 1}$ forms a complete and orthonormal basis in $H = L^2(\Omega)$, it follows that the state space H can be decomposed as

$$H = H_u \oplus H_s, \quad (21)$$

where $H_u = PH = \text{span}\{\phi_1, \phi_2, \dots, \phi_l\}$ and $H_s = (I - P)H = \text{span}\{\phi_{l+1}, \phi_{l+2}, \dots\}$ with $P \in \mathcal{L}(H)$ the projection operator [40]. Hence, system (16) can be decomposed into the following two sub-systems:

$$\begin{cases} {}^C D_t^\alpha z_u(t) = A_u z_u(t) + PBu(t), \\ z_{0u} = Pz_0, \end{cases} \quad (22)$$

and

$$\begin{cases} {}^C D_t^\alpha z_s(t) = A_s z_s(t) + (I - P)Bu(t), \\ z_{0s} = (I - P)z_0, \end{cases} \quad (23)$$

where A_s and A_u are the restrictions of A on H_s and H_u , respectively, and are such that $\sigma(A_s) = \sigma_s(A)$, $\sigma(A_u) = \sigma_u(A)$, and A_u is a bounded operator on H_u .

Our next result asserts that stabilization of system (16) is equivalent to the one of system (22).

Theorem 5. *Let the spectrum $\sigma(A)$ of A satisfy the above spectrum decomposition assumptions (20) for some $\xi > 0$ and A_s be a symmetric uniformly elliptical operator. If system (22) is strongly stabilizable by the control*

$$u(t) = D_u z_u(t) \quad (24)$$

with $D_u \in \mathcal{L}(H, U)$ such that

$$\|z_u(t)\| \leq C t^{-\mu}, \quad \mu, C > 0, \quad (25)$$

then system (16) is strongly stabilizable using the feedback control $v(t) = D_u z_u(t)$.

Proof. Using the fact that system (22) is strongly stabilizable by control (24), and inequality (25) yields

$$\|z_u(t)\| \longrightarrow 0 \text{ as } t \longrightarrow +\infty \quad (26)$$

and

$$\|u(t)\| \leq C \|D_u\| t^{-\mu}, \quad (27)$$

the unique weak solution of system (23) can be written in the space H_s as

$$z_s(t) = \sum_{p=l+1}^{+\infty} E_\alpha(\lambda_p t^\alpha) \langle z_{0s}, \phi_p \rangle \phi_p + \sum_{p=l+1}^{+\infty} \int_0^t (t-s)^{\alpha-1} E_{\alpha,\alpha}(\lambda_p(t-s)^\alpha) \langle (I-P)Bu(s), \phi_p \rangle \phi_p \, ds$$

since A_s is a symmetric uniformly elliptical operator [38]. Using the spectrum decomposition relation (20), Lemma 2, and Lemma 3, one has that

$$E_\alpha(\lambda_p t^\alpha) \leq E_\alpha(-\xi t^\alpha) \quad \text{for all } p \geq l+1 \quad (28)$$

and

$$E_{\alpha,\alpha}(\lambda_p(t-s)^\alpha) \leq E_{\alpha,\alpha}(-\xi(t-s)^\alpha) \quad \text{for all } p \geq l+1. \quad (29)$$

Then, feeding system (23) by the same control $u(t) = D_u z_u(t)$ and using (27)–(29), it follows that

$$\begin{aligned} \|z_s(t)\| &\leq E_\alpha(-\xi t^\alpha) \|z_{0s}\| + C \|D_u\| \|I - P\| \|B\| \int_0^t (t-s)^{\alpha-1} s^{-\mu} E_{\alpha,\alpha}(-\xi(t-s)^\alpha) \, ds \\ &\leq E_\alpha(-\xi t^\alpha) \|z_{0s}\| + C \|D_u\| \|I - P\| \|B\| \sum_{n=1}^{+\infty} \int_0^t \frac{(-\xi)^n (t-s)^{\alpha n + \alpha - 1} s^{-\mu} \, ds}{\Gamma(\alpha n + \alpha)} \\ &\leq E_\alpha(-\xi t^\alpha) \|z_{0s}\| + C \|D_u\| \|I - P\| \|B\| \sum_{n=1}^{+\infty} \frac{(-\xi)^n t^{\alpha n + \alpha - \mu}}{\Gamma(\alpha n + \alpha - \mu - 1) \Gamma(1 - \mu)^{-1}} \\ &\leq E_\alpha(-\xi t^\alpha) \|z_{0s}\| + C \Gamma(1 - \mu) \|D_u\| \|I - P\| \|B\| t^{\alpha - \mu} E_{\alpha,\alpha - \mu + 1}(-\xi t^\alpha). \end{aligned}$$

Lemma 4 implies that

$$\|z_s(t)\| \leq \frac{M_1}{1 + \xi t^\alpha} \|z_{s0}\| + C \Gamma(1 - \mu) \|D_u\| \|I - P\| \|B\| \frac{M_2 t^{\alpha - \mu}}{1 + \xi t^\alpha}$$

for some $M_1, M_2 > 0$. Therefore,

$$\|z_s(t)\| \longrightarrow 0 \text{ as } t \longrightarrow +\infty. \quad (30)$$

On the other hand, we have that

$$\|z(t)\| = \|z_s(t) + z_u(t)\| \leq \|z_s(t)\| + \|z_u(t)\|. \quad (31)$$

Combining (26), (30), and (31), we deduce the strong stabilization of system (16). \square

4. Conclusions and Future Work

We investigated the stability problem of infinite dimensional time fractional differential linear systems under Caputo derivatives of order $\alpha \in (0, 1)$, where the state space is the Hilbert space $L^2(\Omega)$. We proved necessary and sufficient conditions for exponential stability and obtained a characterization for the asymptotic stability, which is guaranteed if the system dynamics are symmetric and uniformly elliptical. Moreover, some stabilization criteria were also proved. Finally, we investigated the strong stabilization of the system via a decomposition method; an explicit feedback control was obtained. Illustrative examples were given, showing the effectiveness of the theoretical results. As future work, we intend to extend our work to the class of infinite dimensional time fractional differential nonlinear systems. Various other questions are still open and deserve further investigations, such as studying boundary stability and gradient stability for time fractional differential linear systems or considering the more recent notion of Λ -fractional derivative [41], and thus obtaining a geometrical interpretation.

Author Contributions: Each author equally contributed to this paper, and read and approved the final manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by Moulay Ismail University (H.Z.); by Hassan II Academy of Science and Technology, project N 630/2016 (A.B.); and by The Portuguese Foundation for Science and Technology, R&D unit CIDMA, within project UIDB/04106/2020 (D.F.M.T.).

Acknowledgments: This research is part of the first author's Ph.D. project, which was carried out at Moulay Ismail University, Meknes, and began during a one-month visit of Zitane to the R&D Unit CIDMA, Department of Mathematics, University of Aveiro, Portugal, June 2019. The hospitality of the host institution is here gratefully acknowledged. The authors are strongly grateful to three anonymous referees for their suggestions and invaluable comments.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Rahimy, M. Applications of fractional differential equations. *Appl. Math. Sci. (Ruse)* **2010**, *4*, 2453–2461.
2. Kilbas, A.A.; Srivastava, H.M.; Trujillo, J.J. *Theory and Applications of Fractional Differential Equations*; Elsevier Science B.V.: Amsterdam, The Netherlands, 2006.
3. Diethelm, K. *The Analysis of Fractional Differential Equations*; Lecture Notes in Mathematics; Springer: Berlin, Germany, 2010; Volume 2004. doi:10.1007/978-3-642-14574-2. [[CrossRef](#)]
4. Hilfer, R. *Applications of Fractional Calculus in Physics*; World Scientific Publishing Co., Inc.: River Edge, NJ, USA, 2000. doi:10.1142/9789812817747. [[CrossRef](#)]
5. Sabatier, J.; Agrawal, O.P.; Machado, J.A.T. *Advances in Fractional Calculus*; Springer: Dordrecht, The Netherlands, 2007. doi:10.1007/978-1-4020-6042-7. [[CrossRef](#)]
6. Gabano, J.D.; Poinot, T. Fractional modelling and identification of thermal systems. *Signal Process.* **2011**, *91*, 531–541. [[CrossRef](#)]
7. Ichise, M.; Nagayanagi, Y.; Kojima, T. An analog simulation of non-integer order transfer functions for analysis of electrode processes. *J. Electroanal. Chem. Interfacial Electrochem.* **1971**, *33*, 253–265. [[CrossRef](#)]
8. Rosa, S.; Torres, D.F.M. Optimal control and sensitivity analysis of a fractional order TB model. *Stat. Optim. Inf. Comput.* **2019**, *7*, 617–625. doi:10.19139/soic.v7i3.836. [[CrossRef](#)]
9. Silva, C.J.; Torres, D.F.M. Stability of a fractional HIV/AIDS model. *Math. Comput. Simul.* **2019**, *164*, 180–190. doi:10.1016/j.matcom.2019.03.016. [[CrossRef](#)]

10. Bagley, R.L.; Calico, R.A. Fractional order state equations for the control of viscoelastically damped structures. *J. Guid. Control Dyn.* **1991**, *14*, 304–311. [\[CrossRef\]](#)
11. Caputo, M. Linear models of dissipation whose Q is almost frequency independent II. *Fract. Calc. Appl. Anal.* **2008**, *11*, 4–14. Reprinted from *Geophys. J. R. Astr. Soc.* **1967**, *13*, 529–539. [\[CrossRef\]](#)
12. Malinowska, A.B.; Torres, D.F.M. *Introduction to the Fractional Calculus of Variations*; Imperial College Press: London, UK, 2012; doi:10.1142/p871. [\[CrossRef\]](#)
13. Almeida, R.; Pooseh, S.; Torres, D.F.M. *Computational Methods in the Fractional Calculus of Variations*; Imperial College Press: London, UK, 2015. doi:10.1142/p991. [\[CrossRef\]](#)
14. Mahmoud, M.S.; Karaki, B.J. Improved stability analysis and control design of reset systems. *IET Control Theory Appl.* **2018**, *12*, 2328–2336. doi:10.1049/iet-cta.2018.5410. [\[CrossRef\]](#)
15. Rocha, D.; Silva, C.J.; Torres, D.F.M. Stability and optimal control of a delayed HIV model. *Math. Methods Appl. Sci.* **2018**, *41*, 2251–2260. doi:10.1002/mma.4207. [\[CrossRef\]](#)
16. Sontag, E.D. Stability and feedback stabilization. In *Mathematics of Complexity and Dynamical Systems*; Springer: New York, NY, USA, 2012; Volumes 1–3, pp. 1639–1652. doi:10.1007/978-1-4614-1806-1_105. [\[CrossRef\]](#)
17. Pritchard, A.J.; Zabczyk, J. Stability and stabilizability of infinite-dimensional systems. *SIAM Rev.* **1981**, *23*, 25–52. doi:10.1137/1023003. [\[CrossRef\]](#)
18. Curtain, R.F.; Zwart, H. *An Introduction to Infinite-Dimensional Linear Systems Theory*; Texts in Applied Mathematics; Springer: New York, NY, USA, 1995; Volume 21. doi:10.1007/978-1-4612-4224-6. [\[CrossRef\]](#)
19. Triggiani, R. On the stabilizability problem in Banach space. *J. Math. Anal. Appl.* **1975**, *52*, 383–403. doi:10.1016/0022-247X(75)90067-0. [\[CrossRef\]](#)
20. Balakrishnan, A.V. Strong stabilizability and the steady state Riccati equation. *Appl. Math. Optim.* **1981**, *7*, 335–345. doi:10.1007/BF01442125. [\[CrossRef\]](#)
21. Wojtak, W.; Silva, C.J.; Torres, D.F.M. Uniform asymptotic stability of a fractional tuberculosis model. *Math. Model. Nat. Phenom.* **2018**, *13*, 9. doi:10.1051/mmnp/2018015. [\[CrossRef\]](#)
22. Zhang, F.; Li, C.; Chen, Y. Asymptotical stability of nonlinear fractional differential system with Caputo derivative. *Int. J. Differ. Equ.* **2011**, *2011*, 635165. doi:10.1155/2011/635165. [\[CrossRef\]](#)
23. Matignon, D. *Stability Results for Fractional Differential Equations with Applications to Control Processing*; Computational Engineering in Systems Applications: Lille, France, 1996; Volume 2, pp. 963–968.
24. Qian, D.; Li, C.; Agarwal, R.P.; Wong, P.J.Y. Stability analysis of fractional differential system with Riemann-Liouville derivative. *Math. Comput. Modelling* **2010**, *52*, 862–874. doi:10.1016/j.mcm.2010.05.016. [\[CrossRef\]](#)
25. Li, Y.; Chen, Y.; Podlubny, I. Stability of fractional-order nonlinear dynamic systems: Lyapunov direct method and generalized Mittag-Leffler stability. *Comput. Math. Appl.* **2010**, *59*, 1810–1821. doi:10.1016/j.camwa.2009.08.019. [\[CrossRef\]](#)
26. Matar, M.M.; Abu Skhail, E.S. On stability of nonautonomous perturbed semilinear fractional differential systems of order $\alpha \in (1, 2)$. *J. Math.* **2018**, *2018*, 1723481. doi:10.1155/2018/1723481. [\[CrossRef\]](#)
27. Ge, F.; Chen, Y.; Kou, C. *Regional Analysis of Time-Fractional Diffusion Processes*; Springer: Cham, Switzerland, 2018; doi:10.1007/978-3-319-72896-4. [\[CrossRef\]](#)
28. Zitane, H.; Larhrissi, R.; Boutoulout, A. On the fractional output stabilization for a class of infinite dimensional linear systems. In *Recent Advances in Modeling, Analysis and Systems Control: Theoretical Aspects and Applications*; Springer: Cham, Switzerland, 2020; pp. 241–259.
29. Zitane, H.; Larhrissi, R.; Boutoulout, A. Fractional output stabilization for a class of bilinear distributed systems. *Rend. Circ. Mat. Palermo* **2019**, *2*, 1–16. doi:10.1007/s12215-019-00429-w. [\[CrossRef\]](#)
30. Zitane, H.; Larhrissi, R.; Boutoulout, A. Riemann Liouville fractional spatial derivative stabilization of bilinear distributed systems. *J. Appl. Nonlinear Dyn.* **2019**, *8*, 447–461. doi:10.5890/JAND.2019.09.008. [\[CrossRef\]](#)
31. Zhou, Y.; Jiao, F. Existence of mild solutions for fractional neutral evolution equations. *Comput. Math. Appl.* **2010**, *59*, 1063–1077. doi:10.1016/j.camwa.2009.06.026. [\[CrossRef\]](#)
32. Mainardi, F.; Paradisi, P.; Gorenflo, R. Probability distributions generated by fractional diffusion equations. *arXiv* **2007**, arXiv:0704.0320.
33. Erdélyi, A.; Magnus, W.; Oberhettinger, F.; Tricomi, F.G. *Higher Transcendental Functions*; Robert, E., Ed.; Krieger Publishing Co., Inc.: Melbourne, Australia, 1981; Volume III.

34. Joshi, S.; Mittal, E.; Pandey, R.M. On Euler type integrals involving extended Mittag-Leffler functions. *Bol. Soc. Parana. Mat.* **2020**, *38*, 125–134.
35. Mainardi, F. On some properties of the Mittag-Leffler function $E_{\alpha}(-t^{\alpha})$, completely monotone for $t > 0$ with $0 < \alpha < 1$. *Discrete Contin. Dyn. Syst. Ser. B* **2014**, *19*, 2267–2278. doi:10.3934/dcdsb.2014.19.2267. [[CrossRef](#)]
36. Schneider, W.R. Completely monotone generalized Mittag-Leffler functions. *Exposition. Math.* **1996**, *14*, 3–16.
37. Podlubny, I. *Fractional Differential Equations*; Academic Press, Inc.: San Diego, CA, USA, 1999.
38. Sakamoto, K.; Yamamoto, M. Initial value/boundary value problems for fractional diffusion-wave equations and applications to some inverse problems. *J. Math. Anal. Appl.* **2011**, *382*, 426–447. doi:10.1016/j.jmaa.2011.04.058. [[CrossRef](#)]
39. Courant, R.; Hilbert, D. *Methods of Mathematical Physics*; Interscience Publishers, Inc.: New York, NY, USA, 1953; Volume I.
40. Kato, T. *Perturbation Theory for Linear Operators*; Springer: New York, NY, USA, 1966.
41. Lazopoulos, K.A.; Lazopoulos, A.K. On the Mathematical Formulation of Fractional Derivatives. *Progr. Fract. Differ. Appl.* **2019**, *5*, 261–267. doi:10.18576/pfda/050402. [[CrossRef](#)]



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).