



Article

On Hybrid Contractions in the Context of Quasi-Metric Spaces

Andreea Fulga ^{1,*}, Erdal Karapınar ^{2,3,*} and Gabriela Petrușel ^{4,†}¹ Department of Mathematics and Computer Science, Transilvania University of Brasov, Brasov 500036, Romania² Department of Medical Research, China Medical University Hospital, China Medical University, Taichung 40402, Taiwan³ Department of Mathematics, Çankaya University, Ankara 06790, Turkey⁴ Faculty of Business, Babeș-Bolyai University, Cluj-Napoca 400084, Romania; gabi.petrusel@tbs.ubbcluj.ro

* Correspondence: afulga@unitbv.ro (A.F.); karapinar@mail.cmuh.org.tw or erdalkarapinar@yahoo.com (E.K.)

† These authors contributed equally to this work.

Received: 1 April 2020; Accepted: 27 April 2020; Published: 29 April 2020



Abstract: In this manuscript, we will investigate the existence of fixed points for mappings that satisfy some hybrid type contraction conditions in the setting of quasi-metric spaces. We provide examples to assure the validity of the given results. The results of this paper generalize several known theorems in the recent literature.

Keywords: contractions; hybrid contractions; quasi-metric spaces; metric spaces

1. Introduction and Preliminaries

Roughly speaking, a quasi-metric is a distance function that is not symmetry but satisfies both the triangle inequality and self-distance property. The notion of quasi-metric was first introduced by Wilson in 1930s [1]. This is a subject of intensive research not only in the setting of topology [2–4] and functional analysis, but also several qualitative sciences, such as theoretical computer science [5–8], biology [9], and many other qualitative disciplines. In particular, as it is mentioned in [10], the notion of quasi-metric plays crucial roles in several distinct branches of mathematics, such as in the existence and uniqueness of iterated function systems' attractor (fractal), in the existence and uniqueness of Hamilton-Jacobi equations, and so on.

Another crucial notion that has no metric counterpart is that of an engaged partial order. Each partial order can be associated with a quasi-metric, and vice versa. Consequently, quasi-metric not only generalizes the concept of the metric, but also partial orders. This is a crucial fact for both the theoretical computer science applications and also has significance in the framework of biology [9].

For the sake of the completeness, we shall give the formal definition of quasi-metric. Throughout the paper, X is a nonempty set A distance function $q : X \times X \rightarrow [0, \infty)$ is called a quasi-metric on X if

$$(q_1) \quad q(u, v) = 0 \Leftrightarrow u = v;$$

$$(q_2) \quad q(u, w) \leq q(u, v) + q(v, w), \text{ for all } u, v, w \in X.$$

In addition, the pair (X, q) is called a quasi-metric space.

In what follows, we indicate the close relation between a standard metric and a quasi-metric. Given q be a quasi-metric on X , it is clear that the function $q_* : X \times X \rightarrow [0, \infty)$ defined by $q_*(u, v) = q(v, u)$ forms also a quasi-metric and it is also called the dual (conjugate) of q . The functions $d_1, d_2 : X \times X \rightarrow [0, \infty)$, where

$$\begin{aligned} d_1(v, u) &= q(u, v) + q_*(u, v), \\ d_2(v, u) &= \max \{q(u, v), q_*(u, v)\} \end{aligned}$$

form standard metrics on X .

We will provide an overview of quasi-metric spaces, presenting the notions of convergence, completeness, and continuity.

Let $\{u_n\}$ be a sequence in X , and $u \in X$, where (X, q) a quasi-metric space. We say that:

1. $\{u_n\}$ converges to u if and only if

$$\lim_{n \rightarrow \infty} q(u_n, u) = \lim_{n \rightarrow \infty} q(u, u_n) = 0. \quad (1)$$

2. $\{u_n\}$ is left-Cauchy if and only if for every $\epsilon > 0$ there exists a positive integer $k = k(\epsilon)$ such that $q(u_n, u_m) < \epsilon$ for all $n \geq m > k$.
3. $\{u_n\}$ is right-Cauchy if and only if for every $\epsilon > 0$ there exists a positive integer $k = k(\epsilon)$ such that $q(u_n, u_m) < \epsilon$ for all $m \geq n > k$.
4. $\{u_n\}$ is Cauchy if and only if it is left-Cauchy and right-Cauchy.

We would remark here that, in a quasi-metric space (X, q) , the limit for a convergent sequence is unique. Indeed, if $u_n \rightarrow u$, for all $v \in X$, we have

$$\lim_{n \rightarrow \infty} q(u_n, v) = q(u, v) \text{ and } \lim_{n \rightarrow \infty} q(v, u_n) = q(v, u).$$

A quasi-metric space (X, q) is said to be: complete (respectively, left-complete or right-complete) if and only if each Cauchy sequence (respectively, left-Cauchy sequence or right-Cauchy sequence) in X is convergent. Notice, in this context, that “right completeness” is equivalent to “Smyth completeness” [11]. See also [12].

A mapping $T : X \rightarrow X$ is continuous provided that, for any sequence $\{u_n\}$ in X such that $u_n \rightarrow u \in X$, the sequence $\{Tu_n\}$ converges to Tu , that is,

$$\lim_{n \rightarrow \infty} q(Tu_n, Tu) = \lim_{n \rightarrow \infty} q(Tu, Tu_n) = 0 \quad (2)$$

If $T : X \rightarrow X$, then the fixed point set of T is $\mathcal{F}_T(X) := \{\chi \in X : T\chi = \chi\}$.

A mapping $\zeta : [0, \infty) \times [0, \infty) \rightarrow \mathbb{R}$ is called an *extended simulation function* if the following axioms are fulfilled:

- (z_d) $\zeta(t, s) < s - t$ for all $t, s > 0$;
- (z₀) $\zeta(t, 0) \leq 0$ for every $t \geq 0$ and $\zeta(t, 0) = 0 \Leftrightarrow t = 0$.

Notice that the axiom (z_d) implies that $\zeta(t, t) < 0$ for all $t > 0$. Let us denote by \mathcal{Z} the family of all extended simulation functions $\zeta : [0, \infty) \times [0, \infty) \rightarrow \mathbb{R}$.

A function $\varphi : [0, \infty) \rightarrow [0, \infty)$ is called a *comparison function* [13] if:

- (c₁) φ is increasing;
- (c₂) $\lim_{n \rightarrow \infty} \varphi^n(t) = 0$, for $t \in [0, \infty)$.

Proposition 1. If φ is a comparison function, then:

- (i) each φ^k is also a comparison function, for all $k \in \mathbb{N}$;
- (ii) φ is continuous at 0;
- (iii) $\varphi(0) = 0$ and $\varphi(t) < t$ for all $t > 0$.

A function $\psi : [0, \infty) \rightarrow [0, \infty)$ is called a *c-comparison function* [13,14] if:

- (cc₁) ψ is increasing;
- (cc₂) $\sum_{n=0}^{\infty} \psi^n(t) < \infty$, for all $t \in (0, \infty)$.

We denote by Ψ the family of *c-comparison functions*. In some papers, instead of a *c-comparison function*, the term of strong comparison function is used. See [13].

Remark 1. Any *c-comparison function* is a comparison function.

Let $\alpha : X \times X \rightarrow [0, \infty)$ be a function. We say that a mapping $T : X \rightarrow X$ is α -orbital admissible [15] if for each $u \in X$ we have

$$\alpha(u, Tu) \geq 1 \Rightarrow \alpha(Tu, T^2u) \geq 1.$$

Lemma 1. Let $T : X \rightarrow X$ be an α -orbital admissible function. If there exists $u_0 \in X$ such that $\alpha(u_0, Tu_0) \geq 1$ and $\alpha(Tu_0, u_0) \geq 1$, then the sequence $(u_n)_{n \in \mathbb{N}}$, defined by $u_n = Tu_{n-1}$, $n \in \mathbb{N}$ satisfies the following relations:

$$\alpha(u_n, u_{n+1}) \geq 1 \text{ and } \alpha(u_{n+1}, u_n) \geq 1, \text{ for all } n \in \mathbb{N}_0.$$

We say that the set X is *regular* with respect to mapping $\alpha : X \times X \rightarrow [0, \infty)$ if the following condition is satisfied: if $\{u_n\}$ is a sequence in X such that $\alpha(u_{n+1}, u_n) \geq 1$ and $\alpha(u_n, u_{n+1}) \geq 1$ for any $n \in \mathbb{N}$ and $u_n \rightarrow u \in X$ as $n \rightarrow \infty$, then there exists a subsequence $\{u_{n(i)}\}$ of $\{u_n\}$ such that

$$\alpha(u_{n(i)}, u) \geq 1 \text{ and } \alpha(u, u_{n(i)}) \geq 1,$$

for each i .

In this manuscript, we will investigate the existence of fixed points for mappings that satisfy some hybrid type contraction conditions in the setting of quasi-metric spaces. We provide examples to assure the validity of the given results. The results of this paper generalize several known theorems in the recent literature, see [13,14,16–25].

2. Main Results

We start with the formal definition of hybrid almost contraction of type \mathbb{I} .

Definition 1. Let (X, q) be a quasi-metric space. We say that the mapping $T : X \rightarrow X$ is a hybrid almost contraction of type \mathbb{I} , if there exist $\zeta \in \mathcal{Z}$, $\psi \in \Psi$, $p \geq 0$, $L \geq 0$ and $a_1, a_2, a_3 \in [0, 1]$ with $a_1 + a_2 > 0$, $a_1 + a_2 + a_3 = 1$, such that, for all distinct $u, v \in X$, we have

$$\frac{1}{2} \min \{q(u, Tu), q(v, Tv)q(Tv, v)\} \leq q(u, v) \text{ implies} \quad (3)$$

$$\zeta(\alpha(u, v)q(Tu, Tv), \psi(I_p(u, v) + L\mathcal{N}(u, v))) \geq 0,$$

where

$$I_p(u, v) = \begin{cases} [a_1(q(u, v))^p + a_2(q(u, Tu))^p + a_3(q(v, Tv))^p]^{1/p}, & \text{for } p > 0, \\ (q(u, v))^{a_1} \cdot (q(u, Tu))^{a_2} \cdot (q(v, Tv))^{a_3} & \text{for } p = 0 \end{cases}$$

and

$$\mathcal{N}(u, v) = \min \{q(u, Tv), q(v, Tu)\}.$$

Theorem 1. Let (X, q) be a complete quasi-metric space and $\alpha : X \times X \rightarrow [0, \infty)$ be a mapping such that:

- (i) $u = Tu$ implies $\alpha(u, v) > 0$ for every $v \in X$;
- (ii) $v = Tv$ implies $\alpha(u, v) > 0$ for every $u \in X$.

Suppose that $T : X \rightarrow X$ is an hybrid almost contraction of type II and

- (C₁) T is α -orbital admissible;
- (C₂) there exists $u_0 \in X$ such that $\alpha(u_0, Tu_0) \geq 1$ and $\alpha(Tu_0, u_0) \geq 1$;
- (C₃) T is continuous.

Then, T has a fixed point.

Proof. Let the sequence $\{u_n\}$ in X be defined by

$$u_1 = Tu_0, u_2 = Tu_1, \dots, u_n = Tu_{n-1} = T^{n-1}u_0$$

where $u_0 \in X$ is the point such that, from (C₂), $\alpha(u_0, Tu_0) \geq 1$ and $\alpha(Tu_0, u_0) \geq 1$. Indubitably, for all $n \in \mathbb{N}$, we have $u_{n+1} \neq u_n$. As a matter of fact, if we suppose that there is $N_0 \in \mathbb{N}$ such that $u_{N_0} = u_{N_0+1}$, from the manner in which the sequence $\{u_n\}$ was defined, we get

$$u_{N_0} = Tu_{N_0} = u_{N_0+1}$$

so that the fixed point of T is u_{N_0} and the proof is completed. Thus, choosing $u = u_{n-1}$ respectively $v = u_n$ and since $\frac{1}{2} \min \{q(u_{n-1}, Tu_{n-1}), q(u_n, Tu_n), q(Tu_{n-1}, u_n)\} \leq \frac{1}{2} q(u_{n-1}, Tu_{n-1}) < q(u_{n-1}, u_n)$ holds for any $n \in \mathbb{N}$, by (3), we get

$$\zeta(\alpha(u_{n-1}, u_n)q(Tu_{n-1}, Tu_n), \psi(I_p(u_{n-1}, u_n) + L\mathcal{N}(u_{n-1}, u_n))) \geq 0. \quad (4)$$

In other words, taking into account (z_d) ,

$$0 \leq \psi(I_p(u_{n-1}, u_n) + L\mathcal{N}(u_{n-1}, u_n)) - \alpha(u_{n-1}, u_n)q(Tu_{n-1}, Tu_n). \quad (5)$$

However, T is an α -orbital admissible and, on the strength of Lemma (1), the above inequality yields

$$q(Tu_{n-1}, Tu_n) \leq \alpha(u_{n-1}, u_n)q(Tu_{n-1}, Tu_n) \leq \psi(I_p(u_{n-1}, u_n) + L\mathcal{N}(u_{n-1}, u_n)). \quad (6)$$

Since

$$\begin{aligned} \mathcal{N}(u_{n-1}, u_n) &= \min \{q(u_{n-1}, Tu_n), q(u_n, Tu_{n-1})\} \\ &= \min \{q(u_{n-1}, u_n), q(u_n, u_n)\} = 0, \end{aligned} \quad (7)$$

the inequality (6) becomes

$$q(Tu_{n-1}, Tu_n) \leq \psi(I_p(u_{n-1}, u_n)). \quad (8)$$

In addition, by taking $u = u_n$, respectively, $v = u_{n-1}$, we have

$$\frac{1}{2} \min \{q(u_n, Tu_n), q(u_{n-1}, Tu_{n-1}), q(Tu_{n-1}, u_{n-1})\} \leq \frac{1}{2} \min \{q(u_n, u_{n+1}), q(u_{n-1}, u_n), q(u_n, u_{n-1})\} < q(u_n, u_{n-1}).$$

As a consequence, (3) becomes

$$\zeta(\alpha(u_n, u_{n-1})q(Tu_n, Tu_{n-1}), \psi(I_p(u_n, u_{n-1}) + L\mathcal{N}(u_n, u_{n-1}))) \geq 0, \quad (9)$$

or, taking into account (z_d) ,

$$0 \leq \psi(I_p(u_n, u_{n-1}) + L\mathcal{N}(u_n, u_{n-1})) - \alpha(u_n, u_{n-1})q(Tu_n, Tu_{n-1}).$$

By Lemma (1), the above inequality yields

$$\begin{aligned} q(u_{n+1}, u_n) &= q(Tu_n, Tu_{n-1}) \leq \alpha(u_n, u_{n-1})q(Tu_n, Tu_{n-1}) \\ &\leq \psi(I_p(u_n, u_{n-1}) + L\mathcal{N}(u_n, u_{n-1})). \end{aligned} \quad (10)$$

However,

$$\begin{aligned} \mathcal{N}(u_n, u_{n-1}) &= \min \{q(u_n, Tu_{n-1}), q(u_{n-1}, Tu_n)\} \\ &= \min \{q(u_n, u_n), q(u_{n-1}, u_{n+1})\} = 0, \end{aligned} \quad (11)$$

and then we get

$$q(Tu_n, Tu_{n-1}) \leq \psi(I_p(u_n, u_{n-1})). \quad (12)$$

From this point of the proof, we will consider the two cases separately: $p > 0$ and $p = 0$.

Case 1. For the case $p > 0$,

$$\begin{aligned} I_p(u_{n-1}, u_n) &= [a_1(q(u_{n-1}, u_n))^p + a_2(q(u_{n-1}, Tu_{n-1}))^p + a_3(q(u_n, Tu_n))^p]^{1/p} \\ &= [a_1(q(u_{n-1}, u_n))^p + a_2(q(u_{n-1}, u_n))^p + a_3(q(u_n, u_{n+1}))^p]^{1/p} \\ &= [(a_1 + a_2)(q(u_{n-1}, u_n))^p + a_3(q(u_n, u_{n+1}))^p]^{1/p} \end{aligned}$$

and the inequality (6) becomes

$$q(u_n, u_{n+1}) = q(Tu_{n-1}, Tu_n) \leq \psi([(a_1 + a_2)(q(u_{n-1}, u_n))^p + a_3(q(u_n, u_{n+1}))^p]^{1/p}). \quad (13)$$

Onward, being a c -comparison function, ψ satisfies (iii) by Proposition 1 that is $\psi(t) < t$ for any $t > 0$, we obtain

$$\begin{aligned} q(u_n, u_{n+1}) &\leq \psi([(a_1 + a_2)(q(u_{n-1}, u_n))^p + a_3(q(u_n, u_{n+1}))^p]^{1/p}) \\ &< [(a_1 + a_2)(q(u_{n-1}, u_n))^p + (1 - a_1 - a_2)(q(u_n, u_{n+1}))^p]^{1/p}, \end{aligned}$$

which is equivalent with

$$(a_1 + a_2)[q(u_n, u_{n+1})]^p < (a_1 + a_2)[q(u_{n-1}, u_n)]^p,$$

or (since $a_1 + a_2 > 0$)

$$q(u_n, u_{n+1}) < q(u_{n-1}, u_n). \quad (14)$$

Using the fact that $\psi \in \Psi$ is increasing, by (13), we have

$$q(u_n, u_{n+1}) < \psi(q(u_{n-1}, u_n)) < \psi^2(q(u_{n-2}, u_{n-1})) < \dots < \psi^n(q(u_0, u_1)) \quad (15)$$

Let now $l \geq 1$. By using (15) and the triangle inequality, we get

$$\begin{aligned} q(u_n, u_{n+l}) &\leq q(u_n, u_{n+1}) + \dots + q(u_{n+l-1}, u_{n+l}) \\ &\leq \sum_{j=n}^{n+l-1} \psi^j(q(u_0, u_1)) \\ &\leq \sum_{j=n}^{\infty} \psi^j(q(u_0, u_1)). \end{aligned} \quad (16)$$

Letting $n \rightarrow \infty$ in the above inequality, we derive that $\sum_{j=n}^{\infty} \psi^j(q(u_0, u_1)) \rightarrow 0$. Hence, $q(u_n, u_{n+l}) \rightarrow 0$ as $n \rightarrow \infty$. Thus, $\{u_n\}$ is a right-Cauchy sequence in (X, q) .

Similarly, since

$$\begin{aligned} I_p(u_n, u_{n-1}) &= [a_1(q(u_n, u_{n-1}))^p + a_2(q(u_n, Tu_n))^p + a_3(q(u_{n-1}, Tu_{n-1}))^p]^{1/p} \\ &= [a_1(q(u_n, u_{n-1}))^p + a_2(q(u_n, u_{n+1}))^p + a_3(q(u_{n-1}, u_n))^p]^{1/p}, \end{aligned}$$

the inequality (12) becomes

$$\begin{aligned} q(u_{n+1}, u_n) &\leq \psi(I_p(u_n, u_{n-1})) < I_p(u_n, u_{n-1}) \\ &= [a_1(q(u_n, u_{n-1}))^p + a_2(q(u_n, u_{n+1}))^p + a_3(q(u_{n-1}, u_n))^p]^{1/p}. \end{aligned} \quad (17)$$

Taking into account (14), we get

$$\begin{aligned} (q(u_{n+1}, u_n))^p &< a_1(q(u_n, u_{n-1}))^p + a_2(q(u_n, u_{n+1}))^p + a_3(q(u_{n-1}, u_n))^p \\ &= a_1(q(u_n, u_{n-1}))^p + a_2(q(u_n, u_{n+1}))^p + (1 - a_1 - a_2)(q(u_{n-1}, u_n))^p \\ &< a_1(q(u_n, u_{n-1}))^p + (1 - a_1)(q(u_{n-1}, u_n))^p, \text{ for any } n \in \mathbb{N}. \end{aligned}$$

We are able to examine it with the following cases.

- a. If $q(u_n, u_{n-1}) < q(u_{n-1}, u_n)$ for any $n \in \mathbb{N}$, the above inequality becomes

$$(q(u_{n+1}, u_n))^p < (q(u_{n-1}, u_n))^p,$$

and then, together with (15),

$$q(u_{n+1}, u_n) < q(u_{n-1}, u_n) < \psi^{n-1}(q(u_0, u_1)), \forall n \geq 1. \quad (18)$$

From the triangle inequality and (18), for all $l \geq 1$, we get that

$$\begin{aligned} q(u_{n+l}, u_n) &\leq q(u_{n+l}, u_{n+l-1}) + \dots + q(u_{n+1}, u_n) \\ &\leq \sum_{j=n}^{n+l-1} \psi^j(q(u_0, u_1)) \\ &\leq \sum_{j=n}^{\infty} \psi^j(q(u_0, u_1)) \rightarrow 0 \text{ as } n \rightarrow \infty. \end{aligned}$$

- b. If, for any $n \in \mathbb{N}$, $q(u_n, u_{n-1}) \leq q(u_{n-1}, u_n)$, we have

$$q(u_{n+1}, u_n) < q(u_n, u_{n-1})$$

and, from (17), regarding $\psi \in \Psi$, we get that

$$q(u_{n+1}, u_n) < \psi(q(u_n, u_{n-1})) < \dots < \psi^n(q(u_1, u_0)). \quad (19)$$

Again, by triangle inequality,

$$\begin{aligned} q(u_{n+l}, u_n) &\leq q(u_{n+l}, u_{n+l-1}) + \dots + q(u_{n+1}, u_n) \\ &\leq \sum_{j=n}^{n+l-1} \psi^j(q(u_1, u_0)) \\ &\leq \sum_{j=n}^{\infty} \psi^j(q(u_1, u_0)) \rightarrow 0 \text{ as } n \rightarrow \infty. \end{aligned}$$

- c. If $q(u_i, u_{i-1}) \leq q(u_{i-1}, u_i)$ for some $i \in \mathbb{N}$ and $q(u_k, u_{k-1}) > q(u_{k-1}, u_k)$ for some $k \in \mathbb{N}$, then we have for $l \in \mathbb{N}$

$$\begin{aligned} q(u_{n+l}, u_n) &\leq q(u_{n+l}, u_{n+l-1}) + \dots + q(u_{n+1}, u_n) \\ &\leq \sum_{j=n}^{\infty} \psi^j(q(u_1, u_0)) + \sum_{j=n}^{\infty} \psi^j(q(u_0, u_1)) \rightarrow 0 \text{ as } n \rightarrow \infty. \end{aligned}$$

Therefore, we proved that $\{u_n\}$ is a left-Cauchy in (X, q) .

Thus, being left and right Cauchy, the sequence $\{u_n\}$ is a Cauchy in complete quasi-metric space (X, q) , which implies that there is $u^* \in X$ such that

$$\lim_{n \rightarrow \infty} q(u_n, u^*) = \lim_{n \rightarrow \infty} q(u^*, u_n) = 0. \quad (20)$$

Using the continuity of T and (q1), we have

$$\lim_{n \rightarrow \infty} q(u_n, Tu^*) = \lim_{n \rightarrow \infty} q(Tu_{n-1}, Tu^*) = 0,$$

$$\lim_{n \rightarrow \infty} q(Tu^*, u_n) = \lim_{n \rightarrow \infty} q(Tu^*, Tu_{n-1}) = 0$$

and so

$$\lim_{n \rightarrow \infty} q(u_n, Tu^*) = \lim_{n \rightarrow \infty} q(Tu^*, u_n) = 0. \quad (21)$$

It follows from (20) and (21) that $Tu^* = u^*$, that is, u^* is a fixed point of T .

Case 2. In the case $p = 0$, we have

$$\begin{aligned} I_p(u_{n-1}, u_n) &= (q(u_{n-1}, u_n))^{a_1} \cdot (q(u_{n-1}, Tu_{n-1}))^{a_2} \cdot (q(u_n, Tu_n))^{a_3} \\ &= (q(u_{n-1}, u_n))^{a_1} \cdot (q(u_{n-1}, u_n))^{a_2} \cdot (q(u_n, u_{n+1}))^{a_3}. \end{aligned}$$

Replacing in (6) and taking into account (7), we get

$$\begin{aligned} q(u_n, u_{n+1}) &= q(Tu_{n-1}, Tu_n) \leq \alpha(u_{n-1}, u_n) q(Tu_{n-1}, Tu_n) \leq \psi(I_p(u_{n-1}, u_n)) \\ &< I_p(u_{n-1}, u_n) = (q(u_{n-1}, u_n))^{a_1+a_2} \cdot (q(u_n, u_{n+1}))^{a_3} \end{aligned} \quad (22)$$

and we deduce that

$$(q(u_n, u_{n+1}))^{a_1+a_2} < (q(u_{n-1}, u_n))^{a_1+a_2}.$$

Thus, taking into account $a_1 + a_2 > 0$, we have

$$q(u_n, u_{n+1}) < q(u_{n-1}, u_n) \quad (23)$$

and, from (22), since $\psi \in \Psi$ we are able to say that, for any $n \in \mathbb{N}$,

$$q(u_n, u_{n+1}) \leq \psi(q(u_{n-1}, u_n)) < \dots < \psi^n(q(u_0, u_1)). \quad (24)$$

Following the above lines and using the triangle inequality, we obtain that the sequence u_n is right Cauchy. Likewise, because

$$\begin{aligned} I_p(u_n, u_{n-1}) &= (q(u_n, u_{n-1}))^{a_1} \cdot (q(u_n, Tu_n))^{a_2} \cdot (q(u_{n-1}, Tu_{n-1}))^{a_3} \\ &= (q(u_n, u_{n-1}))^{a_1} \cdot (q(u_n, u_{n+1}))^{a_2} \cdot (q(u_{n-1}, u_n))^{a_3}, \end{aligned}$$

taking into account (11) and (23), we have

$$\begin{aligned} q(u_{n+1}, u_n) &= q(Tu_n, Tu_{n-1}) \leq \alpha(u_n, u_{n-1}) q(Tu_n, Tu_{n-1}) \leq \psi(I_p(u_n, u_{n-1})) \\ &< I_p(u_n, u_{n-1}) = (q(u_n, u_{n-1}))^{a_1} \cdot (q(u_n, u_{n+1}))^{a_2} \cdot (q(u_{n-1}, u_n))^{a_3} \\ &< (q(u_n, u_{n-1}))^{a_1} \cdot (q(u_{n-1}, u_n))^{a_2+a_3} \\ &\leq (\max\{q(u_n, u_{n-1}), q(u_{n-1}, u_n)\})^{a_1+a_2+a_3} \\ &= \max\{q(u_n, u_{n-1}), q(u_{n-1}, u_n)\}. \end{aligned}$$

We must examine two cases.

If $\max\{q(u_n, u_{n-1}), q(u_{n-1}, u_n)\} = q(u_{n-1}, u_n)$, then since $q(u_{n-1}, u_n) > 0$, we get that

$$q(u_{n+1}, u_n) \leq \psi(q(u_{n-1}, u_n)),$$

and recursively

$$q(u_{n+1}, u_n) \leq \psi^n(q(u_0, u_1)). \quad (25)$$

If $\max\{q(u_n, u_{n-1}), q(u_{n-1}, u_n)\} = q(u_n, u_{n-1})$, we have

$$q(u_{n+1}, u_n) \leq \psi(q(u_n, u_{n-1})) < \dots < \psi^n(q(u_1, u_0)). \quad (26)$$

Therefore, by combining (25) with (26), we derive (due to (c_2)) that

$$\lim_{n \rightarrow \infty} q(u_{n+1}, u_n) = \lim_{n \rightarrow \infty} \max\{\psi^n(q(u_0, u_1)), \psi^n(q(u_1, u_0))\} = 0.$$

Again, using the triangle inequality, and the above inequalities for all $l \geq 1$, we get

$$\begin{aligned} q(u_{n+l}, u_n) &\leq q(u_{n+l}, u_{n+l-1}) + \dots + q(u_{n+1}, u_n) \\ &\leq \sum_{j=n}^{\infty} \psi^j(q(u_1, u_0)) + \sum_{j=n}^{\infty} \psi^j(q(u_0, u_1)) \rightarrow 0 \text{ as } n \rightarrow \infty, \end{aligned}$$

that is, the sequence $\{u_n\}$ is left Cauchy, so that is a Cauchy sequence in a complete quasi-metric space (X, q) . Thus, there is $u^* \in X$ such that

$$\lim_{n \rightarrow \infty} q(u^*, u_n) = 0 = \lim_{n \rightarrow \infty} q(u^*, u_n). \quad (27)$$

Of course, using (q_1) and the continuity of T , we have $Tu^* = u^*$. \square

Corollary 1. Let (X, q) be a complete quasi-metric space, a function $\alpha : X \times X \rightarrow [0, \infty)$ and a mapping $T : X \rightarrow X$ such that there exist $\zeta \in \mathcal{Z}$ and $\psi \in \Psi$ such that, for $p \geq 0$, $L \geq 0$ and $a_1, a_2, a_3 \in [0, 1)$ with $a_1 + a_2 > 0$ and $a_1 + a_2 + a_3 = 1$, we have

$$\zeta(\alpha(u, v)q(Tu, Tv), \psi(I_p(u, v) + L\mathcal{N}(u, v))) \geq 0, \text{ for all distinct } u, v \in X. \quad (28)$$

Suppose also that the following assumptions hold:

- (i) $u = Tu$ implies $\alpha(u, v) > 0$ for every $v \in X$;
- (ii) $v = Tv$ implies $\alpha(u, v) > 0$ for every $u \in X$;
- (i) T is α -orbital admissible;
- (ii) there exists $u_0 \in X$ such that $\alpha(u_0, Tu_0) \geq 1$ and $\alpha(Tu_0, u_0) \geq 1$;
- (iv) T is continuous.

Then, T has a fixed point.

Remark 2. Of course, in particular letting $L = 0$ in the above Corollary, we find Theorem 2.1. in [16].

Corollary 2. Let (X, q) be a complete quasi-metric space and a mapping $T : X \rightarrow X$ such that there exist $\zeta \in \mathcal{Z}$ and $\psi \in \Psi$ such that, for $p \geq 0$, $L \geq 0$ and $a_1, a_2, a_3 \in [0, 1)$ with $a_1 + a_2 > 0$ and $a_1 + a_2 + a_3 = 1$, we have

$$\zeta(q(Tu, Tv), \psi(I_p(u, v) + L\mathcal{N}(u, v))) \geq 0, \text{ for all distinct } u, v \in X. \quad (29)$$

Then, T has a fixed point.

Proof. Let $\alpha(u, v) = 1$ in Corollary 1. \square

Corollary 3. Let (X, q) be a complete quasi-metric space, a function $\alpha : X \times X \rightarrow [0, \infty)$ and a continuous mapping $T : X \rightarrow X$ such that there exist $\psi \in \Psi$ such that, for $p \geq 0$ and $a_1, a_2, a_3 \in [0, 1)$ with $a_1 + a_2 > 0$ and $a_1 + a_2 + a_3 = 1$, we have

$$\alpha(u, v)q(Tu, Tv) \leq \psi(I_p(u, v)), \text{ for all distinct } u, v \in X. \quad (30)$$

Suppose that there exists $u_0 \in X$ such that $\alpha(u_0, Tu_0) \geq 1$ and $\alpha(Tu_0, u_0) \geq 1$. Then, T has a fixed point.

Proof. Let $\zeta(t, s) = \psi(s) - t$ in Corollary 1. \square

Moreover, it easy to see that Theorem 1 is a generalization of Theorem 2.1 in [18] in the context of quasi-metric space. Indeed, if we take $L = 0$ and $p = 0$ in Corollary 3, we find:

Corollary 4. Let (X, q) be a complete quasi-metric space, a function $\alpha : X \times X \rightarrow [0, \infty)$, and a continuous mapping $T : X \rightarrow X$ such that there exists $\psi \in \Psi$ such that, for $a_1, a_2, a_3 \in [0, 1)$ with $a_1 + a_2 > 0$ and $a_1 + a_2 + a_3 = 1$, we have

$$\alpha(u, v)q(Tu, Tv) \leq \psi((q(u, v))^{a_1} \cdot (q(u, Tu))^{a_2} \cdot (q(v, Tv))^{a_3}), \text{ for all distinct } u, v \in X. \quad (31)$$

Suppose that there exists $u_0 \in X$ such that $\alpha(u_0, Tu_0) \geq 1$ and $\alpha(Tu_0, u_0) \geq 1$. Then, T has a fixed point.

Inspired by the example of [10], we consider the following:

Example 1. Let the set $X = [1, \infty)$ and the quasi-metric $q : X \times X \rightarrow [0, \infty)$ given by

$$q(u, v) = \begin{cases} \ln v - \ln u, & \text{if } u \leq v \\ \frac{1}{3}(\ln u - \ln v), & \text{if } u > v \end{cases}.$$

(see Example 4.1 in [10].) Let the mapping $T : X \rightarrow X$, be defined by

$$Tu = \begin{cases} 1, & \text{if } u \in [1, 2] \\ e^{u-2}, & \text{if } u \in (2, \infty) \end{cases}$$

and the function $\alpha : X \times X \rightarrow [0, \infty)$ be defined by

$$\alpha(u, v) = \begin{cases} 2, & \text{if } u, v \in [1, 2) \\ 3, & \text{if } u = 1, v = 2 \text{ or } u = 2, v = 1 \\ 6, & \text{if } u = 3, v = 2 \\ 0, & \text{otherwise} \end{cases}.$$

Since the mapping T is continuous and for $u = 2$, $\alpha(2, T2) = \alpha(2, 1) = 3$ and $\alpha(T2, 2) = \alpha(1, 2) = 3$, we have that the assumptions $(C_2), (C_3)$ are satisfied. Moreover, for any $u \in [1, 2)$, we have

$$\alpha(u, Tu) = \alpha(u, 1) = 2 \Rightarrow \alpha(T1, T^2 1) = \alpha(1, 1) = 2$$

and

$$\alpha(2, T2) = \alpha(2, 1) = 3 \Rightarrow \alpha(T2, T^2 2) = \alpha(1, 1) = 2,$$

so that T is α -orbital admissible.

Choosing $\psi(t) = \frac{1}{3}t$, $p = 2$, $a_1 = a_2 = a_3 = \frac{1}{3}$ and $L = 24$, we have the following cases:

Case 1. If $u, v \in [1, 2]$, then $q(u, v) = q(1, 1) = 0$ and (3) holds for every $\zeta \in \mathcal{Z}$.

Case 2. If $u = 3, v = 2$, then we have

$$\begin{aligned} q(3, T3) &= q(3, e) = \frac{1}{3} \ln \frac{3}{e}, & q(2, T2) &= q(2, 1) = \frac{1}{3} \ln 2, & q(T2, 2) &= q(1, 2) = \ln 2, \\ q(3, 2) &= \frac{1}{3} \ln \frac{3}{2}, & q(T3, T2) &= q(e, 1) = \frac{1}{3}, \\ q(3, T2) &= q(3, 1) = \frac{1}{3} \ln 3, & q(2, T3) &= q(2, e) = \ln \frac{e}{2}. \end{aligned}$$

Thus, we have

$$\frac{1}{2} \min \{q(3, T3), q(2, T2), q(T2, 2)\} = \frac{1}{6} \ln \frac{3}{e} < \frac{1}{3} \ln \frac{3}{2} = q(3, 2)$$

and

$$\alpha(3,2)q(T3,T2) = \frac{6}{3} < \frac{1}{3} \left[\sqrt{\frac{1}{3}} \left(\left(\frac{1}{3} \ln \frac{3}{2} \right)^2 + \left(\frac{1}{3} \ln \frac{3}{e} \right)^2 + \left(\frac{1}{3} \ln 2 \right)^2 \right)^{1/2} + 24 \ln \frac{e}{2} \right] = \psi(I_p(3,2) + L\mathcal{N}(3,2)),$$

so that T is a hybrid almost contraction for any $\zeta \in \mathcal{Z}$.

The other cases are not interesting, while $\alpha(u,v) = 0$. (Consequently, the mapping T has two fixed points, $u_1 = 1$ and $u_2 \in (3,4)$.)

On the other hand, since

$$\begin{aligned} \alpha(3,2)q(T3,T2) &= 2 > \left(\frac{1}{3} \ln \frac{3}{2} \right)^\gamma \left(\frac{1}{3} \ln \frac{3}{e} \right)^\beta \left(\frac{1}{3} \ln 2 \right)^{1-\gamma-\beta} \\ &> \psi \left((q(3,2))^\gamma (q(3,T3))^\beta, (q(2,T2))^{1-\gamma-\beta} \right) \end{aligned}$$

for every $\gamma, \beta \in (0,1)$ and $\psi \in \Psi$, Theorem 2.1 in [18] can not be applied.

In particular, for the case $p = 0$, the continuity condition of T can be replaced with the regularity condition of the space X .

Theorem 2. Let (X, q) be a complete quasi-metric space, a function $\alpha : X \times X \rightarrow [0, \infty)$ and a mapping $T : X \rightarrow X$ such that there exist $\zeta \in \mathcal{Z}$, $\psi \in \Psi$, $L \geq 0$ and $a_1, a_2, a_3 \in [0, 1]$ with $a_1 + a_2 + a_3 = 1$, such that, for all distinct $u, v \in X$, we have

$$\begin{aligned} \frac{1}{2} \min \{q(u, Tu), q(v, Tv), q(Tv, v)\} &\leq q(u, v) \text{ implies} \\ \zeta(\alpha(u, v)q(Tu, Tv), \psi((q(u, v))^{a_1} \cdot (q(u, Tu))^{a_2} \cdot (q(v, Tv))^{a_3} + L\mathcal{N}(u, v))) &\geq 0. \end{aligned} \quad (32)$$

Suppose also that

- (i) $u = Tu$ implies $\alpha(u, v) > 0$ for every $v \in X$;
- (ii) $v = Tv$ implies $\alpha(u, v) > 0$ for every $u \in X$;
- (C₁) T is α -orbital admissible;
- (C₂) there exists $u_0 \in X$ such that $\alpha(u_0, Tu_0) \geq 1$ and $\alpha(Tu_0, u_0) \geq 1$;
- (C₃) X is regular with respect to the mapping α .

Then, T has a fixed point.

Proof. From the above theorem, there exists $u^* \in X$ such that (27) hold. In what follows, we claim that

$$\begin{aligned} \frac{1}{2} \min \left\{ q(u^*, Tu^*), q(u_{n(i)}, Tu_{n(i)}), q(Tu_{n(i)}, u_{n(i)}) \right\} &\leq q(u^*, u_{n(i)}) \quad \text{or} \\ \frac{1}{2} \min \left\{ q(u_{n(i)-1}, Tu_{n(i)-1}), q(u^*, Tu^*), q(Tu^*, u^*) \right\} &\leq q(u_{n(i)-1}, u^*). \end{aligned} \quad (33)$$

Indeed, using the method of Reductio ad Absurdum, we assume that that there exists $k \in \mathbb{N}$ such that

$$\begin{aligned} \frac{1}{2} \min \{q(u^*, Tu^*), q(u_k, Tu_k), q(Tu_k, u_k)\} &> q(u^*, u_k) \quad \text{and} \\ \frac{1}{2} \min \{q(u_{k-1}, Tu_{k-1}), q(u^*, Tu^*), q(Tu^*, u^*)\} &> q(u_{k-1}, u^*) \end{aligned}$$

Therefore, we have

$$\begin{aligned}
q(u_{k-1}, u_k) &\leq q(u_{k-1}, u^*) + q(u^*, u_k) \\
&< \frac{1}{2} \min \{q(u_{k-1}, Tu_{k-1}), q(u^*, Tu^*), q(Tu^*, u^*)\} + \frac{1}{2} \min \{q(u^*, Tu^*), q(u_{k-1}, Tu_{k-1}), q(Tu_{k-1}, u_{k-1})\} \\
&< \frac{1}{2} [\min \{q(u_{k-1}, u_k), q(u^*, Tu^*), q(Tu^*, u^*)\} + \min \{q(u^*, Tu^*), q(u_{k-1}, u_k), q(u_k, u_{k-1})\}] \\
&\leq \frac{1}{2} [q(u_{k-1}, u_k) + q(u_{k-1}, u_k)] \\
&= q(u_{k-1}, u_k),
\end{aligned}$$

which is a contradiction.

In the alternative hypothesis, if the space X is regular with respect to mapping α , we have $\alpha(u^*, u_{n(i)}) \geq 1$, where $\{u_{n(i)}\}$ is a sub-sequence of $\{u_n\}$, for $i \in \mathbb{N}$. We will suppose by *reductio ad absurdum* that $u^* \neq Tu^*$. Then, for $u = u^*$ and $v = u_{n(i)}$ in (3), we get

$$\zeta(\alpha(u^*, u_{n(i)})q(Tu^*, Tu_{n(i)}), \psi(\mathcal{A}_p(u^*, u_{n(i)}))) \geq 0.$$

Taking into account the properties of function ζ , ψ , and α , the above relation becomes

$$\begin{aligned}
q(Tu^*, u^*) &\leq q(Tu^*, Tu_n) + q(Tu_n, u^*) \leq \alpha(u^*, u_n)q(Tu^*, Tu_{n(i)}) + q(u_{n(i)+1}, u^*) \\
&\leq \psi((q(u^*, u_{n(i)}))^{a_1} \cdot (q(u^*, Tu^*))^{a_2} \cdot (q(u_{n(i)}, Tu_{n(i)}))^{a_3} + \mathcal{N}(u^*, u_{n(i)})) + q(u_{n(i)+1}, u^*),
\end{aligned}$$

Letting $i \rightarrow \infty$, we have

$$0 < q(Tu^*, u^*) < \lim_{i \rightarrow \infty} \psi((q(u^*, u_{n(i)}))^{a_1} \cdot (q(u^*, Tu^*))^{a_2} \cdot (q(u_{n(i)}, Tu_{n(i)}))^{a_3} + \mathcal{N}(u^*, u_{n(i)})) + q(u_{n(i)+1}, u^*)$$

and, since ψ is continuous in 0, $\psi(0) = 0$, we get $q(Tu^*, u^*) = 0$. \square

Corollary 5. Let (X, q) be a complete quasi-metric space and $T : X \rightarrow X$ be a given mapping. Assume that there exist $L \geq 0$, $\zeta \in \mathcal{Z}$ and $\psi \in \Psi$ such that, for all distinct $u, v \in X$, we have

$$\begin{aligned}
\frac{1}{2} \min \{q(u, Tu), q(v, Tv)q(Tv, v)\} &\leq q(u, v) \text{ implies} \\
\zeta(q(Tu, Tv), \psi(I_p(u, v) + L\mathcal{N}(u, v))) &\geq 0,
\end{aligned}$$

for all distinct $u, v \in X$. Then, T has a fixed point.

Proof. It is sufficient to take $\alpha(u, v) = 1$ for $u, v \in X$ in Theorem 5. \square

Corollary 6. Let (X, q) be a complete quasi-metric space and $T : X \rightarrow X$ be a given mapping. Assume that there exist $L \geq 0$, $\zeta \in \mathcal{Z}$ and $\psi \in \Psi$ such that, for all distinct $u, v \in X$, we have

$$\frac{1}{2} \min \{q(u, Tu), q(v, Tv)q(Tv, v)\} \leq q(u, v) \text{ implies } q(Tu, Tv) \leq kI_p(u, v)$$

for all distinct $u, v \in X$. Then, T has a fixed point.

Proof. It is sufficient to take $L = 0$, $\zeta(t, s) = k_1s - t$, $\psi(u) = k_2u$ with $k_1, k_2 \in (0, 1)$ and $k = k_1k_2$ in Corollary 5. \square

Corollary 7. Let (X, q) be a complete quasi-metric space and $T : X \rightarrow X$ a continuous mapping such that

$$\begin{aligned} \frac{1}{2} \min \{q(u, Tu), q(v, Tv)q(Tv, v)\} &\leq q(u, v) \text{ implies} \\ q(Tu, Tv) &\leq \frac{\kappa}{\sqrt{3}} \cdot \sqrt{(q(u, v))^2 + (q(u, Tu))^2 + (q(v, Tv))^2} \end{aligned} \quad (34)$$

for all distinct $u, v \in X$ and some $\kappa \in (0, 1)$. Then, T has a fixed point in X .

Proof. Let $p = 2$ and $a_1 = a_2 = a_3 = \frac{1}{3}$ in Corollary 6. \square

In the next theorem, we involve a Jaggi type expression with the hybrid contractions.

Definition 2. Let (X, q) be a quasi-metric space. A mapping $T : X \rightarrow X$ is called a hybrid almost contraction of type \mathbb{J} , if there exist $\zeta \in \mathcal{Z}$ and $\psi \in \Psi$ such that, for $p \geq 0$, $L \geq 0$ and $a_1, a_2 > 0$ with $a_1 + a_2 < 1$, we have

$$\begin{aligned} \frac{1}{2} \min \{q(u, Tu), q(v, Tv)q(Tv, v)\} &\leq q(u, v) \text{ implies} \\ \zeta(\alpha(u, v)q(Tu, Tv), \psi(J_p(u, v) + L\mathcal{N}(u, v))) &\geq 0, \end{aligned} \quad (35)$$

for all distinct $u, v \in X$, where

$$J_p(u, v) = \begin{cases} [a_1(q(u, v))^p + a_2(\frac{q(u, Tu) \cdot q(v, Tv)}{q(u, v)})^p]^{1/p}, & \text{for } p > 0 \\ (q(u, v))^{a_1} \cdot (q(u, Tu))^{a_1} \cdot (q(v, Tv))^{1-a_1-a_2}, & \text{for } p = 0 \end{cases}$$

and

$$\mathcal{N}(u, v) = \min \{q(u, Tv), q(v, Tu)\}.$$

Theorem 3. Let (X, q) be a complete quasi-metric space and $\alpha : X \times X \rightarrow [0, \infty)$ be a mapping such that:

- (i) $u = Tu$ implies $\alpha(u, v) > 0$ for every $v \in X$;
- (ii) $v = Tv$ implies $\alpha(u, v) > 0$ for every $u \in X$.

Suppose that $T : X \rightarrow X$ is a hybrid almost contraction of type \mathbb{J} such that the following assumptions hold:

- (i) T is α -orbital admissible;
- (ii) there exists $u_0 \in X$ such that $\alpha(u_0, Tu_0) \geq 1$ and $\alpha(Tu_0, u_0) \geq 1$;
- (iii) there exists $\Delta > 0$ such that, $(a_1 + a_2\Delta^{2p})^{1/p} \leq 1$ (where $p > 0$) and

$$\frac{1}{\Delta} q(u, v) \leq q(v, u) \leq \Delta q(u, v), \text{ for all } u, v \in X;$$

- (iv) T is continuous.

Then, T has a fixed point.

Proof. We will consider only the case $p > 0$ because, for $p = 0$, the expression is similar to the one in Theorem 1. By verbatim of the first lines in the proof of Theorem 1, starting from a point u_0 , we are able to build a sequence $\{u_n\} \subset X$. Onward, as in the proof of Theorem 1, we suppose that $u_{n+1} \neq u_n$ for all $n \in \mathbb{N}$ and from (35), we have $\frac{1}{2} \min \{q(u_{n-1}, Tu_{n-1}), q(u_n, Tu_n), q(Tu_n, u_n)\} \leq q(u_{n-1}, u_n)$, which implies

$$\zeta(\alpha(u_{n-1}, u_n)q(Tu_{n-1}, Tu_n), \psi(J_p(u_{n-1}, u_n) + L\mathcal{N}(u_{n-1}, u_n))) \geq 0.$$

By the axiom (z_d) , Lemma 1 and taking into account (7), this inequality becomes

$$\begin{aligned} q(u_n, u_{n+1}) &\leq \alpha(u_{n-1}, u_n) q(Tu_{n-1}, Tu_n) \leq \psi(\mathcal{I}_p(u_{n-1}, u_n) + L\mathcal{N}(u_{n-1}, u_n)) < \mathcal{I}_p(u_{n-1}, u_n) \\ &= [a_1(q(u_{n-1}, u_n))^p + a_2(\frac{q(u_{n-1}, Tu_{n-1}) \cdot q(u_n, Tu_n)}{q(u_{n-1}, u_n)})^p]^{1/p} \\ &= [a_1(q(u_{n-1}, u_n))^p + a_2(\frac{q(u_{n-1}, u_n) \cdot (q(u_n, u_{n+1}))}{q(u_{n-1}, u_n)})^p]^{1/p} \\ &= [a_1(q(u_{n-1}, u_n))^p + a_2(q(u_n, u_{n+1}))^p]^{1/p}. \end{aligned} \quad (36)$$

Thereupon,

$$q(u_n, u_{n+1}) < \left(\frac{a_1}{1 - a_2}\right)^{1/p} q(u_{n-1}, u_n) < q(u_{n-1}, u_n)$$

and then, from (36), we have $q(u_n, u_{n+1}) < \psi(q(u_{n-1}, u_n))$. Since $\psi \in \Psi$, recursively, we get

$$q(u_n, u_{n+1}) < \psi(q(u_{n-1}, u_n)) < \dots < \psi^n(q(u_0, u_1)). \quad (37)$$

In order to prove that $\{u_n\}$ is a right-Cauchy sequence, let $l \in \mathbb{N}$. From (37) and the triangle inequality, we get that

$$\begin{aligned} q(u_n, u_{n+l}) &\leq q(u_n, u_{n+1}) + \dots + q(u_{n+l-1}, u_{n+l}) \\ &\leq \sum_{j=n}^{n+l-1} \psi^j(q(u_0, u_1)) \\ &\leq \sum_{j=n}^{\infty} \psi^j(q(u_0, u_1)) \rightarrow 0, \text{ as } n \rightarrow \infty. \end{aligned}$$

We conclude that $\{q_n\}$ is a right-Cauchy sequence in (X, q) .

Substituting in (35) $u = u_n$ and $v = u_{n-1}$ and since $\frac{1}{2} \min \{q(u_n, Tu_n, q(u_{n-1}, Tu_{n-1}), q(Tu_{n-1}, u_{n-1}))\} \leq q(u_n, u_{n-1})$, we have (taking into account (11))

$$\begin{aligned} q(u_{n+1}, u_n) &\leq \alpha(u_n, u_{n-1}) q(Tu_n, Tu_{n-1}) \leq \psi(\mathcal{I}_p(u_n, u_{n-1}) + L\mathcal{N}(u_n, u_{n-1})) < \mathcal{I}_p(u_n, u_{n-1}) \\ &= [a_1(q(u_n, u_{n-1}))^p + a_2(\frac{q(u_n, u_{n+1}) \cdot (q(u_{n-1}, u_n))}{q(u_n, u_{n-1})})^p]^{1/p} \end{aligned}$$

i.e.,

$$(q(u_{n+1}, u_n))^p < a_1(q(u_n, u_{n-1}))^p + a_2(\frac{q(u_n, u_{n+1}) \cdot (q(u_{n-1}, u_n))}{q(u_n, u_{n-1})})^p.$$

On one hand, we have already proved that $q(u_n, u_{n+1}) < q(u_{n-1}, u_n)$. On the other hand, by (iii), there exists a positive constant Δ such that $q(u_{n-1}, u_n) \leq \Delta q(u_n, u_{n-1})$ for $n \in \mathbb{N}$. Thus, we have

$$\begin{aligned} (q(u_{n+1}, u_n))^p &< a_1(q(u_n, u_{n-1}))^p + a_2(\frac{(q(u_{n-1}, u_n))^2}{q(u_n, u_{n-1})})^p \\ &< a_1(q(u_n, u_{n-1}))^p + a_2(\frac{(\Delta \cdot q(u_n, u_{n-1}))^2}{q(u_n, u_{n-1})})^p \\ &= (a_1 + a_2 \Delta^{2p}) \cdot (q(u_n, u_{n-1}))^p, \end{aligned}$$

which is equivalent to the next inequality

$$q(u_{n+1}, u_n) < (a_1 + a_2 \Delta^{2p})^{1/p} q(u_n, u_{n-1}) \leq q(u_n, u_{n-1}).$$

Thus,

$$q(u_{n+1}, u_n) < \psi(q(u_n, u_{n-1})) < \dots < \psi^n(q(u_1, u_0)) \quad (38)$$

Again, considering triangle inequality, together with (38), for $l \in \mathbb{N}$, we have

$$\begin{aligned} q(u_{n+l}, u_n) &\leq q(u_{n+l}, u_{n+l-1}) + \dots + q(u_{n-1}, u_n) \\ &\leq \sum_{j=n}^{\infty} \psi^j(q(u_0, u_1)) \rightarrow 0, \text{ as } n \rightarrow \infty. \end{aligned}$$

Analogously, we deduce that $\{u_n\}$ is left-Cauchy, so that it is a Cauchy sequence in complete quasi-metric space.

Thus, there exists $u^* \in X$ such that

$$\lim_{n \rightarrow \infty} q(u_n, u^*) = \lim_{n \rightarrow \infty} q(u^*, u_n) = 0. \quad (39)$$

Under the assumption (iv), from the continuity of T and (q_1) , we have

$$\lim_{n \rightarrow \infty} q(u_n, Tu^*) = \lim_{n \rightarrow \infty} q(Tu_{n-1}, Tu^*) = 0,$$

$$\lim_{n \rightarrow \infty} q(Tu^*, u_n) = \lim_{n \rightarrow \infty} q(Tu^*, Tu_{n-1}) = 0$$

so that

$$\lim_{n \rightarrow \infty} q(u_n, Tu^*) = \lim_{n \rightarrow \infty} q(Tu^*, u_n) = 0. \quad (40)$$

Hence, $Tu^* = u^*$ that is, u^* is a fixed point of T . \square

The following is a special case for $p = 0$.

Corollary 8. Let (X, q) be a complete quasi-metric space, a function $\alpha : X \times X \rightarrow [0, \infty)$ and a mapping $T : X \rightarrow X$ such that there exist $\zeta \in \mathcal{Z}$ and $\psi \in \Psi$ such that, for $p \geq 0$, $L \geq 0$ and $a_1, a_2 \in [0, 1)$ with $a_1 + a_2 < 1$, we have

$$\zeta(\alpha(u, v)q(Tu, Tv), \psi(\mathcal{I}_p(u, v) + L\mathcal{N}(u, v))) \geq 0, \text{ for all distinct } u, v \in X. \quad (41)$$

Suppose also that the following assumptions hold:

- (i) $u = Tu$ implies $\alpha(u, v) > 0$ for every $v \in X$;
- (ii) $v = Tv$ implies $\alpha(u, v) > 0$ for every $u \in X$;
- (i) T is α -orbital admissible;
- (ii) there exists $u_0 \in X$ such that $\alpha(u_0, Tu_0) \geq 1$ and $\alpha(Tu_0, u_0) \geq 1$;
- (iii) there exists $\Delta > 0$ such that, $(a_1 + a_2\Delta^{2p})^{1/p} \leq 1$ (where $p > 0$) and

$$\frac{1}{\Delta}q(u, v) \leq q(v, u) \leq \Delta q(u, v), \text{ for all } u, v \in X;$$

- (iv) T is continuous.

Then, T has a fixed point.

Example 2. Let $X = [0, 1]$ and the function

$$q(u, v) = \begin{cases} u - v, & \text{for } u \geq v \\ 2(v - u), & \text{for } u < v \end{cases}$$

It is easy to see that the pair (X, q) forms a quasi-metric space.

Let the map $T : X \rightarrow X$ defined by

$$Tu = \begin{cases} \frac{1}{8}, & \text{for } u \in [0, \frac{1}{2}] \\ \frac{u}{4}, & \text{for } u \in [\frac{1}{2}, 1] \end{cases}$$

and choose $\zeta(u, v) = \frac{1}{2}v - u$ and $\psi(t) = \frac{1}{2}t$. For $p = 2$, $L = 0$, $\Delta = 2$, $a_1 = \frac{1}{4}$ and $a_2 = \frac{1}{32}$ because $(a_1 + a_2 \cdot \Delta^{2p})^{1/p} = \frac{1}{4} + \frac{1}{32} \cdot 2^4 = \frac{3}{4} \leq 1$, the assumption (iii) is satisfied. In this case, (41) becomes

$$\alpha(u, v)q(u, v) \leq \mathcal{I}_p(u, v) = \frac{1}{4} \sqrt{\frac{1}{4}(q(u, v))^2 + \frac{1}{32} \left(\frac{q(u, Tu)q(v, Tv)}{q(u, v)} \right)^2}. \quad (42)$$

Define $\alpha : X \times X \rightarrow [0, \infty)$ such that

$$\alpha(u, v) = \begin{cases} 3, & \text{for } u, v \in [0, \frac{1}{2}] \\ 1, & \text{for } u = 1, v = 0 \\ 0, & \text{otherwise} \end{cases}$$

It is easy to see that T is α -admissible. Indeed, we have

$$\alpha(u, v) = 3 \Rightarrow \alpha(Tu, Tv) = \alpha(1/8, 1/8) = 3, \text{ for } u, v \in [0, \frac{1}{2}]$$

and

$$\alpha(1, 0) = 1 \Rightarrow \alpha(T1, T0) = \alpha(1/4, 1/8) = 3.$$

On the other hand, for $q_0 = 0$,

$$\alpha(0, T0) = \alpha(T0, 0) = \alpha(0, 0) = 3,$$

so that the presumptions (i), (ii), and (iv) are satisfied. Of course, if $u, v \in [0, \frac{1}{2}]$, we have $q(Tu, Tv) = q(\frac{1}{8}, \frac{1}{8}) = 0$ and (41) is verified. For $u = 1$ and $v = 0$, we have $q(T1, T0) = \frac{1}{4} - \frac{1}{8} = \frac{1}{8}$, $q(0, T0) = q(0, 1/8) = 2(1/8 - 0) = 1/4$, $q(1, T1) = q(1, 1/4) = 3/4$ and

$$\begin{aligned} \alpha(1, 0)q(T1, T0) &= \frac{1}{8} \leq \frac{1}{4} \sqrt{\frac{1}{4} + \frac{1}{32} \left(\frac{3}{16} \right)^2} \\ &= \frac{1}{4} \sqrt{\frac{1}{4}(q(1, 0))^2 + \frac{1}{32} \left(\frac{q(0, T0)q(1, T1)}{q(1, 0)} \right)^2} \end{aligned} \quad (43)$$

The other cases are not interesting since $\alpha(u, v) = 0$ and the condition (42) is fulfilled trivially. Thus, the presumptions of Theorem 8 are provided and $u = \frac{1}{8}$ is the fixed point of T .

Corollary 9. Let (X, q) be a complete quasi-metric space and T be a continuous self-mapping on X . Suppose that there exist $\zeta \in \mathcal{Z}$, $\psi \in \Psi$ such that

$$\zeta(q(Tu, Tv), \psi(\mathcal{I}_p(u, v))) \geq 0,$$

for each distinct $u, v \in X$. If there exists $\Delta > 0$ such that $(a_1 + a_2 \cdot \Delta^{2p})^{1/p} \leq 1$ for $p > 0$, and $\frac{1}{\Delta}q(u, v) \leq q(v, u) \leq \Delta q(u, v)$ for all $u, v \in X$, then T has a fixed point.

Proof. It is sufficient to take $L = 0$ and $\alpha(u, v) = 1$ for $u, v \in X$ in Corollary 8. \square

Corollary 10. Let (X, q) be a complete quasi-metric space and T be a self-mapping on X . Suppose that there exists $\Delta > 0$ such that $(a_1 + a_2 \cdot \Delta^{2p})^{1/p} \leq 1$ for $p > 0$, and $\frac{1}{\Delta}q(u, v) \leq q(v, u) \leq \Delta q(u, v)$ for all $u, v \in X$. The mapping T has a fixed point provided that

$$q(Tu, Tv) \leq c \cdot \mathcal{I}_p(u, v)$$

for each distinct $u, v \in X$ and some $c \in (0, 1)$.

Proof. We set $\zeta(t, s) = c_1s - t$, $\psi(u) = c_2u$ with $c_1, c_2 \in [0, 1]$ and $c = c_1 + c_2$ in Corollary 9. \square

Remark 3. Letting $p = 0$ in Corollary 10, we find Theorem 2.2. in [20].

Example 3. Let (X, q) be the quasi-metric space, where $X = [1, \infty)$ and

$$q(u, v) = \begin{cases} u - v, & \text{for } u \geq v \\ 2(v - u), & \text{for } u < v \end{cases}$$

Let

$$Tu = \begin{cases} u^3 - 8u^2 + 19u - 9, & \text{for } u \in [1, 5] \\ \ln(u^2 - 24) + u + 6, & \text{for } u \in (5, \infty). \end{cases}$$

Consider the function ζ be arbitrary in \mathcal{Z} , $\psi \in \Psi$ with $\psi(t) = \frac{t}{\sqrt{3}}$ and $\alpha : X \times X \rightarrow [0, \infty)$ such that

$$\alpha(u, v) = \begin{cases} u^2 + 1, & \text{for } (u, v) \in \{(3, 3), (3, 4), (4, 3), (3, 1), (1, 3)\} \\ 1, & \text{for } (u, v) = (2, 1) \\ 0, & \text{otherwise.} \end{cases}$$

It is easily verified that T is α -orbital admissible. Whereas $T1 = T3 = T4 = 3$, taking into account the definition of function α , we have that the inequality (41) holds for every pair $(u, v) \in X^2 \setminus \{(2, 1)\}$. For the case $u = 2$ and $v = 1$, choosing $a_1 = \frac{1}{2}$, $a_2 = \frac{1}{48}$ and $p = 2$, we find that axiom (iii) holds. On the other hand,

$$\begin{aligned} \mathcal{I}_p(2, 1) &= \left[\frac{1}{2}(q(2, 1))^2 + \frac{1}{48} \left(\frac{q(2, T2) \cdot q(1, T1)}{q(2, 1)} \right)^2 \right]^{1/2} \\ &= \sqrt{\frac{1}{2} + \frac{1}{48} \cdot \left(\frac{q(2, 5) \cdot q(1, 3)}{q(2, 1)} \right)^2} = \sqrt{\frac{25}{2}} \end{aligned}$$

and

$$\alpha(2, 1)q(T2, T1) = q(5, 3) = 2 < \sqrt{\frac{25}{2}} = \psi(\mathcal{I}_p(2, 1)).$$

Consequently, by Theorem 8, we have that the mapping T has a fixed point in X .

On the other hand, we can observed that, for $u = 1$ and $v = 5$,

$$q(T1, T(4.5)) = q(2, 5.625) = 7.25, \quad q(1, T1) = q(1, 2) = 2, \quad q(4.5, T(4.5)) = q(4.5, 5.625) = 1.125,$$

so that, since

$$q(T1, T(4.5)) > \lambda(q(1, T1))^\alpha (q(4.5, T(4.5)))^{1-\alpha}$$

for any $\lambda \in [0, 1)$ and $\alpha \in (0, 1)$, Theorem 2.2 in [20] can not be applied.

Corollary 11. Let (X, q) be a complete quasi-metric space and $T : X \rightarrow X$ a continuous mapping. Then, T has a fixed point provided that

$$q(Tu, Tv) \leq k_1 \cdot q(u, v) + k_2 \cdot \frac{q(u, Tu)q(v, Tv)}{q(u, v)} \quad (44)$$

for each $u, v \in X$ and $k_1, k_2 \in (0, 1)$ with $k_1 + k_2 < 1$

Proof. Let $p = 1$ and $k_i = c \cdot a_i$, for $i \in \{1, 2\}$ in Corollary 10. \square

Author Contributions: All authors contributed equally and significantly in writing this article. Writing—original draft, A.F. and E.K.; writing—review and editing, A.F., G.P., and E.K. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Acknowledgments: The authors thank the anonymous referees for their remarkable comments, suggestions, and ideas that helped to improve this paper.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Wilson, W.A. On quasi-metric spaces. *Am. J. Math.* **1931**, *53*, 675–684.
2. Künzi, H.-P.A. Nonsymmetric distances and their associated withologies: about the origins of basic ideas in the area of asymmetric topology. In *Handbook of the History of General Topology*; volume 3 of Hist. Topol.; Kluwer Acad. Publ.: Dordrecht, The Netherlands, 2001; pp. 853–968.
3. Künzi, H.-P.A.; Vajner, V. Weighted quasi-metrics. In *Papers on General Topology and Applications (Flushing, NY, 1992)*; New York Acad. Sci.: New York, NY, USA, 1994; pp. 64–77.
4. Künzi, H.-P.A.; Romaguera, S. Quasi-Metric Spaces, Quasi-Metric Hyperspaces and Uniform Local Compactness. *Rend. Istit. Mat. Univ. Trieste Suppl.* **1999**, *XXX*, 133–144.
5. Romaguera, S.; Schellekens, M. On the structure of the dual complexity space: the general case. *Extracta Math.* **1998**, *13*, 249–253.
6. Romaguera, S.; Schellekens, M. Quasi-metric properties of complexity spaces. *Topology Appl.* **1999**, *98*, 311–322.
7. Romaguera, S.; Schellekens, M. Duality and quasi-normability for complexity spaces. *Appl. Gen. Topol.* **2002**, *3*, 91–112.
8. Romaguera, S.; Schellekens, M.P. Weightable quasi-metric semigroups and semilattices. *Electr. Notes Theor. Comput. Sci.* **2001**, *40*, 347–358.
9. Stojmirović, A. Quasi-metrics, Similarities and Searches: aspects of geometry of protein datasets. *arXiv* **2018**, arXiv:0810.5407.
10. Secelean, N.A.; Mathew, S.; Wardowski, D. New fixed point results in quasi-metric spaces and applications in fractals theory. *Adv. Differ. Equ.* **2019**, *2019*, 177.
11. Romaguera, S.; Tirado, P. A characterization of Smyth complete quasi-metric spaces via Caristi's fixed point theorem. *Fixed Point Theory Appl.* **2015**, *2015*, 183.

12. Romaguera, S.; Tirado, P. The Meir-Keeler fixed point theorems for quasi-metric spaces and some consequences. *Symmetry* **2019**, *11*, 741.
13. Rus, I.A. *Generalized Contractions and Applications*; Cluj University Press: Cluj-Napocca, Romania, 2001.
14. Bianchini, R.M.; Grandolfi, M. Trasformazioni di tipo contrattivo generalizzato in uno spazio metrico. *Atti Acad. Naz. Lincei, VII. Ser. Rend. Cl. Sci. Fis. Mat. Natur.* **1968**, *45*, 212–216.
15. Popescu, O. Some new fixed point theorems for α -Geraghty contractive type maps in metric spaces. *Fixed Point Theory Appl.* **2014**, *2014*, 190.
16. Agarwal, R.P.; Karapinar, E. Interpolative Rus-Reich-Ciric type contractions via simulation functions. *An. St. Univ. Ovidius Constanta Ser. Mat.* **2020**, in press.
17. Aydi, H.; Chen, C.M.; Karapinar, E. Interpolative Ciric-Reich-Rus type contractions via the Branciari distance. *Mathematics* **2019**, *7*, 84.
18. Aydi, H.; Karapinar, E.; Roldán López de Hierro, A.F. ω Interpolative Ciric-Reich-Rus type contractions. *Mathematics* **2019**, *7*, 57.
19. Jaggi, D.S. Some unique fixed point theorems. *Indian J. Pure Appl. Math.* **1977**, *8*, 223–230.
20. Karapinar, E. Revisiting the Kannan type contractions via interpolation. *Adv. Theory Nonlinear Anal. Appl.* **2018**, *2*, 85–87.
21. Karapinar, E.; Alqahtani, O.; Aydi, H. On Interpolative Hardy-Rogers type contractions. *Symmetry* **2019**, *11*, 8.
22. Karapinar, E.; Agarwal, R.P.; Aydi, H. Interpolative Reich-Rus-Ciric type contractions on partial metric spaces. *Mathematics* **2018**, *6*, 256.
23. Khojasteh, F.; Shukla, S.; Radenović, S. A new approach to the study of fixed point theorems via simulation functions. *Filomat* **2015**, *29*, 1189–1194.
24. Petruşel, A.; Rus, I.A. Fixed point theory in terms of a metric and of an order relation. *Fixed Point Theory* **2019**, *20*, 601–622.
25. Rus, I.A.; Petruşel, A.; Petruşel, G. *Fixed Point Theory*; Cluj University Press: Cluj-Napocca, Romania, 2008.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).