

Article

Group Invariant Solutions and Conserved Quantities of a (3+1)-Dimensional Generalized Kadomtsev–Petviashvili Equation

Innocent Simbanefayi ¹ and Chaudry Masood Khalique ^{1,2,3,*} 

¹ International Institute for Symmetry Analysis and Mathematical Modelling, Department of Mathematical Sciences, North-West University, Mafikeng Campus, Private Bag X 2046, Mmabatho 2735, South Africa; isimbanefayi@yahoo.com

² College of Mathematics and Systems Science, Shandong University of Science and Technology, Qingdao 266590, China

³ Department of Mathematics and Informatics, Azerbaijan University, Jeyhun Hajibeyli str., 71, Baku AZ1007, Azerbaijan

* Correspondence: Masood.Khalique@nwu.ac.za

Received: 13 May 2020; Accepted: 10 June 2020; Published: 20 June 2020



Abstract: In this work, we investigate a (3+1)-dimensional generalised Kadomtsev–Petviashvili equation, recently introduced in the literature. We determine its group invariant solutions by employing Lie symmetry methods and obtain elliptic, rational and logarithmic solutions. The solutions derived in this paper are the most general since they contain elliptic functions. Finally, we derive the conserved quantities of this equation by employing two approaches—the general multiplier approach and Ibragimov’s theorem. The importance of conservation laws is explained in the introduction. It should be pointed out that the investigation of higher dimensional nonlinear partial differential equations is vital to our perception of the real world since they are more realistic models of natural and man-made phenomena.

Keywords: (3+1)-dimensional generalised KP equation; invariant solutions; multiplier method; Ibragimov’s conservation theorem; conserved quantities

1. Introduction

The study of nonlinear partial differential equations (NLPDEs) and their solutions has become a subject of much interest in the past few decades. Modelling natural phenomena such as the behaviour of water waves is often an interdisciplinary research area connecting such areas as mathematics, physics and engineering. Along with the progress in modelling nonlinear phenomena came a myriad of methods designed to derive the exact solutions of these models. Some of these methods are Lie group analysis [1–5], homogeneous balance technique [6], the ansatz method [7], the Hirota bilinear method [8], the (G'/G) -expansion technique [9,10], the Kudryashov method [11,12], the simplest equation method [13], the Jacobi elliptic function expansion technique [14], and the F -expansion method [15], bifurcation method [16] just to mention a few.

The (3+1)-dimensional Kadomtsev–Petviashvili (KP) equations [17]

$$(u_t + 6uu_x + u_{xxx})_x \pm 3u_{yy} \pm 3u_{zz} = 0 \quad (1)$$

have their origins in the classic 1970 work of two Soviet physicists Kadomtsev and Petviashvili [18]. It models waves in scenarios where the ratio between the depth of water and the wavelength is very small coupled with insubstantial nonlinear restoring forces. It is also a generalisation of the

Korteweg–de Vries (KdV) equation, named after the Dutch mathematicians Korteweg and de Vries [19]. The highest order term u_{xxx} caters for weak dispersion, while the term uu_x describes the nonlinearity aspect of a wave and the last two terms of (1) describe diffractive divergence [20]. The sign “ $\pm$ ” in the last two terms corresponds to either positive or negative dispersion. Since then, several researchers have studied equations of the form (1), see for instance [21–26]. Investigations into (1) have ranged from establishing the stability of their solitons [23], determining their integrability properties [24], Painlevé analysis [25] and deriving their exact multiple wave solutions [26]. The (3+1)-dimensional generalised KP equation

$$u_{tx} + u_{ty} + 3u_x u_{xy} + 3u_{xx} u_y + u_{xxx} y - u_{zz} = 0 \quad (2)$$

to the best of our knowledge was introduced in [27], where the Plücker relation for determinants was used to obtain one Wronskian solution. Furthermore, in the same work, the Jacobi identity for determinants was used to establish a Grammian solution for (2). Multiple solitons and multiple singular solitons for Equation (2) were derived in [28] using a simplified form of Hirota’s method. In their work, the researchers in [28] contrast their results with those obtained in [27] in terms of the spatial variable z . Later in [29], an extra term u_{tz} , was added to (2), thus introducing a new form of the (3+1)-dimensional generalised KP equation, which we denote here by (3+1)-D gKPe, and it reads

$$u_{tx} + u_{ty} + u_{tz} + 3u_x u_{xy} + 3u_{xx} u_y + u_{xxx} y - u_{zz} = 0. \quad (3)$$

In their work [29], the authors showed that addition of the new term affects the dispersion relations significantly. Furthermore, they applied Hirota’s direct method to determine the multiple soliton solutions of (3). In [30], the Equation (3) was studied and exact solutions were obtained through the use of Hirota’s bilinear method and an extended homoclinic test approach.

In this work, we seek to derive the exact solutions of the (3+1)-D gKPe (3) by making use of its Lie point symmetries and direct integration. We will not employ adhoc methods as often is the case when treating NLPDEs of this nature. Furthermore, we will for the first time derive the conserved quantities of Equation (3) by invoking two approaches.

Lie group theory was discovered by a Norwegian mathematician, Marius Sophus Lie (1842–1899), around the middle of nineteenth century. Lie perceived that the seemingly different methods for finding exact solutions of differential equations were, in reality, all special cases of a broad integration approach; the theory of transformation groups. This theory is an analog of Galois theory and has an enormous impact on mathematics and mathematical physics today. It is contemplated to be one of the most significant techniques to determine the exact solutions of differential equations [1–5].

Conservation laws are essential to our comprehension of the physical world and are considered to be basic laws of nature, with wide application in physics, and in other fields for instance chemistry, engineering, and so on. In classical physics, three types of such laws are conservation of energy, linear momentum and angular momentum. Conservation laws are used in establishing the integrability of differential equations (DEs) and are also used in determining whether the solution of a DE exists and is unique. Moreover, one can also use conservation laws in checking the authenticity of numerical solution methods [31–44].

We want to emphasize here that a study of higher dimensional NLPDEs is critical to our understanding of the world around us since they are more realistic models of natural and man-made phenomena [45].

2. Exact Solutions of the (3+1)-D gKPe

In this section, we determine the exact solutions of the (3+1)-D gKPe (3) by utilizing its Lie point symmetries and direct integration. Roughly speaking, a Lie point symmetry of a PDE is a local group of transformations acting on the independent and dependent variables of the PDE that maps every

solution of the PDE to another solution of the same PDE. That is to say, it maps the solution set of the PDE to itself.

It can be seen, using for example, MathLie package [46], that Equation (3) has a finite-dimensional Lie subalgebra L_6 , which is generated by the vector fields

$$\begin{aligned} X_1 &= \frac{\partial}{\partial t}, \quad X_2 = \frac{\partial}{\partial x}, \quad X_3 = \frac{\partial}{\partial y}, \quad X_4 = \frac{\partial}{\partial z}, \\ X_5 &= -t \frac{\partial}{\partial t} + z \frac{\partial}{\partial x} + z \frac{\partial}{\partial y} + (2t + z) \frac{\partial}{\partial z}, \\ X_6 &= 15t \frac{\partial}{\partial t} + 3(4t + x) \frac{\partial}{\partial x} + 3(3y - 2z) \frac{\partial}{\partial y} - 3(4t - z) \frac{\partial}{\partial z} \\ &\quad + (4x + 4y - 3u) \frac{\partial}{\partial u}, \end{aligned} \quad (4)$$

and the two infinite-dimensional subalgebras generated by

$$X_{F_1} = F_1(t) \frac{\partial}{\partial u}, \quad X_{F_2} = F_2(t + z) \frac{\partial}{\partial u}, \quad (5)$$

where F_1 and F_2 are arbitrary functions of their arguments.

2.1. Invariant Solutions under the Symmetries $X_1, \dots, X_4$

Firstly, we invoke the translation symmetries and reduce the (3+1)-D gKPe (3) to a fourth order ordinary differential equation (ODE). Thus, using the four translation symmetries X_1, X_2, X_3 and X_4 of Equation (3), we can obtain the following invariant solution:

$$u(t, x, y, z) = U(p), \quad p = \gamma x + \alpha y + \beta z - vt, \quad (6)$$

where γ, α, β and v are constants. This reduces the Equation (3) to

$$\alpha \gamma^3 U'''' + 6\alpha \gamma^2 U' U'' - (\alpha v + \beta^2 + \beta v + \gamma v) U'' = 0. \quad (7)$$

Integrating (7) once with respect to p yields

$$\alpha \gamma^3 U''' + 3\alpha \gamma^2 U'^2 - (\alpha v + \beta^2 + \beta v + \gamma v) U' + K_0 = 0 \quad (8)$$

with K_0 a constant. Now letting

$$U'(p) = \gamma \phi(p), \quad \omega = -\frac{\alpha v + \beta^2 + \beta v + \gamma v}{\alpha \gamma^3} \quad \text{and} \quad K_1 = \frac{K_0}{\alpha \gamma^4},$$

Equation (8) becomes

$$\phi'' + 3\phi^2 - \omega \phi + K_1 = 0. \quad (9)$$

This ODE can be integrated easily. Multiplying (9) by ϕ' and integrating once with respect to p gives

$$\phi'^2 = -(2\phi^3 - \omega \phi^2 + 2K_1 \phi + 2K_2) \quad (10)$$

with K_2 an integration constant. If the algebraic equation $\phi^3 - \frac{1}{2}\omega \phi^2 + K_1 \phi + K_2 = 0$ has the roots $\lambda_1 \geq \lambda_2 \geq \lambda_3$, then we have

$$\phi'^2 = -2(\phi - \lambda_1)(\phi - \lambda_2)(\phi - \lambda_3). \quad (11)$$

Equation (11) has the well-known solution expressed in terms of the Jacobi elliptic function, that is,

$$\phi(p) = \lambda_2 + (\lambda_1 - \lambda_2) \operatorname{cn}^2 \left(\sqrt{\frac{\lambda_1 - \lambda_3}{2}} p \middle| M^2 \right), \quad M^2 = \frac{\lambda_1 - \lambda_2}{\lambda_1 - \lambda_3}. \quad (12)$$

Here $\operatorname{cn}(p|M^2)$ is the Jacobi elliptic cosine function. Furthermore, pertinent to note is that when $M^2 \rightarrow 1$, $\operatorname{cn}(p|M^2) \rightarrow \operatorname{sech}(p)$ and when $M^2 \rightarrow 0$, $\operatorname{cn}(p|M^2) \rightarrow \cos(p)$ [47]. A comparison of Equations (9) and (10) reveals that

$$\omega = 2(\lambda_1 + \lambda_2 + \lambda_3), \quad K_1 = \lambda_1\lambda_2 + \lambda_2\lambda_3 + \lambda_1\lambda_3, \quad K_2 = -\lambda_1\lambda_2\lambda_3.$$

Reverting to original variables, the solution of the (3+1)-D gKPe (3) is thus

$$u(t, x, y, z) = \frac{\{\lambda_2 - (1 - M^2)\lambda_1\}p}{M^2} + \frac{(\lambda_1 - \lambda_2) \operatorname{dn}(Ap|M^2) E(\operatorname{am}(Ap|M^2)|M^2)}{AM^2 \sqrt{\operatorname{dn}(Ap|M^2)^2}}, \quad (13)$$

where $E(p|M^2)$ is the elliptic integral of the second kind, $\operatorname{am}(p|M^2)$ is the amplitude function, $\operatorname{dn}(p|M^2)$ is the delta amplitude function, $\operatorname{cn}(p|M^2)$ is the Jacobi cosine function and $A = \sqrt{(\lambda_1 - \lambda_3)/2}$ [47,48]. Figure 1 depicts the profile of solution (13) for suitable parameter values.




Figure 1. Profiles of solution (13).

Figure 1 depicts the coexistence of bright and dark solitons. It is well-known that bright soliton profile are characterized by hyperbolic secant. The bright soliton solution usually takes a bell-shaped figure and propagate undistorted without any change in shape for arbitrarily long distances. However, dark soliton solutions, configured also as topological optical solitons, are given by hyperbolic tangent.

Important to note is that Equation (10) is reminiscent of the ODE obtained in the quintessential work of Korteweg and de Vries in [49]. This ODE is associated with long waves propagating along a rectangular canal. The ODE (10) describes stationary waves and by imposing certain constraints such as having the fluid undisturbed at infinity, Korteweg and de Vries obtained negative and positive solitary waves as well as cnoidal wave solutions [49,50].

2.2. Invariant Solution under the Symmetry X_5

We now turn our attention to the point symmetry X_5 . The usual computations yield the group invariant solution $u(t, x, y, z) = U(\xi)$, where $\xi = t(t + z)$. This, substituted into (3) gives the ODE

$$\xi U'' + U' = 0, \quad (14)$$

whose solution is $U(\xi) = C_1 + C_2 \ln(\xi)$, and hence the group invariant solution of the (3+1)-D gKPe (3) under the symmetry X_5 is

$$u(t, x, y, z) = C_1 + C_2 \ln \{t(t+z)\} \quad (15)$$

with C_1, C_2 integration constants. The corresponding graphical representation of solution (15) is given in Figure 2.




Figure 2. Profiles of solution (15).

The profiles portrayed in Figure 2 are typical of a grey singular soliton solution, more so the distinctive asymptotic form observable for $\lim_{t \rightarrow 0} u(t, x, y, z) = -\infty$.

2.3. Invariant Solution under the Symmetry X_6

Finally, we compute the group invariant solution under X_6 . By following the usual procedure we obtain the group invariant solution

$$u(t, x, y, z) = t^{-1/5} U(\xi) - \frac{2}{9}t + \frac{2}{3}x + \frac{1}{3}y + \frac{1}{3}z, \quad (16)$$

where $\xi = t^{-1/5}(x - t)$. Substituting (16) into (3) and simplifying ultimately yields the second-order ODE

$$\xi U'' + 2U' = 0. \quad (17)$$

Equation (17) can be solved much like Equation (14), which in turn gives the invariant solution of the (3+1)-D gKPe (3) under the group generated by X_6 :

$$u(t, x, y, z) = C_1 + C_2 t^{1/5}(x - t)^{-1} \quad (18)$$

with C_1, C_2 integration constants. In Figure 3, solution (18) is depicted.




Figure 3. Profiles of solution (18).

The density plot coupled with the two dimensional profile in Figure 3 depicts a singular soliton-like wave with a singularity in the spatial domain at $0 \leq x \leq 20$.

3. Conserved Quantities of the (3+1)-D gKPe

In this section, we compute the conserved quantities of the (3+1)-D gKPe (3). In our work we use the multiplier approach and the conservation theorem due to Ibragimov to derive the conserved quantities of (3). However, first we provide salient features on both these approaches.

3.1. Multiplier Approach

The multiplier method is one of the most robust and preferred methods for deriving conserved quantities of DEs [4,36–41]. This method attempts to mitigate the shortcomings of Noether's theorem [31], which requires amongst other things, the existence of a variational principle or a Lagrangian before the theorem can be applied. We begin by providing a concise basis of the method.

3.1.1. Preliminaries

Let G be a system of m PDEs of order k , having n independent variables $x = (x^1, x^2, \dots, x^n)$ and m field variables $\Phi = ((\Phi^1, \Phi^2, \dots, \Phi^m))$:

$$G = \left(G^1(x, \Phi, \Phi_{(1)}, \Phi_{(2)}, \dots, \Phi_{(k)}), \dots, G^m(x, \Phi, \Phi_{(1)}, \Phi_{(2)}, \dots, \Phi_{(k)}) \right), \quad (19)$$

where $\Phi_{(1)}, \Phi_{(2)}, \dots, \Phi_{(k)}$ denote the derivatives of Φ with respect to the variables x up to the k -th derivative. A local conserved quantity $T^i(x, \Phi, \Phi_{(1)}, \Phi_{(2)}, \dots, \Phi_{(l)})$ of system (19) is a continuity equation

$$D_i T^i|_{\varepsilon} = 0, \quad (20)$$

valid for the solution space ε of system (19).

In general, local nontrivial conserved quantities emanate from the divergence identity

$$D_{x^1} T^1 + D_{x^2} T^2 + \dots + D_{x^n} T^n = \Lambda_{\alpha}(x, \Phi, \Phi_{(1)}, \Phi_{(2)}, \dots, \Phi_{(r)}) G(\Phi). \quad (21)$$

Here, $\Lambda_{\alpha}(x, \Phi, \Phi_{(1)}, \Phi_{(2)}, \dots, \Phi_{(r)})$ is a series of conservation law multipliers which are dependent on x, Φ and the derivatives of Φ , up to some arbitrary order $r < k$. The relationship (21) brings to light the pre-eminent interrelation between conserved quantities T^i and multipliers Λ_{α} . A determining condition to derive a set of multipliers $\Lambda_{\alpha}(x, \Phi, \Phi_{(1)}, \Phi_{(2)}, \dots, \Phi_{(r)})$ for system (19) is that

$$\frac{\delta}{\delta \Phi^{\alpha}} (\Lambda_{\alpha} G) = 0, \quad \alpha = 1, \dots, m, \quad (22)$$

where $\delta/\delta \Phi^{\alpha}$ is the Euler–Lagrange operator given by

$$\frac{\delta}{\delta \Phi_i^{\alpha}} = \frac{\partial}{\partial \Phi_i^{\alpha}} + \sum_{s \geq 1} (-1)^s D_{j_1} \dots D_{j_s} \frac{\partial}{\partial \Phi_{ij_1 j_2 \dots j_s}^{\alpha}}, \quad i = 1, \dots, n, \quad \alpha = 1, \dots, m. \quad (23)$$

The condition (22) is requisite and adequate for Λ to be a multiplier. A more rigorous and detailed treatment of the theoretical justification of the multiplier approach including proofs of the formulas utilised in this section can be found in [33].

3.1.2. Application of the Method

In order to determine conserved quantities of Equation (3), we begin by computing the first order multipliers

$$\Lambda = \Lambda(t, x, u, u_x, u_y, u_z),$$

by utilising condition (22), that is,

$$\frac{\delta}{\delta u} (\Lambda E) = 0, \quad (24)$$

where

$$E \equiv u_{tx} + u_{ty} + u_{tz} + 3u_x u_{xy} + 3u_{xx} u_y + u_{xxx} y - u_{zz} = 0.$$

The Euler operator (23) is given by

$$\begin{aligned} \frac{\delta}{\delta u} = & \frac{\partial}{\partial u} - D_x \frac{\partial}{\partial u_x} - D_y \frac{\partial}{\partial u_y} + D_t D_x \frac{\partial}{\partial u_{tx}} + D_t D_y \frac{\partial}{\partial u_{ty}} + D_t D_z \frac{\partial}{\partial u_{tz}} \\ & + D_x^2 \frac{\partial}{\partial u_{xx}} + D_x D_y \frac{\partial}{\partial u_{xy}} + D_z^2 \frac{\partial}{\partial u_{zz}} + D_x^3 D_y \frac{\partial}{\partial u_{xxx} y} \end{aligned}$$

and D_t , D_x , D_y and D_z are total derivative operators [4]. Expanding (24) and splitting on derivatives of u , we obtain the following system of nine multiplier determining equations:

$$\begin{aligned} \Lambda_{ty} + \Lambda_{tz} - \Lambda_{zz} &= 0, \quad \Lambda_{tu_x} = 0, \quad \Lambda_{yu_x} = 0, \quad \Lambda_{zu_x} = 0, \quad \Lambda_{u_x u_x} = 0, \\ \Lambda_x &= 0, \quad \Lambda_u = 0, \quad \Lambda_{u_y} = 0, \quad \Lambda_{u_z} = 0. \end{aligned} \quad (25)$$

The solution algorithm of system (25) is similar to that of obtaining Lie point symmetries. However, with the aid of Gem, a Maple based package [44], this can be expedited. We thus have the multiplier

$$\Lambda = C u_x \quad (26)$$

with C an integration constant. The conserved quantities of (3) are obtained via the divergence identity

$$D_t T^t + D_x T^x + D_y T^y + D_z T^z = E \Lambda,$$

where T^t is a conserved density and T^x, T^y, T^z are spatial fluxes [36]. Thus, after some calculations, we obtain the following conservation law:

$$\begin{aligned} T^t &= \frac{1}{2} u_x^2 + \frac{1}{2} u_y u_x + \frac{1}{2} u_z u_x, \\ T^x &= u u_x u_{xy} + 2 u_x^2 u_y + \frac{1}{2} u_y u_{xxx} - \frac{1}{2} u_{xy} u_{xx} + \frac{1}{2} u_{xxy} u_x + \frac{1}{2} u u_{ty} + \frac{1}{2} u u_{tz} \\ &\quad + \frac{1}{2} u u_{xxy} - \frac{1}{2} u u_{zz}, \\ T^y &= -u u_x u_{xx} - \frac{1}{2} u u_{tx} - \frac{1}{2} u u_{xxx}, \\ T^z &= -\frac{1}{2} u u_{tx} + \frac{1}{2} u u_{xz} - \frac{1}{2} u z u_x. \end{aligned}$$

Since the derivatives of u in the multiplier (26) are of a lower order than both leading derivatives of Equation (3), that is, u_{tx} and u_{xxx} , the conserved quantities derived here are low-order conservation laws [36].

3.2. Ibragimov's Approach

We now determine the conservation laws of (3) by applying a theorem due to Ibragimov [42,43]. As in the case of multiplier method, this theorem does not demand availability of a Lagrangian and is established on a concept of an adjoint equation. Thus, it applies to an arbitrary differential equation, irrespective of whether or not it comes from a variational principle.

3.2.1. Preliminaries

The gist of Ibragimov's method is that every infinitesimal generator is associated with a conserved quantity, notwithstanding the absence of traditional Lagrangians which are envisaged in Noether's theorem [31]. Below we outline the method in detail.

Consider a system of NLPDEs (19) and its adjoint equations given by

$$G^*(x, \Phi, \Psi, \Phi_{(1)}, \Psi_{(1)}, \Phi_{(2)}, \Psi_{(2)}, \dots, \Phi_{(k)}, \Psi_{(k)}) = \frac{\delta}{\delta \Phi^\alpha} (\Psi G_\beta) \quad (27)$$

where $\delta/\delta \Phi^\alpha$ is the Euler–Lagrange operator (23) and m novel field variables $\Psi = (\Psi^1, \dots, \Psi^m)$.

Theorem 1. Consider a system of m Equations (19). The adjoint system given by (27), inherits the symmetries of the system (19). Namely, if the system (19) admits a point transformation group with a generator $X = \xi^i \partial/\partial t + \eta^\alpha \partial/\partial \Phi^\alpha$, then the adjoint system (27) admits the operator X extended to the variables Ψ^α by the formula

$$Y = \xi^i \frac{\partial}{\partial x^i} + \eta^\alpha \frac{\partial}{\partial \Phi^\alpha} + \eta_*^\alpha \frac{\partial}{\partial \Psi^\alpha} \quad (28)$$

with appropriately chosen $\eta_*^\alpha = \eta_*^\alpha(x, \Phi, \Psi)$.

The functions ξ^i and η^α are infinitesimal generator coefficients dependent on x and Φ . In [43], the coefficients η_*^α in (28) are given by

$$\eta_*^\alpha = -[\lambda_\beta^\alpha \Psi^\beta + \Psi^\alpha D_i(\xi^i)], \quad (29)$$

where λ_β^α can be computed by utilising the equation

$$X(G_\alpha) = \lambda_\alpha^\beta G_\beta. \quad (30)$$

We can obtain a conserved vector, for instance, for a third-order Lagrangian by applying the formula

$$\begin{aligned} C^i = & \xi^i \mathcal{L} + W^\alpha \left[\frac{\partial \mathcal{L}}{\partial \Phi_i^\alpha} - D_j \frac{\partial \mathcal{L}}{\partial \Phi_{ij}^\alpha} + D_j D_k \left(\frac{\partial \mathcal{L}}{\partial \Phi_{ijk}^\alpha} \right) + \dots \right] \\ & + D_j (W^\alpha) \left[\frac{\partial \mathcal{L}}{\partial \Phi_{ij}^\alpha} - D_k \frac{\partial \mathcal{L}}{\partial \Phi_{ijk}^\alpha} + \dots \right] + D_j D_k (W^\alpha) \frac{\partial \mathcal{L}}{\partial \Phi_{ijk}^\alpha} + \dots, \end{aligned} \quad (31)$$

where $\mathcal{L}$ is the Lagrangian of the system G and G^* that is defined as

$$\mathcal{L} = \Psi^\alpha G_\alpha \quad (32)$$

and W^α is the Lie characteristic function given by

$$W^\alpha = \eta^\alpha - \xi^j \Phi_j^\alpha, \quad \alpha = 1, \dots, m. \quad (33)$$

The reader is referred to [42,43] for a more comprehensive discussion of this method.

3.2.2. Application of the Method

To begin, we define the adjoint equation of (3) as [43]

$$\begin{aligned} F^* &= \frac{\delta}{\delta u} \{v(u_{tx} + u_{ty} + u_{tz} + 3u_x u_{xy} + 3u_{xx} u_y + u_{xxx} u_y - u_{zz})\} \\ &= \left(\frac{\partial}{\partial u} - D_x \frac{\partial}{\partial u_x} - D_y \frac{\partial}{\partial u_y} + D_t D_x \frac{\partial}{\partial u_{tx}} + D_t D_y \frac{\partial}{\partial u_{ty}} + D_t D_z \frac{\partial}{\partial u_{tz}} \right. \\ &\quad \left. + D_x^2 \frac{\partial}{\partial u_{xx}} + D_x D_y \frac{\partial}{\partial u_{xy}} + D_z^2 \frac{\partial}{\partial u_{zz}} + D_x^3 D_y \frac{\partial}{\partial u_{xxx}} \right) \{v(u_{tx} + u_{ty} \\ &\quad + u_{tz} + 3u_x u_{xy} + 3u_{xx} u_y + u_{xxx} u_y - u_{zz})\} \\ &= v_{tx} + v_{ty} + v_{tz} + 6v_x u_{xy} + 3u_x v_{xy} + 3u_y v_{xx} + v_{xxx} u_y - v_{zz} = 0. \end{aligned} \quad (34)$$

We have introduced a new variable $v = v(t, x, y, z)$. According to [43], the Equation (3) considered together with its adjoint (34) has a Lagrangian $\mathcal{L}$, given by

$$\mathcal{L} = vF = v(u_{tx} + u_{ty} + u_{tz} + 3u_x u_{xy} + 3u_{xx} u_y + u_{xxx} u_y - u_{zz}). \quad (35)$$

Notice how $\delta \mathcal{L} / \delta u = F^*$ and $\delta \mathcal{L} / \delta v = F$. The Lagrangian (35) is equivalent to the second order Lagrangian

$$\mathcal{L} = v(u_{tx} + u_{ty} + u_{tz} + 3u_x u_{xy} + 3u_{xx} u_y - u_{zz}) + v_{xx} u_{xy}. \quad (36)$$

Now, Equation (34) admits all the symmetries (4) and (5) of (3) extended to the new variable $v(t, x, y, z)$. That is, the generators (4) and (5) become

$$Y = \xi^1 \frac{\partial}{\partial t} + \xi^2 \frac{\partial}{\partial x} + \xi^3 \frac{\partial}{\partial y} + \xi^4 \frac{\partial}{\partial z} + \eta \frac{\partial}{\partial u} + \eta^* \frac{\partial}{\partial v} \quad (37)$$

with

$$\eta^* = \eta^*(t, x, y, z, u, v) = - \left\{ \lambda + D_t(\xi^1) + D_x(\xi^2) + D_y(\xi^3) + D_z(\xi^4) \right\} v. \quad (38)$$

The parameter λ is determined by using

$$X'(F) = \lambda F, \quad (39)$$

where X' is the generator (4) prolonged to all the derivatives in (3), that is,

$$\begin{aligned} X' &= X + \zeta_x \frac{\partial}{\partial u_x} + \zeta_y \frac{\partial}{\partial u_y} + \zeta_{tx} \frac{\partial}{\partial u_{tx}} + \zeta_{ty} \frac{\partial}{\partial u_{ty}} + \zeta_{tz} \frac{\partial}{\partial u_{tz}} + \zeta_{xx} \frac{\partial}{\partial u_{xx}} \\ &\quad + \zeta_{xy} \frac{\partial}{\partial u_{xy}} + \zeta_{zz} \frac{\partial}{\partial u_{zz}} + \zeta_{xxx} \frac{\partial}{\partial u_{xxx}}. \end{aligned} \quad (40)$$

Here $X = \xi^1 \partial / \partial t + \xi^2 \partial / \partial x + \xi^3 \partial / \partial y + \xi^4 \partial / \partial z + \eta \partial / \partial u$ and $\xi^1, \dots, \xi^4$, and η are functions of (t, x, y, z, u) . Furthermore, $\zeta_x, \zeta_y, \zeta_{tx}, \zeta_{ty}, \zeta_{tz}, \zeta_{xx}, \zeta_{xy}, \zeta_{zz}$ and ζ_{xxx} are coefficient functions, each given by the following formulae:

$$\begin{aligned}
\zeta_x &= D_x(\eta) - u_t D_x(\zeta^1) - u_x D_x(\zeta^2) - u_y D_x(\zeta^3) - u_z D_x(\zeta^4), \\
\zeta_y &= D_y(\eta) - u_t D_y(\zeta^1) - u_x D_y(\zeta^2) - u_y D_y(\zeta^3) - u_z D_y(\zeta^4), \\
\zeta_{tx} &= D_x(\zeta_t) - u_{tt} D_x(\zeta^1) - u_{tx} D_x(\zeta^2) - u_{ty} D_x(\zeta^3) - u_{tz} D_x(\zeta^4), \\
\zeta_{ty} &= D_y(\zeta_t) - u_{tt} D_y(\zeta^1) - u_{tx} D_y(\zeta^2) - u_{ty} D_y(\zeta^3) - u_{tz} D_y(\zeta^4), \\
\zeta_{tz} &= D_z(\zeta_t) - u_{tt} D_z(\zeta^1) - u_{tx} D_z(\zeta^2) - u_{ty} D_z(\zeta^3) - u_{tz} D_z(\zeta^4), \\
\zeta_{xx} &= D_x(\zeta_x) - u_{xt} D_x(\zeta^1) - u_{xx} D_x(\zeta^2) - u_{xy} D_x(\zeta^3) - u_{xz} D_x(\zeta^4), \\
\zeta_{xy} &= D_y(\zeta_x) - u_{xt} D_y(\zeta^1) - u_{xx} D_y(\zeta^2) - u_{xy} D_y(\zeta^3) - u_{xz} D_y(\zeta^4), \\
\zeta_{zz} &= D_z(\zeta_x) - u_{zt} D_z(\zeta^1) - u_{zx} D_z(\zeta^2) - u_{zy} D_z(\zeta^3) - u_{zz} D_z(\zeta^4), \\
\zeta_{xxx} &= D_x(\zeta_{xx}) - u_{xxt} D_x(\zeta^1) - u_{xxx} D_x(\zeta^2) - u_{xxy} D_x(\zeta^3) - u_{xxz} D_x(\zeta^4), \\
\zeta_{xxxy} &= D_y(\zeta_{xxx}) - u_{xxtt} D_y(\zeta^1) - u_{xxx} D_y(\zeta^2) - u_{xxxy} D_y(\zeta^3) - u_{xxxz} D_y(\zeta^4).
\end{aligned} \tag{41}$$

See the full expansions of (41) in Appendix A. Now using Equations (3), (39) and (40) we compute the values of the parameter λ corresponding to each of the vector fields in (4) and (5).

Cases $X_1, \dots, X_4$

For the time translation symmetry X_1 , we have $\zeta^1 = 1$ and $\zeta^2 = \zeta^3 = \zeta^4 = \eta = 0$. Thus, it is easy to see that

$$\zeta_x = \zeta_y = \zeta_{tx} = \zeta_{ty} = \zeta_{tz} = \zeta_{xx} = \zeta_{xy} = \zeta_{zz} = \zeta_{xxxy} = 0.$$

Consequently, we have $X_1'(F) = 0F$, that is $\lambda = 0$. From (38), we obtain $\eta^* = 0$ and the new generator (4) retains the form of X_1 , i.e., $Y_1 = \partial/\partial t$. Since the generator coefficients are all constants, the translation symmetries will retain their form and this leads us to conclude that $Y_2 = \partial/\partial x$, $Y_3 = \partial/\partial y$ and $Y_4 = \partial/\partial z$.

Case X_5

Here, we have the generator coefficients $\zeta^1 = -t$, $\zeta^2 = x$, $\zeta^3 = z$, $\zeta^4 = 2t + z$ and $\eta = 0$ from which we can establish the following:

$$\begin{aligned}
\zeta_x &= 0, \quad \zeta_y = 0, \quad \zeta_{tx} = u_{tx} - 2u_{xz}, \quad \zeta_{ty} = u_{ty} - 2u_{yz}, \quad \zeta_{tz} = -2u_{zz} - u_{tx} - u_{ty}, \\
\zeta_{xx} &= 0, \quad \zeta_{xy} = 0, \quad \zeta_{zz} = -2u_{xz} - 2u_{yz} - 2u_{zz}, \quad \zeta_{xxxy} = 0.
\end{aligned}$$

Consequently, we can verify that $X_5'(F) = 0F$, that is, $\lambda = 0$. From (38), we can further establish that $\eta^* = 0$. The generator for the adjoint Equation (34) is thus $Y_5 = -t\partial/\partial t + z\partial/\partial x + z\partial/\partial y + (2t + z)\partial/\partial z$.

Case X_6

The vector field X_6 has the coefficients $\zeta^1 = 15t$, $\zeta^2 = 12t + 3x$, $\zeta^3 = 9y - 6z$, $\zeta^4 = -12t + 3z$ and $\eta = 4x + 4y - 3u$. The reckoning of the coefficient functions (41) yields

$$\begin{aligned}
\zeta_x &= 4 - 6u_x, \quad \zeta_y = 4 - 12u_y, \quad \zeta_{tx} = -21u_{tx} - 12u_{xx} + 12u_{xz}, \\
\zeta_{ty} &= -27u_{ty} - 12u_{xy} + 12u_{yz}, \quad \zeta_{tz} = -21u_{tz} - 12u_{xz} + 12u_{zz} + 6u_{ty}, \\
\zeta_{xx} &= -9u_{xx}, \quad \zeta_{xy} = -15u_{xy}, \quad \zeta_{zz} = -9u_{zz} + 12u_{yz}, \quad \zeta_{xxxy} = -21u_{xxxy}.
\end{aligned}$$

Now, from Equations (39)–(41), we have

$$\begin{aligned}
X_6'(F) &= -21u_{tx} - 21u_{ty} - 21u_{tz} - 63u_x u_{xy} - 63u_{xx} u_y - 21u_{xxxy} + 21u_{zz} \\
&= -21(u_{tx} + u_{ty} + u_{tz} + 3u_x u_{xy} + 3u_{xx} u_y + u_{xxxy} - u_{zz}) \\
&= -21F.
\end{aligned}$$

We can thus see that $\lambda = -21$. Consequently,

$$\begin{aligned}\eta^* &= -(-21 + D_t(15t) + D_x(12t + 3x) + D_y(9y - 6z) + D_z(-12t + 3z))v \\ &= -7v.\end{aligned}$$

We now have the vector field:

$Y_6 = 15t\partial t + (12t + 3x)\partial x + (9y - 6z)\partial y + (3z - 12t)\partial z + (4x + 4y - 3u)\partial u - 9v\partial v$, corresponding to (34).

Cases X_7, X_8

Finally, considering the infinite-dimensional vector fields X_7 and X_8 we have, for (34), $Y_7 = F(t)\partial u$ and $Y_8 = F(t + z)\partial u$, respectively.

To compute the conservation laws of (3), we utilise the formula [43]

$$C^i = \zeta^i \mathcal{L} + W^\alpha \left(\frac{\partial \mathcal{L}}{\partial u_i^\alpha} - D_k \frac{\partial \mathcal{L}}{\partial u_{ik}^\alpha} \right) + D_k (W^\alpha) \frac{\partial \mathcal{L}}{\partial u_{ik}^\alpha}, \quad (42)$$

where W^α is the Lie characteristic function given by $W^\alpha = \eta^\alpha - \zeta^j u_j^\alpha$. The reckoning of (42) gives the following conserved quantities:

$$\begin{aligned}T_1 &= -vu_{zz} + 3vu_xu_{xy} + 3vu_yu_{xx} + vu_{xxy} + \frac{1}{2}v_zu_t + \frac{1}{2}v_yu_t + \frac{1}{2}v_xu_t + \frac{1}{2}vu_{tz} \\ &\quad + \frac{1}{2}vu_{ty} + \frac{1}{2}vu_{tx}, \\ X_1 &= \frac{3}{2}v_yu_xu_t + 3u_yv_xu_t + \frac{3}{2}vu_{xy}u_t + \frac{3}{4}v_{xxy}u_t + \frac{1}{2}v_tu_t - \frac{3}{2}vu_xu_{ty} - \frac{1}{4}v_{xx}u_{ty} \\ &\quad - 3vu_yu_{tx} - \frac{1}{2}v_{xy}u_{tx} + \frac{1}{2}v_xu_{txy} + \frac{1}{4}v_yu_{txx} - \frac{3}{4}vu_{txxy} - \frac{1}{2}vu_{tt}, \\ Y_1 &= \frac{3}{2}u_xv_xu_t - \frac{3}{2}vu_{xx}u_t + \frac{1}{4}v_{xxx}u_t + \frac{1}{2}v_tu_t - \frac{3}{2}vu_xu_{tx} - \frac{1}{4}v_{xx}u_{tx} + \frac{1}{4}v_xu_{txx} \\ &\quad - \frac{1}{4}vu_{txxx} - \frac{1}{2}vu_{tt}, \\ Z_1 &= -v_zu_t + \frac{1}{2}v_tu_t + vu_{tz} - \frac{1}{2}vu_{tt}; \\ T_2 &= -\frac{1}{2}u_{xz}v - \frac{1}{2}u_{xy}v - \frac{1}{2}u_{xx}v + \frac{1}{2}u_xv_y + \frac{1}{2}u_xv_z + \frac{1}{2}u_xv_x, \\ X_2 &= 3u_xu_{xy}v - u_{zz}v + \frac{1}{4}u_{xxy}v + u_{tz}v + u_{ty}v + \frac{1}{2}u_{tx}v + \frac{1}{2}v_tu_x + \frac{3}{2}u_x^2v_y + 3u_xu_yv_x \\ &\quad + \frac{3}{4}u_xv_{xxy} - \frac{1}{2}u_{xx}v_{xy} - \frac{1}{4}v_{xx}u_{xy} + \frac{1}{2}v_xu_{xxy} + \frac{1}{4}u_{xxx}v_y, \\ Y_2 &= -3u_{xx}u_xv - \frac{1}{4}u_{xxx}v - \frac{1}{2}u_{tx}v + \frac{1}{2}v_tu_x + \frac{3}{2}u_x^2v_x + \frac{1}{4}u_xv_{xxx} - \frac{1}{4}u_{xx}v_{xx} \\ &\quad + \frac{1}{4}u_{xxx}v_x, \\ Z_2 &= u_{xz}v - \frac{1}{2}u_{tx}v + \frac{1}{2}v_tu_x - u_xv_z;\end{aligned}$$

$$\begin{aligned}
T_3 &= -\frac{1}{2}u_{yz}v - \frac{1}{2}u_{yy}v - \frac{1}{2}u_{xy}v + \frac{1}{2}u_yv_x + \frac{1}{2}u_yv_z + \frac{1}{2}u_yv_y, \\
X_3 &= -\frac{3}{2}u_yu_{xy}v - \frac{3}{2}u_xu_{yy}v - \frac{3}{4}u_{xxy}v - \frac{1}{2}u_{ty}v + \frac{1}{2}v_tu_y + 3u_y^2v_x + \frac{3}{2}u_xu_yv_y + \frac{3}{4}u_yv_{xxy} \\
&\quad - \frac{1}{2}u_{xy}v_{xy} + \frac{1}{2}v_xu_{xyy} - \frac{1}{4}u_{yy}v_{xx} + \frac{1}{4}v_yu_{xxy}, \\
Y_3 &= -u_{zz}v + \frac{3}{2}u_xu_{xy}v + \frac{3}{2}u_{xx}u_yv + \frac{3}{4}u_{xxy}v + u_{tz}v + \frac{1}{2}u_{ty}v + u_{tx}v + \frac{1}{2}v_tu_y + \frac{3}{2}u_xu_yv_x \\
&\quad - \frac{1}{4}v_{xx}u_{xy} + \frac{1}{4}v_xu_{xxy} + \frac{1}{4}u_yv_{xxx}, \\
Z_3 &= u_{yz}v - \frac{1}{2}u_{ty}v + \frac{1}{2}v_tu_y - u_yv_z;
\end{aligned}$$

$$\begin{aligned}
T_4 &= \frac{1}{2}u_zv_z - \frac{1}{2}vu_{zz} + \frac{1}{2}u_zv_y - \frac{1}{2}vu_{yz} + \frac{1}{2}u_zv_x - \frac{1}{2}vu_{xz}, \\
X_4 &= \frac{3}{2}u_zv_yu_x - \frac{3}{2}vu_{yz}u_x + 3u_zu_yv_x - 3vu_yu_{xz} + \frac{3}{2}vu_zu_{xy} - \frac{1}{2}u_{xz}v_{xy} + \frac{1}{2}v_xu_{xyz} - \frac{1}{4}u_{yz}v_{xx} \\
&\quad + \frac{1}{4}v_yu_{xxz} + \frac{3}{4}u_zv_{xxy} - \frac{3}{4}vu_{xxyz} + \frac{1}{2}u_zv_t - \frac{1}{2}vu_{tz}, \\
Y_4 &= \frac{3}{2}u_zu_xv_x + \frac{1}{4}u_{xxz}v_x - \frac{3}{2}vu_xu_{xz} - \frac{3}{2}vu_zu_{xx} - \frac{1}{4}u_{xz}v_{xx} + \frac{1}{4}u_zv_{xxx} - \frac{1}{4}vu_{xxxz} + \frac{1}{2}u_zv_t \\
&\quad - \frac{1}{2}vu_{tz}, \\
Z_4 &= -u_zv_z + 3vu_xu_{xy} + 3vu_yu_{xx} + vu_{xxx} + \frac{1}{2}u_zv_t + \frac{1}{2}vu_{tz} + vu_{ty} + vu_{tx};
\end{aligned}$$

$$\begin{aligned}
T_5 &= -\frac{1}{2}vu_z + tv_zu_z + \frac{1}{2}zv_zu_z + tv_yu_z + \frac{1}{2}zv_yu_z + tv_xu_z + \frac{1}{2}zv_xu_z - \frac{1}{2}zv_u_{zz} - \frac{1}{2}vu_y \\
&\quad + \frac{1}{2}zv_zu_y + \frac{1}{2}zu_yv_y - tv_u_{yz} - zv_u_{yz} - \frac{1}{2}zv_u_{yy} - \frac{1}{2}vu_x + \frac{1}{2}zv_zu_x + \frac{1}{2}zv_yu_x + \frac{1}{2}zu_yv_x \\
&\quad + \frac{1}{2}zu_xv_x - tv_u_{xz} - zv_u_{xz} - zv_u_{xy} - 3tv_u_{xxy} - \frac{1}{2}zv_u_{xx} - 3tv_u_{xy}u_{xx} - tv_u_{xxx} \\
&\quad - \frac{1}{2}tv_zu_t - \frac{1}{2}tv_yu_t - \frac{1}{2}tv_xu_t - \frac{1}{2}tv_u_{tz} - \frac{1}{2}tv_u_{ty} - \frac{1}{2}tv_u_{tx}, \\
X_5 &= 3zv_xu_y^2 + \frac{3}{2}zv_yu_xu_y + 6tu_zv_xu_y + 3zu_zv_xu_y + 3zu_xv_xu_y - 6tv_u_{xz}u_y - 3zv_u_{xz}u_y \\
&\quad - \frac{3}{2}zv_u_{xy}u_y + \frac{3}{4}zv_{xxy}u_y - 3tv_xu_tu_y + \frac{1}{2}zv_tu_y + 3tv_u_{tx}u_y + \frac{3}{2}zv_yu_x^2 - vu_z - zv_u_{zz} \\
&\quad + 3tu_zv_yu_x + \frac{3}{2}zu_zv_yu_x - 3tv_u_{yz}u_x - \frac{3}{2}zv_u_{yz}u_x - \frac{3}{2}zv_u_{yy}u_x + 3tv_u_{zxy} + \frac{3}{2}zv_u_{zxy} \\
&\quad + 3zv_u_{xxy} - tu_{xz}v_{xy} - \frac{1}{2}zu_{xz}v_{xy} - \frac{1}{2}zu_{xy}v_{xy} + tv_xu_{xyz} + \frac{1}{2}zv_xu_{xyz} + \frac{1}{2}zv_xu_{xyy} \\
&\quad - \frac{1}{2}zv_{xy}u_{xx} - \frac{1}{2}tu_{yz}v_{xx} - \frac{1}{4}zu_{yz}v_{xx} - \frac{1}{4}zu_{yy}v_{xx} - \frac{1}{4}zu_{xy}v_{xx} + \frac{1}{2}tv_yu_{xxz} \\
&\quad + \frac{1}{4}zv_yu_{xxz} + \frac{1}{4}zv_yu_{xxy} + \frac{1}{2}zv_xu_{xxy} + \frac{3}{2}tu_zv_{xxy} + \frac{3}{4}zu_zv_{xxy} + \frac{3}{4}zu_xv_{xxy} - \frac{3}{2}tv_u_{xxyz} \\
&\quad - \frac{3}{4}zv_u_{xxyz} - \frac{3}{4}zv_u_{xxyy} + \frac{1}{4}zv_yu_{xxx} + \frac{1}{4}zv_u_{xxx} + \frac{1}{2}vu_t - \frac{3}{2}tv_yu_xu_t - \frac{3}{2}tv_u_{xy}u_t \\
&\quad - \frac{3}{4}tv_{xxy}u_t + tu_zv_t + \frac{1}{2}zu_zv_t + \frac{1}{2}zu_xv_t - \frac{1}{2}tu_tv_t - tv_u_{tz} + \frac{1}{2}zv_u_{tz} + \frac{1}{2}zv_u_{ty} \\
&\quad + \frac{3}{2}tv_u_{xxy}u_t + \frac{1}{4}tv_{xx}u_{ty} + \frac{1}{2}zv_u_{tx} + \frac{1}{2}tv_{xy}u_{tx} - \frac{1}{2}tv_xu_{txy} - \frac{1}{4}tv_yu_{txx} + \frac{3}{4}tv_u_{txxy} + \frac{1}{2}tv_u_{tt},
\end{aligned}$$

$$\begin{aligned}
Y_5 = & \frac{3}{2}zv_xu_x^2 + 3tu_zv_xu_x + \frac{3}{2}zu_zv_xu_x + \frac{3}{2}zu_yv_xu_x - 3tvu_{xz}u_x - \frac{3}{2}zvu_{xz}u_x \\
& + \frac{3}{2}zvu_{xy}u_x - 3zvu_{xx}u_x + \frac{1}{4}zv_{xxx}u_x - \frac{3}{2}tv_xu_tu_x + \frac{1}{2}zv_tu_x + \frac{3}{2}tvu_{tx}u_x \\
& - vu_z - zvu_{zz} - 3tvu_zu_{xx} - \frac{3}{2}zvu_zu_{xx} + \frac{3}{2}zvu_yu_{xx} - \frac{1}{2}tu_{xz}v_{xx} - \frac{1}{4}zu_{xz}v_{xx} \\
& - \frac{1}{4}zu_{xy}v_{xx} - \frac{1}{4}zu_{xx}v_{xx} + \frac{1}{2}tv_xu_{xxz} + \frac{1}{4}zv_xu_{xxz} + \frac{1}{4}zv_xu_{xxy} + \frac{1}{4}zv_xu_{xxx} \\
& + \frac{1}{2}tu_zv_{xxx} + \frac{1}{4}zu_zv_{xxx} + \frac{1}{4}zu_yv_{xxx} - \frac{1}{2}tvu_{xxxz} - \frac{1}{4}zvu_{xxxz} + \frac{3}{4}zvu_{xxxy} \\
& - \frac{1}{4}zvu_{xxxx} + \frac{1}{2}vu_t + \frac{3}{2}tvu_{xx}u_t - \frac{1}{4}tv_{xxx}u_t + tu_zv_t + \frac{1}{2}zu_zv_t + \frac{1}{2}zu_yv_t \\
& - \frac{1}{2}tu_tv_t - tvu_{tz} + \frac{1}{2}zvu_{tz} + \frac{1}{2}zvu_{ty} + \frac{1}{2}zvu_{tx} + \frac{1}{4}tv_{xx}u_{tx} - \frac{1}{4}tv_xu_{txx} \\
& + \frac{1}{4}tvu_{txxx} + \frac{1}{2}tvu_{tt},
\end{aligned}$$

$$\begin{aligned}
Z_5 = & -2tu_zv_z - zu_zv_z - zu_yv_z - zu_xv_z + tu_tv_z + vu_y + zvu_{yz} + vu_x + zvu_{xz} \\
& + 6tvu_xu_{xy} + 3zvu_xu_{xy} + 6tvu_yu_{xx} + 3zvu_yu_{xx} + 2tvu_{xxxy} + zvu_{xxxxy} \\
& + \frac{1}{2}vu_t + tu_zv_t + \frac{1}{2}zu_zv_t + \frac{1}{2}zu_yv_t + \frac{1}{2}zu_xv_t - \frac{1}{2}tu_tv_t + \frac{1}{2}zvu_{tz} + 2tvu_{ty} \\
& + \frac{1}{2}zvu_{ty} + 2tvu_{tx} + \frac{1}{2}zvu_{tx} + \frac{1}{2}tvu_{tt};
\end{aligned}$$

$$\begin{aligned}
T_6 = & -3vu_z - 9tu_{zz}v - \frac{3}{2}zu_{zz}v - 3u_yv + 6tu_{yz}v - \frac{9}{2}yu_{yz}v + \frac{3}{2}zu_{yz}v - \frac{9}{2}yu_{yy}v \\
& + 3zvu_{yy} - 3u_xv - \frac{3}{2}xu_{xz}v - \frac{3}{2}zu_{xz}v - 6tu_{xy}v - \frac{3}{2}xu_{xy}v - \frac{9}{2}yu_{xy}v + 3zu_{xy}v \\
& + 45tu_xu_{xy}v - 6tu_{xx}v - \frac{3}{2}xu_{xx}v + 45tu_yu_{xx}v + 15tu_{xxxy}v + \frac{15}{2}tu_{tz}v + \frac{15}{2}tu_{ty}v \\
& + \frac{15}{2}tvu_{tx} + 4v - 2xv_z - 2yv_z + \frac{3}{2}uv_z - 6tu_zv_z + \frac{3}{2}zu_zv_z + \frac{9}{2}yv_zu_y - 3zv_zu_y \\
& - 2xv_y - 2yv_y + \frac{3}{2}uv_y - 6tu_zv_y + \frac{3}{2}zu_zv_y + \frac{9}{2}yu_yv_y - 3zu_yv_y + 6tv_zu_x \\
& + \frac{3}{2}xv_zu_x + 6tv_yu_x + \frac{3}{2}xv_yu_x - 2xv_x - 2yv_x + \frac{3}{2}uv_x - 6tu_zv_x + \frac{3}{2}zu_zv_x \\
& + \frac{9}{2}yu_yv_x - 3zu_yv_x + 6tu_xv_x + \frac{3}{2}xu_xv_x + \frac{15}{2}tv_zu_t + \frac{15}{2}tv_yu_t + \frac{15}{2}tv_xu_t,
\end{aligned}$$

$$\begin{aligned}
X_6 = & 27yv_xu_y^2 - 18zv_xu_y^2 + 12vu_y - 36vu_xu_y + \frac{27}{2}yv_yu_xu_y - 9zv_yu_xu_y - 12xv_xu_y \\
& - 12yv_xu_y + 9uv_xu_y - 36tu_zv_xu_y + 9zu_zv_xu_y + 36tu_xv_xu_y + 9xu_xv_xu_y \\
& + 36tvu_{xz}u_y - 9zv_{xz}u_y - \frac{27}{2}yv_{xy}u_y + 9zv_{xy}u_y - 3v_{xx}u_y + \frac{27}{4}yv_{xy}u_y \\
& - \frac{9}{2}zv_{xy}u_y + 45tv_xu_tu_y + \frac{9}{2}yv_tu_y - 3zv_tu_y - 45tv_{tx}u_y + 18tv_yu_x^2 + \frac{9}{2}xv_yu_x^2 \\
& + 6vu_z - 12tvu_{zz} - 3xv_{zz} - 6xv_yu_x - 6yv_yu_x + \frac{9}{2}uv_yu_x - 18tu_zv_yu_x \\
& + \frac{9}{2}zu_zv_yu_x + 18tv_{yz}u_x - \frac{9}{2}zv_{yz}u_x - \frac{27}{2}yv_{yy}u_x + 9zv_{yy}u_x - 6xv_{xy} \\
& - 6yv_{xy} + \frac{9}{2}uv_{xy} - 18tv_zu_{xy} + \frac{9}{2}zv_zu_{xy} + 36tv_{xz}u_{xy} + 9xv_{xz}u_{xy} \\
& + \frac{15}{2}v_xu_{xy} - 3u_xv_{xy} + 6tu_{xz}v_{xy} - \frac{3}{2}zu_{xz}v_{xy} - \frac{9}{2}yu_{xy}v_{xy} + 3zu_{xy}v_{xy} + 2v_{xy} \\
& - 6tv_xu_{xyz} + \frac{3}{2}zv_xu_{xyz} + \frac{9}{2}yv_xu_{xyy} - 3zv_xu_{xyy} + \frac{9}{4}v_yu_{xx} - 6tv_{xy}u_{xx} - \frac{3}{2}xv_{xy}u_{xx} \\
& + 3tu_{yz}v_{xx} - \frac{3}{4}zu_{yz}v_{xx} - \frac{9}{4}yu_{yy}v_{xx} + \frac{3}{2}zu_{yy}v_{xx} - 3tu_{xy}v_{xx} - \frac{3}{4}xu_{xy}v_{xx} + v_{xx} \\
& - 3tv_yu_{xxz} + \frac{3}{4}zv_yu_{xxz} - \frac{27}{2}vu_{xxy} + \frac{9}{4}yv_yu_{xxy} - \frac{3}{2}zv_yu_{xxy} + 6tv_xu_{xxy} \\
& + \frac{3}{2}xv_xu_{xxy} - 3xv_{xxy} - 3yv_{xxy} + \frac{9}{4}uv_{xxy} - 9tu_zv_{xxy} + \frac{9}{4}zu_zv_{xxy} + 9tu_xv_{xxy} \\
& + \frac{9}{4}xu_xv_{xxy} + 9tv_{xxyz} - \frac{9}{4}zv_{xxyz} - \frac{27}{4}yv_{xxyy} + \frac{9}{2}zv_{xxyy} + 3tv_yu_{xxx} \\
& + \frac{3}{4}xv_yu_{xxx} + 3tv_{xxxy} + \frac{3}{4}xv_{xxxy} - 9vu_t + \frac{45}{2}tv_yu_xu_t + \frac{45}{2}tv_{xy}u_t + \frac{45}{4}tv_{xxy}u_t \\
& - 2xv_t - 2yv_t + \frac{3}{2}uv_t - 6tu_zv_t + \frac{3}{2}zu_zv_t + 6tu_xv_t + \frac{3}{2}xu_xv_t + \frac{15}{2}tu_tv_t + 18tv_{tz} \\
& + 3xv_{tz} - \frac{3}{2}zv_{tz} + 12tv_{ty} + 3xv_{ty} - \frac{9}{2}yv_{ty} + 3zv_{ty} - \frac{45}{2}tv_{tx}u_{ty} \\
& - \frac{15}{4}tv_{xx}u_{ty} + 6tv_{tx} + \frac{3}{2}xv_{tx} - \frac{15}{2}tv_{xy}u_{tx} + \frac{15}{2}tv_xu_{txy} + \frac{15}{4}tv_yu_{txx} \\
& - \frac{45}{4}tv_{txxy} - \frac{15}{2}tv_{tt},
\end{aligned}$$

$$\begin{aligned}
Y_6 = & -9vu_x^2 + 18tv_xu_x^2 + \frac{9}{2}xv_xu_x^2 - 6xv_xu_x - 6yv_xu_x + \frac{9}{2}uv_xu_x - 18tu_zv_xu_x \\
& + \frac{9}{2}zu_zv_xu_x + \frac{27}{2}yu_yv_xu_x - 9zu_yv_xu_x + 18tvu_{xz}u_x - \frac{9}{2}zv_{xz}u_x + \frac{27}{2}yvu_{xy}u_x \\
& - 9zv_{xy}u_x - 36tvu_{xx}u_x - 9xv_{xx}u_x - \frac{3}{2}v_{xx}u_x + 3tv_{xxx}u_x + \frac{3}{4}xv_{xxx}u_x \\
& + \frac{45}{2}tv_xu_tu_x + 6tv_tu_x + \frac{3}{2}xv_tu_x - \frac{45}{2}tvu_{tx}u_x + 6vu_z - 9yv_{zz} + 6zv_{zz} \\
& + 6xv_{xx} + 6yv_{xx} - \frac{9}{2}uv_{xx} + 18tvu_{zx} - \frac{9}{2}zv_{zx} + \frac{27}{2}yvu_{yx} \\
& - 9zv_{yx} + \frac{9}{4}v_{xx} + 3tu_{xz}v_{xx} - \frac{3}{4}zu_{xz}v_{xx} - \frac{9}{4}yu_{xy}v_{xx} + \frac{3}{2}zu_{xy}v_{xx} \\
& - 3tu_{xx}v_{xx} - \frac{3}{4}xu_{xx}v_{xx} + v_{xx} - 3tv_{xx} + \frac{3}{4}zv_{xx} + \frac{9}{4}yv_{xx} - \frac{3}{2}zv_{xx} \\
& - 3vu_{xxx} + 3tv_{xxx} + \frac{3}{4}xv_{xxx} - xv_{xxx} - yv_{xxx} + \frac{3}{4}uv_{xxx} - 3tu_{zx}v_{xxx} \\
& + \frac{3}{4}zu_{zx}v_{xxx} + \frac{9}{4}yu_{yz}v_{xxx} - \frac{3}{2}zu_{yz}v_{xxx} + 3tvu_{xxxz} - \frac{3}{4}zv_{xxxz} + \frac{27}{4}yvu_{xxxxy} \\
& - \frac{9}{2}zv_{xxxxy} - 3tvu_{xxxxy} - \frac{3}{4}xv_{xxxxy} - 9vu_t - \frac{45}{2}tvu_{xx}u_t + \frac{15}{4}tv_{xxx}u_t - 2xv_t \\
& - 2yv_t + \frac{3}{2}uv_t - 6tu_zv_t + \frac{3}{2}zu_zv_t + \frac{9}{2}yu_yv_t - 3zu_yv_t + \frac{15}{2}tu_tv_t + 6tv_{tz} \\
& + 9yv_{tz} - \frac{15}{2}zv_{tz} + \frac{9}{2}yvu_{ty} - 3zv_{ty} - 6tv_{tx} - \frac{3}{2}xv_{tx} + 9yv_{tx} \\
& - 6zv_{tx} - \frac{15}{4}tv_{xx}u_{tx} + \frac{15}{4}tv_{xx}u_{tx} \\
& - \frac{15}{4}tv_{txxx} - \frac{15}{2}tv_{ttx},
\end{aligned}$$

$$\begin{aligned}
Z_6 = & 12vu_z + 12tv_zu_z - 3zv_zu_z - 6tv_tu_z + \frac{3}{2}zv_tu_z + 4xv_z + 4yv_z - 3uv_z \\
& - 6vu_y - 9yv_zu_y + 6zv_zu_y + 9yv_{yz} - 6zv_{yz} - 6vu_x - 12tv_zu_x \\
& - 3xv_zu_x + 12tvu_{xz} + 3xv_{xz} - 36tvu_{xz}u_{xy} + 9zv_{xz}u_{xy} - 36tvu_{yx} \\
& + 9zv_{yx}u_{xx} - 12tvu_{xxx} + 3zv_{xxx} - 9vu_t - 15tv_zu_t - 2xv_t - 2yv_t \\
& + \frac{3}{2}uv_t + \frac{9}{2}yu_yv_t - 3zu_yv_t + 6tu_xv_t + \frac{3}{2}xu_xv_t + \frac{15}{2}tu_tv_t + 9tv_{tz} + \frac{3}{2}zv_{tz} \\
& - 12tv_{ty} - \frac{9}{2}yvu_{ty} + 6zv_{ty} - 18tv_{tx} - \frac{3}{2}xv_{tx} + 3zv_{tx} - \frac{15}{2}tv_{ttx};
\end{aligned}$$

$$T_{F_1} = \frac{1}{2}F_1(t)v_x - \frac{1}{2}F_1(t)v_y - \frac{1}{2}F_1(t)v_z,$$

$$X_{F_1} = -\frac{3}{2}F_1(t)u_{xy}v + \frac{1}{2}F_1'(t)v - \frac{3}{2}F_1(t)u_xv_y - 3F_1(t)u_yv_x - \frac{3}{4}F_1(t)v_{xy} - \frac{1}{2}F_1(t)v_t,$$

$$Y_{F_1} = \frac{3}{2}F_1(t)u_{xx}v + \frac{1}{2}F_1'(t)v - \frac{3}{2}F_1(t)u_xv_x - \frac{1}{4}F_1(t)v_{xx} - \frac{1}{2}F_1(t)v_t,$$

$$Z_{F_1} = \frac{1}{2}F_1'(t)v + F_1(t)v_z - \frac{1}{2}F_1(t)v_t;$$

$$\begin{aligned}
T_{F_2} &= \frac{1}{2}F_2'v - \frac{1}{2}v_xF_2(t+z) - \frac{1}{2}v_yF_2(t+z) - \frac{1}{2}v_zF_2(t+z), \\
X_{F_2} &= -\frac{3}{2}F_2(t+z)u_{xy}v + \frac{1}{2}F_2'v - \frac{3}{2}u_xv_yF_2(t+z) - 3u_yv_xF_2(t+z) \\
&\quad - \frac{3}{4}F_2(t+z)v_{xxy} - \frac{1}{2}v_tF_2(t+z), \\
Y_{F_2} &= \frac{3}{2}u_{xx}F_2(t+z)v + \frac{1}{2}F_2'v - \frac{3}{2}u_xv_xF_2(t+z) - \frac{1}{4}v_{xxx}F_2(t+z) - \frac{1}{2}v_tF_2(t+z), \\
Z_{F_2} &= -\frac{1}{2}F_2'v + v_zF_2(t+z) - \frac{1}{2}v_tF_2(t+z).
\end{aligned}$$

4. Concluding Remarks

In this work, we used Lie symmetry methods to obtain analytical solutions of the (3+1)-dimensional generalised KP Equation (3). The first solution contains an elliptic integral of the second kind, an amplitude function, a delta amplitude function and a Jacobi cosine function. It is common knowledge that these functions degenerate to trigonometric or hyperbolic functions depending on the behaviour of the parameter M^2 . Thus, the solution (13) is to our knowledge the most general solution of (3) that has been obtained to date. Some of the solutions obtained were presented graphically. Furthermore, we obtained logarithmic and rational solutions. Finally, we computed the conserved quantities of Equation (3) using the multiplier method as well as Ibragimov's conservation theorem. The former method yielded a local low-order conserved quantity, while the later method yielded eight conservation laws with each conserved vector corresponding to an infinitesimal generator.

Author Contributions: Conceptualization, I.S. and C.M.K.; methodology, C.M.K.; software, I.S.; validation, C.M.K.; writing—original draft preparation, I.S.; writing—review and editing, C.M.K. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Acknowledgments: The authors thank North-West University for its continued support and sincerely thank the reviewers for their positive suggestions, which helped to improve the paper enormously.

Conflicts of Interest: The authors declare no conflict of interest.

Appendix A. Prolongation Coefficients: Full Expansions

$$\begin{aligned}
\zeta_x &= \eta_x + u_x\eta_u - u_tu_x\zeta_u^1 - u_t\zeta_x^1 - u_x^2\zeta_u^2 - u_x\zeta_x^2 - u_xu_y\zeta_u^3 - u_y\zeta_x^3 - u_xu_z\zeta_u^4 \\
&\quad - u_z\zeta_x^4, \\
\zeta_y &= \eta_y + \eta_uu_y - u_t\zeta_u^1u_y - u_t\zeta_y^1 - \zeta_u^2u_xu_y - u_x\zeta_y^2 - \zeta_u^4u_yu_z - u_z\zeta_y^4 - u_y^2\zeta_u^3 \\
&\quad - u_y\zeta_y^3, \\
\zeta_{tx} &= u_tu_x\eta_{uu} - u_t\zeta_{uu}^2u_x^2 - \zeta_{tu}^2u_x^2 - 2u_{tx}\zeta_u^2u_x - u_{tt}\zeta_u^1u_x - u_{ty}\zeta_u^3u_x - u_{tz}\zeta_u^4u_x \\
&\quad - u_t^2\zeta_{uu}^1u_x - u_yu_t\zeta_{uu}^3u_x - u_zu_t\zeta_{uu}^4u_x - u_t\zeta_{xu}^2u_x + \eta_{tu}u_x - u_t\zeta_{tu}^1u_x \\
&\quad - u_z\zeta_{tu}^4u_x - \zeta_{tx}^2u_x + u_{tx}\eta_u - u_{xx}u_t\zeta_u^2 - 2u_tu_{tx}\zeta_u^1 - u_{xy}u_t\zeta_u^3 - u_yu_{tx}\zeta_u^3 \\
&\quad - u_{xz}u_t\zeta_u^4 - u_zu_{tx}\zeta_u^4 - u_{tx}\zeta_x^2 - u_{tt}\zeta_x^1 - u_{ty}\zeta_x^3 - u_{tz}\zeta_x^4 + u_t\eta_{xu} - u_t^2\zeta_{xu}^1 \\
&\quad - u_yu_t\zeta_{xu}^3 - u_zu_t\zeta_{xu}^4 - u_{xx}\zeta_t^2 - u_{tx}\zeta_t^1 - u_{xy}\zeta_t^3 - u_{xz}\zeta_t^4 + \eta_{tx} - u_t\zeta_{tx}^1 \\
&\quad - u_y\zeta_{tx}^3 - u_z\zeta_{tx}^4 - u_y\zeta_{tu}^3u_x,
\end{aligned}$$

$$\begin{aligned}
\zeta_{ty} &= u_t \eta_{uu} u_y - u_t \zeta_{uu}^3 u_y^2 - \zeta_{tu}^3 u_y^2 - u_{tx} \zeta_u^2 u_y - u_{tt} \zeta_u^1 u_y - 2u_{ty} \zeta_u^3 u_y - u_{tz} \zeta_u^4 u_y \\
&\quad - u_x u_t \zeta_{uu}^2 u_y - u_t^2 \zeta_{uu}^1 u_y - u_z u_t \zeta_{uu}^4 u_y - u_t \zeta_{yu}^3 u_y + \eta_{tu} u_y - u_x \zeta_{tu}^2 u_y \\
&\quad - u_z \zeta_{tu}^4 u_y - \zeta_{ty}^3 u_y + u_{ty} \eta_u - u_{xy} u_t \zeta_u^2 - u_x u_{ty} \zeta_u^2 - 2u_t u_{ty} \zeta_u^1 - u_{yy} u_t \zeta_u^3 \\
&\quad - u_{yz} u_t \zeta_u^4 - u_t \zeta_{tu}^1 u_y - u_z u_{ty} \zeta_u^4 - u_{tx} \zeta_y^2 - u_{tt} \zeta_y^1 - u_{ty} \zeta_y^3 - u_{tz} \zeta_y^4 + u_t \eta_{yu} \\
&\quad - u_x u_t \zeta_{yu}^2 - u_t^2 \zeta_{yu}^1 - u_z u_t \zeta_{yu}^4 - u_{xy} \zeta_t^2 - u_{ty} \zeta_t^1 - u_{yy} \zeta_t^3 - u_{yz} \zeta_t^4 + \eta_{ty} \\
&\quad - u_x \zeta_{ty}^2 - u_t \zeta_{ty}^1 - u_z \zeta_{ty}^4, \\
\zeta_{tz} &= u_t \eta_{uu} u_z - u_t \zeta_{uu}^4 u_z^2 - \zeta_{tu}^4 u_z^2 - u_{tx} \zeta_u^2 u_z - u_{tt} \zeta_u^1 u_z - u_{ty} \zeta_u^3 u_z - 2u_{tz} \zeta_u^4 u_z \\
&\quad - u_x u_t \zeta_{uu}^2 u_z - u_t^2 \zeta_{uu}^1 u_z - u_y u_t \zeta_{uu}^3 u_z - u_t \zeta_{zu}^4 u_z + \eta_{tu} u_z - u_x \zeta_{tu}^2 u_z \\
&\quad - u_t \zeta_{tu}^1 u_z - u_y \zeta_{tu}^3 u_z - \zeta_{tz}^4 u_z + u_{tz} \eta_u - u_{xz} u_t \zeta_u^2 - u_x u_{tz} \zeta_u^2 - 2u_t u_{tz} \zeta_u^1 \\
&\quad - u_{yz} u_t \zeta_u^3 - u_y u_{tz} \zeta_u^3 - u_{zz} u_t \zeta_u^4 - u_{tx} \zeta_z^2 - u_{tt} \zeta_z^1 - u_{ty} \zeta_z^3 - u_{tz} \zeta_z^4 + u_t \eta_{zu} \\
&\quad - u_x u_t \zeta_{zu}^2 - u_t^2 \zeta_{zu}^1 - u_y u_t \zeta_{zu}^3 - u_{xz} \zeta_t^2 - u_{tz} \zeta_t^1 - u_{yz} \zeta_t^3 - u_{zz} \zeta_t^4 + \eta_{tz} \\
&\quad - u_x \zeta_{tz}^2 - u_t \zeta_{tz}^1 - u_y \zeta_{tz}^3, \\
\zeta_{xx} &= \eta_{uu} u_x^2 - \zeta_{uu}^2 u_x^3 - u_t \zeta_{uu}^1 u_x^2 - u_y \zeta_{uu}^3 u_x^2 - u_z \zeta_{uu}^4 u_x^2 - 2\zeta_{xu}^2 u_x^2 - 3u_{xx} \zeta_u^2 u_x \\
&\quad - 2u_{tx} \zeta_u^1 u_x - 2u_{xy} \zeta_u^3 u_x - 2u_{xz} \zeta_u^4 u_x + 2\eta_{xu} u_x - 2u_t \zeta_{xu}^1 u_x - 2u_y \zeta_{xu}^3 u_x \\
&\quad - 2u_z \zeta_{xu}^4 u_x - \zeta_{xx}^2 u_x + u_{xx} \eta_u - u_{xx} u_t \zeta_u^1 - u_y u_{xx} \zeta_u^3 - u_z u_{xx} \zeta_u^4 - 2u_{xx} \zeta_x^2 \\
&\quad - 2u_{tx} \zeta_x^1 - 2u_{xy} \zeta_x^3 - 2u_{xz} \zeta_x^4 + \eta_{xx} - u_t \zeta_{xx}^1 - u_y \zeta_{xx}^3 - u_z \zeta_{xx}^4, \\
\zeta_{xy} &= u_x \eta_{uu} u_y - u_x \zeta_{uu}^3 u_y^2 - \zeta_{xu}^3 u_y^2 - u_{xx} \zeta_u^2 u_y - u_{tx} \zeta_u^1 u_y - 2u_{xy} \zeta_u^3 u_y - u_{xz} \zeta_u^4 u_y \\
&\quad - u_x^2 \zeta_{uu}^2 u_y - u_x u_t \zeta_{uu}^1 u_y - u_z u_x \zeta_{uu}^4 u_y - u_x \zeta_{yu}^3 u_y + \eta_{xu} u_y - u_x \zeta_{xu}^2 u_y \\
&\quad - u_t \zeta_{xu}^1 u_y - u_z \zeta_{xu}^4 u_y - \zeta_{xy}^3 u_y + u_{xy} \eta_u - 2u_x u_{xy} \zeta_u^2 - u_{xy} u_t \zeta_u^1 - u_x u_{ty} \zeta_u^1 \\
&\quad - u_{yy} u_x \zeta_u^3 - u_{yz} u_x \zeta_u^4 - u_z u_{xy} \zeta_u^4 - u_{xx} \zeta_y^2 - u_{tx} \zeta_y^1 - u_{xy} \zeta_y^3 - u_{xz} \zeta_y^4 \\
&\quad + u_x \eta_{yu} - u_x^2 \zeta_{yu}^2 - u_x u_t \zeta_{yu}^1 - u_z u_x \zeta_{yu}^4 - u_{xy} \zeta_x^2 - u_{ty} \zeta_x^1 - u_{yy} \zeta_x^3 - u_{yz} \zeta_x^4 \\
&\quad + \eta_{xy} - u_x \zeta_{xy}^2 - u_t \zeta_{xy}^1 - u_z \zeta_{xy}^4, \\
\zeta_{zz} &= \eta_{uu} u_z^2 - \zeta_{uu}^4 u_z^3 - u_x \zeta_{uu}^2 u_z^2 - u_t \zeta_{uu}^1 u_z^2 - u_y \zeta_{uu}^3 u_z^2 - 2\zeta_{zu}^4 u_z^2 - 2u_{xz} \zeta_u^2 u_z \\
&\quad - 2u_{tz} \zeta_u^1 u_z - 2u_{yz} \zeta_u^3 u_z - 3u_{zz} \zeta_u^4 u_z + 2\eta_{zu} u_z - 2u_x \zeta_{zu}^2 u_z - 2u_t \zeta_{zu}^1 u_z \\
&\quad - 2u_y \zeta_{zu}^3 u_z - \zeta_{zz}^4 u_z + u_{zz} \eta_u - u_{zz} u_x \zeta_u^2 - u_{zz} u_t \zeta_u^1 - u_{zz} u_y \zeta_u^3 - 2u_{xz} \zeta_z^2 \\
&\quad - 2u_{tz} \zeta_z^1 - 2u_{yz} \zeta_z^3 - 2u_{zz} \zeta_z^4 + \eta_{zz} - u_x \zeta_{zz}^2 - u_t \zeta_{zz}^1 - u_y \zeta_{zz}^3, \\
\zeta_{xxx} &= u_y \eta_{uuuu} u_x^3 - u_y \zeta_{uuuu}^1 u_x^4 - \zeta_{yuuu}^2 u_x^4 - 4u_{xy} \zeta_{uuuu}^2 u_x^3 - u_{ty} \zeta_{uuuu}^1 u_x^3 - u_{yy} \zeta_{uuuu}^3 u_x^3 \\
&\quad - u_{yz} \zeta_{uuuu}^4 u_x^3 - u_y u_t \zeta_{uuuu}^1 u_x^3 - u_y^2 \zeta_{uuuu}^3 u_x^3 - u_z u_y \zeta_{uuuu}^4 u_x^3 + \eta_{yuuu} u_x^3 \\
&\quad - u_t \zeta_{yuuu}^1 u_x^3 - u_y \zeta_{yuuu}^3 u_x^3 - u_z \zeta_{yuuu}^4 u_x^3 - 3u_y \zeta_{xuuu}^2 u_x^3 - 3\zeta_{xyuu}^2 u_x^3 \\
&\quad - 6u_{xy} \zeta_{uuu}^2 u_x^2 - 3u_{txy} \zeta_{uuu}^1 u_x^2 - 3u_{xyy} \zeta_{uuu}^3 u_x^2 - 3u_{xyz} \zeta_{uuu}^4 u_x^2 + 3u_{xy} \eta_{uuu} u_x^2 \\
&\quad - 6u_y u_{xx} \zeta_{uuu}^2 u_x^2 - 3u_{xy} u_t \zeta_{uuu}^1 u_x^2 - 3u_y u_{tx} \zeta_{uuu}^3 u_x^2 - 6u_y u_{xy} \zeta_{uuu}^4 u_x^2 \\
&\quad - 3u_y u_{xz} \zeta_{uuu}^4 u_x^2 - 3u_z u_{xy} \zeta_{uuu}^4 u_x^2 - 6u_{xx} \zeta_{uuu}^2 u_x^2 - 3u_{tx} \zeta_{uuu}^1 u_x^2 - 3u_{xy} \zeta_{uuu}^3 u_x^2 \\
&\quad - 3u_{xz} \zeta_{uuu}^4 u_x^2 - 9u_{xy} \zeta_{xuu}^2 u_x^2 - 3u_{ty} \zeta_{xuu}^1 u_x^2 - 3u_{yy} \zeta_{xuu}^3 u_x^2 - 3u_{yz} \zeta_{xuu}^4 u_x^2 \\
&\quad + 3u_y \eta_{xuuu} u_x^2 - 3u_y u_t \zeta_{xuuu}^1 u_x^2 - 3u_y^2 \zeta_{xuuu}^3 u_x^2 - 3u_z u_y \zeta_{xuuu}^4 u_x^2 + 3\eta_{xyuu} u_x^2 \\
&\quad - 3u_t \zeta_{xyuu}^1 u_x^2 - 3u_y \zeta_{xyuu}^3 u_x^2 - 3u_z \zeta_{xyuu}^4 u_x^2 - 3u_y \zeta_{xxuu}^2 u_x^2 - 3\zeta_{xxyu}^2 u_x^2 \\
&\quad - 4u_{xxx} \zeta_u^2 u_x - u_{xxx} \zeta_u^1 u_x - 3u_{txxy} \zeta_u^1 u_x - u_{txxx} \zeta_u^1 u_x - 3u_{xxyy} \zeta_u^3 u_x
\end{aligned}$$

$$\begin{aligned}
& -u_{xxxy}\tilde{\zeta}_u^3 u_x - 3u_{xxyz}\tilde{\zeta}_u^4 u_x - u_{xxxz}\tilde{\zeta}_u^4 u_x + 3u_{xxy}\eta_{uu} u_x - 12u_{xy}u_{xx}\tilde{\zeta}_{uu}^2 u_x \\
& - 4u_y u_{xxx}\tilde{\zeta}_{uu}^2 u_x - 3u_{xxy}u_t\tilde{\zeta}_{uu}^1 u_x - 3u_{xx}u_{ty}\tilde{\zeta}_{uu}^1 u_x - 6u_{xy}u_{tx}\tilde{\zeta}_{uu}^1 u_x \\
& - 3u_y u_{txx}\tilde{\zeta}_{uu}^1 u_x - 6u_{xy}^2\tilde{\zeta}_{uu}^3 u_x - 3u_{yy}u_{xx}\tilde{\zeta}_{uu}^3 u_x - 6u_y u_{xxy}\tilde{\zeta}_{uu}^3 u_x \\
& - 6u_{xz}u_{xy}\tilde{\zeta}_{uu}^4 u_x - 3u_{yz}u_{xx}\tilde{\zeta}_{uu}^4 u_x - 3u_y u_{xxz}\tilde{\zeta}_{uu}^4 u_x - 3u_z u_{xxy}\tilde{\zeta}_{uu}^4 u_x \\
& + 3u_y u_{xx}\eta_{uuu} u_x - 3u_y u_{xx}u_t\tilde{\zeta}_{uuu}^1 u_x - 3u_y^2 u_{xx}\tilde{\zeta}_{uuu}^3 u_x - 3u_z u_y u_{xx}\tilde{\zeta}_{uuu}^4 u_x \\
& - 4u_{xxx}\tilde{\zeta}_{yu}^2 u_x - 3u_{txx}\tilde{\zeta}_{yu}^1 u_x - 3u_{xxy}\tilde{\zeta}_{yu}^3 u_x - 3u_{xxz}\tilde{\zeta}_{yu}^4 u_x + 3u_{xx}\eta_{yuu} u_x \\
& - 3u_{xx}u_t\tilde{\zeta}_{yuu}^1 u_x - 3u_y u_{xx}\tilde{\zeta}_{yuu}^3 u_x - 3u_z u_{xx}\tilde{\zeta}_{yuu}^4 u_x - 9u_{xxy}\tilde{\zeta}_{xu}^2 u_x \\
& - 6u_{txy}\tilde{\zeta}_{xu}^1 u_x - 6u_{xyy}\tilde{\zeta}_{xu}^3 u_x - 6u_{yz}\tilde{\zeta}_{xu}^4 u_x + 6u_{xy}\eta_{xuu} u_x - 9u_y u_{xx}\tilde{\zeta}_{xuu}^2 u_x \\
& - 6u_{xy}u_t\tilde{\zeta}_{xuu}^1 u_x - 6u_y u_{tx}\tilde{\zeta}_{xuu}^1 u_x - 12u_y u_{xy}\tilde{\zeta}_{xuu}^3 u_x - 6u_y u_{xz}\tilde{\zeta}_{xuu}^4 u_x \\
& - 6u_z u_{xy}\tilde{\zeta}_{xuu}^4 u_x - 9u_{xx}\tilde{\zeta}_{xyu}^2 u_x - 6u_{tx}\tilde{\zeta}_{xyu}^1 u_x - 6u_{xy}\tilde{\zeta}_{xyu}^3 u_x - 6u_{xz}\tilde{\zeta}_{xyu}^4 u_x \\
& - 6u_{xy}\tilde{\zeta}_{xxu}^2 u_x - 3u_{ty}\tilde{\zeta}_{xxu}^1 u_x - 3u_{yy}\tilde{\zeta}_{xxu}^3 u_x - 3u_{yz}\tilde{\zeta}_{xxu}^4 u_x + 3u_y\eta_{xuu} u_x \\
& - 3u_y u_t\tilde{\zeta}_{xxuu}^1 u_x - 3u_y^2\tilde{\zeta}_{xxuu}^3 u_x - 3u_z u_y\tilde{\zeta}_{xxuu}^4 u_x + 3\eta_{xxyu} u_x - 3u_t\tilde{\zeta}_{xxyu}^1 u_x \\
& - 3u_y\tilde{\zeta}_{xxyu}^3 u_x - 3u_z\tilde{\zeta}_{xxyu}^4 u_x - u_y\tilde{\zeta}_{xxxu}^2 u_x - \tilde{\zeta}_{xxxy}^2 u_x + u_{xxx}\eta_y u \\
& - 6u_{xx}u_{xxy}\tilde{\zeta}_u^2 u_x - 4u_{xy}u_{xxx}\tilde{\zeta}_u^2 u_x - u_{xxx}u_t\tilde{\zeta}_u^1 u_x - u_{xxx}u_{ty}\tilde{\zeta}_u^1 u_x - 3u_{xxy}u_{tx}\tilde{\zeta}_u^1 u_x \\
& - 3u_{xx}u_{txy}\tilde{\zeta}_u^1 u_x - 3u_{xy}u_{txx}\tilde{\zeta}_u^1 u_x - 3u_{xyy}u_{xx}\tilde{\zeta}_u^3 u_x - 6u_{xy}u_{xxy}\tilde{\zeta}_u^3 u_x - u_{yy}u_{xxx}\tilde{\zeta}_u^3 u_x \\
& - u_y u_{xxx}\tilde{\zeta}_u^3 u_x - 3u_{xyz}u_{xx}\tilde{\zeta}_u^4 u_x - 3u_{xy}u_{xxz}\tilde{\zeta}_u^4 u_x - 3u_{xz}u_{xxy}\tilde{\zeta}_u^4 u_x - u_{yz}u_{xxx}\tilde{\zeta}_u^4 u_x \\
& - u_z u_{xxy}\tilde{\zeta}_u^4 u_x + 3u_{xy}u_{xx}\eta_{uu} + u_y u_{xxx}\eta_{uu} - 3u_y u_{xx}\tilde{\zeta}_{uu}^2 u_x - 3u_{xy}u_{xx}u_t\tilde{\zeta}_{uu}^1 u_x \\
& - u_y u_{xxx}u_t\tilde{\zeta}_{uu}^1 u_x - 3u_y u_{xx}u_{tx}\tilde{\zeta}_{uu}^1 u_x - 6u_y u_{xy}u_{xx}\tilde{\zeta}_{uu}^3 u_x - u_y^2 u_{xxx}\tilde{\zeta}_{uu}^3 u_x \\
& - 3u_y u_{xz}u_{xx}\tilde{\zeta}_{uu}^4 u_x - 3u_z u_{xy}u_{xx}\tilde{\zeta}_{uu}^4 u_x - u_z u_y u_{xxx}\tilde{\zeta}_{uu}^4 u_x + u_{xxx}\eta_{yu} - 3u_{xx}^2\tilde{\zeta}_{yu}^2 u_x \\
& - u_{xxx}u_t\tilde{\zeta}_{yu}^1 u_x - 3u_{xx}u_{tx}\tilde{\zeta}_{yu}^1 u_x - 3u_{xy}u_{xx}\tilde{\zeta}_{yu}^3 u_x - u_y u_{xxx}\tilde{\zeta}_{yu}^3 u_x - 3u_{xz}u_{xx}\tilde{\zeta}_{yu}^4 u_x \\
& - u_z u_{xxx}\tilde{\zeta}_{yu}^4 u_x - 3u_{xxy}\tilde{\zeta}_x^2 u_x - u_{xxx}\tilde{\zeta}_x^2 u_x - 3u_{txxy}\tilde{\zeta}_x^1 u_x - u_{txxx}\tilde{\zeta}_x^1 u_x - 3u_{xxyy}\tilde{\zeta}_x^3 u_x \\
& - u_{xxyy}\tilde{\zeta}_x^3 u_x - 3u_{xxy}\tilde{\zeta}_x^4 u_x - u_{xxxz}\tilde{\zeta}_x^4 u_x + 3u_{xxy}\eta_{xu} - 9u_{xy}u_{xx}\tilde{\zeta}_{xu}^2 u_x - 3u_y u_{xxx}\tilde{\zeta}_{xu}^2 u_x \\
& - 3u_{xxy}u_t\tilde{\zeta}_{xu}^1 u_x - 3u_{xx}u_{ty}\tilde{\zeta}_{xu}^1 u_x - 6u_{xy}u_{tx}\tilde{\zeta}_{xu}^1 u_x - 3u_y u_{txx}\tilde{\zeta}_{xu}^1 u_x - 6u_{xy}^2\tilde{\zeta}_{xu}^3 u_x \\
& - 3u_{yy}u_{xx}\tilde{\zeta}_{xu}^3 u_x - 6u_y u_{xxy}\tilde{\zeta}_{xu}^3 u_x - 6u_{xz}u_{xy}\tilde{\zeta}_{xu}^4 u_x - 3u_{yz}u_{xx}\tilde{\zeta}_{xu}^4 u_x - 3u_y u_{xxz}\tilde{\zeta}_{xu}^4 u_x \\
& - 3u_z u_{xxy}\tilde{\zeta}_{xu}^4 u_x + 3u_y u_{xx}\eta_{xuu} - 3u_y u_{xx}u_t\tilde{\zeta}_{xuu}^1 u_x - 3u_y^2 u_{xx}\tilde{\zeta}_{xuu}^3 u_x - 3u_z u_y u_{xx}\tilde{\zeta}_{xuu}^4 u_x \\
& - 3u_{xxx}\tilde{\zeta}_{xy}^2 u_x - 3u_{txx}\tilde{\zeta}_{xy}^1 u_x - 3u_{xxy}\tilde{\zeta}_{xy}^3 u_x - 3u_{xxz}\tilde{\zeta}_{xy}^4 u_x + 3u_{xx}\eta_{xyu} - 3u_{xx}u_t\tilde{\zeta}_{xyu}^1 u_x \\
& - 3u_y u_{xx}\tilde{\zeta}_{xyu}^3 u_x - 3u_z u_{xx}\tilde{\zeta}_{xyu}^4 u_x - 3u_{xxy}\tilde{\zeta}_{xx}^2 u_x - 3u_{txy}\tilde{\zeta}_{xx}^1 u_x - 3u_{xyy}\tilde{\zeta}_{xx}^3 u_x - 3u_{xyz}\tilde{\zeta}_{xx}^4 u_x \\
& + 3u_{xy}\eta_{xuu} - 3u_y u_{xx}\tilde{\zeta}_{xxu}^2 u_x - 3u_{xy}u_t\tilde{\zeta}_{xxu}^1 u_x - 3u_y u_{tx}\tilde{\zeta}_{xxu}^1 u_x - 6u_y u_{xy}\tilde{\zeta}_{xxu}^3 u_x \\
& - 3u_y u_{xz}\tilde{\zeta}_{xxu}^4 u_x - 3u_z u_{xy}\tilde{\zeta}_{xxu}^4 u_x - 3u_{xx}\tilde{\zeta}_{xxy}^2 u_x - 3u_{tx}\tilde{\zeta}_{xxy}^1 u_x - 3u_{xy}\tilde{\zeta}_{xxy}^3 u_x \\
& - 3u_{xz}\tilde{\zeta}_{xxy}^4 u_x - u_{xy}\tilde{\zeta}_{xxx}^2 u_x - u_{ty}\tilde{\zeta}_{xxx}^1 u_x - u_{yy}\tilde{\zeta}_{xxx}^3 u_x - u_{yz}\tilde{\zeta}_{xxx}^4 u_x + u_y\eta_{xxxu} - u_y u_t\tilde{\zeta}_{xxxu}^1 u_x \\
& - u_y^2\tilde{\zeta}_{xxxu}^3 u_x - u_z u_y\tilde{\zeta}_{xxxu}^4 u_x + \eta_{xxy} - u_t\tilde{\zeta}_{xxy}^1 u_x - u_y\tilde{\zeta}_{xxy}^3 u_x - u_z\tilde{\zeta}_{xxy}^4 u_x.
\end{aligned}$$

References

1. Ovsiannikov, L.V. *Group Analysis of Differential Equations*; Academic Press: New York, NY, USA, 1982.
2. Ibragimov, N.H. *CRC Handbook of Lie Group Analysis of Differential Equations*; CRC Press: Boca Raton, FL, USA, 1994; Volume 1.
3. Stephani, H. *Differential Equations: Their Solution Using Symmetries*; Cambridge University Press: New York, NY, USA, 1989.
4. Olver, P.J. *Applications of Lie Groups to Differential Equations*, 2nd ed.; Springer: Berlin, Germany, 1993.
5. Hydon, P.E. *Symmetry Methods for Differential Equations*; Cambridge University Press: Cambridge, UK, 2000.
6. Zhou, Y.; Wang, M.; Li, Z. Application of a homogeneous balance method to exact solutions of nonlinear equations in mathematical physics. *Phys. Lett. A* **1996**, *216*, 67–75.

7. Hu, J.; Zhang, H. A new method for finding exact traveling wave solutions to nonlinear partial differential equations. *Phys. Lett. A* **2001**, *286*, 175–179.
8. Hirota, R. *The Direct Method in Soliton Theory*; Cambridge University Press: Cambridge, UK, 2004.
9. Wang, M.; Li, X.; Zhang, J. The (G'/G) –expansion method and travelling wave solutions for linear evolution equations in mathematical physics. *Phys. Lett. A* **2008**, *372*, 417–423.
10. Khalique, C.M.; Moleleki, L.D. A (3+1)-dimensional generalized BKP-Boussinesq equation: Lie group approach. *Results Phys.* **2019**, *13*, 102239.
11. Kudryashov, N.A. One method for finding exact solutions of nonlinear differential equations. *Commun. Nonlinear Sci. Numer. Simul.* **2012**, *17*, 2248–2253.
12. Motsepa, T.; Khalique, C.M. Closed-form solutions and conserved vectors of the (3+1)-dimensional negative-order KdV equation. *Adv. Math. Model. Appl.* **2020**, *5*, 7–18.
13. Kudryashov, N.A. Simplest equation method to look for exact solutions of nonlinear differential equations. *Chaos Soliton Fract.* **2005**, *24*, 1217–1231.
14. Simbanefayi, I.; Khalique, C.M. Travelling wave solutions and conservation laws for the Korteweg-de Vries-Benjamin-Bona-Mahony equation. *Results Phys.* **2018**, *8*, 57–63.
15. Zhou, Y.; Wang, M.; Wang, Y. Periodic wave solutions to a coupled KdV equations with variable coefficients. *Phys. Lett. A* **2003**, *308*, 31–36.
16. Zhang, L.; Khalique, C.M. Classification and bifurcation of a class of second-order ODEs and its application to nonlinear PDEs. *Discret. Contin. Dynam. Syst. S* **2018**, *11*, 777–790.
17. Ma, W.X. Comment on the (3+1)-dimensional Kadomtsev–Petviashvili equations. *Commun. Nonlinear Sci. Numer. Simulat.* **2011**, *16*, 2663–2666.
18. Kadomtsev, B.B.; Petviashvili, V.I. On the stability of solitary waves in weakly dispersing media. *Sov. Phys. Dokl.* **1970**, *192*, 753–756.
19. You, F.; Xia, T.; Chen, D. Decomposition of the generalized KP, cKP and mKP and their exact solutions. *Phys. Lett. A* **2008**, *372*, 3184–3194.
20. Kuznetsov, E.A.; Turitsyn, S.K. Two- and three-dimensional solitons in weakly dispersive media. *Zh. Ebp. Teor. Fa.* **1982**, *82*, 1457–1463.
21. Ablowitz, M.J.; Segur, H. On the evolution of packets of water waves. *J. Fluid Mech.* **1979**, *92*, 691–715.
22. Infeld, E.; Rowlands, G. Three-dimensional stability of Korteweg–de Vries waves and solitons II. *Acta Phys. Polon. A* **1979**, *56*, 329–332.
23. Senatorski, A.; Infeld, E. Simulations of two-dimensional Kadomtsev–Petviashvili soliton dynamics in three-dimensional space. *Phys. Rev. Lett.* **1996**, *77*, 2855–2858.
24. Alagesan, T.; Uthayakumar, A.; Porsezian, K. Painlevé analysis and Bäcklund transformation for a three-dimensional Kadomtsev–Petviashvili equation. *Chaos Soliton Fract.* **1997**, *8*, 893–895.
25. Xu, G.; Li, Z. Symbolic computation of the Painlevé test for nonlinear partial differential equations using Maple. *Comput. Phys. Commun.* **2004**, *161*, 65–75.
26. Ma, W.X.; Fan, E. Linear superposition principle applying to Hirota bilinear equations. *Comput. Math. Appl.* **2011**, *61*, 950–959.
27. Ma, W.X.; Abdeljabbar, A.; Asaad, M.G. Wronskian and Grammian solutions to a (3+1)-dimensional generalized KP equation. *Appl. Math. Comput.* **2011**, *217*, 10016–10023.
28. Wazwaz, A.M. Multiple-soliton solutions for a (3 + 1)-dimensional generalized KP equation, *Commun. Nonlinear Sci. Numer. Simulat.* **2012**, *17*, 491–495.
29. Wazwaz, A.M.; El-Tantawy, S.A. A new (3+1)-dimensional generalized Kadomtsev–Petviashvili equation. *Nonlinear Dyn.* **2016**, *84*, 1107–1112.
30. Liu, J.G.; Tian, Y.; Zeng, Z.F. New exact periodic solitary-wave solutions for the new (3+1)-dimensional generalized Kadomtsev–Petviashvili equation in multi-temperature electron plasmas. *AIP Adv.* **2017**, *7*, 2158–3226.
31. Noether, E. Invariante variationsprobleme. *Nachr. V. D. Ges. D. Wiss. Zu Göttingen* **1918**, *2*, 235–257.
32. Bessel-Hagen, E. Über die Erhaltungssätze der Elektrodynamik. *Math. Ann.* **1921**, *84*, 258–276.
33. Bluman, G.W.; Cheviakov, A.F.; Anco, S.C. *Applications of Symmetry Methods to Partial Differential Equations*. Springer: New York, NY, USA, 2010.
34. Leveque, R.J. *Numerical Methods for Conservation Laws*, 2nd ed.; Birkhäuser-Verlag: Basel, Switzerland, 1992.

35. Sarlet, W. Comment on 'Conservation laws of higher order nonlinear PDEs and the variational conservation laws in the class with mixed derivatives'. *J. Phys. A Math. Theor.* **2010**, *43*, 458001.
36. Anco, S.C. Generalization of Noether's Theorem in Modern Form to Non-variational Partial Differential Equations. In *Recent Progress and Modern Challenges in Applied Mathematics, Modeling and Computational Science*; Melnik, R., Makarov, R., Belair, J., Eds.; Fields Institute Communications, Springer: New York, NY, USA, 2017; Volume 79.
37. Johnpillai, A.G.; Khalique, C.M.; Mahomed, F.M. Travelling wave group-invariant solutions and conservation laws for θ -equation. *Malays. J. Math. Sci.* **2019**, *13*, 13–29.
38. Motsepa, T.; Abudiab, M.; Khalique, C.M. A Study of an extended generalized (2+1)-dimensional Jaulent-Miodek equation. *Int. J. Nonlin. Sci. Numer. Simul.* **2018**, *19*, 391–395.
39. Khalique, C.M.; Abdallah, S.A. Coupled Burgers equations governing polydisperse sedimentation: A Lie symmetry approach. *Results Phys.* **2020**, *16*, 102967.
40. Bruzón, M.S.; Gandarias, M.L. Traveling wave solutions of the K(m, n) equation with generalized evolution. *Math. Meth. Appl. Sci.* **2018**, *41*, 5851–5857.
41. Khalique, C.M.; Adeyemo, O.D. A study of (3+1)-dimensional generalized Korteweg-de Vries-Zakharov-Kuznetsov equation via Lie symmetry approach. *Results Phys.* **2020**, in press.
42. Ibragimov, N.H. Integrating factors, adjoint equations and Lagrangians. *J. Math. Anal. Appl.* **2006**, *318*, 742–757.
43. Ibragimov, N.H. A new conservation theorem. *J. Math. Anal. Appl.* **2007**, *333*, 311–328.
44. Cheviakov, A.F. Computation of fluxes of conservation laws. *J. Eng. Math.* **2010**, *66*, 153–173.
45. Wazwaz, A.M. Exact soliton and kink solutions for new (3+1)-dimensional nonlinear modified equations of wave propagation. *Open Eng.* **2017**, *7*, 169–174.
46. Baumann, G. *Symmetry Analysis of Differential Equations with Mathematica®*; Springer-Verlag: New York, NY, USA, 2000.
47. Gradshteyn, I.S.; Ryzhik, I.M. *Table of Integrals, Series, and Products*, 7th ed.; Academic Press: New York, NY, USA, 2007.
48. Billingham, J.; King, A.C. *Wave Motion*; Cambridge University Press: Cambridge, UK, 2000.
49. Korteweg, D.J.; de Vries, G. On the change of form of long waves advancing in a rectangular canal, and on a new type of long stationary waves. *Phil Mag.* **1895**, *39*, 422–443.
50. Drazin, P.G.; Johnson, R.S. *Soliton: An Introduction*; Cambridge University Press: Cambridge, UK, 1989.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).