

Article

A p -Ideal in BCI-Algebras Based on Multipolar Intuitionistic Fuzzy Sets

Jeong-Gon Lee ^{1,*} , Mohammad Fozouni ² , Kul Hur ³ and Young Bae Jun ⁴¹ Division of Applied Mathematics, Wonkwang University, 460, Iksan-daero, Iksan-Si, Jeonbuk 54538, Korea² Department of Mathematics, Faculty of Sciences and Engineering, Gonbad Kavous University, Gonbad Kavous P.O. 163, Iran; fozouni@hotmail.com³ Department of Applied Mathematics, Wonkwang University, 460, Iksan-daero, Iksan-Si, Jeonbuk 54538, Korea; kulhur@wku.ac.kr⁴ Department of Mathematics Education, Gyeongsang National University, Jinju 52828, Korea; skywine@gmail.com

* Correspondence: jukolee@wku.ac.kr

Received: 21 April 2020; Accepted: 25 May 2020; Published: 17 June 2020



Abstract: In 2020, Kang, Song and Jun introduced the notion of multipolar intuitionistic fuzzy set with finite degree, which is a generalization of intuitionistic fuzzy set, and they applied it to BCK/BCI-algebras. In this paper, we used this notion to study p -ideals of BCI-algebras. The notion of k -polar intuitionistic fuzzy p -ideals in BCI-algebras is introduced, and several properties were investigated. An example to illustrate the k -polar intuitionistic fuzzy p -ideal is given. The relationship between k -polar intuitionistic fuzzy ideal and k -polar intuitionistic fuzzy p -ideal is displayed. A k -polar intuitionistic fuzzy p -ideal is found to be k -polar intuitionistic fuzzy ideal, and an example to show that the converse is not true is provided. The notions of p -ideals and k -polar (\in, \in) -fuzzy p -ideal in BCI-algebras are used to study the characterization of k -polar intuitionistic p -ideal. The concept of normal k -polar intuitionistic fuzzy p -ideal is introduced, and its characterization is discussed. The process of eliciting normal k -polar intuitionistic fuzzy p -ideal using k -polar intuitionistic fuzzy p -ideal is provided.

Keywords: multipolar intuitionistic fuzzy set with finite degree k ; k -polar (\in, \in) -fuzzy ideal; k -polar intuitionistic fuzzy ideal; k -polar intuitionistic fuzzy p -ideal

MSC: 06F35; 03G25; 08A72

1. Introduction

BCI-algebras were introduced by Iséki [1] as the algebraic counterpart of the BCI-logic. BCI-algebras are a generalization of BCK-algebras, and they originated from two sources: set theory and propositional calculi. See the books [2,3] for more information on BCK/BCI-algebras. Fuzzy sets were first introduced by Zadeh [4], in which the membership degree is represented by only one function—the truth function. Intuitionistic fuzzy sets, which were introduced by Atanassov (see [5,6]), are a generalization of fuzzy sets. As an extension of the bipolar fuzzy set, Chen et al. [7] introduced an m -polar fuzzy set in 2014, and then this concept was applied to certain algebraic structures as BCK/BCI algebras, graph theory and decision making problem. For BCK/BCI-algebras, see [8–10], for graph theory, see [11–14] and see [15–18] for decision making problems. Al-Masarwah and Ahmad discussed the notion of m -polar fuzzy sets with applications in BCK/BCI-algebras. They introduced the notions of m -polar fuzzy subalgebras and m -polar fuzzy (closed, commutative) ideals and gave characterizations of m -polar fuzzy subalgebras and m -polar fuzzy (commutative) ideals. They considered relations

between m -polar fuzzy subalgebras, m -polar fuzzy ideals and m -polar fuzzy commutative ideals (see [8]). Using the notion of multipolar fuzzy point, Mohseni Takallo et al. [9] studied p -ideals of BCI-algebras. In [19], Kang et al. introduced the notion of multipolar intuitionistic fuzzy set with finite degree as a generalization of intuitionistic fuzzy set, and applied it to BCK/BCI-algebras. They introduced the concepts of a k -polar intuitionistic fuzzy subalgebra and a (closed) k -polar intuitionistic fuzzy ideal in a BCK/BCI-algebra, and investigated their relations and characterizations. In a BCI-algebra, they considered the relationship between a k -polar intuitionistic fuzzy ideal and a closed k -polar intuitionistic fuzzy ideal, and discussed the characterization of a closed k -polar intuitionistic fuzzy ideal. They consulted conditions for a k -polar intuitionistic fuzzy ideal to be a closed k -polar intuitionistic fuzzy ideal in a BCI-algebra. The aim of this manuscript was to use Kang et al.'s notion so called multipolar intuitionistic fuzzy set for studying p -ideal in BCI-algebras. This is a generalization of multipolar fuzzy p -ideals of BCI-algebras which is studied in [9]. We introduce the concept of k -polar intuitionistic fuzzy p -ideals in BCI-algebras, and then we study several properties. We first give an example to illustrate the k -polar intuitionistic fuzzy p -ideal. We consider the relationship between k -polar intuitionistic fuzzy ideal and k -polar intuitionistic fuzzy p -ideal. We first prove that every k -polar intuitionistic fuzzy p -ideal is a k -polar intuitionistic fuzzy ideal, and then give an example to show that the converse is not true in general. We use the notion of p -ideals in BCI-algebras to study the characterization of k -polar intuitionistic fuzzy p -ideal. We also use the notion of k -polar (\in, \in) -fuzzy p -ideal in BCI-algebras to study the characterization of k -polar intuitionistic fuzzy p -ideal. We define the concept of normal k -polar intuitionistic fuzzy p -ideal, and discuss its characterization. We look at the process of eliciting normal k -polar intuitionistic fuzzy p -ideal from a given k -polar intuitionistic fuzzy p -ideal.

2. Preliminaries

If a set U has a special element 0 and a binary operation $*$ satisfying the conditions:

- (I) $(\forall \omega, v, \tau \in U) (((\omega * v) * (\omega * \tau)) * (\tau * v) = 0)$,
- (II) $(\forall \omega, v \in U) ((\omega * (\omega * v)) * v = 0)$,
- (III) $(\forall \omega \in U) (\omega * \omega = 0)$,
- (IV) $(\forall \omega, v \in U) (\omega * v = 0, v * \omega = 0 \Rightarrow \omega = v)$,

then it is said that U is a BCI-algebra. If a BCI-algebra U satisfies the following identity:

- (V) $(\forall \omega \in U) (0 * \omega = 0)$,

then U is called a BCK-algebra.

Any BCK/BCI-algebra U satisfies the following conditions:

$$(\forall \omega \in U) (\omega * 0 = \omega), \tag{1}$$

$$(\forall \omega, v, \tau \in U) ((\omega * v) * \tau = (\omega * \tau) * v). \tag{2}$$

A subset I of a BCI-algebra U is called

- a subalgebra of U if $\omega * v \in I$ for all $\omega, v \in I$.
- an ideal of U if it satisfies:

$$0 \in I, \tag{3}$$

$$(\forall \omega \in U) (\forall v \in I) (\omega * v \in I \Rightarrow \omega \in I). \tag{4}$$

- a p -ideal of U (see [20]) if it satisfies Equation (3) and

$$(\forall \omega, v, \tau \in U) ((\omega * \tau) * (v * \tau) \in I, v \in I \Rightarrow \omega \in I). \tag{5}$$

Let $\{b_i \mid i \in \Gamma\}$ be a family of real numbers where Γ is any index set and we define

$$\bigvee \{b_i \mid i \in \Gamma\} := \begin{cases} \max\{b_i \mid i \in \Gamma\} & \text{if } \Gamma \text{ is finite,} \\ \sup\{b_i \mid i \in \Gamma\} & \text{otherwise.} \end{cases}$$

$$\bigwedge \{b_i \mid i \in \Gamma\} := \begin{cases} \min\{b_i \mid i \in \Gamma\} & \text{if } \Gamma \text{ is finite,} \\ \inf\{b_i \mid i \in \Gamma\} & \text{otherwise.} \end{cases}$$

If $\Gamma = \{1, 2\}$, we will also use $b_1 \vee b_2$ and $b_1 \wedge b_2$ instead of $\bigvee \{b_i \mid i \in \Gamma\}$ and $\bigwedge \{b_i \mid i \in \Gamma\}$, respectively.

Let k be a natural number and $[0, 1]^k$ denote the k -Cartesian product of $[0, 1]$, that is,

$$[0, 1]^k = [0, 1] \times [0, 1] \times \cdots \times [0, 1]$$

in which $[0, 1]$ is repeated k times. The order “ \leq ” in $[0, 1]^k$ is given by the pointwise order.

By a k -polar fuzzy set on a set U (see [7]), we mean a function $\widehat{\xi} : U \rightarrow [0, 1]^k$ where k is a natural number. The membership value of every element $z \in U$ is denoted by

$$\widehat{\xi}(z) = \left((\text{proj}_1 \circ \widehat{\xi})(z), (\text{proj}_2 \circ \widehat{\xi})(z), \dots, (\text{proj}_k \circ \widehat{\xi})(z) \right),$$

where $\text{proj}_i : [0, 1]^k \rightarrow [0, 1]$ is the i -th projection for all $i = 1, 2, \dots, k$ and \circ is the composition of functions.

A k -polar fuzzy set $\widehat{\xi}$ on a BCK/BCI-algebra U is called a k -polar fuzzy ideal of U (see [8]) if the following conditions are valid.

$$(\forall z \in U) \left(\widehat{\xi}(0) \geq \widehat{\xi}(z) \right), \tag{6}$$

$$(\forall z, x \in U) \left(\widehat{\xi}(z) \geq \widehat{\xi}(z * x) \wedge \widehat{\xi}(x) \right). \tag{7}$$

By a k -polar fuzzy point on a set U , we mean a k -polar fuzzy set $\widehat{\xi}$ on U of the form

$$\widehat{\xi}(x) = \begin{cases} \widehat{r} = (r_1, r_2, \dots, r_k) \in (0, 1]^k & \text{if } x = z, \\ \widehat{0} = (0, 0, \dots, 0) & \text{if } x \neq z, \end{cases} \tag{8}$$

and it is denoted by $z_{\widehat{r}}$ where z is a given element of U . We say that z is the support of $z_{\widehat{r}}$ and \widehat{r} is the value of $z_{\widehat{r}}$.

We say that a k -polar fuzzy point $z_{\widehat{r}}$ is contained in a k -polar fuzzy set $\widehat{\xi}$, denoted by $z_{\widehat{r}} \in \widehat{\xi}$, if $\widehat{\xi}(z) \geq \widehat{r}$, that is, $(\text{proj}_i \circ \widehat{\xi})(z) \geq r_i$ for all $i = 1, 2, \dots, k$.

A k -polar fuzzy set $\widehat{\xi}$ on a BCI-algebra U is called a k -polar (\in, \in) -fuzzy p -ideal of U (see [9]) if it satisfies

$$(\forall z \in U) (\forall \widehat{r} \in [0, 1]^k) \left(z_{\widehat{r}} \in \widehat{\xi} \Rightarrow 0_{\widehat{r}} \in \widehat{\xi} \right), \tag{9}$$

$$(\forall z, x, y \in U) (\forall \widehat{r}, \widehat{t} \in [0, 1]^k) \left(((z * y) * (x * y))_{\widehat{r}} \in \widehat{\xi}, x_{\widehat{t}} \in \widehat{\xi} \Rightarrow z_{\inf\{\widehat{r}, \widehat{t}\}} \in \widehat{\xi} \right). \tag{10}$$

It is easy to show that Condition (10) is equivalent to the following condition.

$$(\forall z, x, y \in U) \left(\widehat{\xi}(z) \geq \widehat{\xi}((z * y) * (x * y)) \wedge \widehat{\xi}(x) \right). \tag{11}$$

A multipolar intuitionistic fuzzy set with finite degree k (briefly, k -pIF set) over a set U (see [19]) is a mapping

$$(\widehat{\xi}, \widehat{\varrho}) : U \rightarrow [0, 1]^k \times [0, 1]^k, z \mapsto (\widehat{\xi}(z), \widehat{\varrho}(z)) \tag{12}$$

where $\widehat{\xi} : U \rightarrow [0, 1]^k$ and $\widehat{\varrho} : U \rightarrow [0, 1]^k$ are k -polar fuzzy sets over a set U such that $\widehat{\xi}(z) + \widehat{\varrho}(z) \leq \widehat{1}$ for all $z \in U$, that is, $(\text{proj}_i \circ \widehat{\xi})(z) + (\text{proj}_i \circ \widehat{\varrho})(z) \leq 1$ for all $z \in U$ and $i = 1, 2, \dots, k$. We know that if the multipolar intuitionistic fuzzy set has degree 1, then it is an intuitionistic fuzzy set. So, the intuitionistic fuzzy set is a special case of the multipolar intuitionistic fuzzy set. From this point of view, multipolar intuitionistic fuzzy set is a generalization of intuitionistic fuzzy set.

Given a k -pIF set $(\widehat{\xi}, \widehat{\varrho})$ over a set U , we consider the sets

$$U(\widehat{\xi}, \widehat{t}) := \{z \in U \mid \widehat{\xi}(z) \geq \widehat{t}\} \text{ and } L(\widehat{\varrho}, \widehat{s}) := \{z \in U \mid \widehat{\varrho}(z) \leq \widehat{s}\}, \tag{13}$$

where $\widehat{t} = (t_1, t_2, \dots, t_k) \in [0, 1]^k$ and $\widehat{s} = (s_1, s_2, \dots, s_k) \in [0, 1]^k$ with $\widehat{t} + \widehat{s} \leq \widehat{1}$, which is called a k -polar upper (resp., lower) level set of $(\widehat{\xi}, \widehat{\varrho})$ where "+" is the componentwise operation in $[0, 1]^k$, that is, $t_i + s_i \leq 1$ for all $i = 1, 2, \dots, k$. It is clear that $U(\widehat{\xi}, \widehat{t}) = \bigcap_{i=1}^k U(\widehat{\xi}, t_i)$ and $L(\widehat{\varrho}, \widehat{s}) = \bigcap_{i=1}^k L(\widehat{\varrho}, s_i)$ where

$$U(\widehat{\xi}, t_i) = \{z \in U \mid (\text{proj}_i \circ \widehat{\xi})(z) \geq t_i\} \text{ and } L(\widehat{\varrho}, s_i) = \{z \in U \mid (\text{proj}_i \circ \widehat{\varrho})(z) \leq s_i\}.$$

A k -pIF set $(\widehat{\xi}, \widehat{\varrho})$ over U is called a k -polar intuitionistic fuzzy ideal (briefly, k -pIF ideal) of U (see [19]) if it satisfies the conditions

$$(\forall z \in U)(\widehat{\xi}(0) \geq \widehat{\xi}(z), \widehat{\varrho}(0) \leq \widehat{\varrho}(z)), \tag{14}$$

that is, $(\text{proj}_i \circ \widehat{\xi})(0) \geq (\text{proj}_i \circ \widehat{\xi})(z)$ and $(\text{proj}_i \circ \widehat{\varrho})(0) \leq (\text{proj}_i \circ \widehat{\varrho})(z)$ for $i = 1, 2, \dots, k$. and

$$(\forall z, x \in U) \begin{pmatrix} \widehat{\xi}(z) \geq \widehat{\xi}(z * x) \wedge \widehat{\xi}(x) \\ \widehat{\varrho}(z) \leq \widehat{\varrho}(z * x) \vee \widehat{\varrho}(x) \end{pmatrix}. \tag{15}$$

3. k -Polar Intuitionistic Fuzzy p -Ideals

In this section, let U be a BCI-algebra unless otherwise stated.

Definition 1. A k -pIF set $(\widehat{\xi}, \widehat{\varrho})$ over U is called a k -polar intuitionistic fuzzy p -ideal (briefly, k -pIF p -ideal) of U if it satisfies Condition (14) and

$$(\forall z, x, y \in U) \begin{pmatrix} \widehat{\xi}(z) \geq \widehat{\xi}((z * x) * (y * x)) \wedge \widehat{\xi}(y) \\ \widehat{\varrho}(z) \leq \widehat{\varrho}((z * x) * (y * x)) \vee \widehat{\varrho}(y) \end{pmatrix}. \tag{16}$$

Example 1. Let $U = \{0, x, a, b\}$ be a set with a binary operation $*$ which is given in Table 1.

Table 1. Cayley table for the binary operation “*”.

| * | 0 | x | a | b |
|---|---|---|---|---|
| 0 | 0 | x | a | b |
| x | x | 0 | b | a |
| a | a | b | 0 | x |
| b | b | a | x | 0 |

Then, U is a BCI-algebra (see [2]). Let $(\widehat{\xi}, \widehat{\varrho})$ be a 4-polar intuitionistic fuzzy set over U given by

$$(\widehat{\xi}, \widehat{\varrho}) : U \rightarrow [0, 1]^4 \times [0, 1]^4,$$

$$z \mapsto \begin{cases} ((0.8, 0.67, 0.9, 0.56), (0.19, 0.15, 0.07, 0.28)) & \text{if } z = 0, \\ ((0.7, 0.57, 0.7, 0.56), (0.19, 0.24, 0.07, 0.35)) & \text{if } z = x, \\ ((0.5, 0.37, 0.4, 0.32), (0.37, 0.44, 0.39, 0.58)) & \text{if } z = a, \\ ((0.5, 0.37, 0.4, 0.32), (0.37, 0.44, 0.39, 0.58)) & \text{if } z = b. \end{cases}$$

It is routine to check that $(\widehat{\xi}, \widehat{\varrho})$ is a 4-polar intuitionistic fuzzy p -ideal of U .

Theorem 1. Let I be a subset of U and let $(\widehat{\xi}_I, \widehat{\varrho}_I)$ be a k -pIF set on U defined by

$$\widehat{\xi}_I : U \rightarrow [0, 1]^k, z \mapsto \begin{cases} \widehat{1} & \text{if } z \in I, \\ \widehat{0} & \text{otherwise} \end{cases}$$

$$\widehat{\varrho}_I : U \rightarrow [0, 1]^k, z \mapsto \begin{cases} \widehat{0} & \text{if } z \in I, \\ \widehat{1} & \text{otherwise} \end{cases}$$

Then, $(\widehat{\xi}_I, \widehat{\varrho}_I)$ is a k -pIF ideal p -ideal of U if and only if I is a p -ideal of U .

Proof. Straightforward. \square

In the following theorem, we look at the relationship between k -pIF ideal and k -pIF p -ideal.

Theorem 2. Every k -pIF p -ideal is a k -pIF ideal.

Proof. Let $(\widehat{\xi}, \widehat{\varrho})$ be a k -pIF p -ideal of U . If we put $x = 0$ in (16) and use (1), then

$$\begin{aligned} (\text{proj}_i \circ \widehat{\xi})(z) &\geq \min\{(\text{proj}_i \circ \widehat{\xi})((z * 0) * (x * 0)), (\text{proj}_i \circ \widehat{\xi})(x)\} \\ &= \min\{(\text{proj}_i \circ \widehat{\xi})(z * x), (\text{proj}_i \circ \widehat{\xi})(x)\} \end{aligned}$$

and

$$\begin{aligned} (\text{proj}_i \circ \widehat{\varrho})(z) &\leq \max\{(\text{proj}_i \circ \widehat{\varrho})((z * 0) * (x * 0)), (\text{proj}_i \circ \widehat{\varrho})(x)\} \\ &= \max\{(\text{proj}_i \circ \widehat{\varrho})(z * x), (\text{proj}_i \circ \widehat{\varrho})(x)\} \end{aligned}$$

for all $z, x \in U$. Therefore $(\widehat{\xi}, \widehat{\varrho})$ is a k -pIF ideal of U . \square

In the following example, we find that the converse of Theorem 2 is not true.

Example 2. Let $U = \{0, x, b, c, d\}$ be a set with a binary operation $*$, which is given in Table 2.

Table 2. Cayley table for the binary operation “ $*$ ”.

| $*$ | 0 | x | b | c | d |
|-----|---|---|---|---|---|
| 0 | 0 | 0 | d | c | b |
| x | x | 0 | d | c | b |
| b | b | b | 0 | d | c |
| c | c | c | b | 0 | d |
| d | d | d | c | b | 0 |

Then, U is a BCI-algebra (see [2]). Define a 3-polar intuitionistic fuzzy set $(\widehat{\xi}, \widehat{\varrho})$ on U as follows:

$$(\widehat{\xi}, \widehat{\varrho}) : U \rightarrow [0, 1]^3 \times [0, 1]^3,$$

$$z \mapsto \begin{cases} ((0.6, 0.7, 0.9), (0.2, 0.25, 0.07)) & \text{if } z = 0, \\ ((0.6, 0.5, 0.7), (0.3, 0.25, 0.17)) & \text{if } z = x, \\ ((0.2, 0.3, 0.4), (0.6, 0.45, 0.27)) & \text{if } z = b, \\ ((0.5, 0.4, 0.6), (0.4, 0.35, 0.37)) & \text{if } z = c, \\ ((0.2, 0.3, 0.4), (0.6, 0.45, 0.27)) & \text{if } z = d. \end{cases}$$

It is easy to confirm that $(\widehat{\xi}, \widehat{\varrho})$ is a 3-polar intuitionistic fuzzy ideal of U . But it is not a 3-polar intuitionistic fuzzy p -ideal of U since

$$(\text{proj}_2 \circ \widehat{\xi})(x) = 0.5 < 0.7 = \min\{(\text{proj}_2 \circ \widehat{\xi})((x * b) * (0 * b)), (\text{proj}_2 \circ \widehat{\xi})(0)\}$$

and/or

$$(\text{proj}_3 \circ \widehat{\varrho})(x) = 0.17 > 0.07 = \max\{(\text{proj}_3 \circ \widehat{\varrho})((x * b) * (0 * b)), (\text{proj}_3 \circ \widehat{\varrho})(0)\}.$$

Proposition 1. Every k -pIF p -ideal $(\widehat{\xi}, \widehat{\varrho})$ of U satisfies the following inequalities.

$$(\forall z \in U)(\widehat{\xi}(z) \geq \widehat{\xi}(0 * (0 * z)), \widehat{\varrho}(z) \leq \widehat{\varrho}(0 * (0 * z))). \tag{17}$$

Proof. If we change y to z and x to 0 in Equation (16), then

$$\begin{aligned} (\text{proj}_i \circ \widehat{\xi})(z) &\geq \min\{(\text{proj}_i \circ \widehat{\xi})((z * z) * (0 * z)), (\text{proj}_i \circ \widehat{\xi})(0)\} \\ &= \min\{(\text{proj}_i \circ \widehat{\xi})(0 * (0 * z)), (\text{proj}_i \circ \widehat{\xi})(0)\} \\ &= (\text{proj}_i \circ \widehat{\xi})(0 * (0 * z)) \end{aligned}$$

and

$$\begin{aligned} (\text{proj}_i \circ \widehat{\varrho})(z) &\leq \max\{(\text{proj}_i \circ \widehat{\varrho})((z * z) * (0 * z)), (\text{proj}_i \circ \widehat{\varrho})(0)\} \\ &= \max\{(\text{proj}_i \circ \widehat{\varrho})(0 * (0 * z)), (\text{proj}_i \circ \widehat{\varrho})(0)\} \\ &= (\text{proj}_i \circ \widehat{\varrho})(0 * (0 * z)) \end{aligned}$$

for all $z \in U$. \square

Proposition 2. Every k -pIF p -ideal $(\widehat{\xi}, \widehat{\varrho})$ of U satisfies the following inequalities.

$$(\forall z, x, y \in U) \left(\begin{array}{l} \widehat{\xi}(z * x) \leq \widehat{\xi}((z * y) * (x * y)) \\ \widehat{\varrho}(z * x) \geq \widehat{\varrho}((z * y) * (x * y)) \end{array} \right). \tag{18}$$

Proof. Let $(\widehat{\xi}, \widehat{\varrho})$ be a k -pIF p -ideal of U . Then, it is a k -pIF ideal of U by Theorem 2. For any $z, x, y \in U$, we have $((z * y) * (x * y)) * (z * x) = 0$. Hence

$$\begin{aligned} &(\text{proj}_i \circ \widehat{\xi})((z * y) * (x * y)) \\ &\geq \min\{(\text{proj}_i \circ \widehat{\xi})(((z * y) * (x * y)) * (z * x)), (\text{proj}_i \circ \widehat{\xi})(z * x)\} \\ &= \min\{(\text{proj}_i \circ \widehat{\xi})(0), (\text{proj}_i \circ \widehat{\xi})(z * x)\} = (\text{proj}_i \circ \widehat{\xi})(z * x) \end{aligned}$$

and

$$\begin{aligned} & (\text{proj}_i \circ \widehat{\varrho})((z * y) * (x * y)) \\ & \leq \max\{(\text{proj}_i \circ \widehat{\varrho})(((z * y) * (x * y)) * (z * x)), (\text{proj}_i \circ \widehat{\varrho})(z * x)\} \\ & = \max\{(\text{proj}_i \circ \widehat{\varrho})(0), (\text{proj}_i \circ \widehat{\varrho})(z * x)\} = (\text{proj}_i \circ \widehat{\varrho})(z * x) \end{aligned}$$

for all $z, x, y \in U$. \square

We provide conditions for a k -pIF ideal to be a k -pIF p -ideal.

Theorem 3. Let $(\widehat{\xi}, \widehat{\varrho})$ be a k -pIF ideal of U satisfying the condition

$$(\forall z, x, y \in U) \left(\begin{array}{l} \widehat{\xi}(z * x) \geq \widehat{\xi}((z * y) * (x * y)) \\ \widehat{\varrho}(z * x) \leq \widehat{\varrho}((z * y) * (x * y)) \end{array} \right). \tag{19}$$

Then, it is a k -pIF p -ideal of U .

Proof. Using Equations (15) and (19), we have that

$$\widehat{\xi}(z) \geq \widehat{\xi}(z * x) \wedge \widehat{\xi}(x) \geq \widehat{\xi}((z * y) * (x * y)) \wedge \widehat{\xi}(x)$$

and

$$\widehat{\varrho}(z) \leq \widehat{\varrho}(z * x) \vee \widehat{\varrho}(x) \leq \widehat{\varrho}((z * y) * (x * y)) \vee \widehat{\varrho}(x)$$

for all $z, x, y \in U$. Therefore $(\widehat{\xi}, \widehat{\varrho})$ is a k -pIF p -ideal of U . \square

Lemma 1. Every k -pIF ideal $(\widehat{\xi}, \widehat{\varrho})$ of U satisfies the following inequalities.

$$(\forall z \in U)(\widehat{\xi}(z) \leq \widehat{\xi}(0 * (0 * z)), \widehat{\varrho}(z) \geq \widehat{\varrho}(0 * (0 * z))). \tag{20}$$

Proof. For any $z, x \in U$, we obtain

$$\widehat{\xi}(0 * (0 * z)) \geq \widehat{\xi}((0 * (0 * z)) * z) \wedge \widehat{\xi}(z) = \widehat{\xi}((0 * z) * (0 * z)) \wedge \widehat{\xi}(z) = \widehat{\xi}(0) \wedge \widehat{\xi}(z) = \widehat{\xi}(z)$$

and

$$\widehat{\varrho}(0 * (0 * z)) \leq \widehat{\varrho}((0 * (0 * z)) * z) \vee \widehat{\varrho}(z) = \widehat{\varrho}((0 * z) * (0 * z)) \vee \widehat{\varrho}(z) = \widehat{\varrho}(0) \vee \widehat{\varrho}(z) = \widehat{\varrho}(z)$$

by Equations (2), (3), (14) and (15). \square

Theorem 4. Let $(\widehat{\xi}, \widehat{\varrho})$ be a k -pIF set over U . If $(\widehat{\xi}, \widehat{\varrho})$ satisfies the following inequalities

$$(\forall z \in U)(\widehat{\xi}(z) \geq \widehat{\xi}(0 * (0 * z)), \widehat{\varrho}(z) \leq \widehat{\varrho}(0 * (0 * z))). \tag{21}$$

Proof. For any $z, x, y \in U$ and $i = 1, 2, \dots, k$, we have

$$\begin{aligned} (\text{proj}_i \circ \widehat{\xi})((z * y) * (x * y)) & \leq (\text{proj}_i \circ \widehat{\xi})(0 * (0 * (z * y) * (x * y))) \\ & = (\text{proj}_i \circ \widehat{\xi})((0 * x) * (0 * y)) \\ & = (\text{proj}_i \circ \widehat{\xi})(0 * (0 * (z * y))) \\ & \leq (\text{proj}_i \circ \widehat{\xi})(z * x), \end{aligned}$$

and

$$\begin{aligned} (\text{proj}_i \circ \widehat{q})((z * y) * (x * y)) &\geq (\text{proj}_i \circ \widehat{q})(0 * (0 * (z * y) * (x * y))) \\ &= (\text{proj}_i \circ \widehat{q})((0 * x) * (0 * y)) \\ &= (\text{proj}_i \circ \widehat{q})(0 * (0 * (z * y))) \\ &\geq (\text{proj}_i \circ \widehat{q})(z * x), \end{aligned}$$

which imply that $\widehat{\xi}((z * y) * (x * y)) \leq \widehat{\xi}(z * x)$ and $\widehat{q}((z * y) * (x * y)) \geq \widehat{q}(z * x)$ for all $z, x, y \in U$. Therefore $(\widehat{\xi}, \widehat{q})$ is a k -pIF p -ideal of U by Theorem 3. \square

We consider characterizations of a k -pIF p -ideal.

Theorem 5. Given a k -pIF set $(\widehat{\xi}, \widehat{q})$ over U , the following assertions are equivalent.

- (i) $(\widehat{\xi}, \widehat{q})$ is a k -pIF p -ideal of U .
- (ii) The k -polar upper and lower level sets $U(\widehat{\xi}, \widehat{r})$ and $L(\widehat{q}, \widehat{q})$ are p -ideals of U for all $(\widehat{r}, \widehat{q}) \in [0, 1]^k \times [0, 1]^k$ with $U(\widehat{\xi}, \widehat{r}) \neq \emptyset \neq L(\widehat{q}, \widehat{q})$.

Proof. Assume that $(\widehat{\xi}, \widehat{q})$ is a k -pIF p -ideal of U . It is clear that $0 \in U(\widehat{\xi}; \widehat{r})$ and $0 \in L(\widehat{q}; \widehat{q})$ for any $\widehat{r} = (r_1, r_2, \dots, r_k) \in (0, 1]^k$ and $\widehat{q} = (q_1, q_2, \dots, q_k) \in (0, 1]^k$. Let $z, x, y, b, c, d \in U$ be such that $(z * y) * (x * y) \in U(\widehat{\xi}; \widehat{r})$, $x \in U(\widehat{\xi}; \widehat{r})$, $(b * d) * (c * d) \in L(\widehat{q}; \widehat{q})$ and $c \in L(\widehat{q}; \widehat{q})$. Then, $(\text{proj}_i \circ \widehat{\xi})((z * y) * (x * y)) \geq r_i$, $(\text{proj}_i \circ \widehat{\xi})(x) \geq r_i$, $(\text{proj}_i \circ \widehat{q})((b * d) * (c * d)) \leq q_i$ and $(\text{proj}_i \circ \widehat{q})(c) \leq q_i$. It follows from Equations (16) that

$$(\text{proj}_i \circ \widehat{\xi})(z) \geq \min\{(\text{proj}_i \circ \widehat{\xi})((z * y) * (x * y)), (\text{proj}_i \circ \widehat{\xi})(x)\} \geq r_i$$

and

$$(\text{proj}_i \circ \widehat{q})(b) \leq \max\{(\text{proj}_i \circ \widehat{q})((b * d) * (c * d)), (\text{proj}_i \circ \widehat{q})(c)\} \leq q_i$$

for $i = 1, 2, \dots, k$. Hence $z \in U(\widehat{\xi}; \widehat{r})$ and $b \in L(\widehat{q}; \widehat{q})$ and therefore $U(\widehat{\xi}; \widehat{r})$ and $L(\widehat{q}; \widehat{q})$ are p -ideals of U .

Conversely, suppose that the k -polar upper and lower level sets $U(\widehat{\xi}, \widehat{r})$ and $L(\widehat{q}, \widehat{q})$ are p -ideals of U for all $(\widehat{r}, \widehat{q}) \in [0, 1]^k \times [0, 1]^k$ with $U(\widehat{\xi}, \widehat{r}) \neq \emptyset \neq L(\widehat{q}, \widehat{q})$. If $\widehat{\xi}(0) < \widehat{\xi}(b)$ for some $b \in U$, then $b \in U(\widehat{\xi}; \widehat{r})$ and $0 \notin U(\widehat{\xi}; \widehat{r})$ where $\widehat{r} := \widehat{\xi}(b)$. This is a contradiction, and so $\widehat{\xi}(0) \geq \widehat{\xi}(z)$ for all $z \in U$. If $\widehat{q}(0) > \widehat{q}(c)$ for some $c \in U$, then $(\text{proj}_i \circ \widehat{q})(0) > (\text{proj}_i \circ \widehat{q})(c)$ for $i = 1, 2, \dots, k$. If we take $q_i := (\text{proj}_i \circ \widehat{q})(c)$ for $i = 1, 2, \dots, k$, then $c \in L(\widehat{q}, \widehat{q})^i$ and $0 \notin L(\widehat{q}, \widehat{q})^i$ for $i = 1, 2, \dots, k$. Thus $c \in \bigcap_{i=1}^k L(\widehat{q}, \widehat{q})^i = L(\widehat{q}, \widehat{q})$ and $0 \notin L(\widehat{q}, \widehat{q})$, which is a contradiction; hence $\widehat{q}(0) \leq \widehat{q}(z)$ for all $z \in U$. Now, suppose that there exist $b, c, d \in U$ such that $\widehat{\xi}(b) < \widehat{\xi}((b * d) * (c * d)) \wedge \widehat{\xi}(c) > \widehat{q}(b) > \widehat{q}((b * d) * (c * d)) \vee \widehat{q}(c)$. If we take

$$\widehat{r} := \widehat{\xi}((b * d) * (c * d)) \wedge \widehat{\xi}(c)$$

and

$$\widehat{q} := \widehat{q}((b * d) * (c * d)) \vee \widehat{q}(c),$$

then

$$(b * d) * (c * d) \in U(\widehat{\xi}; \widehat{r}) \text{ and } c \in U(\widehat{\xi}; \widehat{r})$$

or

$$(b * d) * (c * d) \in L(\widehat{q}, \widehat{q}) \text{ and } c \in L(\widehat{q}, \widehat{q}).$$

Since $U(\widehat{\xi}; \widehat{r})$ and $L(\widehat{\varrho}; \widehat{q})$ are p -ideals of U by assumption, it follows that $b \in U(\widehat{\xi}; \widehat{r})$ or $b \in L(\widehat{\varrho}; \widehat{q})$. Hence $\widehat{\xi}(b) \geq \widehat{r} = \widehat{\xi}((b * d) * (c * d)) \wedge \widehat{\xi}(c)$ or $\widehat{\varrho}(b) \leq \widehat{q} = \widehat{\varrho}((b * d) * (c * d)) \vee \widehat{\varrho}(c)$, which is a contradiction. Thus $\widehat{\xi}(z) \geq \widehat{\xi}((z * y) * (x * y)) \wedge \widehat{\xi}(x)$ and $\widehat{\varrho}(z) \leq \widehat{\varrho}((z * y) * (x * y)) \vee \widehat{\varrho}(x)$ for all $z, x, y \in U$; therefore $(\widehat{\xi}, \widehat{\varrho})$ is a k -pIF p -ideal of U . \square

Given a k -pIF set $(\widehat{\xi}, \widehat{\varrho})$ over U and $(\widehat{t}, \widehat{s}) \in (0, 1]^k \times [0, 1)^k$, we consider the sets:

$$R_{(\widehat{\xi}, \widehat{t})}(U) := \{z \in U \mid \widehat{\xi}(z) + \widehat{t} > \widehat{1}\}$$

and

$$R_{(\widehat{\varrho}, \widehat{s})}(U) := \{z \in U \mid \widehat{\varrho}(z) + \widehat{s} < \widehat{1}\}.$$

Then, $R_{(\widehat{\xi}, \widehat{t})}(U) = \bigcap_{i=1}^k R_{(\widehat{\xi}, \widehat{t})}(U)^i$ and $R_{(\widehat{\varrho}, \widehat{s})}(U) = \bigcap_{i=1}^k R_{(\widehat{\varrho}, \widehat{s})}(U)^i$ where

$$R_{(\widehat{\xi}, \widehat{t})}(U)^i := \{z \in U \mid (\text{proj}_i \circ \widehat{\xi})(z) + t_i > 1\}$$

and

$$R_{(\widehat{\varrho}, \widehat{s})}(U)^i := \{z \in U \mid (\text{proj}_i \circ \widehat{\varrho})(z) + s_i < 1\}$$

for $i = 1, 2, \dots, k$.

Theorem 6. Given a k -pIF set $(\widehat{\xi}, \widehat{\varrho})$ over U , the following assertions are equivalent.

- (i) $(\widehat{\xi}, \widehat{\varrho})$ is a k -pIF p -ideal of U .
- (ii) The sets $R_{(\widehat{\xi}, \widehat{t})}(U)$ and $R_{(\widehat{\varrho}, \widehat{s})}(U)$ are p -ideals of U for all $(\widehat{t}, \widehat{s}) \in (0, 1]^k \times [0, 1)^k$ with $R_{(\widehat{\xi}, \widehat{t})}(U) \neq \emptyset \neq R_{(\widehat{\varrho}, \widehat{s})}(U)$.

Proof. Assume that $(\widehat{\xi}, \widehat{\varrho})$ is a k -pIF p -ideal of U . It is clear that $0 \in R_{(\widehat{\xi}, \widehat{t})}(U)$ and $0 \in R_{(\widehat{\varrho}, \widehat{s})}(U)$. Let $z, x, y, b, c, d \in U$ be such that $(z * y) * (x * y) \in R_{(\widehat{\xi}, \widehat{t})}(U)$, $x \in R_{(\widehat{\xi}, \widehat{t})}(U)$, $(b * d) * (c * d) \in R_{(\widehat{\varrho}, \widehat{s})}(U)$ and $c \in R_{(\widehat{\varrho}, \widehat{s})}(U)$. Then, $\widehat{\xi}((z * y) * (x * y)) + \widehat{t} > \widehat{1}$, $\widehat{\xi}(x) + \widehat{t} > \widehat{1}$, $\widehat{\varrho}((b * d) * (c * d)) + \widehat{s} < \widehat{1}$ and $\widehat{\varrho}(c) + \widehat{s} < \widehat{1}$. It follows that

$$\begin{aligned} (\text{proj}_i \circ \widehat{\xi})(z) + t_i &\geq \min\{(\text{proj}_i \circ \widehat{\xi})((z * y) * (x * y)), (\text{proj}_i \circ \widehat{\xi})(x)\} + t_i \\ &= \min\{(\text{proj}_i \circ \widehat{\xi})((z * y) * (x * y)) + t_i, (\text{proj}_i \circ \widehat{\xi})(x) + t_i\} > 1 \end{aligned}$$

and

$$\begin{aligned} (\text{proj}_i \circ \widehat{\varrho})(b) + s_i &\leq \max\{(\text{proj}_i \circ \widehat{\varrho})((b * d) * (c * d)), (\text{proj}_i \circ \widehat{\varrho})(c)\} + s_i \\ &= \max\{(\text{proj}_i \circ \widehat{\varrho})((b * d) * (c * d)) + s_i, (\text{proj}_i \circ \widehat{\varrho})(c) + s_i\} < 1 \end{aligned}$$

for all $i = 1, 2, \dots, k$. Hence $z \in \bigcap_{i=1}^k R_{(\widehat{\xi}, \widehat{t})}(U)^i = R_{(\widehat{\xi}, \widehat{t})}(U)$ and $b \in \bigcap_{i=1}^k R_{(\widehat{\varrho}, \widehat{s})}(U)^i = R_{(\widehat{\varrho}, \widehat{s})}(U)$; therefore $R_{(\widehat{\xi}, \widehat{t})}(U)$ and $R_{(\widehat{\varrho}, \widehat{s})}(U)$ are p -ideals of U for all $(\widehat{t}, \widehat{s}) \in (0, 1]^k \times [0, 1)^k$.

Conversely suppose that (ii) is valid. If $\widehat{\xi}(0) < \widehat{\xi}(z)$ or $\widehat{\varrho}(0) > \widehat{\varrho}(b)$ for some $z, b \in U$, then $\widehat{\xi}(0) + \widehat{t} \leq \widehat{1} < \widehat{\xi}(z) + \widehat{t}$ or $\widehat{\varrho}(0) + \widehat{s} \geq \widehat{1} > \widehat{\varrho}(b) + \widehat{s}$ for some $(\widehat{t}, \widehat{s}) \in (0, 1]^k \times [0, 1)^k$. Thus $0 \notin R_{(\widehat{\xi}, \widehat{t})}(U)$ or $0 \notin R_{(\widehat{\varrho}, \widehat{s})}(U)$ which is a contradiction. Hence $(\widehat{\xi}, \widehat{\varrho})$ satisfies Condition (14). Suppose that $\widehat{\xi}(b) < \widehat{\xi}((b * d) * (c * d)) \wedge \widehat{\xi}(c)$ for some $b, c \in U$. Then, $\widehat{\xi}(b) + \widehat{t} \leq \widehat{1} < (\widehat{\xi}((b * d) * (c * d)) \wedge \widehat{\xi}(c)) + \widehat{t} = (\widehat{\xi}((b * d) * (c * d)) + \widehat{t}) \wedge (\widehat{\xi}(c) + \widehat{t})$ for some $\widehat{t} \in (0, 1]^k$. It follows that $(b * d) * (c * d) \in R_{(\widehat{\xi}, \widehat{t})}(U)$ and

$c \in R_{(\widehat{\xi}, \widehat{t})}(U)$, which implies that $b \in R_{(\widehat{\xi}, \widehat{t})}(U)$ since $R_{(\widehat{\xi}, \widehat{t})}(U)$ is a p -ideal of U ; hence $\widehat{\xi}(b) + \widehat{t} > \widehat{1}$, which is a contradiction. If $\widehat{\varrho}(z) > \widehat{\varrho}((z * y) * (x * y)) \vee \widehat{\varrho}(x)$ for some $z, x \in U$, then

$$\widehat{\varrho}(z) + \widehat{s} \geq \widehat{1} > (\widehat{\xi}((z * y) * (x * y)) \vee \widehat{\xi}(x)) + \widehat{s} = (\widehat{\xi}((z * y) * (x * y)) + \widehat{s}) \vee (\widehat{\xi}(x) + \widehat{s})$$

for some $\widehat{s} \in [0, 1)^k$. Thus $(z * y) * (x * y) \in R_{(\widehat{\varrho}, \widehat{s})}(U)$ and $x \in R_{(\widehat{\varrho}, \widehat{s})}(U)$. Since $R_{(\widehat{\varrho}, \widehat{s})}(U)$ is a p -ideal of U , it follows that $z \in R_{(\widehat{\varrho}, \widehat{s})}(U)$, that is, $\widehat{\varrho}(z) + \widehat{s} < \widehat{1}$. This is a contradiction. This shows that $(\widehat{\xi}, \widehat{\varrho})$ satisfies Condition (16); therefore $(\widehat{\xi}, \widehat{\varrho})$ is a k -pIF p -ideal of U . \square

The following theorem shows the characterization of k -pIF p -ideal using k -polar (\in, \in) -fuzzy p -ideal.

Theorem 7. A k -pIF set $(\widehat{\xi}, \widehat{\varrho})$ over U is a k -pIF p -ideal of U if and only if $\widehat{\xi}$ and $\widehat{\varrho}^c$ are k -polar (\in, \in) -fuzzy p -ideals of U where $\widehat{\varrho}^c = 1 - \widehat{\varrho}$, i.e., $(\text{proj}_i \circ \widehat{\varrho})^c = 1 - (\text{proj}_i \circ \widehat{\varrho})$ for $i = 1, 2, \dots, k$.

Proof. Let $(\widehat{\xi}, \widehat{\varrho})$ be a k -pIF p -ideal of U . It is clear that $\widehat{\xi}$ is a k -polar (\in, \in) -fuzzy p -ideal of U . Let $z, x, y \in U$. Then,

$$(\text{proj}_i \circ \widehat{\varrho})^c(0) = 1 - (\text{proj}_i \circ \widehat{\varrho})(0) \geq 1 - (\text{proj}_i \circ \widehat{\varrho})(z) = (\text{proj}_i \circ \widehat{\varrho})^c(z)$$

and

$$\begin{aligned} (\text{proj}_i \circ \widehat{\varrho})^c(z) &= 1 - (\text{proj}_i \circ \widehat{\varrho})(z) \geq 1 - \max\{(\text{proj}_i \circ \widehat{\varrho})((z * y) * (x * y)), (\text{proj}_i \circ \widehat{\varrho})(x)\} \\ &= \min\{1 - (\text{proj}_i \circ \widehat{\varrho})((z * y) * (x * y)), 1 - (\text{proj}_i \circ \widehat{\varrho})(x)\} \\ &= \min\{(\text{proj}_i \circ \widehat{\varrho})^c((z * y) * (x * y)), (\text{proj}_i \circ \widehat{\varrho})^c(x)\}. \end{aligned}$$

Thus $\widehat{\varrho}^c$ is a k -polar (\in, \in) -fuzzy p -ideal of U .

Conversely, suppose that $\widehat{\xi}$ and $\widehat{\varrho}^c$ are k -polar (\in, \in) -fuzzy p -ideals of U . For any $z, x \in U$, we have $(\text{proj}_i \circ \widehat{\xi})(0) \geq (\text{proj}_i \circ \widehat{\xi})(z)$, $(\text{proj}_i \circ \widehat{\xi})(z) \geq \min\{(\text{proj}_i \circ \widehat{\xi})((z * y) * (x * y)), (\text{proj}_i \circ \widehat{\xi})(x)\}$, $1 - (\text{proj}_i \circ \widehat{\varrho})(0) = (\text{proj}_i \circ \widehat{\varrho})^c(0) \geq (\text{proj}_i \circ \widehat{\varrho})^c(z) = 1 - (\text{proj}_i \circ \widehat{\varrho})(z)$, i.e., $(\text{proj}_i \circ \widehat{\varrho})(0) \leq (\text{proj}_i \circ \widehat{\varrho})(z)$ and

$$\begin{aligned} 1 - (\text{proj}_i \circ \widehat{\varrho})(z) &= (\text{proj}_i \circ \widehat{\varrho})^c(z) \geq \min\{(\text{proj}_i \circ \widehat{\varrho})^c((z * y) * (x * y)), (\text{proj}_i \circ \widehat{\varrho})^c(x)\} \\ &= \min\{1 - (\text{proj}_i \circ \widehat{\varrho})((z * y) * (x * y)), 1 - (\text{proj}_i \circ \widehat{\varrho})(x)\} \\ &= 1 - \max\{(\text{proj}_i \circ \widehat{\varrho})((z * y) * (x * y)), (\text{proj}_i \circ \widehat{\varrho})(x)\}, \end{aligned}$$

that is, $(\text{proj}_i \circ \widehat{\varrho})(z) \leq \max\{(\text{proj}_i \circ \widehat{\varrho})((z * y) * (x * y)), (\text{proj}_i \circ \widehat{\varrho})(x)\}$; therefore $(\widehat{\xi}, \widehat{\varrho})$ is a k -pIF p -ideal of U . \square

The following corollary is an immediate consequence of Theorem 7.

Corollary 1. Let $(\widehat{\xi}, \widehat{\varrho})$ be a k -pIF set over U . Then, $(\widehat{\xi}, \widehat{\varrho})$ is a k -pIF p -ideal of U if and only if the necessary operator $\square(\widehat{\xi}, \widehat{\varrho}) = (\widehat{\xi}, \widehat{\xi}^c)$ and the possibility operator $\diamond(\widehat{\xi}, \widehat{\varrho}) = (\widehat{\varrho}^c, \widehat{\varrho})$ of $(\widehat{\xi}, \widehat{\varrho})$ are k -pIF p -ideals of U .

Definition 2. A k -pIF p -ideal $(\widehat{\xi}, \widehat{\varrho})$ of U is said to be normal if there exists $z, x \in U$ such that $\widehat{\xi}(z) = \widehat{1}$ and $\widehat{\varrho}(x) = \widehat{0}$.

Example 3. Consider the BCI-algebra $U = \{0, x, a, b\}$, which is given in Example 1. Let $(\widehat{\xi}, \widehat{\varrho})$ be a 3-polar intuitionistic fuzzy set over U given by

$$(\widehat{\xi}, \widehat{\varrho}) : U \rightarrow [0, 1]^3 \times [0, 1]^3,$$

$$z \mapsto \begin{cases} ((1.00, 1.00, 1.00), (0.00, 0.00, 0.00)) & \text{if } z = 0, \\ ((0.72, 0.57, 1.00), (0.00, 0.24, 0.35)) & \text{if } z = x, \\ ((0.52, 0.37, 0.32), (0.37, 0.44, 0.58)) & \text{if } z = a, \\ ((0.52, 0.37, 0.32), (0.37, 0.44, 0.58)) & \text{if } z = b. \end{cases}$$

It is routine to check that $(\widehat{\xi}, \widehat{\varrho})$ is a normal 3-polar intuitionistic fuzzy p -ideal of U .

It is clear that if a k -pIF p -ideal $(\widehat{\xi}, \widehat{\varrho})$ of U is normal, then $\widehat{\xi}(0) = \hat{1}$ and $\widehat{\varrho}(0) = \hat{0}$, that is, $(\text{proj}_i \circ \widehat{\xi})(0) = 1$ and $(\text{proj}_i \circ \widehat{\varrho})(0) = 0$ for all $i = 1, 2, \dots, k$.

Lemma 2. A k -pIF p -ideal $(\widehat{\xi}, \widehat{\varrho})$ of U is normal if and only if $\widehat{\xi}(0) = \hat{1}$ and $\widehat{\varrho}(0) = \hat{0}$.

Proof. Straightforward. \square

In the following theorem we look at the process of eliciting normal k -pIF p -ideal from a given k -pIF p -ideal.

Theorem 8. If $(\widehat{\xi}, \widehat{\varrho})$ is k -pIF p -ideal of U , then the k -pIF set $(\widehat{\xi}, \widehat{\varrho})^+ = (\widehat{\xi}^+, \widehat{\varrho}^+)$ on U defined by

$$\begin{aligned} \widehat{\xi}^+ : U &\rightarrow [0, 1]^k, z \mapsto \hat{1} + \widehat{\xi}(z) - \widehat{\xi}(0), \\ \widehat{\varrho}^+ : U &\rightarrow [0, 1]^k, z \mapsto \widehat{\varrho}(z) - \widehat{\varrho}(0) \end{aligned} \tag{22}$$

is a normal k -pIF p -ideal of U containing $(\widehat{\xi}, \widehat{\varrho})$.

Proof. Assume that $(\widehat{\xi}, \widehat{\varrho})$ is a k -pIF p -ideal of U . Then, $(\widehat{\xi}, \widehat{\varrho})$ is a k -pIF ideal of U by Theorem 2. For any $z, x \in U$, we have

$$\begin{aligned} (\text{proj}_i \circ \widehat{\xi})(0) &= 1 + (\text{proj}_i \circ \widehat{\xi})(0) - (\text{proj}_i \circ \widehat{\xi})(0) = 1 \geq (\text{proj}_i \circ \widehat{\xi})(z), \\ (\text{proj}_i \circ \widehat{\varrho})(0) &= (\text{proj}_i \circ \widehat{\varrho})(0) - (\text{proj}_i \circ \widehat{\varrho})(0) = 0 \leq (\text{proj}_i \circ \widehat{\varrho})(z), \end{aligned}$$

$$\begin{aligned} (\text{proj}_i \circ \widehat{\xi})^+(z) &= 1 + (\text{proj}_i \circ \widehat{\xi})(z) - (\text{proj}_i \circ \widehat{\xi})(0) \\ &\geq 1 + \min\{(\text{proj}_i \circ \widehat{\xi})((z * y) * (x * y)), (\text{proj}_i \circ \widehat{\xi})(x)\} - (\text{proj}_i \circ \widehat{\xi})(0) \\ &= \min\{1 + (\text{proj}_i \circ \widehat{\xi})((z * y) * (x * y)) - (\text{proj}_i \circ \widehat{\xi})(0), 1 + (\text{proj}_i \circ \widehat{\xi})(x) - (\text{proj}_i \circ \widehat{\xi})(0)\} \\ &= \min\{(\text{proj}_i \circ \widehat{\xi})^+((z * y) * (x * y)), (\text{proj}_i \circ \widehat{\xi})^+(x)\} \end{aligned}$$

and

$$\begin{aligned} (\text{proj}_i \circ \widehat{\varrho})^+(z) &= (\text{proj}_i \circ \widehat{\varrho})(z) - (\text{proj}_i \circ \widehat{\varrho})(0) \\ &\leq \max\{(\text{proj}_i \circ \widehat{\varrho})((z * y) * (x * y)), (\text{proj}_i \circ \widehat{\varrho})(x)\} - (\text{proj}_i \circ \widehat{\varrho})(0) \\ &= \max\{(\text{proj}_i \circ \widehat{\varrho})((z * y) * (x * y)) - (\text{proj}_i \circ \widehat{\varrho})(0), (\text{proj}_i \circ \widehat{\varrho})(x) - (\text{proj}_i \circ \widehat{\varrho})(0)\} \\ &= \max\{(\text{proj}_i \circ \widehat{\varrho})^+((z * y) * (x * y)), (\text{proj}_i \circ \widehat{\varrho})^+(x)\} \end{aligned}$$

for all for $i = 1, 2, \dots, k$. Hence $(\widehat{\xi}, \widehat{\varrho})^+$ is a k -pIF p -ideal of U and it is normal by Lemma 2. It is clear that $(\widehat{\xi}, \widehat{\varrho})$ is contained in $(\widehat{\xi}, \widehat{\varrho})^+$. \square

Theorem 9. Let $(\widehat{\xi}, \widehat{\varrho})$ be a k -pIF p -ideal of U . Then, $(\widehat{\xi}, \widehat{\varrho})$ is normal if and only if $(\widehat{\xi}, \widehat{\varrho})^+ = (\widehat{\xi}, \widehat{\varrho})$, that is, $\widehat{\xi}^+ = \widehat{\xi}$ and $\widehat{\varrho}^+ = \widehat{\varrho}$.

Proof. The sufficiency is clear. Assume that $(\widehat{\xi}, \widehat{\varrho})$ is normal. Then,

$$\begin{aligned} (\text{proj}_i \circ \widehat{\xi})^+(z) &= 1 + (\text{proj}_i \circ \widehat{\xi})(z) - (\text{proj}_i \circ \widehat{\xi})(0) = (\text{proj}_i \circ \widehat{\xi})(z) \\ (\text{proj}_i \circ \widehat{\varrho})^+(z) &= (\text{proj}_i \circ \widehat{\varrho})(z) - (\text{proj}_i \circ \widehat{\xi})(0) = (\text{proj}_i \circ \widehat{\xi})(z) \end{aligned}$$

for all $z \in U$ by Lemma 2. This completes the proof. \square

Corollary 2. Let $(\widehat{\xi}, \widehat{\varrho})$ be a k -pIF p -ideal of U . If $(\widehat{\xi}, \widehat{\varrho})$ is normal, then $((\widehat{\xi}, \widehat{\varrho})^+)^+ = (\widehat{\xi}, \widehat{\varrho})$.

Theorem 10. Let $(\widehat{\xi}, \widehat{\varrho})$ be a non-constant normal k -pIF p -ideal of U , which is maximal in the poset of normal k -pIF p -ideals under set inclusion. Then, $\widehat{\xi}$ and $\widehat{\varrho}$ have the values $\widehat{0}$ and $\widehat{1}$ only.

Proof. Since $(\widehat{\xi}, \widehat{\varrho})$ is normal, we have $\widehat{\xi}(0) = \widehat{1}$ and $\widehat{\varrho}(0) = \widehat{0}$ by Lemma 2. Let $z, x \in U$ be such that $\widehat{\xi}(z) \neq \widehat{1}$ and $\widehat{\varrho}(x) \neq \widehat{0}$. It is sufficient to show that $\widehat{\xi}(z) = \widehat{0}$ and $\widehat{\varrho}(x) = \widehat{1}$. If $\widehat{\xi}(z) \neq \widehat{0}$ and $\widehat{\varrho}(x) \neq \widehat{1}$, then there exists $b, c \in U$ such that $\widehat{0} < \widehat{\xi}(b) < \widehat{1}$ and $\widehat{0} < \widehat{\varrho}(c) < \widehat{1}$. Let $(\widehat{\xi}, \widehat{\varrho})_* = (\widehat{\xi}_*, \widehat{\varrho}_*)$ be a k -pIF set on U given by

$$\widehat{\xi}_* : U \rightarrow [0, 1]^k, z \mapsto \frac{1}{2} (\widehat{\xi}(z) + \widehat{\xi}(b)).$$

and

$$\widehat{\varrho}_* : U \rightarrow [0, 1]^k, z \mapsto \frac{1}{2} (\widehat{\varrho}(z) + \widehat{\varrho}(c)).$$

It is clear that $(\widehat{\xi}, \widehat{\varrho})_*$ is well-defined. For any $z, x \in U$, we have

$$\widehat{\xi}_*(0) = \frac{1}{2} (\widehat{\xi}(0) + \widehat{\xi}(b)) = \frac{1}{2} (\widehat{1} + \widehat{\xi}(b)) \geq \frac{1}{2} (\widehat{\xi}(z) + \widehat{\xi}(b)) = \widehat{\xi}_*(z),$$

$$\widehat{\varrho}_*(0) = \frac{1}{2} (\widehat{\varrho}(0) + \widehat{\varrho}(c)) = \frac{1}{2} (\widehat{0} + \widehat{\varrho}(c)) \leq \frac{1}{2} (\widehat{\varrho}(z) + \widehat{\varrho}(c)) = \widehat{\varrho}_*(z),$$

$$\begin{aligned} \widehat{\xi}_*(z) &= \frac{1}{2} (\widehat{\xi}(z) + \widehat{\xi}(b)) \geq \frac{1}{2} ((\widehat{\xi}(z * x) \wedge \widehat{\xi}(x)) + \widehat{\xi}(b)) \\ &= \frac{1}{2} ((\widehat{\xi}(z * x) + \widehat{\xi}(b)) \wedge (\widehat{\xi}(x) + \widehat{\xi}(b))) \\ &= \frac{1}{2} (\widehat{\xi}(z * x) + \widehat{\xi}(b)) \wedge \frac{1}{2} (\widehat{\xi}(x) + \widehat{\xi}(b)) \\ &= \widehat{\xi}_*(z * x) \wedge \widehat{\xi}_*(x) \end{aligned}$$

and

$$\begin{aligned} \widehat{\varrho}_*(z) &= \frac{1}{2} (\widehat{\varrho}(z) + \widehat{\varrho}(c)) \leq \frac{1}{2} ((\widehat{\varrho}(z * x) \vee \widehat{\varrho}(x)) + \widehat{\varrho}(c)) \\ &= \frac{1}{2} ((\widehat{\varrho}(z * x) + \widehat{\varrho}(c)) \vee (\widehat{\varrho}(x) + \widehat{\varrho}(c))) \\ &= \frac{1}{2} (\widehat{\varrho}(z * x) + \widehat{\varrho}(c)) \vee \frac{1}{2} (\widehat{\varrho}(x) + \widehat{\varrho}(c)) \\ &= \widehat{\varrho}_*(z * x) \vee \widehat{\varrho}_*(x). \end{aligned}$$

Hence $(\widehat{\xi}, \widehat{\varrho})$ is a k -pIF ideal of U . We have

$$\widehat{\xi}_*(z) = \frac{1}{2} (\widehat{\xi}(z) + \widehat{\xi}(b)) \geq \frac{1}{2} (\widehat{\xi}(0 * (0 * z)) + \widehat{\xi}(b)) = \widehat{\xi}_*(0 * (0 * z))$$

and

$$\widehat{q}_*(z) = \frac{1}{2}(\widehat{q}(z) + \widehat{q}(c)) \leq \frac{1}{2}(\widehat{q}(0 * (0 * z)) + \widehat{q}(c)) = \widehat{q}_*(0 * (0 * z))$$

for all $z \in U$. Hence $(\widehat{\xi}, \widehat{q})_*$ is a k -pIF p -ideal of U by Theorem 4. Now, we get

$$\widehat{\xi}_*^+(z) = \widehat{1} + \widehat{\xi}_*(z) - \widehat{\xi}_*(0) = \widehat{1} + \frac{1}{2}(\widehat{\xi}(z) + \widehat{\xi}(b)) - \frac{1}{2}(\widehat{\xi}(0) + \widehat{\xi}(b)) = \frac{1}{2}(\widehat{1} + \widehat{\xi}(z)),$$

and

$$\widehat{q}_*^+(z) = \widehat{q}_*(z) - \widehat{q}_*(0) = \frac{1}{2}(\widehat{q}(z) + \widehat{q}(c)) - \frac{1}{2}(\widehat{q}(0) + \widehat{q}(c)) = \frac{1}{2}\widehat{q}(z),$$

and so $\widehat{\xi}_*^+(0) = \frac{1}{2}(\widehat{1} + \widehat{\xi}(0)) = \widehat{1}$ and $\widehat{q}_*^+(z) = \frac{1}{2}\widehat{q}(0) = \widehat{0}$. Hence $(\widehat{\xi}, \widehat{q})_*$ is normal. Note that

$$\widehat{\xi}_*^+(0) = \widehat{1} > \widehat{\xi}_*^+(b) = \frac{1}{2}(\widehat{1} + \widehat{\xi}(b)) > \widehat{\xi}(b)$$

and

$$\widehat{q}_*^+(0) = \widehat{0} < \widehat{q}_*^+(c) = \frac{1}{2}(\widehat{0} + \widehat{q}(c)) < \widehat{q}(c).$$

Hence $(\widehat{\xi}, \widehat{q})_*^+$ is non-constant and $(\widehat{\xi}, \widehat{q})$ is not maximal, which is a contradiction; therefore $\widehat{\xi}$ and \widehat{q} have the values $\widehat{0}$ and $\widehat{1}$ only. \square

4. Conclusions and Future Works

As a generalization of intuitionistic fuzzy set, Kang et al. [19] introduced the notion of multipolar intuitionistic fuzzy set with finite degree, and then they applied the notion to BCK/BCI-algebras. In this manuscript, we used Kang et al.'s multipolar intuitionistic fuzzy set to study p -ideal in BCI-algebras. We introduced the notion of k -polar intuitionistic fuzzy p -ideals (see Definition 1) in BCI-algebras, and then we studied several properties (See Proposition 1, Proposition 2). We gave an example to illustrate the k -polar intuitionistic fuzzy p -ideal (see Example 1), and considered the relationship between k -polar intuitionistic fuzzy ideal and k -polar intuitionistic fuzzy p -ideal. We have shown that every k -polar intuitionistic fuzzy p -ideal is a k -polar intuitionistic fuzzy ideal (see Theorem 2), and then provided an example to show that the converse is not true in general (see Example 2). We used the notion of p -ideals in BCI-algebras to study the characterization of k -polar intuitionistic fuzzy p -ideal (see Theorem 1, Theorem 5 and Theorem 6), and also used the notion of k -polar (\in, \in) -fuzzy p -ideal in BCI-algebras to study the characterization of k -polar intuitionistic fuzzy p -ideal (see Theorem 7). We defined the concept of normal k -polar intuitionistic fuzzy p -ideal (see Definition 2), and discussed its characterization (see Lemma 2 and Theorem 9). We looked at the process of eliciting normal k -polar intuitionistic fuzzy p -ideal from a given k -polar intuitionistic fuzzy p -ideal (see Theorem 8). Our goal in the future is to apply the ideas and results of this paper to other forms of ideals, filters, etc. in BCK/BCI-algebras. We will also apply the ideas and results of this paper to other algebraic structures, for example, MV-algebras, EQ-algebras, equality algebras, hoops, etc.

Author Contributions: Created and conceptualized ideas, J.-G.L. and Y.B.J.; writing—original draft preparation, Y.B.J.; writing—review and editing, M.F. and K.H.; funding acquisition, J.-G.L. All authors have read and agreed to the published version of the manuscript.

Funding: This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (2018R1D1A1B07049321).

Acknowledgments: We would like to thank the guest editor and the anonymous reviewers for their very careful reading and valuable comments/suggestions.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Iséki, K. An algebra related with a propositional calculus. *Proc. Jpn. Acad.* **1966**, *42*, 26–29. [[CrossRef](#)]
2. Huang, Y. *BCI-Algebra*; Science Press: Beijing, China, 2006.
3. Meng, J.; Jun, Y.B. *BCK-Algebras*; Kyungmoonsa Co.: Seoul, Korea, 1994.
4. Zadeh, L.A. Fuzzy sets. *Inform. Control* **1965**, *8*, 338–353. [[CrossRef](#)]
5. Atanassov, K.T. Intuitionistic fuzzy sets. VII ITKR Session, Sofia, 20–23 June 1983 (Deposited in Centr. Sci.-Techn. Library of the Bulg. Acad. of Sci., 1697/84) (in Bulgarian). *Repr. Int. Bioautomation* **2016**, *20*, S1–S6.
6. Atanassov, K.T. Intuitionistic fuzzy sets. *Fuzzy Sets Syst.* **1986**, *20*, 87–96. [[CrossRef](#)]
7. Chen, J.; Li, S.; Ma, S.; Wang, X. m -polar fuzzy sets: An extension of bipolar fuzzy sets. *Sci. World J.* **2014**, *2014*, 416530. [[CrossRef](#)] [[PubMed](#)]
8. Al-Masarwah, A.; Ahmad, A.G. m -polar fuzzy ideals of BCK/BCI-algebras. *J. King Saud Univ. Sci.* **2019**, *31*, 1220–1226. [[CrossRef](#)]
9. Mohseni Takallo, M.; Ahn, S.S.; Borzooei, R.A.; Jun, Y.B. Multipolar fuzzy p -ideals of BCI-algebras. *Mathematics* **2019**, *7*, 1094. [[CrossRef](#)]
10. Al-Masarwah, A.; Ahmad, A.G. m -polar (α, β) -fuzzy ideals in BCK/BCI-algebras. *Symmetry* **2019**, *11*, 44. [[CrossRef](#)]
11. Akram, M.; Sarwar, M. New applications of m -polar fuzzy competition graphs. *New Math. Nat. Comput.* **2018**, *14*, 249–276. [[CrossRef](#)]
12. Akram, M.; Adeel, A. m -polar fuzzy graphs and m -polar fuzzy line graphs. *J. Discret. Math. Sci. Cryptogr.* **2017**, *20*, 1597–1617. [[CrossRef](#)]
13. Akram, M.; Waseem, N.; Davvaz, B. Certain types of domination in m -polar fuzzy graphs. *J. Mult. Valued Log. Soft Comput.* **2017**, *29*, 619–646.
14. Sarwar, M.; Akram, M. Representation of graphs using m -polar fuzzy environment. *Ital. J. Pure Appl. Math.* **2017**, *38*, 291–312.
15. Akram, M.; Waseem, N.; Liu, P. Novel approach in decision making with m -polar fuzzy ELECTRE-I. *Int. J. Fuzzy Syst.* **2019**, *21*, 1117–1129. [[CrossRef](#)]
16. Akram, M.; Ali, G.; Alshehri, N.O. A New Multi-Attribute Decision-Making Method Based on m -Polar Fuzzy Soft Rough Sets. *Symmetry* **2017**, *9*, 271. [[CrossRef](#)]
17. Adeel, A.; Akram, M.; Koam, A.N.A. Group decision-making based on m -polar fuzzy linguistic TOPSIS method. *Symmetry* **2019**, *11*, 735. [[CrossRef](#)]
18. Adeel, A.; Akram, M.; Ahmed, I.; Nazar, K. Novel m -polar fuzzy linguistic ELECTRE-I method for group decision-making. *Symmetry* **2019**, *11*, 471. [[CrossRef](#)]
19. Kang, K.T.; Song, S.Z.; Jun, Y.B. Multipolar intuitionistic fuzzy set with finite degree and its application in BCK/BCI-algebras. *Mathematics* **2020**, *8*, 177. [[CrossRef](#)]
20. Zhang, X.H.; Hao, J.; Bhatti, S.A. On p -ideals of a BCI-algebra. *Punjab Univ. J. Math. (Lahore)* **1994**, *27*, 121–128.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).