

Remarks on the Local Irregularity Conjecture

Jelena Sedlar ^{1,2,†}  and Riste Škrekovski ^{2,3,*,†} 

¹ Faculty of Civil Engineering, Architecture and Geodesy, University of Split, 21000 Split, Croatia; jsedlar@gradst.hr or jelena.sedlar@gmail.com

² Faculty of Information Studies, University of Ljubljana, 8000 Novo Mesto, Slovenia

³ Faculty of Mathematics and Physics, University of Ljubljana, 1000 Ljubljana, Slovenia

* Correspondence: riste.skrekovski@fmf.uni-lj.si or skrekovski@gmail.com

† These authors contributed equally to this work.

Abstract: A locally irregular graph is a graph in which the end vertices of every edge have distinct degrees. A locally irregular edge coloring of a graph G is any edge coloring of G such that each of the colors induces a locally irregular subgraph of G . A graph G is colorable if it allows a locally irregular edge coloring. The locally irregular chromatic index of a colorable graph G , denoted by $\chi'_{\text{irr}}(G)$, is the smallest number of colors used by a locally irregular edge coloring of G . The local irregularity conjecture claims that all graphs, except odd-length paths, odd-length cycles and a certain class of cacti are colorable by three colors. As the conjecture is valid for graphs with a large minimum degree and all non-colorable graphs are vertex disjoint cacti, we study rather sparse graphs. In this paper, we give a cactus graph B which contradicts this conjecture, i.e., $\chi'_{\text{irr}}(B) = 4$. Nevertheless, we show that the conjecture holds for unicyclic graphs and cacti with vertex disjoint cycles.

Keywords: locally irregular edge coloring; local irregularity conjecture; unicyclic graph; cactus graph



Citation: Sedlar, J.; Škrekovski, R. Remarks on the Local Irregularity Conjecture. *Mathematics* **2021**, *9*, 3209. <https://doi.org/10.3390/math9243209>

Academic Editor: Elena Guardo

Received: 16 November 2021

Accepted: 10 December 2021

Published: 12 December 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

All graphs mentioned in this paper are considered to be simple and finite. An edge coloring of a graph is *neighbor sum distinguishing* if any two neighboring vertices differ in the sum of the colors of the edges incident to them. This notion was first introduced in [1] and the following conjecture was proposed there.

Conjecture 1 (1-2-3 Conjecture). *Every graph G without isolated edges admits a neighbor-sum-distinguishing edge coloring with the colors $\{1, 2, 3\}$.*

This conjecture attracted a lot of interest [2–6] and for a survey we refer the reader to [7]. The best upper bound is that every graph without isolated edges admits a neighbor-sum-distinguishing edge coloring with five colors [8], but the 1-2-3 Conjecture remains open.

This variant of edge coloring and the 1-2-3 Conjecture motivated the introduction of similar variants of edge coloring. A *locally irregular* graph is any graph in which the two end vertices of every edge differ in degree. A *locally irregular k -edge coloring*, or *k -liec* for short, is any edge coloring of G with k colors such that every color induces a locally irregular subgraph of G . This variant of edge coloring was introduced in [9]. A third related edge coloring variant is the *neighbor multiset-distinguishing edge coloring*, where neighboring vertices must have assigned distinct multisets of colors on incident edges. In [10], it was established that every graph without isolated edges admits the neighbor multiset-distinguishing edge coloring with four colors. Notice that every locally irregular edge coloring is also a neighbor multiset-distinguishing edge coloring, but the reverse does not have to hold. In order to see that the reverse does not hold, consider the following graph with its edge coloring: G is a 2-path abc and edges ab and bc are colored by 1 and 2, respectively. This is obviously not a locally irregular edge coloring but it is a neighbor

multiset-distinguishing edge coloring as $\{1\}$, $\{1, 2\}$, $\{1, 2\}$ are the (multi)sets of colors at vertices a , b , c , respectively.

In this paper, we focus our attention on locally irregular edge colorings exclusively, and we say a graph is *colorable* if it admits such a coloring. The *locally irregular chromatic index* of a colorable graph G , denoted by $\chi'_{\text{irr}}(G)$, is the smallest k , such that G admits a k -liec. In [9], the family of graphs \mathfrak{T} has been defined as follows:

- \mathfrak{T} contains the triangle K_3 ;
- if G is a graph from \mathfrak{T} , then a graph H obtained from G by identifying a vertex $v \in V(G)$ of degree 2, which belongs to a triangle of G , with an end-vertex of an even length path or with an end vertex of an odd-length path such that the other end vertex of that path is identified with a vertex of a triangle, also belongs to \mathfrak{T} .

Note that every graph $G \in \mathfrak{T}$ has an odd size. A *cactus graph* is any graph in which cycles are pairwise edge disjoint. Notice that \mathfrak{T} is a special family of cacti. Additionally, if we imagine triangles to be vertices and paths attached to vertices of a triangle as edges, we might informally say that G has tree-like structure. For the sake of simplicity, we define a broader family \mathfrak{T}' as the family obtained from \mathfrak{T} by introducing to it all odd-length paths and all odd-length cycles. Notice that \mathfrak{T}' is a subclass of vertex-disjoint cactus graphs. It was established in [9] that a connected graph G is not colorable if and only if $G \in \mathfrak{T}'$. Additionally, the following conjecture on the irregular chromatic index was proposed.

Conjecture 2 (Local Irregularity Conjecture). *For every connected graph $G \notin \mathfrak{T}'$, it holds that $\chi'_{\text{irr}}(G) \leq 3$.*

Let us mention some of the results related to Conjecture 2. For general graphs it was first established $\chi'_{\text{irr}}(G) \leq 328$ [11], then it was lowered to $\chi'_{\text{irr}}(G) \leq 220$ [12]. For some special classes of graphs Conjecture 2 is shown to hold, namely for trees [13], graphs with minimum degree at least 10^{10} [14], k -regular graphs where $k \geq 10^7$ [9].

In this paper we will show that every unicyclic graph G which does not belong to \mathfrak{T}' admits a 3-liec, thus establishing that Conjecture 2 holds for unicyclic graphs. We will further extend this result to cactus graphs with vertex-disjoint cycles. Finally, we will provide an example of a colorable graph B with $\chi'_{\text{irr}}(B) = 4$ showing that Conjecture 2 does not hold in general.

2. Revisiting the Trees

Since a unicyclic graph is obtained from a tree by adding a single edge to it, we first need to introduce the notation and several important results for trees from [13]. Additionally, we will establish several auxiliary results for trees, which will be useful throughout the paper.

First, a *shrub* is any tree rooted at a leaf. The only edge in a shrub G incident to the root we will call the *root edge* of G . An *almost locally irregular k -edge coloring* of a shrub G , or *k -aliec* for short, is an edge coloring of G which is either k -liec or a coloring in which only the root edge is not locally irregular (notice that in this case the root edge is an isolated edge of its color, i.e., it is not adjacent to any other edge of the same color). A *proper k -aliec* is k -aliec which is not a k -liec. The following results for trees were established in [13].

Theorem 1. *Every shrub admits a 2-aliec.*

Theorem 2. *For every colorable tree T , it holds that $\chi'_{\text{irr}}(T) \leq 3$. Moreover, $\chi'_{\text{irr}}(T) \leq 2$ if $\Delta(T) \geq 5$.*

If an edge coloring uses at most three colors, we will denote those colors by a, b, c . A 1-liec (resp. 2-liec, 3-liec) of a graph G will be denoted by $\phi_a(G)$ (resp. $\phi_{a,b}(G)$, $\phi_{a,b,c}(G)$). A 2-aliec of a shrub G will be denoted by $\phi_{a,b}(G)$ where a is the color of the root edge in G . Let a, b, c, d be four colors, if $\phi_{a,b}(G)$ is a 2-liec of G in colors a and b , then 2-liec

$\phi_{c,d}(G)$ of G in colors c and d is obtained from $\phi_{a,b}(G)$ by replacing colors a and b for c and d respectively, i.e., $\phi_{c,d}(e) = c$ if and only if $\phi_{a,b}(e) = a$. Particularly, 2-(a)lic $\phi_{b,a}(G)$ is called the *inversion* of the 2-(a)lic $\phi_{a,b}(G)$, where colors a and b are replaced. Moreover, let $\phi_{a,b,c}^i$ be an edge coloring of a graph G_i , for $i = 1, \dots, k$, and let these graphs be pairwise edge disjoint, i.e., $E(G_i) \cap E(G_j) = \emptyset$ for any two distinct i and j . For a graph G such that $E(G) = \cup_{i=1}^k E(G_i)$, by $\sum_{i=1}^k \phi_{a,b,c}^i$ we will denote the edge coloring of G such that an edge e is colored by $\phi_{a,b,c}^i(e)$ if and only if $e \in E(G_i)$.

For any color of the edge coloring $\phi_{a,b,c}$, say a , we define the a -degree of a vertex $v \in V(G)$ as the number of edges incident to v which are colored by a . The a -degree of a vertex v is denoted by $d_G^a(v)$. Assume that a vertex $v \in V(G)$ has k neighbors w_1, \dots, w_k such that each vw_i is colored by a . Then, the sequence $d_G^a(w_1), \dots, d_G^a(w_k)$ is called the a -sequence of the vertex v . We usually assume that neighbors of v are denoted so that the a -sequence is non-increasing.

Throughout the paper we will use the technique of finding a 2-lic for trees introduced in [13]. Namely, if T is a tree with maximum degree 5 or more, then T admits a 2-lic according to Theorem 2. Otherwise, if the maximum degree of T is at most 4, let v be a vertex from T and w_1, \dots, w_k all the neighbors of v for $k \leq 4$. Notice that T consists of k shrubs T_i starting at v , let T_i denote a shrub with the root edge vw_i and let $\phi_{a,b}^i$ denote a 2-lic of T_i which exists according to Theorem 1. Recall that $\phi_{a,b}^i(vw_i) = a$ for every $i \leq k$. The coloring $\phi_{a,b} = \sum_{i=1}^k \phi_{a,b}^i$ is called a *shrubs-based* edge coloring of T . We say that a shrub-based coloring $\phi_{a,b}$ is *inversion resistant* if neither $\phi_{a,b}$ is a 2-lic of T nor any of the colorings which can be obtained from $\phi_{a,b}$ by color inversion in some of the shrubs T_i . Let us now introduce the following lemma which stems from the technique used in [13].

Lemma 1. Let T be a tree with $\Delta(T) \leq 4$ and v a vertex from T of degree k . Let T_1, \dots, T_k be all the shrubs of T rooted at v and let $\phi_{a,b}^i$ be a 2-lic of T_i . If $\phi_{a,b}^i$ is a 2-lic of T_i for every $i = 1, \dots, k$, then the shrub-based coloring $\phi_{a,b} = \sum_{i=1}^k \phi_{a,b}^i$ can be inversion resistant in two cases only:

- If $d_T(v) = 3$ and the a -sequence of v by $\phi_{a,b}$ is 3, 2, 2;
- If $d_T(v) = 4$ and the a -sequence of v by $\phi_{a,b}$ is 4, 3, 3, 2.

Proof. If $d_T(v) = 1$, then the shrub-based coloring of T equals $\phi_{a,b}^1$, which is 2-lic. If $d_T(v) = 2$, then $\phi_{a,b}^1 + \phi_{b,a}^2$ would be a 2-lic of T .

If $d_T(v) = 3$, then the a -degree of v by $\phi_{a,b}$ is 3, and inverting colors in one of the shrubs would decrease the a -degree of v to 2. Therefore, the a -sequence of v by $\phi_{a,b}$ must contain 3 and 2. Considering the two possibilities 4, 3, 2, and 3, 3, 2, we see that $\phi_{a,b}^1 + \phi_{a,b}^2 + \phi_{b,a}^3$ would be 2-lic in both of them. The only remaining possibility is 3, 2, 2, and it is inversion resistant.

Finally, assume $d_T(v) = 4$. By a similar consideration as above, we see that the a -sequence of v by $\phi_{a,b}$ must contain 4, 3, and 2. Therefore, we must consider the possibilities 4, 4, 3, 2, then 4, 3, 3, 2, then 4, 3, 2, 2. It is easily seen that only in the case 4, 3, 3, 2, the shrub-based coloring $\phi_{a,b}$ is inversion resistant. \square

A *spidey* is a tree with radius at most two which consists of a central vertex u of degree at least 3 and the remaining vertices have degree at most 2 and are at distance at most 2 from u . Notice that every spidey is locally irregular, hence it admits a 1-lic. We say that a vertex v of a spidey G is a *short leg* if it is a leaf which is a neighbor of the central vertex of G .

Lemma 2. Let H be a spidey with a short leg v and let K be a tree. Let G be a graph obtained from H and K by identifying the vertex v with a vertex from K . Then, G admits a 3-lic such that all edges of $E(H)$ are colored by a same color.

Proof. Since H is a spidey, H admits a 1-lic, say ϕ_c^0 . Assume first that a tree K is not colorable, i.e., K is an odd-length path. This implies there exists in K an edge e_v incident with v , such that $K - e_v$ is a collection of even paths which, therefore, admits 2-lic $\phi_{a,b}^1$. The edge coloring $\phi_{a,b,c}$ of G defined by

$$\phi_{a,b,c}(e) = \begin{cases} c & \text{if } e = e_v, \\ \phi_{a,b}^1(e) & \text{if } e \in E(K) \setminus \{e_v\}, \\ \phi_c^0(e) & \text{if } e \in E(H), \end{cases}$$

is a 3-lic of G .

Assume now that K is a colorable tree. If K admits a 2-lic $\phi_{a,b}^1$, then $\phi_c^0 + \phi_{a,b}^1$ is a 3-lic of G with the desired property. So, we may assume K is a colorable tree which does not admit a 2-lic. Theorem 2 implies $\Delta(K) \leq 4$. Let $d_K(v) = k \leq 4$ and let T_1, \dots, T_k be all the shrubs of K rooted at v . By Theorem 1, each shrub T_i admits a 2-alic $\phi_{a,b}^i$, where without loss of generality we may assume that $\phi_{a,b}^i$ is a proper 2-alic if and only if $i \leq l$. We distinguish the following four cases with respect to l .

Case 1: $l \geq 3$. Notice that $l \in \{3, 4\}$ and $l \leq k \leq 4$. If $l = 3$ and $k = 4$, then $\phi_{a,b}^1 + \phi_{a,b}^2 + \phi_{a,b}^3 + \phi_{b,a}^4$ would be a 2-lic of K , a contradiction. Otherwise, the shrub-based coloring $\phi_{a,b} = \sum_{i=1}^k \phi_{a,b}^i$ would be a 2-lic of K , again a contradiction.

Case 2: $l = 2$. If $k = 2$, then $\phi_{a,b}^1 + \phi_{a,b}^2$ is a 2-lic of K , a contradiction. If $k = 3$, then $\phi_{a,b}^1 + \phi_{a,b}^2 + \phi_{b,a}^3$ is a 2-lic of K , a contradiction. If $k = 4$, then let w_1, \dots, w_4 be all the neighbors of v in K . The shrub-based coloring $\phi_{a,b} = \sum_{i=1}^k \phi_{a,b}^i$ is not a 2-lic only if the a -degree of w_3 or w_4 by $\phi_{a,b}$ is 4. Without loss of generality we may assume that a -degree of w_3 by $\phi_{a,b}$ is 4, but then $\phi_{a,b}^1 + \phi_{a,b}^2 + \phi_{a,b}^3 + \phi_{b,a}^4$ is a 2-lic of K , a contradiction.

Case 3: $l = 1$. In this case, T_1 is the only shrub with a proper 2-alic $\phi_{a,b}^1$. Let w_1 be the neighbor of v in T_1 , we define the coloring $\phi'_{a,b,c}$ of K as follows

$$\phi'_{a,b,c}(e) = \begin{cases} c & \text{if } e = vw_1, \\ \sum_{i=1}^k \phi_{a,b}^i(e) & \text{if } e \in E(K) \setminus \{vw_1\}. \end{cases}$$

Notice that $\phi'_{a,b,c}$ is not a lic of K , but $\phi_c^0 + \phi'_{a,b,c}$ is a 3-lic of $G = H + K$ with the desired property that all edges of H are colored by a same color, in this case c .

Case 4: $l = 0$. Notice that in this case Lemma 1 applies on K and v . Therefore, the only cases when K does not admit a 2-lic are: i) $d_K(v) = 3$ and the a -sequence of v by the shrub-based coloring $\phi_{a,b} = \sum_{i=1}^k \phi_{a,b}^i$ is 3, 2, 2, or ii) $d_K(v) = 4$ and the a -sequence of v by $\phi_{a,b}$ is 4, 3, 3, 2. In the first case the coloring $\phi_c^0 + \phi_{c,b}^1 + \phi_{b,a}^2 + \phi_{a,b}^3$ is a 3-lic of G such that $E(H)$ is colored by the same color c , as it is illustrated in Figure 1. In the other case, the coloring $\phi_c^0 + \phi_{a,b}^1 + \phi_{c,b}^2 + \phi_{a,b}^3 + \phi_{b,a}^4$ is a 3-lic of G such that $E(H)$ is colored by a same color, as it is illustrated in Figure 2. \square




Figure 1. A graph $G = H + K$ and a vertex v of degree 3 in K : (a) the coloring $\phi_c^0 + \phi_{a,b}^1 + \phi_{a,b}^2 + \phi_{a,b}^3$ is not a 3-lic of G ; (b) the coloring $\phi_c^0 + \phi_{c,b}^1 + \phi_{b,a}^2 + \phi_{a,b}^3$ is a 3-lic of G .




Figure 2. A graph $G = H + K$ and a vertex v of degree 4 in K : (a) the coloring $\phi_c^0 + \phi_{a,b}^1 + \phi_{a,b}^2 + \phi_{a,b}^3 + \phi_{a,b}^4$ is not a 3-lic of G , (b) the coloring $\phi_c^0 + \phi_{a,b}^1 + \phi_{c,b}^2 + \phi_{a,b}^3 + \phi_{b,a}^4$ is a 3-lic of G .

3. Unicyclic Graphs

In this section, we will establish Conjecture 2 for unicyclic graphs. It is already known that there exist colorable unicyclic graphs which do not admit 2-lic, but require three colors in order for edge coloring to be locally irregular, namely cycles of length $4k + 2$, for $k \in \mathbb{N}$. We will show that such cycles are not an isolated family of unicyclic graphs that require three colors. The main result for unicyclic graphs is established through the following two lemmas in which we will consider separately cases whether the cycle of G is a triangle or not.

Lemma 3. Let G be a unicyclic graph with the unique cycle being a triangle. If $G \notin \mathfrak{T}'$, then $\chi'_{\text{irr}}(G) \leq 3$.

Proof. Let $C = u_1u_2u_3$ be the triangle in G , let T_i denote the connected component of $G - E(C)$ which contains u_i . Since $G \notin \mathfrak{T}'$, there must exist a vertex u_i on C such that T_i is not a pendant even length path, say it is u_1 . Let $G_1 = T_1 + u_1u_2$ and let $G_0 = G - E(G_1)$. First notice that both G_0 and G_1 are trees and that $E(G) = E(G_0) \cup E(G_1)$. Since T_1 is not a pendant path of even length, it follows that G_1 is not an odd length path, hence it is colorable. Let $\phi_{a,b,c}^1$ be a 3-lic of G_1 . Without loss of generality we may assume that $\phi_{a,b,c}^1(u_1u_2) = c$. Let H be the subgraph of G_1 induced by all edges incident to u_1 in G_1 . We may assume $\phi_{a,b,c}^1(e) = c$ for every $e \in E(H)$, namely if $d_H(u_1) = 2$ this follows from the local irregularity of $\phi_{a,b,c}^1$; otherwise, it follows from Lemma 2 applied on H and every component of $G_1 - E(H)$ repeatedly.

Let us now consider the graph G_0 and notice that it is a shrub rooted at u_1 with the root edge u_1u_3 . By Theorem 1 there exists a 2-alic $\phi_{a,b}^0$ of G_0 . If $\phi_{a,b}^0$ is a 2-lic, then

$\phi_{a,b,c} = \phi_{a,b}^0 + \phi_{a,b,c}^1$ is a 3-lic of G . Additionally, if $\phi_{a,b}^0$ is a proper 2-aliec of G_0 , then $G_1 + u_1u_3$ is colorable and let $\phi_{a,b,c}^2$ be a 3-lic of $G_1 + u_1u_3$ such that $\phi_{a,b,c}^2(e) = c$ for every edge $e \in E(H) \cup \{u_1u_3\}$ by Lemma 2. We define the edge coloring $\phi_{a,b,c}$ of G as follows

$$\phi_{a,b,c}(e) = \begin{cases} \phi_{a,b}^0(e) & \text{if } e \in E(G_0) \setminus \{u_1u_3\}, \\ \phi_{a,b,c}^1(e) & \text{if } e \in E(G_1). \end{cases}$$

It is easily seen that $\phi_{a,b,c}$ is a 3-lic of G . \square

Let us now consider unicyclic graphs with larger cycles.

Lemma 4. Let G be a unicyclic graph with the unique cycle being of length at least four. If $G \notin \mathfrak{T}'$, then $\chi'_{\text{irr}}(G) \leq 3$.

Proof. If G is a cycle, then $G \notin \mathfrak{T}'$ implies that G is an even-length cycle and hence admits a 3-lic. So, we may assume G is not a cycle, i.e., at least one vertex from the cycle of G is of degree at least three. Denote the cycle in G by $C = u_1u_2 \cdots u_g$ with $g \geq 4$. Without loss of generality we may assume that u_1 is the vertex with maximum degree among vertices from C . We distinguish the following two cases with respect to $d_G(u_1)$.

Case 1: $d_G(u_1) \geq 4$. Let E_1 denote the set of all edges incident to u_1 in G except the edge u_1u_2 and let H denote the subgraph of G induced by E_1 . The assumption $d_G(u_1) \geq 4$ implies $d_H(u_1) \geq 3$, so H is a spidey in which every leg is short. Let G_0 be the connected component of $G - E_1$ which contains u_2 and let $G_1 = G - E(G_0)$. Let G'_1, \dots, G'_k be all connected components of $G_1 - E(H)$. Each G'_i is a tree, so Lemma 2 can be applied to H and $K = G'_i$, for every $i = 1, \dots, k$. We conclude that there exists a 3-lic $\phi_{a,b,c}^1$ of G_1 such that $\phi_{a,b,c}^1(e) = c$ for every $e \in E(H)$. On the other hand, G_0 is a shrub rooted at u_1 with the root edge u_1u_2 , so G_0 admits 2-aliec $\phi_{a,b}^0$ according to Theorem 1.

If $\phi_{a,b}^0$ is a 2-lic of G_0 , then $\phi_{a,b,c} = \phi_{a,b}^0 + \phi_{a,b,c}^1$ is a 3-lic of G . Additionally, if $\phi_{a,b}^0$ is a proper 2-aliec of G_0 , then $G_1 + u_1u_2$ is colorable and let $\phi_{a,b,c}^2$ be a 3-lic of $G_1 + u_1u_2$ such that $\phi_{a,b,c}^2(e) = c$ for every edge $e \in E(H) \cup \{u_1u_2\}$ by Lemma 2. We define the edge coloring $\phi_{a,b,c}$ of G as follows

$$\phi_{a,b,c}(e) = \begin{cases} \phi_{a,b}^0(e) & \text{if } e \in E(G_0) \setminus \{u_1u_2\}, \\ \phi_{a,b,c}^1(e) & \text{if } e \in E(G_1). \end{cases}$$

It is easily seen that thus defined $\phi_{a,b,c}$ is a 3-lic of G .

Case 2: $d_G(u_1) = 3$. Let E_1 be the set of all edges incident to u_1 in G and H a subgraph of G induced by E_1 . Let G_0 be the connected component of $G - E_1$ which contains u_2 and let $G_1 = G - E(G_0)$. Similarly as in the previous case, there exists a 3-lic $\phi_{a,b,c}^1$ of G_1 such that $\phi_{a,b,c}^1(e) = c$ for every $e \in E(H)$. Notice that $d_G(u_2) \in \{2, 3\}$, since u_1 is the vertex with maximum degree among vertices from C . Now we distinguish two possibilities with regard to $d_G(u_2)$.

If $d_G(u_2) = 2$, then G_0 is a shrub rooted in u_2 with the root edge u_2u_3 . According to Theorem 1, there exists a 2-aliec $\phi_{a,b}^0$ of G_0 . If $\phi_{a,b}^0$ is 2-lic of G_0 , then $\phi_{a,b,c} = \phi_{a,b}^0 + \phi_{a,b,c}^1$ is a 3-lic of G . Otherwise, $\phi_{a,b,c}$ defined by

$$\phi_{a,b,c}(e) = \begin{cases} c & \text{if } e = u_2u_3, \\ \phi_{a,b}^0(e) & \text{if } e \in E(G_0) \setminus \{u_2u_3\}, \\ \phi_{a,b,c}^1(e) & \text{if } e \in E(G_1), \end{cases}$$

is a 3-lic of G .

If $d_G(u_2) = 3$, then consider G_0 to be a tree rooted at u_2 which consists of two shrubs G'_0 and G''_0 , the first with the root edge u_2u_3 and the other with the root edge u_2v_2 , where v_2 is the only neighbor of u_2 which does not belong to the cycle C . Theorem 1 implies that

there exist 2 aliecs $\phi'_{a,b}$ and $\phi''_{a,b}$ of G'_0 and G''_0 , respectively. If both $\phi'_{a,b}$ and $\phi''_{a,b}$ are a 2-lic of the respective shrub, then $\phi_{a,b,c} = \phi'_{a,b} + \phi''_{a,b} + \phi^1_{a,b,c}$ is a 3-lic of G . If both $\phi'_{a,b}$ and $\phi''_{a,b}$ are a proper 2-aliec of the respective shrub, then $\phi_{a,b,c} = \phi'_{a,b} + \phi''_{a,b} + \phi^1_{a,b,c}$ is a 3-lic of G . The only remaining possibility is that precisely one of $\phi'_{a,b}$ and $\phi''_{a,b}$, say $\phi'_{a,b}$, is a proper 2-aliec of the respective shrub. In this case we define the coloring $\phi^0_{a,b,c}$ of G_0 as follows

$$\phi^0_{a,b,c}(e) = \begin{cases} c & \text{if } e = u_2u_3, \\ \phi'_{a,b}(e) & \text{if } e \in E(G'_0) \setminus \{u_2u_3\}, \\ \phi''_{a,b}(e) & \text{if } e \in E(G''_0). \end{cases}$$

Since $|V(C)| \geq 4$, it is easily seen that $\phi_{a,b,c} = \phi^0_{a,b,c} + \phi^1_{a,b,c}$ is a 3-lic of G . \square

The previous two lemmas yield the following result.

Theorem 3. *Let G be a unicyclic graph. If $G \notin \mathcal{T}'$, then $\chi'_{\text{irr}}(G) \leq 3$.*

A natural question that arises is whether the bound $\chi'_{\text{irr}}(G) \leq 3$ is tight, i.e., are there colorable unicyclic graphs which are not 2-colorable. The family of cycles of length $4k + 2$ are such graphs, but this family is not an isolated case, there exist other unicyclic graphs which require three colors, for example, the graph from Figure 3. One can assure infinitely many such graphs, for example, by taking longer threads of suitable parity in the given graph.




Figure 3. A colorable unicyclic graph distinct from cycle which requires 3 colors for locally irregular edge coloring.

4. Cacti with Vertex Disjoint Cycles

In this section, we will extend the result from the previous section to cacti with vertex disjoint cycles. We will also show that the result does not extend to all cacti by providing an example of a cactus graph with four cycles which is colorable, but requires 4 colors for a locally irregular edge coloring. This establishes that Conjecture 2 does not hold in general. We first need to introduce several useful notions in order to deal with cacti.

Let G be a cactus graph with at least two cycles, let C be a cycle in G and let u be a vertex from C . We say that u is a *root vertex* of C if the connected component of $G - E(C)$ which contains u is a cyclic graph. A cycle C of G is a *proper end cycle* if $G - V(C)$ contains at most one cyclic connected component. Every cactus graph with vertex disjoint cycles contains at least two proper end cycles, given it is not a unicyclic graph.

Theorem 4. *Let G be a cactus graph with vertex disjoint cycles. If $G \notin \mathcal{T}'$, then $\chi'_{\text{irr}}(G) \leq 3$.*

Proof. The proof is by induction on the number of cycles in G . If G is a unicyclic graph, then the claim holds by Theorem 3. Assume that the claim holds for all cacti with fewer than p cycles, where $p \geq 2$. Let G be a cactus graph with p cycles. We will show that G admits a 3-lic and this will establish the claim of the theorem. Let C be a proper end cycle of G , u_1 the root vertex of C , and v the only neighbor of u_1 which belongs to the cyclic component of $G - E(C)$. Denote the other neighbors of u_1 by u_2, \dots, u_k so that u_2 and u_3 belong to the cycle C . In what follows, we distinguish two cases.

Case 1: $d_G(u_1) = 3$. Let G_1 be the connected component $G - u_1v$ which does not contain v and let $G_0 = G - E(G_1)$. Let $G'_0 = G_0 + u_1u_2$ and $G'_1 = G_1 - u_1u_2$.

Suppose first that G'_0 is colorable. Then, it admits a 3-lic $\phi_{a,b,c}^0$ where the edges u_1u_2 and u_1v must be colored by a same color, say color c . Notice that G'_1 is a shrub rooted at u_1 with the root edge u_1u_3 . By Theorem 1, G'_1 admits a 2-lic $\phi_{a,b}^1$. If $\phi_{a,b}^1$ is a 2-lic of G'_1 , then $\phi_{a,b,c}^0 + \phi_{a,b}^1$ is a 3-lic of G . Otherwise, if $\phi_{a,b}^1$ is a proper 2-lic of G_1 , then the restriction of $\phi_{a,b}^1$ to $G''_1 = G'_1 - u_1u_3$ is a 2-lic of that graph. Notice that $G''_0 = G'_0 + u_1u_3$ does not belong to \mathfrak{T}' , so it is colorable and by induction hypothesis it admits a 3-lic $\phi_{a,b,c}''$ in which edges u_1u_2 and u_1u_3 must be colored by a same color (say color c) since $d_G(u_1) = 3$. Now, we infer that

$$\phi_{a,b,c}(e) = \begin{cases} \phi_{a,b}^1(e) & \text{if } e \in E(G'_1), \\ \phi_{a,b,c}''(e) & \text{if } e \in E(G''_0), \end{cases}$$

is a 3-lic of G .

Suppose now that G'_0 is not colorable. Assume first G_1 is not colorable. Notice that G_1 is a unicyclic graph, so if the cycle of G_1 is a triangle, then the assumption that G'_0 and G_1 are not colorable would imply $G \in \mathfrak{T}$, a contradiction. Otherwise, if G_1 is a unicyclic graph on a larger cycle, then it is not colorable only if it is an odd-length cycle. In this case let w be the only neighbor of u_3 distinct from u_1 , let $G''_0 = G'_0 + \{u_1u_3, u_3w\}$ and $G''_1 = G - E(G''_0)$. Notice that by induction hypothesis G''_0 is colorable and admits a 3-lic $\phi_{a,b,c}^0$ for which we may assume $\phi_{a,b,c}^0(u_3w) = a$ and $\phi_{a,b,c}^0(u_1u_2) \in \{a, b\}$. Additionally, notice that G''_1 is an even length path, so it admits a 2-lic $\phi_{b,c}^1$ where we may assume that the edge of G''_1 incident to u_2 is colored by c . Then $\phi_{a,b,c}^0 + \phi_{b,c}^1$ is a 3-lic of G .

Suppose now that G_1 is colorable. Since G'_0 is not colorable, the edge u_1u_2 of G'_0 must belong to an even length path hanging at a vertex of a triangle in G'_0 , so the graph $G_0 = G'_0 - u_1u_2$ contains an odd-length path hanging at a vertex of a triangle, which means $G_0 \notin \mathfrak{T}'$, so it is colorable. Therefore, by induction hypothesis G_0 admits a 3-lic $\phi_{a,b,c}^0$. Since u_1 is a leaf in G_0 , we may assume that $\phi_{a,b,c}^0(u_1v) = c$. By Theorem 3, G_1 admits 3-lic $\phi_{a,b,c}^1$. Since the degree of u_1 in G_1 equals two, we may assume that the colors of edges u_1u_2 and u_1u_3 are from $\{a, b\}$. Therefore, $\phi_{a,b,c}^0 + \phi_{a,b,c}^1$ is a 3-lic of G .

Case 2: $d_G(u_1) \geq 4$. Let H be the subgraph of G induced by the set of all edges incident to u_1 in G . Denote the connected components of $G - E(H)$ in the following way, let G'_0 be the component which contains v and G'_1 the component which contains u_2 and u_3 . Additionally, let $G_1 = G'_1 + u_1u_2$ and $G_0 = G'_0 + u_1v$. We may assume G_0 is colorable, as otherwise G would contain a proper end cycle which is a triangle with the root vertex of degree 3, which would reduce to the previous case. Let $G_2 = G - (E(G_0) \cup E(G_1) \cup E(H))$ and $H' = H - \{u_1u_2, u_1v\}$.

Suppose first that the tree $H' + G_2$ is not colorable. This implies that it is an odd-length path. Notice that $H' + G_2$, as a shrub rooted at u_3 , admits a proper 2-lic $\phi_{a,b}^2$, and since it is proper we have $\phi_{a,b}^2(u_1u_4) = b$. Since G_0 is colorable, by induction hypothesis it admits a 3-lic $\phi_{a,b,c}^0$ where we may assume $\phi_{a,b,c}^0(u_1v) = a$. Since G_1 is a shrub rooted at u_1 with the root edge u_1u_2 , it admits a 2-lic $\phi_{a,b}^1$. If $\phi_{a,b}^1$ is a proper 2-lic of G_1 , then

$$\phi_{a,b,c}(e) = \begin{cases} c & \text{if } e = u_1u_2 \text{ or } u_1u_3, \\ (\phi_{a,b,c}^0 + \phi_{a,b}^1 + \phi_{a,b}^2)(e) & \text{if } e \in E(G) \setminus \{u_1u_2, u_1u_3\}, \end{cases}$$

is a 3-lic of G . Otherwise, if $\phi_{a,b}^1$ is a 2-lic of G_1 , then let us consider the graph $G''_0 = G_0 + u_1u_3$. It is colorable by the same argument as G_0 , so it admits a 3-lic $\phi_{a,b,c}''$ in which u_1u_3 and u_1v must be colored by a same color, say c . Then

$$\phi_{a,b,c}(e) = \begin{cases} \phi_{a,b,c}''(e) & \text{if } e \in E(G''_0), \\ (\phi_{a,b}^1 + \phi_{a,b}^2)(e) & \text{if } e \in E(G) \setminus E(G''_0), \end{cases}$$

is a 3-lic of G .

Suppose now that $H' + G_2$ is a colorable tree, so it admits a 3-lic $\phi_{a,b,c}^2$. We may assume that $\phi_{a,b,c}^2(e) = c$ for every $e \in E(H')$, as this follows either from $d_{H'}(u_1) = 2$ or from Lemma 2 applied to H' as a spidey and every connected component of G_2 as K . As for G_0 , recall that it is colorable, so by induction hypothesis, it has a 3-lic $\phi_{a,b,c}^0$. Since u_1 is a leaf in G_0 , we may assume $\phi_{a,b,c}^0(u_1v) = a$. Let us now consider the graph G_1 . Recall that it is a shrub rooted at u_1 with the root edge u_1u_2 . Hence, by Theorem 1 the graph G_1 admits a 2-alic $\phi_{a,b}^1$. If $\phi_{a,b}^1$ is a 2-lic of G_1 , then $\phi_{a,b,c}^0 + \phi_{b,a}^1 + \phi_{a,b,c}^2$ is a 3-lic of G . Otherwise, we define $H'' = H' + u_1u_2$, and notice that H'' is a spidey. According to Lemma 2, the graph $G_2'' = H'' + G_2$ admits a 3-lic $\phi_{a,b,c}''$ such that $\phi_{a,b,c}''(e) = c$ for every $e \in E(H'')$. We conclude that

$$\phi_{a,b,c}(e) = \begin{cases} \phi_{a,b,c}''(e) & \text{if } e \in E(G_2''), \\ (\phi_{a,b,c}^0 + \phi_{a,b}^1)(e) & \text{if } e \in E(G) \setminus E(G_2''), \end{cases}$$

is a 3-lic of G . \square

Let us now consider the so called *bow-tie* graph B shown in Figure 4. This is a cactus graph with four cycles, but in which cycles are not vertex disjoint. This graph is colorable and admits the 4-lic shown in Figure 4, but it does not admit a k -lic for $k \leq 3$ since the two end vertices of the cut edge must have the degree three in the color of that edge. Hence, for the bow-tie graph B it holds that $\chi'_{\text{irr}}(B) = 4$. We conclude that Conjecture 2 does not hold in general.




Figure 4. The bow-tie graph B and a 4-lic of it.

The consideration of the bow-tie graph gives rise to the following questions: are there any other graphs for which Conjecture 2 does not hold, do all colorable cacti admit a 4-lic, what is the tight upper bound on $\chi'_{\text{irr}}(G)$ of general graphs? We believe the following conjecture holds, which is a weaker form of the Local Irregularity Conjecture.

Conjecture 3. Every connected graph G which does not belong to \mathfrak{T}' satisfies $\chi'_{\text{irr}}(G) \leq 4$.

5. Concluding Remarks

It was established in [9] that a graph does not admit a locally irregular edge coloring if and only if it belongs to the family \mathfrak{T}' . Additionally, it was further conjectured that all other graphs admit a locally irregular 3-edge coloring, see Conjecture 2. As the conjecture is valid for graphs with large minimum degree and all non-colorable graphs are vertex disjoint cacti, we took direction to study cacti.

In this paper we first consider trees to obtain an auxiliary result, then we considered unicyclic graphs and we established that every unicyclic graph $G \notin \mathfrak{T}'$ admits a locally irregular 3-edge coloring as Conjecture 2 claims. Moreover, we have remarked that the bound 3 is tight also on a family of unicyclic graphs, distinct from cycles of the length $4k + 2$, for which it was previously known.

We further extended the result to cacti with vertex disjoint cycles, showing that Conjecture 2 also holds for this class of graphs. Then we gave the so called bow-tie graph B in which cycles are not vertex disjoint, which is colorable and admits a 4-lic, but it does not

admit a locally irregular 3-edge coloring. This graph is a counterexample for Conjecture 2. So the conjecture does not hold in general, but possibly this is a lonely counterexample. We concluded the paper by conjecturing that all colorable graphs admits a locally irregular 4-edge coloring.

Author Contributions: Conceptualization, J.S. and R.Š.; methodology, J.S. and R.Š.; investigation, J.S. and R.Š.; writing—original draft preparation, J.S.; software, R.Š. All authors have read and agreed to the published version of the manuscript.

Funding: Both authors acknowledge partial support of the Slovenian research agency ARRS program P1-0383, ARRS projects J1-1692 and J1-3002. The first author also the support of Project KK.01.1.1.02.0027, a project co-financed by the Croatian Government and the European Union through the European Regional Development Fund—the Competitiveness and Cohesion Operational Programme.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Karoński, M.; Łuczak, T.; Thomason, A. Edge weights and vertex colours. *J. Combin. Theory Ser. B* **2004**, *91*, 151–157. [CrossRef]
2. Addario-Berry, L.; Dalal, K.; McDiarmid, C.; Reed, B.A.; Thomason, A. Vertex-Colouring Edge-Weightings. *Combinatorica* **2007**, *27*, 1–12. [CrossRef]
3. Kalkowski, M. A Note on 1,2-Conjecture. Manuscript. Ph.D. Thesis, Adam Mickiewicz University, Poznan, Poland 2009.
4. Przybyło, J. The 1-2-3 Conjecture almost holds for regular graphs. *J. Combin. Theory Ser. B* **2021**, *147*, 183–200. [CrossRef]
5. Przybyło, J.; Woźniak, M. On a 1,2 Conjecture. *Discrete Math. Theor. Comput. Sci.* **2010**, *12*, 101–108.
6. Wang, T.; Yu, Q. On vertex-coloring 13-edge-weighting. *Front. Math. China* **2008**, *3*, 581–587. [CrossRef]
7. Seamone, B. The 1-2-3 Conjecture and Related Problems: A Survey, Technical Report. 2012. Available online: <http://arxiv.org/abs/1211.5122> (accessed on 15 September 2021).
8. Kalkowski, M.; Karoński, M.; Pfender, F. Vertex-coloring edge-weightings: towards the 1-2-3-conjecture. *J. Combin. Theory Ser. B* **2010**, *100*, 347–349. [CrossRef]
9. Baudon, O.; Bensmail, J.; Przybyło, J.; Woźniak, M. On decomposing regular graphs into locally irregular subgraphs. *Eur. J. Combin.* **2015**, *49*, 90–104. [CrossRef]
10. Addario-Berry, L.; Aldred, R.E.L.; Dalal, K.; Reed, B.A. Vertex colouring edge partitions. *J. Combin. Theory Ser. B* **2005**, *94*, 237–244. [CrossRef]
11. Bensmail, J.; Merker, M.; Thomassen, C. Decomposing graphs into a constant number of locally irregular subgraphs. *Eur. J. Combin.* **2017**, *60*, 124–134. [CrossRef]
12. Lužar, B.; Przybyło, J.; Soták, R. New bounds for locally irregular chromatic index of bipartite and subcubic graphs. *J. Comb. Optim.* **2018**, *36*, 1425–1438. [CrossRef]
13. Baudon, O.; Bensmail, J.; Sopena, É. On the complexity of determining the irregular chromatic index of a graph. *J. Discret. Algorithms* **2015**, *30*, 113–127. [CrossRef]
14. Przybyło, J. On decomposing graphs of large minimum degree into locally irregular subgraphs. *Electron. J. Combin.* **2016**, *23*, 2–31. [CrossRef]