



Lower-Estimates on the Hochschild (Co)Homological Dimension of Commutative Algebras and Applications to Smooth Affine Schemes and Quasi-Free Algebras

Anastasis Kratsios 问

Article

Department of Mathematics, ETH Zürich, Rämistrasse 101, 8092 Zurich, Switzerland; anastasis.kratsios@math.ethz.ch; Tel.: +41-078-933-99-19

Abstract: The Hochschild cohomological dimension of any commutative k-algebra is lower-bounded by the least-upper bound of the flat-dimension difference and its global dimension. Our result is used to show that for a smooth affine scheme *X* satisfying Pointcaré duality, there must exist a vector bundle with section *M* and suitable *n* which the module of algebraic differential *n*-forms $\Omega^n(X, M)$. Further restricting the notion of smoothness, we use our result to show that most *k*-algebras fail to be smooth in the quasi-free sense. This consequence, extends the currently known results, which are restricted to the case where $k = \mathbb{C}$.

Keywords: hochschild cohomology; homological dimension theory; non-commutative geometry; quasi-free algebras; pointcaré duality; higher differential forms



Citation: Kratsios, A.

Lower-Estimates on the Hochschild (Co)Homological Dimension of Commutative Algebras and Applications to Smooth Affine Schemes and Quasi-Free Algebras. *Mathematics* 2021, 9, 251. https://doi.org/10.3390/ math9030251

Academic Editor: Sonia Pérez-Díaz Received: 12 January 2021 Accepted: 25 January 2021 Published: 27 January 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https://creativecommons.org/licenses/by/4.0/).

1. Introduction

Non-commutative geometry is a rapidly developing area of contemporary mathematical research that studies non-commutative algebras using formal geometric tools. The field traces its most evident origins back to the results of [1], which show that any compact Hausdorff space can be fully reconstructed, and largely understood, from its associated C^* -algebra of functions C(X). However, the trend of understanding geometric properties via algebraic dual theories is echoed throughout mathematics; with notable examples coming from the duality between finitely generated algebras and affine schemes (see [2]), the description of any smooth manifold M through its commutative algebra $C^{\infty}(M)$, and ultimately culminating with the work of [3,4] describing the duality relationship between algebra and geometry in full generality.

Though a large portion of the interest in non-commutative geometry stems from its connections with physics, see [5–7]. A. Connes largely made these connections through the cyclic cohomology theory of [8], a generalized de Rham cohomology theory for non-commutative spaces, which closely tied through the Connes complex to one of the central tools of non-commutative geometry and the central object of study of this paper, namely *Hochschild (co)homology*.

Hochschild (co)homology, originally introduced in [9], is a cohomology theory for non-commutative *k*-algebras. Since its introduction, it has become a key tool and object of study in non-commutative geometry since the results of [10] (and more recently generalized in [11] to characteristic *p* fields); which identifies the Hochschild homology of commutative *k*-algebras over a characteristic 0 field *k*, to the module of Khäler differentials over their associated affine scheme. Likewise, the result identifies Hochschild's cohomology theory with the modules of derivations and, therefore, with the tangential structure over the commutative algebra's associated affine scheme. Likewise, in these cases, Pointcaré dualitylike results can also be entirely formulated between these structures and the Hochschild (co)homology theories as shown in [12]. This article focuses on a fundamental non-commutative geometric invariant derived from the Hochschild (co)homology, namely its (co)homological *dimension*. We focus on the interplay between this (co)homological invariant of commutative *k*-algebras over general commutative rings k, and its implications on various notions of smoothness of its associated dual non-commutative space; such as the quasi-freeness (or formal smoothness) of [13,14], or more generally, the vanishing of their higher modules of differential forms as seen in [12].

The relationship between the Hochschild (co)homology theory and smoothness has seen study in the case where k is a field in [15,16]. However, the general case is still far from understood and this is likely due to it requiring a more subtle treatment offered by the less-standard tools of relative homological-algebraic (see [17,18] for example). Indeed, this paper proposes a set of lower-estimates of this invariant, which can be easily computed from local data of any commutative k-algebra over a commutative ring k with unity.

The paper's main results are used to show that for any smooth affine scheme *X* there must exist a vector bundle on *X* with section *M* and a suitably small natural number *n* for which the module of algebraic differential *n*-forms with values in *M*, denoted by $\Omega^n(X, M)$ is non-trivial. Our results are also used to derive simple tests for a *k*-algebras' quasi-freeness. This latter application extends known results of [14] in the special case where $k = \mathbb{C}$. Using this result, we conclude that typical *k*-algebras are not quasi-free. Concrete applications are considered within the scope of arithmetic geometry.

Organization of the Paper

The paper is organized as follows. Section 2 contains the paper's main theorems as well as its non-commutative geometric questions consequences. Each result is followed by examples which unpack the general implications in the context of algebraic geometry. Appendix A contains detailed background material in the relative homological algebraic tools required for the paper's proofs is included after the paper's conclusion. Likewise, the paper's proofs and any auxiliary technical lemma is also relegated to Appendices B–D.

2. Main Result

From here on out, *A* will always be a commutative *k*-algebra. The remainder of this paper will focus on establishing the following result. An analogous statement was made in [14] that all affine algebraic varieties over \mathbb{C} of dimension at greater than 1 fail to have a quasi-free \mathbb{C} -algebra of functions. Once, the assumption that $k = \mathbb{C}$ is relaxed, we find an analogous claim is true; however, the analysis is more delicate. Our principle result is the following.

Theorem 1 (Lower-Bound on Hochschild Cohomological Dimension). Let A be a commutative k-algebra and m be a non-zero maximal ideal in A such that $A_{\mathfrak{m}}$ is has finite $k_{i^{-1}[\mathfrak{m}]}$ -flat dimension and $D(k_{i^{-1}[\mathfrak{m}]})$ is finite. Then:

$$fd_{A_{\mathfrak{m}}}(M_{\mathfrak{m}}) - D(k_{i^{-1}[\mathfrak{m}]}) - fd_{k_{\mathfrak{m}}}(A_{\mathfrak{m}}) \leq HCdim(A|k)$$

Theorem 1 allows for an easily computable lower-bound on the Hochschild cohomological dimension of nearly any commutative *k*-algebra *A*, granted that it is smooth in the classical sense at-least at one point. The next result, obtains an even simpler criterion under the additional assumption that *A* is *k*-flat.

Theorem 2. Let *k* be of finite global dimension, *A* be a *k*-algebra which is flat as a *k*-module. Then *M*:

$$fd_A(M) - D(k) \le HCdim(A|k).$$
(1)

Example 1. Let A be a commutative k-algebra and \mathfrak{m} be a non-zero maximal ideal in A such that $A_{\mathfrak{m}}$ is has finite $k_{i^{-1}[\mathfrak{m}]}$ -flat dimension, $D(k_{i^{-1}[\mathfrak{m}]})$, and A is Cohen-Macaulay at some maximal ideal \mathfrak{m} . Then

$$Krull(A_{\mathfrak{m}}) - D(k_{i^{-1}[\mathfrak{m}]}) - fd_{\mathfrak{m}}(A_{\mathfrak{m}}) \leq HCdim(A|k).$$

Example 2. Let k be of finite global dimension, A be a k-algebra which is flat as a k-module. Then, for every A-module M, if $x_1, ..., x_n$ is a regular sequence in A then:

$$n - D(k) \le HCdim(A|k).$$
⁽²⁾

Furthermore if A is commutative and Cohen-Macaulay at a maximal ideal m then:

$$Krull(A_{\mathfrak{m}}) - D(k) \le HCdim(A|k).$$
 (3)

Next, we consider the implications of our dimension-theoretic formulas within the scope of algebraic geometry from the non-commutative geometric vantage-point.

2.1. Non-Triviality of Higher Differential Forms

The paper's provides a homological argument showing that a smooth affine scheme must have some non-trivial module of higher-differential forms. These begin with the non-triviality of the Hochschild homology modules.

To show our result, we begin by recalling the terminology introduced in [12]. Recall that a *k*-algebra is satisfies *Pointcaré duality in dimension d* if the *dualising module* $\omega_A \triangleq Ext^d_{\mathcal{E}^k_A}(A, A)$ satisfies $Ext^i_{\mathcal{E}^k_A}(A, k) = 0$ for every $i \neq d$ and if in addition $pd_{\mathcal{E}^k_A}(\omega_A) < \infty$. We also recall that an *A*-bimodule *M* is invertible if and only if there exits another *A*-bimodule, which we denote by M^{-1} , for which $M \otimes_A M^{-1} \cong M^{-1} \otimes_A M \cong A$ in $_A Mod_A$.

Corollary 1 (Non-Triviality of Hochschild Homology Modules). Let *k* be a commutative ring and *X* be a *d*-dimensional smooth affine scheme over *k* whose coordinate ring satisfies Pointcaré duality in dimension *d* and is invertible. Then, there is an *A*-bimodule *M* and some $0 \le n \le d - fd_A(M) + D(k)$ satisfying

$$HH_n(A, M) \cong 0.$$

On applying the Hochschild-Kostant-Rosenberg Theorem to Corollary 1, we immediately obtain the claimed result. Recall that $\Omega^n(X, M)$ denotes the algebraic differential *n*-forms on the affine scheme *X* with coefficients in the vector bundle whose section is the k[A]-bimodule *M*.

Corollary 2. Let *k* be a commutative ring and *X* be a *d*-dimensional smooth affine scheme over *k* whose coordinate ring satisfies Pointcaré duality in dimension *d* and is invertible. Then, there exists a some $0 \le n \le d - fd_A(M) + D(k)$ and a vector bundle whose section is the k[A]-module *M* for which the algebraic differential *n*-forms for which

$$\Omega^n(X, M) \not\cong 0.$$

Next, we use Theorem 1 to demonstrate the rarity of commutative quasi-free *k*-algebras.

2.2. Quasi-Free Algebras are Uncommon

Corollary 3 (Krull Dimension-Theoretic Criterion for Quasi-Freeness). Let A be a commutative k-algebra and m be a non-zero maximal ideal in A such that $A_{\mathfrak{m}}$ is has finite $k_{i^{-1}[\mathfrak{m}]}$ -flat dimension, $D(k_{i^{-1}[\mathfrak{m}]})$, and A is Cohen-Macaulay at some maximal ideal m. Then, A is not Quasi-free if

$$Krull(A_{\mathfrak{m}}) \leq 2 + D(k_{i^{-1}[\mathfrak{m}]}) - fd_{\mathfrak{m}}(A_{\mathfrak{m}}).$$

Let us also consider the simpler form implied by Theorem 2.

We unpack Theorem 2 in the context of classical algebraic and arithmetic geometry.

Examples

To build intuition before proceeding, we consider a counter-intuitive consequence. Namely, that most examples of smooth commutative algebras fail to be quasi-free, even when $k \neq \mathbb{C}$. This makes smoothness, in the sense of [14], very rare in the non-commutative category. The following example from arithmetic geometry is of interest.

Let be an affine algebraic \mathbb{C} -variety V(A). For any point x in V(A) the ideal generated by the collection of regular functions on V(A) vanishing at the point x is denoted by $\mathscr{I}(x)$; in fact $\mathscr{I}(x)$ is a maximal ideal in A [19]. Moreover, for any affine-algebraic variety V(A)there exists a point x such that $A_{\mathscr{I}(x)}$ is regular. Since every regular local \mathbb{C} -algebra is Cohen Macaulay at its maximal ideal, then A is Cohen-Macaulay at $\mathscr{I}(x)$. Since \mathbb{C} is a field it is a regular local ring of Krull dimension 0; the Auslander-Buchsbaum-Serre theorem thus implies D(k) = Krull(k) = 0, moreover $A_{\mathscr{I}(x)}$ is a \mathbb{C} -vector space whence it is a \mathbb{C} -free and so is a \mathbb{C} -flat module. Therefore Theorem 2 applies if $Krull(A) \ge 2$. We summarize this finding as follows.

Corollary 5. If X is an affine \mathbb{C} -variety and k[A]'s Krull dimension is greater than 1 then the \mathbb{C} -algebra A is not quasi-free

Remark 1. Corollary 5 implies that any affine algebraic \mathbb{C} -variety which is not a disjoint union of curves or points has a coordinate ring which fails to be quasi-free over \mathbb{C} .

Example 3. The \mathbb{C} -algebra $\mathbb{C}[x_{1,1}, x_{1,2}, x_{2,1}, x_{2,2}]_{(det)}$ is not quasi-free.

Proof. $\mathbb{C}[x_{1,1}, x_{1,2}, x_{2,1}, x_{2,2}]_{(det)}$ is of Krull dimension 4 > 1 [20] therefore Theorem 2 applies. \Box

Corollary 6 (Arithmetic Polynomial-Algebras). *The* \mathbb{Z} *-algebra* $\mathbb{Z}[x_1, ..., x_n]$ *fails to be quasi-free for values of* n > 1.

Proof. Since $\mathbb{Z}[x_1, ..., x_n]$ is Cohen-Macaulay at the maximal ideal $(x_1, ..., x_n, p)$ and is of Krull dimension $n + 1 = Krull(\mathbb{Z}[x_1, ..., x_n])$. Moreover, one computes that $D(\mathbb{Z}) = 1$. Whence by point 2 of Theorem 2: $\mathbb{Z}[x_1, ..., x_n]$ fails to be Quasi-free if $2 \le Krull(\mathbb{Z}[x_1, ..., x_n]) - D(\mathbb{Z}) = (n + 1) - 1 = n$. \Box

The contributions of the paper are now summarized.

3. Conclusions

This paper's main result derived a general lower bound on the Hochschild cohomological dimension of an arbitrary commutative k-algebra A over a general commutative ring k. Theorem 1 derived, the lower-bound for this (co)homological invariant was expressed in terms of other (co)homological dimension-theoretic invariants, namely the flat dimension over A, the global dimension of A, and the flat dimension of A over k; where each quantity was appropriately localized. Examples 1 and 2, built on these results to lower-bound the Hochschild cohomological dimension purely in terms of easily computable quantities, such as the Krull dimension, when A was Cohen-Macaulay. Theorem 2 then expresses a non-localized analog of Theorem 1 wherein no commutativity of A was required.

The paper's results have then been applied the results to purely geometric questions. First, the dimension-theoretic formula was used in Corollary 2 to show infer the nontriviality of certain higher algebraic differential forms of any smooth affine scheme with values in a vector bundle with a non-trivial section. The dual result was also considered in Corollary 1 where dimension-theoretic conditions were obtained for the non-vanishing of some of the Hochschild homology modules under Pointcaré duality in the sense of [12].

Next, using the general (co)homological dimension-theoretic estimates, a result of [14], which showed that most commutative affine k-algebras fail to be smooth in the noncommutative sense formalized by quasi-freeness, was extended from the simple case where k was a field to the general case where k is simply a commutative ring. Specifically, in Corollaries 3 and 4, easily applicable dimension-theoretic tests for the non-quasi-freeness (non-formal smoothness) of a commutative k-algebra over a general ring k were derived. The tools are simple and only require a simple computation involving the Krull dimension of A, the flat-dimension of k at one point, and the base ring's global dimension to identify if A's associated non-commutative space is quasi-free or not.

Author Contributions: All relevant work was carried out by the author. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the ETH Zürich Foundation.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not Applicable.

Acknowledgments: I would like to thank Abraham Broer for the numerous helpful and insightful conversations.

Conflicts of Interest: The author declares no conflict of interest.

Appendix A. Background

This appendix contains the necessary background material for the formulation of this paper's main results. We refer the reader in further reading to the notes of [21].

Appendix A.1. Relative Homological Algebra

The results in this paper are formulated using the *relative homological algebra*, see [17] for example. The theory is analogous to standard homological algebra; see [22] for example, but in this case, one builds the entire theory relative to a suitable subclass of epi(resp. mono)-morphisms. In our case, these are defined as follows.

Definition A1 (\mathscr{E}_A^k -Epimorphism). For any k-algebra A, an epimorphism ϵ in $_A$ Mod is an \mathscr{E}_A^k -epimorphism if and only if ϵ 's underlying morphism of k-modules is a k-split epimorphism in $_k$ Mod. The class of these epimorphisms is denoted \mathscr{E}_A^k .

Definition A2 (\mathscr{E}_{A}^{k} -Exact sequence). An exact sequence of A-modules:

$$.. \xrightarrow{\phi_{i-1}} M_i \xrightarrow{\phi_i} M_{i+1} \xrightarrow{\phi_{i+1}} M_{i+2} \xrightarrow{\phi_{i+2}} ...$$
(A1)

is said to be \mathscr{E}_A^k -exact if and only if for every integer *i* the there exists a morphism of *k*-modules $\psi_i : M_{i+1} \to M_i$ such that:

$$\phi_i = \phi_i \circ \psi_i \circ \phi_i. \tag{A2}$$

In particular, a short exact sequence of A-modules which is \mathscr{E}^k_A -exact is called an \mathscr{E}^k_A -short exact sequence.

Remark A1. Property (A2) is called $\mathcal{E}_{A^e}^k$ -admissibility [18]. Alternatively, it is called $\mathcal{E}_{A^e}^k$ -allowable [23].

Example A1. The augmented bar complex $\hat{CB}_{\star}(A)$ of a k-algebra A is $\mathscr{E}_{A^e}^k$ -exact.

$$0 \to M \xrightarrow{\eta} N \xrightarrow{\epsilon} N' \to 0 \tag{A3}$$

the sequence of k-modules:

$$0 \to Hom_A(P, M) \xrightarrow{\eta^*} Hom_A(P, N) \xrightarrow{\epsilon^*} Hom_A(P, N') \to 0$$
(A4)

is exact.

Remark A2. This definition is equivalent to requiring that P verify the universal property of projective modules only on \mathcal{E}_A^k -epimorphisms [23].

Example A2. $A^{\otimes n+2}$ is $\mathscr{E}^k_{A^e}$ -projective for all $n \in \mathbb{N}$.

 \mathscr{E}_{A}^{k} -projective *A*-modules have analogous properties to projective *A*-modules. For example, \mathscr{E}_{A}^{k} -projective *A*-modules admit the following characterization.

Proposition A1. For any A-module P the following are equivalent:

- \mathscr{E}_A^k -Short exact sequence preservation property P is \mathscr{E}_A^k -projective.
- $\mathscr{E}_{A}^{\vec{k}}$ -lifting property For every \mathscr{E}_{A}^{k} -epimorphism $f : N \to M$ if there exists an A-module morphism $g : P \to M$ then there exists an A-module map $\tilde{f} : P \to N$ such that $f \circ \tilde{f} = g$.

$$\mathfrak{E}_{\pi}: 0 \to M \to N \to P \to 0 \tag{A5}$$

is A-split-exact.

 \mathscr{E}^k_A -free direct summand property There exists a k-module F, an A-module Q and an isomorphism of A-modules $\phi : P \oplus Q \xrightarrow{\cong} A \otimes_k F$.

Remark A3. If *F* is a free *k*-module, some authors call $A \otimes_k F$ an \mathscr{E}^k_A -free module. In fact this gives an alternative proof that $A^e \otimes_k A^{\otimes n} \cong A^{\otimes n+2}$ is $\mathscr{E}^k_{A^e}$ -free for every $n \in \mathbb{N}$.)

Proof. See [23] pages 261 for the equivalence of 1, 2 and 3 and page 277 for the equivalence of 1 and 4. □

For a homological algebraic theory to be possible, one needs enough projective (resp. injective) objects. The next result shows that there are indeed enough \mathscr{E}_A^k -projectives in $_AMod$.

Proposition A2 (Enough \mathscr{E}^k_A -projectives). If A is a k-algebra and M is an A-module then there exists an \mathscr{E}^k_A -epimorphism $\epsilon : P \to M$ where P is an \mathscr{E}^k_A -projective.

Proof. By Proposition A1 $A \otimes_k M$ is \mathscr{E}_A^k -projective. Moreover, the A-map $\zeta : A \otimes_k M \to M$ described on elementary tensors as $(\forall a \otimes_k m \in A \otimes_k M)\zeta(a \otimes_k m) := a \cdot m$ is epi and is k-split by the section $m \mapsto 1 \otimes_k m$. \Box

Since there are enough projective objects, then one can build a resolution of any *A*-module by \mathscr{E}_A^k -projective modules.

Definition A4 (\mathscr{E}_A^k -projective resolution). If M is an A^e -module then a resolution P_* of M is called an \mathscr{E}_A^k -projective resolution of M if and only if each P_i is an \mathscr{E}_A^k -projective module and P_* is an \mathscr{E}_A^k -exact sequence.

Example A3. The augmented bar complex $\hat{CB}_{\star}(A)$ of A is an \mathscr{E}^{k}_{Ae} -projective resolution of A.

Remark A4. A nearly completely analogous argument to Example A3 shows that for any (A, A)-bimodule $M, M \otimes_A CB_*(A)$ is an $\mathcal{E}^k_{A^e}$ -projective resolution of M, see for details [24].

Following [18], the \mathscr{E}_A^k -relative derived functors of the tensor product and the Hom_A -functors are introduced, as follows.

Definition A5. \mathscr{E}^k_A -relative Tor

If N is a right A-module, M is an A-module and P_* is an \mathscr{E}^k_A -projective resolution of N then the k-modules $H_*(P_* \otimes_A M)$ are called the \mathscr{E}^k_A -relative Tor k-modules of N with coefficients in the A-module M and are denoted by $\operatorname{Tor}^n_{\mathscr{E}^k}(N, M)$.

Let H_{\star} (resp. H^{\star}) denote the (co)homology functor from the category of chain (co)complexes on an *A*-module to the category of *A*-modules. The \mathscr{E}_{A}^{k} -relative Tor functors are defined as follows.

Example A4. The \mathscr{E}^k_A -relative Tor functors may differ from the usual (or "absolute") Tor functors. For example consider all the \mathbb{Z} -algebra \mathbb{Z} , any \mathbb{Z} -modules N and M are $\mathscr{E}^{\mathbb{Z}}_{\mathbb{Z}}$ -projective. In particular, this is true for the \mathbb{Z} -modules \mathbb{Z} and $\mathbb{Z}/2\mathbb{Z}$. Therefore $\operatorname{Tor}^n_{\mathscr{E}^{\mathbb{Z}}_{\mathbb{Z}}}(\mathbb{Z}, \mathbb{Z}/2\mathbb{Z})$ vanish for every positive n,

however $\operatorname{Tor}^n_{\mathbb{Z}}(\mathbb{Z}, \mathbb{Z}/2\mathbb{Z})$ does not. For example, $\operatorname{Tor}^1_{\mathbb{Z}}(\mathbb{Z}, \mathbb{Z}/2\mathbb{Z}) \cong \mathbb{Z}/2\mathbb{Z}$ [22].

Similarly there are \mathscr{E}^k_A -relative Ext functors.

Definition A6 (\mathscr{E}_A^k -relative Ext). If N is and M are A-modules and P_{\star} is an \mathscr{E}_A^k -projective resolution of N then the k-modules $H^{\star}(Hom_A(P_{\star}, M))$ are called the \mathscr{E}_A^k -relative Ext k-modules of N with coefficients in the A-module M and are denoted by $\operatorname{Ext}_{\mathscr{E}_A^k}^n(N, M)$.

The \mathscr{E}_A^k -relative homological algebra is indeed well defined, since both the definitions of \mathscr{E}_A^k -relative Ext and \mathscr{E}_A^k -relative Tor are independent of the choice of \mathscr{E}_A^k -projective resolution.

Theorem A1 (\mathscr{E}_A^k -Comparison theorem). If P_* and P'_* are \mathscr{E}_A^k -projective resolutions of an A-module N then for any A-module M there are natural isomorphisms:

$$H^{\star}(Hom_{\mathcal{E}^{k}_{A}}(P_{\star},N)) \stackrel{\equiv}{\to} H^{\star}(Hom_{\mathcal{E}^{k}_{A}}(P'_{\star},N))$$
(A6)

and if P_{\star} and P'_{\star} are \mathscr{E}^k_A -projective resolutions of a right A-module N then:

$$H_{\star}(P_{\star} \otimes_A N) \xrightarrow{=} H_{\star}(P'_{\star} \otimes_A N) \tag{A7}$$

Proof. Nearly identical to the usual comparison theorem, see [23]. \Box

Example A5. The $Ext_{\mathbb{Z}}$ and $\mathscr{E}_{\mathbb{Z}}^{\mathbb{Z}}$ -relative Ext may differ. For example, one easily computes $Ext_{\mathbb{Z}}^{1}(\mathbb{Z},\mathbb{Z}/2\mathbb{Z}) \cong \mathbb{Z}/2\mathbb{Z}$. However, $Ext_{\mathscr{E}_{\mathbb{Z}}}^{1}(\mathbb{Z},\mathbb{Z}/2\mathbb{Z}) \cong 0$.

Analogous to the fact that for any *A*-module *P*, *P* is projective if and only if $Ext_A^1(P, N) \cong 0$ for every *A*-module *N* there is the following result, which can be found in ([18], Chapter IX).

Proposition A3. *P* is an \mathscr{E}_A^k -projective module if and only if for every *A*-module *N*:

$$Ext^{1}_{\mathscr{E}^{k}}(P,N) \cong 0 \tag{A8}$$

Using the theory of relative (co)homology, we are now in-place to review the Hochschild cohomology theory over general *k*-algebras.

Appendix A.2. Hochschild (Co)homological Dimension

Since $CB_*(A)$ is an $\mathscr{E}_{A^e}^k$ -projective resolution of A then Theorem A1 and the definition of the $Ext_{\mathscr{E}_{A^e}^k}^*(A, -)$ functors imply that the Hochschild cohomology of A with coefficients in of [9], denoted by $HH^*(A, N)$, can be expressed using the $Ext_{\mathscr{E}_{A^e}^k}^*$. We maintain this perspective throughout this entire article.

Proposition A4. For every A^e module N there are k-module isomorphisms, natural in N:

$$HH^{\star}(A,N) \stackrel{\cong}{\to} Ext^{\star}_{\mathscr{E}^{k}}(A,N) \tag{A9}$$

Taking short $\mathscr{E}^k_{A^e}$ -exact sequences to isomorphic long exact sequences.

Definition A7 (Hochschild Homology). *The Hochschild homology* $HH_{\star}(A, N)$ *of a k-algebra A with coefficient in the* (A, A)*-bimodule* N *is defined as:*

$$HH_{\star}(A,N) := H_{\star}(P_{\star} \otimes_A N) \tag{A10}$$

where P_{\star} is an $\mathscr{E}^k_{A^e}$ -projective resolution of A.

Following the results of [10], the Hochschild cohomology has become the central tool for obtaining non-commutative algebraic geometric analogues of classical commutative algebraic geometric notions. The one of central focus in this paper, is the Hochschild cohomological dimension,

Definition A8 (Hochschild cohomological dimension). *The Hochschild cohomological dimension of a k-algebra A is defined as:*

$$HCdim(A|k) := \sup_{M \in {}_{A^e}Mod} (\sup\{n \in \mathbb{N}^{\#}|HH^n(A,M) \not\cong 0\}).$$
(A11)

where $\mathbb{N}^{\#}$ is the ordered set of extended natural numbers.

The Hochschild cohomological dimension may be related to the following cohomological dimension.

Definition A9 (\mathscr{E}^k_A -projective dimension). If *n* is a natural number and *M* is an *A*-module then *M* is said to be of \mathscr{E}^k_A -projective dimension at most *n* if and only if there exists a deleted \mathscr{E}^k_A -projective resolution of *M* of length *n*. If no such \mathscr{E}^k_A -projective resolution of *M* exists then *M* is said to be of \mathscr{E}^k_A -projective dimension ∞ . The \mathscr{E}^k_A -projective dimension of *M* is denoted $pd_{\mathscr{E}^k_A}(M)$.

The following is a translation of a classical homological algebraic result into the setting of $\mathscr{E}_{A^e}^k$ -projective dimension, $\Omega^n(A/k)$ and Hochschild cohomology. Here, $\Omega^n(A/k) \triangleq Ker(b'_{n-1})$ and b'_{n-1} is the $(n-1)^{th}$ differential in the augmented Bar resolution of A; see [24] for details on the augmenter Bar complex.

Theorem A2. For every natural number *n*, the following are equivalent:

- $HCdim(A|k) \le n$
- *A is of* $\mathscr{E}^k_{A^e}$ *-projective dimension at most n*
- $\Omega^n(A/k)$ is an $\mathscr{E}^k_{A^e}$ -projective module.
- $HH^{n+1}(A, M)$ vanishes for every (A, A)-bimodule M.

 $Ext_{\mathcal{E}_{Ae}^{k}}^{n+1}(A, M)$ vanishes for every A^{e} -module M.

Proof. $(1 \Rightarrow 4)$ By definition of the Hochschild cohomological dimension. $(4 \Leftrightarrow 5)$ By Proposition A4. $(3 \Rightarrow 2)$ Since $\Omega^n(A/k)$ is $\mathscr{E}^k_{A^e}$ -projective:

$$0 \to \Omega^n(A/k) \to CB_{n-1}(A) \xrightarrow{b'_{n-1}} \dots \xrightarrow{b'_0} A \to 0$$

is a $\mathscr{E}_{A^e}^k$ -projective resolution of A of length n. Therefore $pd_{\mathscr{E}_{A^e}^k}(A) \leq n$. (3 \Leftrightarrow 4) By Proposition A9 there are isomorphism natural in M:

$$(\forall M \in_{A^e} Mod) HH^{1+n}(A, M) \cong Ext^{1+n}_{\mathscr{E}^k_{A^e}}(A, M) \cong Ext^1_{\mathscr{E}^k_{A^e}}(\Omega^n(A/k), M).$$

Therefore for every A^e -module M:

$$Ext^{1}_{\mathscr{E}^{k}_{A^{e}}}(\Omega^{n}(A/k), M) \cong 0 \text{ if and only if } HH^{1+n}(A, M) \cong 0.$$

By Proposition A3 $\Omega^n(A/k)$ is \mathscr{E}^k_A -projective if and only if $Ext^1_{\mathscr{E}^k_{A^e}}(\Omega^n(A/k), M) \cong 0$.

 $(2 \Rightarrow 1)$ If *A* admits an $\mathscr{E}_{A^e}^k$ -projective resolution P_{\star} of length *n* then Theorem A1 implies there are natural isomorphisms of A^e -modules:

$$(\forall M \in_{A^e} Mod) Ext^*_{\mathscr{E}^k_{A^e}}(A, M) \cong H^*(Hom_{A^e}(P_\star, M)).$$
(A12)

Since P_* is of length n all the maps $p_j : P_{j+1} \to P_j$ are the zero maps therefore so are the maps $p_j^* : Hom_{A^e}(P_j) \to Hom_{A^e}(P_{j+1})$. Whence (A12) entails that for all j > n + 1 $Ext^*_{\mathcal{E}^{A^e}_{A^e}}(A, M)$ vanishes. By Proposition A4 this is equivalent to $HH^j(A, M)$ vanishing for all j > n + 1 for all $M \in_{A^e} Mod$. Hence A is of Hochschild cohomological dimension at most n. \Box

Next, the non-commutative geometric object focused on in this paper is reviewed.

Appendix A.3. Quasi-Free Algebras

Many of the properties of an algebra are summarized by its Hochschild cohomological dimension, see [10,17] for example. However, this article focuses on the following non-commutative analogue of smoothness of [13], introduced by [14].

Remark A5. Due to their lifting property, the quasi-free k-algebras are considered a non-commutative analogue to smooth k-algebras; that is k-algebras for which $\Omega_{A|k}$ is a projective A-module.

This notion of smoothness has played a key role in a number of places in noncommutative algebraic geometry, especially in the cyclic (co)homology of [25].

Definition A10 (Quasi-free *k*-algebra). *A k-algebra for which all k-Hochschild extensions of A by an* (*A*, *A*)-*bimodule lift is called a quasi-free k-algebra*.

Corollary A1. For a k-algebra A, the following are equivalent:

- $A \text{ is } HCdim(A|k) \leq 1.$
- $\Omega^1(A/k)$ is a $\mathscr{E}^k_{A^e}$ -projective A^e -module.
- A is quasi-free.

One typically construct quasi-free algebras using Morita equivalences. However, the next proposition, which extends a result of [14] to the case where k need not be a field, may also be used without any such restrictions on k.

Proposition A5. If A is a quasi-free k-algebra and P is an $\mathscr{E}_{A^e}^k$ -projective (A, A)-bimodule then $T_A(P)$ is a quasi-free A-algebra.

Proof. Differed until the appendix. \Box

Example A6. Let
$$n \in \mathbb{N}$$
. The \mathbb{Z} -algebra $T_{\mathbb{Z}} \begin{pmatrix} n \\ \bigoplus \\ i=0 \end{pmatrix}$ is quasi-free.

Proof. Since all free \mathbb{Z} -modules are projective \mathbb{Z} -modules and all projective \mathbb{Z} -modules are $\mathscr{E}_{\mathbb{Z}}^{\mathbb{Z}}$ -projective modules, the free \mathbb{Z} -module $\bigoplus_{i=0}^{n} \mathbb{Z}$ is $\mathscr{E}_{\mathbb{Z}}^{\mathbb{Z}}$ -projective. Whence Proposition A5 implies $T_{\mathbb{Z}}\left(\bigoplus_{i=0}^{n} \mathbb{Z}\right)$ is a quasi-free \mathbb{Z} -algebra. \Box

Example A7. If A is a quasi-free k-algebra then $T_A(\Omega^1(A/k))$ is a quasi-free A-algebra.

Proof. By Corollary A1 if *A* is quasi-free $\Omega^1(A/k)$ must be an $\mathscr{E}^k_{A^e}$ -projective (A, A)-bimodule; whence Proposition A5 applies. \Box

Next, we overview some relevant dimension-theoretic notions and terminology.

Appendix A.4. Classical Cohomological Dimensions

We remind the reader of a few important algebraic invariants which we will require. The reader unfamiliar with certain of these notions from commutative algebra and algebraic geometry is referred to [2,26] or to [19].

Definition A11 (A-Flat Dimension). If A is a commutative ring then the A-flat dimension $fd_A(M)$ of an A-module M is the extended natural number n, defined as the shortest length of a resolution of M by A-flat A-modules. If no such finite n exists n is taken to be ∞ .

We will require the following result, whose proof can be found in [24].

Proposition A6. If *n* is a positive integer and if there exists a regular sequence $x_1, ..., x_n$ in *A* of length *n* then:

$$n = fd_A(A/(x_1, ..., x_n)).$$
(A13)

One more ingredient related to the flat dimension will soon be needed.

Proposition A7. If A is a commutative ring and m is a maximal ideal of A then for any A-module $M fd_{A_{\mathfrak{m}}}(M_{\mathfrak{m}})$ is a lower-bound for $fd_{A}(M)$.

Definition A12. A-Projective Dimension

If A is a commutative ring and M is an A-module then the A-projective dimension $pd_A(M)$ of M is the extended natural number n, defined as the shortest length of a deleted A-projective resolution of M. If no such finite n exists n is taken to be ∞ .

Lemma A1. If A is a commutative ring and M is an A-module then $fd_A(M) \le pd_A(M)$.

Proof. Since all *A*-projective *A*-modules are *A*-flat, then any *A*-projective resolution is a *A*-flat resolution. \Box

Lemma A2. If A is a commutative ring then for any A-module M the following are equivalent:

- The A-projective dimension of M is at most n.
- For every A-module N, the A-module $Ext_{n+1}^A(M, N)$ is trivial.
- For every A-module N and every integer $m \ge n + 1$: $Ext_m^A(M, N) \cong 0$.

Proof. Nearly identical to the proof of Theorem A2, see page 456 of [22] for details. \Box

Definition A13 (Cohen-Macaulay at an Ideal). *A commutative ring A is said to be Cohen-Macaulay at a maximal ideal* m *if and only if either:*

- *Krull*($A_{\mathfrak{m}}$) *is finite and there is an* $A_{\mathfrak{m}}$ *-regular sequence* $x_1, ..., x_d$ *in* $A_{\mathfrak{m}}$ *of maximal length* $d = Krull(A_{\mathfrak{m}})$ *such that* $\{x_1, ..., x_d\} \subseteq \mathfrak{m}$.
- $Krull(A_m)$ is infinite and for every positive integer d there is an A_m -regular sequence $x_1, ..., x_d$ in \mathfrak{m} on A of length d.

Proposition A8 ([24]). *If* A *is a commutative ring which is Cohen Macaulay at the maximal ideal* \mathfrak{m} *and* $Krull(A_{\mathfrak{m}})$ *is finite then:*

$$Krull(A_{\mathfrak{m}}) = fd_{A_{\mathfrak{m}}}(A_{\mathfrak{m}}/(x_1, .., x_n)) \le pd_A(A_{\mathfrak{m}}/(x_1, .., x_n))$$
(A14)

Definition A14. Global Dimension

The *global dimension* D(A) of a ring A, is defined as the supremum of all the A-projective dimensions of its A-modules. That is:

$$D(A) := \sup_{M \in {}_{A}Mod} pd_{A}(M).$$
(A15)

The following modification of the global dimension of a *k*-algebra, does not ignore the influence of *k* on a *k*-algebra *A*, as will be observed in the next section of this paper.

Definition A15. \mathscr{E}^k -Global dimension

The \mathscr{E}^k -global Dimension $D_{\mathscr{E}^k}(A)$ of a k-algebra A is defined as the supremum of all the \mathscr{E}^k_A -projective dimensions of its A-modules. That is:

$$D_{\mathscr{E}^k}(A) := \sup_{M \in {}_A Mod} pd_{\mathscr{E}^k_A}(M).$$
(A16)

Appendix B. Proofs

This appendix contains certain technical lemmas or auxiliary results that otherwise detracted from the overall flow of the paper.

Appendix C. Technical Lemmas

We make use of the following result appearing in a technical note of Hochschild circa 1958, see [27].

Theorem A3 ([27]). *If k is of finite global dimension, A is a k-algebra which is flat as a k-module and M is an A-module then:*

$$pd_A(M) - D(k) \le pd_{\mathcal{E}_A^k}(M) \tag{A17}$$

Proposition A9 (Dimension Shifting). If

$$\dots \xrightarrow{d_{n+1}} P_{n+j}P_n \xrightarrow{d_n} \dots \xrightarrow{d_2} P_1 \xrightarrow{d_1} P_0 \to 0$$
(A18)

is a deleted \mathscr{E}_A^k -projective resolution of an A-module M then for every A-module N and for every positive integer n there are isomorphisms natural in N:

$$Ext^{1}_{\mathscr{C}^{k}_{A}}(Ker(d_{n}), N) \cong Ext^{n+1}_{\mathscr{C}^{k}_{A}}(A, N)$$
(A19)

Proof. By definition the truncated sequence is exact:

$$\dots \xrightarrow{d_{n+j}} P_{n+j} \xrightarrow{d_{n+j-1}} \dots \xrightarrow{d_{n+1}} P_{n+1} \xrightarrow{\eta} Ker(d_n) \to 0,,$$
(A20)

where η is the canonical map satisfying $d_n = ker(d_n) \circ \eta$ (arising from the universal property of $ker(d_n)$). Moreover, since (A30) is \mathscr{E}_A^k -exact, d_n is k-split; whence η must be k-split. Moreover, for every $j \ge n + 1$, d_j was by assumption k-split therefore (A20) is \mathscr{E}_A^k -exact and since for every natural number $m > n P_m$ is by hypothesis \mathscr{E}_A^k -projective then (A20) is an augmented \mathscr{E}_A^k -projective resolution of the A-module $Ker(d_n)$.

For every natural number *m*, relabel:

$$Q_m := P_{m+n} \text{ and } p_m := d_{n+m}. \tag{A21}$$

By Theorem A1, for all $N \in_A Mod$ and all $m \in \mathbb{N}$, we have that:

$$Ext_{\mathscr{E}_{A}^{k}}^{m}(Ker(d_{n}), N) \cong H^{m}(Hom_{A}(Q_{\star}, N))$$

$$= Ker(Hom_{A}(p_{n}, N)) / Im(Hom_{A}(p_{n+1}, N))$$

$$= Ker(Hom_{A}(d_{n+m}, N)) / Im(Hom_{A}(d_{n+m+1}, N))$$

$$= H^{m+n}(Hom_{A}(P_{\star}, N))$$

$$\cong Ext_{\mathscr{E}_{A}^{k}}^{m}(A, N).$$
(A22)

Therefore, the result follows. \Box

Appendix D. Auxiliary Results Proof of Proposition A5. Let

$$0 \to M \to B \xrightarrow{\pi} T_A(P) \to 0 \tag{A23}$$

be a *k*-Hochschild extension of $T_A(P)$ by *M*. We use the universal property of $T_A(P)$ to show that there must exist a lift *l* of (A23).

Let $p : T_A(P) \to A$ be the projection *k*-algebra homomorphism of $T_A(P)$ onto *A*. *p* is *k*-split since the *k*-module inclusion $i : A \to T_A(P)$ is a section of *p*; therefore *p* is an $\mathscr{E}^k_{A^e}$ -epimorphism and

$$0 \to Ker(p \circ \pi) \to B \to A \to 0$$
 (A24)

is a *k*-Hochschild extension of *A* by the (A, A)-bimodule $Ker(p \circ \pi)$. Since *A* is a quasifree *k*-algebra there exists a *k*-algebra homomorphism $l_1 : A \to B$ lifting $p \circ \pi$. Hence *B* inherits the structure of an (A, A)-bimodule and π may be viewed as an (A, A)-bimodule homomorphism. Moreover, l_1 induces an *A*-algebra structure on *B*.

Let $f : P \to T_A(P)$ be the (A, A)-bimodule homomorphism satisfying the universal property of the tensor algebra on the (A, A)-bimodule P. Since $\pi : B \to A$ is an $\mathscr{E}_{A^e}^k$ epimorphism and since P is an $\mathscr{E}_{A^e}^k$ -projective (A, A)-bimodule, Proposition A1 implies that that there exists an (A, A)-bimodule homomorphism $l_2 : P \to B$ satisfying $\pi \circ l_2 = f$.

Since $l_2 : P \to B$ is an (A, A)-bimodule homomorphism to a A-algebra the universal property of the tensor algebra $T_A(P)$ on the (A, A)-bimodule P, see [28], implies there is an A-algebra homomorphism $l : T_A(P) \to B$ whose underlying function satisfies: $l \circ f = l_2$.

Therefore $l \circ \pi \circ l_2 = l_2$; whence $l \circ \pi = 1_{T_A(P)}$; that is l is a A-algebra homomorphism which is a section of π , that is l lifts π . \Box

Appendix D.1. Proof of Theorem 1

Our first lemma is a generalization of the central theorem of [27]; which does not rely on the assumption that A is k-flat.

Lemma A3. If *k* is of finite global dimension and *A* is a *k*-algebra which is of finite flat dimension as a *k*-module, then for every *A*-module *M*:

$$pd_A(M) - D(k) - fd_k(A) \le pd_{\mathcal{E}_A^k}(M)$$
(A25)

The proof of Lemma A3 relies on the following lemma.

Lemma A4. If A is a k-algebra such that $fd_k(A) < \infty$ then:

$$(\forall M \in_k Mod) \ pd_A(A \otimes_k M) - fd_k(A) \le pd_k(M) \tag{A26}$$

Proof. For every *k*-module *M* and every *A*-module *N* there is a convergent third quadrant spectral sequence (see [22], page 667):

$$Ext_{A}^{p}(Tor_{q}^{k}(A,M),N) \xrightarrow{p} Ext_{k}^{p+q}(M,Hom_{A}(A,N)).$$
(A27)

Moreover, the adjunction $-\otimes_k A \dashv Hom_A(A, -)$ extends to a natural isomorphism:

$$(\forall p,q \in \mathbb{N}) Ext_k^{p+q}(M, Hom_A(A, N)) \cong Ext_A^{p+q}(M \otimes_k A, N).$$
(A28)

Therefore there is a convergent third-quadrant spectral sequence:

$$Ext_{A}^{p}(Tor_{q}^{k}(A,M),N) \Longrightarrow_{p} Ext_{A}^{p+q}(M \otimes_{k} A,N).$$
(A29)

If $pd_A(N) < \infty$, then the result is immediate. Therefore assume that: $pd_A(N) < \infty$. If $p + q > fd_k(A) + pd_A(N)$ then either $p > pd_A(N)$ or $q > fd_k(A)$. In the case of th

$$0 \cong E_2^{p,q} \cong E_{\infty}^{p,q} \cong Ext_A^{p+q}(M \otimes_k A, N)$$

and in the latter case

$$0 \cong E_2^{p,q} \cong E_\infty^{p,q} \cong Ext_A^{p+q}(M \otimes_k A, N)$$

also. Therefore

$$(\forall N \in A_M od) \ 0 \cong Ext_A^n (M \otimes_k A, N) \text{ if } n > fd_k(A) + pd_A(N);$$

hence: $pd_A(M \otimes_k A) \leq fd_k(A) + pd_A(M)$.

Finally, the result follows since $fd_k(A)$ is finite and, therefore, can be subtracted unambiguously. \Box

Lemma A5. If A is a k-algebra then for any k-module M there is an \mathscr{E}^k_A -exact sequence:

$$0 \to Ker(a) \to A \otimes_k M \xrightarrow{\alpha} M \to 0 \tag{A30}$$

where α be the map defined on elementary tensors $(a \otimes_k m)$ in $A \otimes_k M$ as $a \otimes_k m \mapsto a \cdot m$.

Proof. α is *k*-split by the map $\beta : M \to A \otimes_k M$ defined on elements $m \in M$ as $m \mapsto 1 \otimes_k m$. Indeed if $m \in M$ then:

$$\alpha \circ \beta(m) = \alpha(1 \otimes_k m) = 1 \cdot m = m.$$
(A31)

Lemma A6. If M and N are A-modules then:

$$pd_A(M) \le pd_A(M \oplus N).$$
 (A32)

Proof.

$$(\forall n \in \mathbb{N})(\forall X \in_A Mod) Ext_A^n(M, X) \oplus Ext_A^n(N, X) \cong Ext_A^n(M \oplus N, X).$$
(A33)

Therefore $Ext_A^n(M \oplus N, X)$ vanishes only if both $Ext_A^n(M, X)$ and $Ext_A^n(N, X)$ vanish. Lemma A2 then implies: $pd_A(M) \le pd(M \oplus N)$. \Box

Proof of Lemma A3 **Proof.**

Case 1: $pd_{\mathcal{E}^k_4}(M) = \infty$

By definition $pd_A(M) \leq \infty$ therefore trivially if $pd_{\mathscr{E}_A^k}(M) = \infty$ then:

$$pd_A(M) \le pd_{\mathscr{E}^k}(M) + D(k). \tag{A34}$$

Since *k*'s global dimension is finite hence (A34) implies:

$$pd_A(M) - D(k) \le \infty = pd_{\mathscr{C}_A^k}(M).$$
(A35)

Case 2: $pd_{\mathscr{E}^k_A}(M) < \infty$

Let $d := pd_{\mathcal{E}^k_A}(M) + D(k) + fd_k(A)$. The proof will proceed by induction on d.

Base: d = 0

Suppose $pd_{\mathscr{E}^k}(M) = 0$.

By Theorem A2 *M* is \mathscr{E}_A^k -projective. Lemma A5 implies there is an \mathscr{E}_A^k -exact sequence:

$$0 \to Ker(a) \to A \otimes_k M \xrightarrow{a} M \to 0.$$
(A36)

Proposition A1 implies that (A36) is *A*-split therefore *M* is a direct summand of the *A*-module $A \otimes_k M$. Hence Lemma A6 implies:

$$pd_A(M) \le pd_A(M \otimes_k A).$$
 (A37)

Lemma A4 together with (A37) imply:

$$pd_A(M) \le pd_A(M \otimes_k A) \le pd_k(M).$$
 (A38)

Definition A15 and (A38) together with the assumption that $pd_{\mathcal{E}^k_A}(M) = 0$ imply:

$$pd_A(M) \le pd_k(M) \le D(k) = D(k) + 0 + 0 = D(k) + pd_{\mathscr{E}^k_A}(M) + fd_k(A).$$
 (A39)

Since *k*'s global dimension and $fd_k(A)$ are finite then (A39) implies:

$$pd_A(M) - D(k) - fd_k(A) \le pd_{\mathcal{E}_A^k}(M).$$
(A40)

Inductive Step: d > 0

Suppose the result holds for all *A*-modules *K* such that $pd_{\mathcal{E}_A^k}(K) + D(k) + fd_k(A) = d$ for some integer d > 0. Again appealing to Lemma A5, there is an \mathcal{E}_A^k -exact sequence:

$$0 \to Ker(a) \to A \otimes_k M \xrightarrow{\alpha} M \to 0. \tag{A41}$$

Proposition A1 implies $A \otimes_k M$ is \mathscr{E}_A^k -projective; whence (A41) implies:

$$pd_{\mathcal{E}^k_A}(Ker(\alpha)) + 1 = pd_{\mathcal{E}^k_A}(M).$$
(A42)

$$pd_{A}(Ker(\alpha)) + 1 \leq pd_{\mathscr{E}^{k}_{A}}(Ker(\alpha)) + 1 + D(k) + fd_{k}(A)$$

$$\leq pd_{\mathscr{E}^{k}_{A}}(M) + D(k) + fd_{k}(A).$$
(A43)

The proof will be completed by demonstrating that: $pd_A(M) \le pd_A(Ker(\alpha)) + 1$. For any $N \in_A Mod Ext^*_A(-, N)$ applied to (A41) gives way to the long exact sequence in homology, particularly the following of its segments are exact:

$$Ext_{A}^{n-1}(A \otimes_{k} M, N) \to Ext_{A}^{n-1}(Ker(a), N) \xrightarrow{d^{n}} Ext_{A}^{n}(M, N) \to Ext_{A}^{n}(A \otimes_{k} M, N)$$
(A44)

Since $A \otimes_k M$ is \mathscr{E}^k_A -projective $pd_{\mathscr{E}^k_A}(A \otimes_k M) = 0$, therefore by the base case of the induction hypothesis $pd_A(A \otimes_k M) \leq pd_{\mathscr{E}_A^k} + D(k) + fd_k(A) = D(k) + d_k(A)$ $fd_k(A)$; thus for every positive integer $n \ge D(k)$ (in particular *d* is at least *n*):

$$(\forall N \in_A Mod) Ext_A^{n-1}(A \otimes_k M, N) \cong 0 \cong Ext_A^n(A \otimes_k M, N);$$
(A45)

whence ∂^n must be an isomorphism. Therefore Lemma A2 implies $pd_A(M)$ is at most equal to $pd_A(Ker(\alpha)) + 1$. Therefore:

$$pd_A(M) \le pd_A(Ker(\alpha)) + 1$$

$$d_A(M) \le pd_A(Ker(\alpha)) + 1 \tag{A46}$$

$$\leq p u_{\mathcal{E}_A^k}(\mathbf{R}(u)) + 1 + D(k) + j u_k(A) \tag{A47}$$

$$\leq pd_{\mathscr{E}^k_A}(M) + D(k) + fd_k(A). \tag{A48}$$

Finally since *k* is of finite global dimension and *A* is of finite *k*-flat dimension then (A48) implies:

$$pd_A(M) - D(k) - fd_k(A) \le pd_{\mathscr{E}_A^k}(M);$$
(A49)

thus concluding the induction.

We will also require the following result.

Remark A6. Let A be a k-algebra, $i : k \to A$ the morphism defining the k-algebra A and m a maximal ideal in A. For legibility the $\mathscr{E}_{A_{\mathfrak{m}}}^{k_{i-1}[\mathfrak{m}]}$ -projective dimension of an $A_{\mathfrak{m}}$ -module N will be abbreviated by $pd_{\mathscr{E}_{\mathfrak{m},k}}(N)$ (instead of writing $pd_{\mathscr{E}_{A_{\mathfrak{m}}}^{k_{i-1}[\mathfrak{m}]}}(N)$).

Lemma A7. If A is a commutative k-algebra and \mathfrak{m} is a non-zero maximal ideal in A then for every A-module M:

$$pd_{\mathscr{E}_{\mathfrak{m},k}}(M_{\mathfrak{m}}) \le pd_{\mathscr{E}_{A}^{k}}(M),\tag{A50}$$

where $i: k \to A$ is the inclusion of k into A.

Proof. Since m is a prime ideal in A, $i^{-1}[m]$ is a maximal ideal in $k_{i^{-1}[m]}$, whence the localized ring $k_{i^{-1}[\mathfrak{m}]}$ is a well-defined sub-ring of $A_{\mathfrak{m}}$. Let

$$\dots \xrightarrow{d_{n+1}} P_n \xrightarrow{d_n} \dots \xrightarrow{d_2} P_1 \xrightarrow{d_1} P_0 \xrightarrow{d_0} M \to 0$$
(A51)

be an \mathscr{E}^k_A -projective resolution of an A-module M. The exactness of localization [26] implies:

$$\dots \xrightarrow{d_{n+1}} P_n \otimes_A A_{\mathfrak{m}} \xrightarrow{d_n \otimes_A A_{\mathfrak{m}}} \dots \xrightarrow{d_2 \otimes_A A_{\mathfrak{m}}} P_1 \otimes_A A_{\mathfrak{m}} \xrightarrow{d_1 \otimes_A A_{\mathfrak{m}}} P_0 \otimes_A A_{\mathfrak{m}} \xrightarrow{d_0 \otimes_A A_{\mathfrak{m}}} M \otimes_A A_{\mathfrak{m}} \to 0$$
(A52)

is exact. It will now be verified that (A52) is a $\mathscr{E}_{\mathfrak{m},k}$ -projective resolution of the $A_{\mathfrak{m}}$ -module $M_{\mathfrak{m}}$.

The $d_n \otimes_A A_{\mathfrak{m}}$ are $k_{i^{-1}[\mathfrak{m}]}$ -split

Since (A51) was k-split then for every $i \in \mathbb{N}$ there existed a k-module homomorphism $s_i : P_{n-1} \to P_n$ (where for convenience write $P_{-1} := M$) satisfying $d_i = d_i \circ s_i \circ d_i$. Since $A_{\mathfrak{m}}$ is a $k_{i^{-1}[\mathfrak{m}]}$ -algebra $A_{\mathfrak{m}}$ may be viewed as a $k_{i^{-1}[\mathfrak{m}]}$ -module therefore the maps: $s_i \otimes_A 1_{A_{\mathfrak{m}}}$ are $k_{i^{-1}[\mathfrak{m}]}$ -module homomorphisms; moreover they must satisfy:

$$d_i \otimes_A \mathbf{1}_{A_{\mathfrak{m}}} = d_i \otimes_A \mathbf{1}_{A_{\mathfrak{m}}} \circ s_i \otimes_A \mathbf{1}_{A_{\mathfrak{m}}} \circ d_i \otimes_A \mathbf{1}_{A_{\mathfrak{m}}}.$$
 (A53)

Therefore (A52) is $k_{i^{-1}[\mathfrak{m}]}$ -split-exact.

The $P_i \otimes_A A_{\mathfrak{m}}$ are $\mathscr{E}_{\mathfrak{m},k}$ -projective

For each $i \in \mathbb{N}$ if P_i is \mathscr{E}_A^k -projective therefore Proposition A1 implies there exists some *A*-module *Q* and some *k*-module *X* satisfying:

$$P_i \oplus Q \cong A \otimes_k X. \tag{A54}$$

Therefore we have that:

$$(P_{i} \otimes_{A} A_{\mathfrak{m}}) \oplus (Q \otimes_{A} A_{\mathfrak{m}}) \cong (P_{i} \otimes_{A} Q) \otimes_{A} A_{\mathfrak{m}}$$

$$\cong (A \otimes_{k} X) \otimes_{A} A_{\mathfrak{m}}$$

$$\cong (A \otimes_{k} X) \otimes_{A} (A_{\mathfrak{m}} \otimes_{k_{i^{-1}[\mathfrak{m}]}} k_{i^{-1}[\mathfrak{m}]})$$
(A55)

Since *A*, *k* and $k_{i^{-1}[\mathfrak{m}]}$ are commutative rings the tensor products $-\otimes_A -, -\otimes_k$ and $-\otimes_{k_{i^{-1}[\mathfrak{m}]}} -$ are symmetric [22], hence (A55) implies:

$$(P_{i} \otimes_{A} A_{\mathfrak{m}}) \oplus (Q \otimes_{A} A_{\mathfrak{m}}) \cong (A \otimes_{k} X) \otimes_{A} (A_{\mathfrak{m}} \otimes_{k_{i^{-1}[\mathfrak{m}]}} k_{i^{-1}[\mathfrak{m}]})$$
$$\cong (A_{\mathfrak{m}} \otimes_{A} A) \otimes_{k_{i^{-1}[\mathfrak{m}]}} (k_{i^{-1}[\mathfrak{m}]} \otimes_{k} X)$$
(A56)

Since *A* is a subring of $A_{\mathfrak{m}}$ then (A56) implies:

$$(P_i \otimes_A A_{\mathfrak{m}}) \oplus (Q \otimes_A A_{\mathfrak{m}}) \cong A_{\mathfrak{m}} \otimes_{k_{i^{-1}[\mathfrak{m}]}} (k_{i^{-1}[\mathfrak{m}]} \otimes_k X).$$
(A57)

 $(k_{i^{-1}[\mathfrak{m}]} \otimes_k X)$ may be viewed as a $k_{i^{-1}[\mathfrak{m}]}$ -module with action $\hat{\cdot}$ defined as:

$$(\forall c \in k) (\forall (c' \otimes_k x) \in k_{i^{-1}[\mathfrak{m}]} \otimes_k X) c^{\cdot}(c' \otimes_k x) := c \cdot c' \otimes x.$$
(A58)

Since $(k_{i^{-1}[\mathfrak{m}]} \otimes_k X)$ is a $k_{i^{-1}[\mathfrak{m}]}$ -module then for each $i \in \mathbb{N}$ $(P_i \otimes_A A_\mathfrak{m})$ is a direct summand of an $A_\mathfrak{m}$ -module of the form $A_\mathfrak{m} \otimes_{k_{i^{-1}[\mathfrak{m}]}} X'$ where X' is a $k_{i^{-1}[\mathfrak{m}]}$ -module, thus Proposition A1 implies that $P_i \otimes_A A_\mathfrak{m}$ is $A_\mathfrak{m}$ -projective.

Hence (A52) is an $\mathscr{E}_{\mathfrak{m},k}$ -projective resolution of $M \otimes_A A_{\mathfrak{m}} \cong M_{\mathfrak{m}}$; whence:

$$pd_{\mathscr{E}_{\mathfrak{m},k}}(M_{\mathfrak{m}}) \le pd_{\mathscr{E}_{A}^{k}}(M).$$
 (A59)

All the homological dimensions discussed to date are related as follows:

Proposition A10. If A is a commutative k-algebra and m be a non-zero maximal ideal in A such that $A_{\mathfrak{m}}$ has finite $k_{i^{-1}[\mathfrak{m}]}$ -flat dimension and $D(k_{i^{-1}[\mathfrak{m}]})$ is finite then there is a string of inequalities:

$$\begin{aligned} fd_{A_{\mathfrak{m}}}(M_{\mathfrak{m}}) - D(k_{i^{-1}[\mathfrak{m}]}) - fd_{k}(A) &\leq pd_{A_{\mathfrak{m}}}(M_{\mathfrak{m}}) - D(k_{i^{-1}[\mathfrak{m}]}) - fd_{k}(A) \\ &\leq pd_{\mathscr{E}_{\mathfrak{m},k}}(M_{\mathfrak{m}}) \\ &\leq pd_{\mathscr{E}_{A}^{k}}(M) \\ &\leq D_{\mathscr{E}^{k}}(A). \end{aligned}$$

Proof.

- By definition: $pd_{\mathscr{E}^k_A}(M) \leq D_{\mathscr{E}^k}(A)$.
- By Lemma A7: $pd_{\mathscr{E}_{\mathfrak{m},k}}(M_{\mathfrak{m}}) \leq pd_{\mathscr{E}_{A}^{k}}(M)$
- Since $A_{\mathfrak{m}}$ is flat as a $k_{i^{-1}[\mathfrak{m}]}$ -module and $D(k_{i^{-1}[\mathfrak{m}]})$ is finite Lemma A3 entails: $pd_{A_{\mathfrak{m}}}(M_{\mathfrak{m}}) - D(k_{i^{-1}[\mathfrak{m}]}) - fd_k(A) \leq pd_{\mathscr{E}_{\mathfrak{m},k}}(M_{\mathfrak{m}})$
- Lemma A1 implies:

$$fd_{A_{\mathfrak{m}}}(M_{\mathfrak{m}}) \le pd_{A_{\mathfrak{m}}}(M_{\mathfrak{m}}).$$
(A60)

Since the global dimension of $k_{i^{-1}[m]}$ was assumed to be finite (A60) implies:

$$fd_{A_{\mathfrak{m}}}(M_{\mathfrak{m}}) - D(k_{i^{-1}[\mathfrak{m}]}) \le pd_{A_{\mathfrak{m}}}(M_{\mathfrak{m}}) - D(k_{i^{-1}[\mathfrak{m}]}).$$
(A61)

Lemma A8. If *A* is a commutative *k*-algebra and *M* and *N* be *A*-modules, then there are natural isomorphisms:

$$Ext^{n}_{\mathscr{E}^{k}_{A}}(M,N) \cong HH^{n}(A,Hom_{k}(M,N)) \cong Ext^{n}_{\mathscr{E}^{k}_{Ae}}(A,Hom_{k}(M,N)).$$
(A62)

Proof.

- For any (A, A)-bimodule $X, X \otimes_A M$ is an (A, A)-bimodule [22] [Cor. 2.53].
- Moreover, there are natural isomorphisms [22]:

$$Hom_{AMod}(X \otimes_A M, N) \xrightarrow{\cong} Hom_{AMod_A}(X, Hom_{kMod}(M, N))$$
 [Thrm. 2.75]. (A63)

In particular (A63) implies that for every *n* in \mathbb{N} there is an isomorphism which is natural in the first input:

$$Hom_{AMod}(A^{\otimes n} \otimes_A M, N) \xrightarrow{\psi_n} Hom_{AMod_A}(A^{\otimes n}, Hom_{kMod}(M, N)).$$
(A64)

whence if $b'_{n+1} : A^{\otimes n+3} \to A^{\otimes n+2}$ is the n^{th} map in the Bar complex (recall Example A3) and for legibility denote $Hom_{AMod_A}(b'_n, Hom_k(M, N))$ by β_n . The naturality of the maps ψ_n imply the following diagram of *k*-modules commutes:

• Therefore for every n in \mathbb{N} :

$$(\psi_{n+2}^{-1} \circ \beta_{n+1} \circ \psi_{n+1}) \circ (\psi_{n+1}^{-1} \circ \beta_n \circ \psi_n) = \beta_{n+1} \circ \beta_n$$

$$=0.$$
(A66)

Whence $< Hom_{AMod}(A^{\otimes \star + 2} \otimes_A M, N), (\psi_{\star+1}^{-1} \circ \beta_{\star} \circ \psi_{\star}) >$ is a chain complex. Moreover, the commutativity of (A65) implies that:

$$(\forall n \in \mathbb{N}) \ H^{n}(Hom_{AMod}(A^{\otimes \star +2} \otimes_{A} M, N)) = Ker(\psi_{\star+1}^{-1} \circ \beta_{\star} \circ \psi_{\star}) / Im(\psi_{n+2}^{-1} \circ \beta_{n+1} \circ \psi_{n+1})$$

$$\cong Ker(\beta_{n}) / Im(\beta_{n+1})$$

$$= H^{n}(Hom_{AMod_{A}}(A^{\otimes \star +2}, Hom_{kMod}(M, N)))$$

$$= HH^{n}(A, Hom_{k}(M, N)).$$

$$(A67)$$

Furthermore Proposition A4 implies there are natural isomorphisms:

$$HH^{n}(A, Hom_{k}(M, N)) \cong Ext^{n}_{\mathscr{E}^{k}_{A^{e}}}(A, Hom_{k}(M, N));$$
(A68)

Whence for all *n* in \mathbb{N} there are natural isomorphisms:

$$H^{n}(Hom_{A}Mod}(A^{\otimes \star +2} \otimes_{A} M, N)) \cong HH^{n}(A, Hom_{k}(M, N)) \cong Ext^{n}_{\mathscr{C}^{k}}(A, Hom_{k}(M, N)).$$
(A69)

• Finally if *M* is an *A*-module then $< Hom_{AMod}(A^{\otimes \star +2} \otimes_A M, N), (\psi_{\star+1}^{-1} \circ \beta_{\star} \circ \psi_{\star}) >$ calculates the \mathscr{E}_A^k -relative Ext groups of *M* with coefficients in *N*; therefore, by ([24], pg. 289), there are natural isomorphisms:

$$H^{n}(Hom_{AMod}(A^{\otimes \star+2} \otimes_{A} M, N)) \cong Ext^{n}_{\mathscr{E}^{k}_{A}}(M, N).$$
(A70)

• Putting it all together, for every n in \mathbb{N} there are natural isomorphisms:

$$Ext^{n}_{\mathscr{C}^{k}_{A^{e}}}(A, Hom_{k}(M, N)) \cong HH^{n}(A, Hom_{k}(M, N)) \cong Ext^{n}_{\mathscr{C}^{k}_{A^{e}}}(A, Hom_{k}(M, N)).$$
(A71)

We may now prove Theorem 1.

Proof of Theorem 1.

• For any *A*-modules *M* and *N* Lemma A8 implied:

$$Ext^{\star}_{\mathscr{O}^{k}_{A}}(N,M) \cong HH^{\star}(A,Hom_{k}(N,M)).$$
(A72)

Therefore taking supremums over all the *A*-modules M, N, of the integers *n* for which (A85) is non-trivial implies:

$$D_{\mathscr{E}^k}(A) = \sup_{M,N \in {}_A Mod} (\sup\{n \in \mathbb{N}^{\#} | Ext^n(M,N) \neq 0\}))$$
(A73)

$$= \sup_{M,N \in A^{Mod}} (\sup\{n \in \mathbb{N}^{\#} | HH^n(A, Hom_k(N, M)) \neq 0\})).$$
(A74)

 $Hom_k(N, M)$ is only a particular case of an A^e -module; therefore taking supremums over *all* A-modules bounds (A87) above as follows:

$$D_{\mathscr{E}^k}(A) = \sup_{M,N \in A^{Mod}} (\sup\{n \in \mathbb{N}^{\#} | HH^*(A, Hom_k(N, M)) \neq 0\}))$$
(A75)

$$\leq \sup_{\tilde{M} \in _{A^c} Mod} (\sup\{n \in \mathbb{N}^{\#} | HH^n(A, \tilde{M}) \neq 0\})).$$
(A76)

The right hand side of (A89) is precisely the definition of the Hochschild cohomological dimension. Therefore

$$D_{\mathscr{E}^k}(A) \le HCdim(A|k) \tag{A77}$$

Proposition A10 applied to (A90), which draws out the conclusion.

• **Case 1:** *Krull*(*A*_m) **is finite**

Since *A* is Cohen-Macaulay at m there is an A_m -regular sequence $x_1, ..., x_d$ in m of length $d := Krull(A_m)$ in A_m . Therefore Proposition A6 implies:

$$Krull(A_{\mathfrak{m}}) = fd_{A_{\mathfrak{m}}}(A_{\mathfrak{m}}/(x_1,..,x_n)).$$
(A78)

Part 1 of Theorem 1 applied to (A78) implies:

$$Krull(A_{\mathfrak{m}}) - D(k_{i^{-1}[\mathfrak{m}]}) - fd_{\mathfrak{m}}(A_{\mathfrak{m}}) = fd_{A_{\mathfrak{m}}}(A_{\mathfrak{m}}) - D(k_{i^{-1}[\mathfrak{m}]}) - fd_{\mathfrak{m}}(A_{\mathfrak{m}}) \le HCdim(A|k).$$
(A79)

Moreover, the *characterization of quasi-freeness* given in Corollary A1 implies that *A* cannot be quasi-free if:

$$2 + D(k_{i^{-1}[\mathfrak{m}]}) - fd_{\mathfrak{m}}(A_{\mathfrak{m}}) \le Krull(A_{\mathfrak{m}}).$$
(A80)

Case 2: Krull(A_m) is infinite
 For every positive integer *d* there exists an A_m-regular sequence x^d₁, ..., x^d_d in m of length *d*. Therefore Proposition A6 implies:

$$(\forall d \in \mathbb{Z}^+) \ d = f d_{A_{\mathfrak{m}}}(A_{\mathfrak{m}}/(x_1^d, .., x_d^d)).$$
(A81)

Therefore part one of Theorem 1 implies:

$$(\forall d \in \mathbb{Z}^+) \ d - D(k_{i^{-1}[\mathfrak{m}]}) - fd_{\mathfrak{m}}(A_{\mathfrak{m}}) = fd_{A_{\mathfrak{m}}}(A_{\mathfrak{m}}/(x_1^d, ..., x_d^d)) - D(k_{i^{-1}[\mathfrak{m}]}) - fd_{\mathfrak{m}}(A_{\mathfrak{m}}) \le HCdim(A|k).$$
(A82)

Since D(k) and $fd_{\mathfrak{m}}(A_{\mathfrak{m}})$ are finite:

$$\infty - D(k_{i^{-1}[\mathfrak{m}]}) - fd_{\mathfrak{m}}(A_{\mathfrak{m}}) = \infty \le HCdim(A|k).$$
(A83)

Since $Krull(A_m)$ is infinite (A83) implies:

$$Krull(A_{\mathfrak{m}}) - D(k_{i^{-1}[\mathfrak{m}]}) - fd_{\mathfrak{m}}(A_{\mathfrak{m}}) = \infty = HCdim(A|k).$$
(A84)

In this case Corollary A1 implies that *A* is not quasi-free.

Appendix D.2. Proof of Theorem 2

Proof of Theorem 2. For any *A*-modules *M* and *N* Lemma *A*8 implied:

$$Ext^{\star}_{\mathscr{E}^{k}_{A}}(N,M) \cong HH^{\star}(A,Hom_{k}(N,M)).$$
(A85)

Therefore taking supremums over all the *A*-modules M, N, of the integers *n* for which (A85) is non-trivial implies:

$$D_{\mathscr{E}^k}(A) = \sup_{M, N \in A Mod} (\sup\{n \in \mathbb{N}^{\#} | Ext^n(M, N) \neq 0\}))$$
(A86)

$$= \sup_{M,N \in AMod} (\sup(\{n \in \mathbb{N}^{\#} | HH^n(A, Hom_k(N, M)) \neq 0\})).$$
(A87)

 $Hom_k(N, M)$ is only a particular case of an A^e -module; therefore taking supremums over *all A*-modules bounds (A87) above as follows:

$$D_{\mathscr{E}^k}(A) = \sup_{M, N \in A Mod} (\sup\{n \in \mathbb{N}^{\#} | HH^*(A, Hom_k(N, M)) \neq 0\}))$$
(A88)

$$\leq \sup_{\tilde{M} \in A^{e} Mod} (\sup\{n \in \mathbb{N}^{\#} | HH^{n}(A, \tilde{M}) \neq 0\})).$$
(A89)

$$D_{\mathscr{E}^k}(A) \le HCdim(A|k) \tag{A90}$$

Proposition A10 applied to (A90) then draws out the conclusion. Proposition A6 implies that:

$$n = fd_A(A/(x_1, ..., x_n)).$$
(A91)

Therefore (1) applied to the *A*-module $A/(x_1, ..., x_n$ together with (A91) imply:

$$n - D(k) = fd_A(A/(x_1, \dots, x_n) \le D_{\mathscr{E}^k} \le HCDim(A/k).$$
(A92)

If $\Omega^1(A/k)$ is generated by a regular sequence $x_1, ..., x_n$ then Proposition A6 implies:

$$n = fd_{A^e}(A \otimes_k A/\Omega^1(A/k)) \tag{A93}$$

However by definition of $\Omega^1(A/k)$ as the kernel of μ_A : $A \otimes_k A/\Omega^1(A/k) \cong A$. Therefore:

 $n = f d_{A^e}(A). \tag{A94}$

Lemma A1 together with Lemma A3 imply:

$$n = fd_{A^e}(A) \le pd_{A^e}(A) \le pd_{\mathcal{E}^k_{A^e}}(A) + D(k).$$
(A95)

Since D(k) is finite then (A95) entails:

$$n - D(k) \le pd_{\mathcal{E}^k_{A^c}}(A).$$
(A96)

By Theorem A2 (A96) is equivalent to:

$$n - D(k) \le HCDim(A). \tag{A97}$$

If *A* is Cohen-Macaulay at one of its maximal ideals \mathfrak{m} then there exists a maximal regular $x_1, ..., x_d$ in $A_{\mathfrak{m}}$ with $d = Krull(A_{\mathfrak{m}})$. Therefore (2) implies:

$$Krull(A_{\mathfrak{m}}) - D(k) = d - D(k) \le D(A_{\mathfrak{m}}) - D(k).$$
(A98)

Since $D(A_{\mathfrak{m}}) \leq D(A)$, then

$$Krull(A_{\mathfrak{m}}) - D(k) \le D(A_{\mathfrak{m}}) - D(k) \le D(A) - D(k).$$
(A99)

Finally (1) applied to (A99) implies:

$$Krull(A_{\mathfrak{m}}) - D(k) \le D(A) - D(k) \le HCDim(A).$$
(A100)

Appendix D.3. Proofs of Consequences

Proof of Corollary 1. Since *X* is a smooth affine scheme its coordinate ring satisfies Pointcaré duality in dimension *d* then Van den Bergh's Theorem ([12]) applied. Hence, we have that for every $M \in_{A^e} Mod$

$$HH^{n}(A,M) \cong HH_{d-n}(A,H_{d-n}(A,\omega_{A}^{-1}\otimes_{A}M).$$
(A101)

$$0 \not\cong HH^n(A, M) \cong HH_{d-n}(A, \omega_A^{-1} \otimes_A M), \tag{A102}$$

for some *A*-bimodule *M* and some $n \ge fd_A(M) - D(k)$. Again by Van den Bergh's theorem we conclude that $HH_m(A, \omega_A^{-1} \otimes_A M) \cong 0$ for any m > d. Hence, (A102) must hold for some *A*-bimodule *M* and some

$$fd_A(M) - D(k) \le n \le d$$

Thus, there exists an *A*-bimodule M' and some non-negative integer n' for which

$$0 \le n' \le d - f d_A(M) + D(k),$$

and $HH_{n'}(A, M') \not\cong 0$; where $M' \triangleq \omega_A^{-1} \otimes_A M$. Relabeling the index we obtain the conclusion. \Box

Proof of Corollary 2. By the Hochschild-Kostant-Rosenberg ([10]) there are isomorphisms of *A*-bimodules

$$HH_n(A, M) \cong \Omega^n(A, M) \triangleq \Omega^n(A) \otimes_A M, \tag{A103}$$

for every *A*-bimodule *M*. In particular, (A103) holds for the *A*-bimodule *M* of Corollary 1. \Box

References

- 1. Gel'and, I.; Neumark, M. On the imbedding of normed rings into the ring of operators in Hilbert space. *C*-algebras* 1943–1993 (*San Antonio TX* 1993) **1994**, 167, 2–19. Corrected reprint of the 1943 original [MR 5, 147]. [CrossRef]
- 2. Hartshorne, R. Algebraic Geometry; Graduate Texts in Mathematics, No. 52; Springer: New York, NY, USA, 1977; p. xvi+496.
- 3. Isbell, J.R. Structure of categories. Bull. Am. Math. Soc. 1966, 72, 619–655. [CrossRef]
- 4. Isbell, J.R. Normal completions of categories. In *Reports of the Midwest Category Seminar*; Springer: Berlin, Germany, 1967; pp. 110–155.
- 5. Seiberg, N.; Witten, E. String theory and noncommutative geometry. J. High Energy Phys. 1999, 032. [CrossRef]
- 6. Lurie, J. On the classification of topological field theories. In *Current Developments in Mathematics*, 2008; International Press: Somerville, MA, USA, 2009; pp. 129–280.
- 7. Hawkins, E. A cohomological perspective on algebraic quantum field theory. Commun. Math. Phys. 2018, 360, 439–479. [CrossRef]
- 8. Connes, A. Noncommutative differential geometry. Inst. Hautes Études Sci. Publ. Math. 1985, 62, 141–144. [CrossRef]
- 9. Hochschild, G. On the cohomology groups of an associative algebra. Ann. Math. 1945, 46, 58–67. [CrossRef]
- 10. Hochschild, G.; Kostant, B.; Rosenberg, A. Differential forms on regular affine algebras. *Trans. Am. Math. Soc.* **1962**, *102*, 383–408. [CrossRef]
- 11. Antieau, B.; Vezzosi, G. A remark on the Hochschild-Kostant-Rosenberg theorem in characteristic *p. Ann. Sc. Norm. Super. Pisa Cl. Sci.* **2020**, 20, 1135–1145. [CrossRef]
- 12. Van den Bergh, M. A relation between Hochschild homology and cohomology for Gorenstein rings. *Proc. Am. Math. Soc.* **1998**, 126, 1345–1348. [CrossRef]
- 13. Grothendieck, A.; Raynaud, M. Revêtements étales et groupe fondamental (SGA 1). arXiv 2002, arXiv:math/0206203
- 14. Cuntz, J.; Quillen, D. Algebra extensions and nonsingularity. J. Am. Math. Soc. 1995, 8, 251–289. [CrossRef]
- 15. Avramov, L.L.; Vigué-Poirrier, M. Hochschild homology criteria for smoothness. Int. Math. Res. Not. 1992, 1992, 17–25. [CrossRef]
- 16. Avramov, L.L.; Iyengar, S. Gaps in Hochschild cohomology imply smoothness for commutative algebras. *Math. Res. Lett.* **2005**, 12, 789–804. [CrossRef]
- 17. Beck, J.M. Triples, Algebras and Cohomology. Reprints in Theory and Applications of Categories. 2003, pp. 1–59. Available online: http://www.tac.mta.ca/tac/reprints/articles/2/tr2abs.html (accessed on 20 November 2020)
- 18. Hilton, P.J.; Stammbach, U. A Course in Homological Algebra. In *Graduate Texts in Mathematics*, 2nd ed.; Springer: New York, NY, USA, 1997; Volume 4, p. xii+364.
- 19. De Jong, A.J. The Stacks Project. Algebr. Stacks. 2013. Available online: https://stacks.math.columbia.edu/ (accessed on 20 November 2020).
- Springer, T.A. Linear Algebraic Groups, 2nd ed.; Modern Birkhäuser Classics; Birkhäuser Boston, Inc.: Boston, MA, USA, 2009; p. xvi+334.

- 21. Keller, B. Introduction to Kontsevich's Quantization Theorem. Preprint. 2004. Available online: https://webusers.imjprg.fr/ bernhard.keller/publ/emalca.pdf (accessed on 20 November 2020)
- 22. Rotman, J.J. An Introduction to Homological Algebra, 2nd ed.; Universitext; Springer: New York, NY, USA, 2009; p. xiv+709.
- 23. MacLane, S. Homology, 1st ed.; Springer: Berlin, Germany; New York, NY, USA, 1967; p. x+422.
- 24. Weibel, C.A. *An Introduction to Homological Algebra*; Cambridge Studies in Advanced Mathematics; Cambridge University Press: Cambridge, UK, 1994; Volume 38, p. xiv+450.
- Connes, A.; Cuntz, J.; Guentner, E.; Higson, N.; Kaminker, J.; Roberts, J.E. Noncommutative Geometry. In *Lectures Given at the C.I.M.E. Summer School Held in Martina Franca*, 3–9 September 2000; Doplicher, S., Longo, R., Eds.; Fondazione CIME/CIME Foundation Subseries; Springer: Berlin, Germamy, 2004; Volume 1831, p. xiv+343. [CrossRef]
- 26. Eisenbud, D. Commutative Algebra; Graduate Texts in Mathematics; Springer: New York, NY, USA, 1995; Volume 150, p. xvi+785.
- 27. Hochschild, G. Note on relative homological dimension. Nagoya Math. J. 1958, 13, 89–94. [CrossRef]
- 28. Bourbaki, N. Commutative Algebra; Elements of Mathematics (Berlin); Springer: Berlin, Germany, 1998; Chapters 1–7. p. xxiv+625.