

Supplementary Table S1. Demographics, clinical, pathological and postoperative complications of the patients

Variable	Number of patients (%)
Age	
≤ 65 y-o	151 (62.1%)
> 65 y-o	92 (37.9%)
Gender	
Male	140 (57.6%)
Female	103 (42.4%)
Primary tumor location	
Right Colon	48 (19.7%)
Left colon	105 (43.2%)
Rectum	90 (37.1%)
T stage	
T1	1 (0.4%)
T2	6 (2.4%)
T3	216 (88.8%)
T4	17 (7.0%)
NA	3 (1.2%)
N stage	
N0	77 (31.7%)
N1	78 (32.1%)
N2	83 (34.2%)
NA	5 (2%)
Distributions of SCLMs	
Unilobar	176 (72.5%)
Bilobar	67 (27.5%)
Number of SCLMs	
1	137 (56.3%)
2-3	73 (30.1%)
≥ 4	33 (13.6%)
Size of SCLMs	
< 3 cm	126 (51.9%)
≥ 3 cm	117 (48.1%)
TBS score	
≤ 3.2	120 (49.3%)
> 3.2	123 (50.7%)
Type of hepatectomy	
Anatomic	34 (14%)
Non-anatomic	209 (86%)
Extension of hepatectomy	
Minor	211 (86.8%)
Major	32 (13.2%)
Postoperative complications (Clavien-Dindo)	
No complications	
I	121 (49.7%)

II	20 (8.3%)
III	48 (19.7%)
IV	35 (14.5%)
V	7 (2.9%)
	12 (4.9%)
Septic complications	
No	157 (64.6%)
Yes	86 (35.4%)
Hepatic complications	
No	184 (75.7)
Yes	59 (24.3%)
Hepatic septic complications	
No	203 (83.5%)
Yes	40 (16.5%)
Comprehensive complication index (median [IQR])	8.70 (0 - 26.20)
Neoadjuvant treatment	
No	210 (86.4%)
Yes	33 (13.6%)
Adjuvant treatment	
No	29 (11.9%)
Yes	214 (88.1%)