

Article

Physical and Mental Disabilities among the Gender-Diverse Population Using the Behavioral Risk Factor Surveillance System, BRFSS (2017–2019): A Propensity-Matched Analysis

Jennifer R. Pharr ^{1,*} and Kavita Batra ²¹ Department of Environmental and Occupational Health, School of Public Health, University of Nevada, Las Vegas, NV 89119, USA² Office of Research, Kirk Kerkorian School of Medicine, University of Nevada, Las Vegas, NV 89102, USA; Kavita.batra@unlv.edu

* Correspondence: jennifer.pharr@unlv.edu

Abstract: This propensity-matched analysis utilized the publicly available Behavioral Risk Factor Surveillance System (2017–2019) data to compare the burden of disabilities among transgender/non-binary (TGNB) and cisgender groups. The groups were matched (1:1 ratio) on demographic variables using Nearest Neighborhood Matching. Categorical variables were compared among groups using a Chi-square analysis to test differences in the proportions. Multivariate logistic regression analysis was fit to predict the likelihood of the physical and mental disabilities among the TGNB group compared with the cisgender group while controlling for healthcare access factors, income, and employment. Survey weights were included in the model to account for the complex survey design. In a weighted sample of 664,103 respondents, only 2827 (0.4%) self-identified as TGNB. In the matched sample, a higher proportion of the TGNB group belonged to the low-income group (39.5% vs. 29.8%, $p < 0.001$), were unable to work (12.5% vs. 8.6%, $p < 0.001$), and delayed care due to cost barriers (19.0% vs. 12.4%, $p < 0.001$). Compared with the cisgender group, the odds of having difficulty making decisions were 1.94 times higher (95% CI: 1.67–2.27) and odds of difficulty walking were 1.38 times higher (95% CI: 1.19, 1.59) among the TGNB group. Additionally, the TGNB group had 59.8% higher adjusted odds ratio (aOR) (aOR 1.598, 95% Confidence interval (CI): 1.256, 2.034) of experiencing difficulty dressing and 83.3% higher odds (aOR 1.833, 95% CI: 1.533, 2.191) in having difficulty doing things alone. The findings of this study advocate for developing policies and interventions to deliver culturally competent care to the TGNB population with disabilities.

Citation: Pharr, J.R.; Batra, K. Physical and Mental Disabilities among the Gender-Diverse Population Using the Behavioral Risk Factor Surveillance System, BRFSS (2017–2019): A Propensity-Matched Analysis. *Healthcare* **2021**, *9*, 1285. <https://doi.org/10.3390/healthcare9101285>

Academic Editor: Alessandro Sartorio

Received: 3 September 2021

Accepted: 24 September 2021

Published: 28 September 2021

Keywords: physical disability; mental disability; transgender; propensity score matching; Behavioral Risk Factor Surveillance System

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Transgender and gender nonbinary (TGNB) persons have gender identities, expressions, or behaviors not traditionally associated with the sex they were assigned at birth [1]. In the United States (U.S.), it is estimated that TGNB persons constitute 6% of the population, or 1.4 million people [2]. Despite the growing population, visibility, and acceptance of gender minorities in the U.S., evidence continues to show wide disparities in social determinants of health (income, employment, education) and health outcomes among this population compared with cisgender (non-transgender) populations [3–11]. The minority stress theory posits that TGNB people are at greater risk of experiencing worse health outcomes due to stigma and its resulting discrimination [12–15]. The stigma that they experience may be at the individual, interpersonal, and societal/structural levels and occurs because their gender identity lies outside of the cisgender, binary norm [12–15]. Identity-related stigmatization results in employment discrimination, public services

discrimination, macro and micro aggressions, and internalized transphobia, which leads to elevated levels of stress [12–15].

Minority stress has been linked to an increased burden of mental health disorders and chronic diseases among sexual and gender minority populations, particularly among TGNB people [12,14]. Studies reported a higher prevalence of mental health disorders such as anxiety, depression, suicidality, and substance use among TGNB persons compared with their cisgender counterparts [10,12–17]. Previous studies explained increased suicidal ideation among TGNB people through gender minority stress and interpersonal factors [16,17]. Additionally, compared with the cisgender people, TGNB people are more likely to have chronic diseases, such as hypertension, hypercholesterolemia, prediabetes, stroke, cardiovascular diseases, asthma, and cancers, while also engaging in more health risk behaviors including smoking and heavy drinking [10,18]. While there is a paucity of research examining disability disparities among TGNB people, a few studies found an elevated risk of mobility, cognitive, and independent living disabilities among TGNB people compared to cisgender men, the odds of which increase with age [5,19–24].

Disability is defined by the Americans with Disabilities Act (ADA) as a “physical or mental impairment that substantially limits one or more major life activities” with major life activities including, but not limited to, self-care, manual tasks, seeing, hearing, walking, standing, lifting, speaking, learning, concentrating, and communicating [25]. Reportedly, sixty-one million American adults have a disability, equaling 25.7% of the adult population [26]. Among transgender people, 39% report having a disability [26,27]. People with disabilities have been stigmatized throughout history [3,19,27]. The stigma of disability contributes to discrimination in many facets of life such as employment, education, and life decisions; stereotypes; negative attitudes; and labeling of people with disabilities [28].

Because people with disabilities are subjected to disability stigma, this could result in TGNB people with disabilities having multiple minority stress. However, the epidemiology of these two minority stressors is relatively unknown in the TGNB population. There is evidence to suggest that sexual minority people are more likely to have disabilities compared with heterosexual people and that this intersection results in poorer health and a great need for community support [29–31]. However, minimal research is available for gender minorities. The little research that has been conducted used uneven samples to draw conclusions. Therefore, the purpose of this study was to use a propensity-matched sample to examine disabilities disparities among TGNB people compared with cisgender people. Authors hypothesize that TGNB people experience a higher burden of physical disabilities compared to their cisgender counterparts after matching on certain demographic variables.

2. Materials and Methods

2.1. Study Design and Data Source

This cross-sectional study utilized publicly available data from the Behavioral Risk Factor Surveillance System (BRFSS) for the years 2017–2019 [32]. The BRFSS is the largest, population-based, and nationwide computer-assisted telephone interview (CATI) survey conducted by the CDC, which collects data from all 50 states and selected territories of the United States (U.S.) [32–34]. The target population of the BRFSS survey is non-institutionalized adults aged 18 years or older [32–34]. The BRFSS collects information about participants’ demographics, health-related risk behaviors, chronic health conditions, and use of preventive services in the core module of the questionnaire [32]. In 2014, the BRFSS expanded the utility of the surveillance data by including an optional module related to sexual orientation and gender identity (SOGI), which was utilized by 28 states and territories in 2017, 30 states and territories in 2018, and 31 states and territories in 2019 [33,34]. To account for the bias resulting from the selection probabilities

and non-response, the BRFSS utilizes a complex sampling weighting methodology, which can be viewed at <https://www.cdc.gov/brfss>. This helps in generating the nationally representative samples [35]. This study was deemed excluded from the ethical review by the University of Nevada, Las Vegas Institutional Review Board, because it involves secondary data analysis of publicly available, deidentified data with no direct involvement of the human subjects.

2.2. Participants

The study was comprised of survey participants of at least 18 years of age and who responded “yes” to the question “Do you consider yourself to be a transgender?” [36–38]. Participants who self-identified as Transgender, male-to-female; Transgender, female-to-male; and gender nonconforming were included in the TGNB (yes) category. Participants who responded “no” to the above question were included in a comparative cisgender (transgender “no”) group. Respondents who responded “don’t know or unsure” or “refused to answer” were excluded from the study. This methodology was consistent with the previous studies, which utilized the BRFSS data [39–41].

2.3. Variables and Measures

Variables included in this study were sociodemographic information (e.g., birth-assigned sex, education, age, race/ethnicity, marital status, employment status and income) and healthcare accessibility information (health insurance, having personal doctor, frequency of medical check-up, and delayed check-up due to cost) [36–38]. We recoded some variables to construct meaningful categories. For instance, unemployed includes those who were out of the labor force (e.g., retired, homemakers, students, and unable to work) and the non-white category includes Blacks, multiracial, Asian, Native Hawaiian/Pacific Islanders, and Alaska Natives/American Indians. The main outcomes for this study were physical and mental disabilities, e.g., difficulty hearing, seeing, performing routine activities such as walking, dressing, bathing, and limited ability to make decisions. We also evaluated the number of days of poor mental and physical health in the past month (i.e., 0 days, 1–13 days, or <14 days). Because some disabilities are associated with poor health, chronic conditions, and lack of access to healthcare, we included healthcare access as a demographic characteristic in this study. Additionally, we included healthcare access as a covariate in this analysis because previous research has identified that financial barriers to healthcare are predictors of disability among SGM populations [5,24].

2.4. Statistical Analysis

The data from the three surveys (2017, 2018, and 2019) were pooled to increase the sample size following the methodology described by the CDC [35]. The final weight was calculated and adjusted according to the proportion of the sample size in each year. To account for the sample size differences in the cisgender and transgender groups and to minimize selection bias, we used the propensity-score-matched (PSM) analysis [10,42,43]. Cisgender (control) and TGNB (case) groups were 1:1 matched on demographic variables, including birth-assigned sex, age, race, marital status, and education. Nearest Neighborhood Matching (NNM) was performed using the matching package [43] and covariate balance was assessed using cobalt [44,45], both in R [46]. Propensity scores were first estimated using a logistic regression and the probability distribution was then compared. Covariance balance in unmatched and matched sample was compared by visually inspecting the Love plots. In addition, standardized mean differences and variance ratio were used as indicators to assess covariance balance [10]. Categorical variables were compared among groups using a Chi-square analysis to test the differences in the proportions. Multivariate logistic regression analysis was fit to predict the likelihood of the physical and mental disabilities among the TGNB group compared with

the cisgender group while controlling for healthcare access factors, income, and employment. Survey weights were included in the model to account for the complex survey design using a survey package in R software. The significance level was set at 5%, and all descriptive analysis of the matched samples were conducted using SPSS version 26 (IBM Corp. Armonk, NY, USA).

3. Results

In a weighted sample of 664,103 respondents, only 2827 (0.4%) self-identified as TGNB. In an unmatched sample, a higher proportion of the TGNB group were non-white (23.2% vs. 20.5%, $p < 0.001$), Hispanic (21.0% vs. 15.8%, $p < 0.001$), aged 18–34 years (47.4% vs. 27.6%, $p < 0.001$), were never married (38.8% vs. 23.5%, $p < 0.001$), and were less educated with only a high school diploma or less (55.1% vs. 41.4%, $p < 0.001$; Table 1). The post-matched sample ($n = 2687$) had an adequate covariate balance, as indicated by absolute standardized mean differences close to zero and variance ratios close to 1, as indicated in Figure 1 and Table 2.

Table 1. Demographic characteristics of the unmatched study population ($N = 664,103$).

Characteristics	Gender Identities		Total <i>n</i> (weighted %)	<i>p</i> Value
	TGNB <i>n</i> (weighted %)	Cisgender <i>n</i> (weighted %)		
All	2827 (0.5)	661,276 (99.5)	664,103	-
Sex assigned at birth				
Male	1456 (51.9)	294,479 (47.7)	295,935 (48.0)	<0.001
Female	1324 (48.1)	366,243 (51.7)	367,567 (52.0)	
Education status				
High school graduate or less	1343 (55.1)	223,742 (41.4)	225,085 (41.4)	<0.001
Attended college	723 (28.9)	183,259 (31.3)	183,982 (31.3)	
College graduate	742 (16.0)	252,304 (27.4)	253,052 (27.3)	
Age (in years)				
18–34	847 (47.4)	99,009 (27.6)	99,856 (27.7)	<0.001
35–54	700 (25.1)	172,366 (32.3)	173,066 (32.2)	
55–74	936 (20.8)	279,048 (31.0)	279,984 (30.9)	
75+	307 (6.7)	101,102 (9.2)	101,409 (9.2)	
Race/ethnicity				
White	1784 (55.8)	500,935 (63.7)	502,719 (63.7)	<0.001
Non-white	680 (23.2)	109,607 (20.5)	110,287 (20.5)	
Hispanic	337 (21.0)	46,450 (15.8)	46,787 (15.9)	
Marital status				
Married	1032 (34.3)	338,983 (51.1)	340,015 (51.0)	<0.001
Divorced	382 (9.8)	92,057 (10.9)	92,439 (10.9)	
Widowed	285 (6.2)	82,717 (7.4)	83,002 (7.3)	
Separated	80 (3.0)	14,374 (2.6)	14,454 (2.6)	
Never married	850 (38.8)	107,597 (23.5)	108,447 (23.6)	
A member of unmarried couple	181 (7.8)	21,651 (4.6)	21,832 (4.7)	

TGNB = transgender, non-binary. Unemployed includes those being out of the labor force (e.g., retired, homemakers, students, and unable to work). Non-white includes Blacks, multiracial, Asian, NH/PI, AI/AN.

Table 2. Summary of covariate balance for matched cohort units ($N = 2687$).

	Means Treated (Trans)	Mean Control (Cis)	SMD	VR	Standard Pair Distance
Age	2.949	2.951	−0.0015	1.001	0.0098
Biological sex	1.569	1.533	0.0386	1.353	0.0507
Race	1.843	1.849	−0.0041	0.992	0.0102
Marital status	2.988	3.004	−0.0083	0.973	0.0146
Education level	2.635	2.630	0.0047	0.998	0.0069

SMD = Standardized Mean Difference; VR = Variance ratio. Matching variables = Age, Biological sex, Race, Marital status, Education level.




Figure 1. Love plot displaying covariate balance in unadjusted and adjusted samples. Legend: Absolute standardized mean differences close to zero and variance ratio close to one is indicative of a good covariate balance.

Upon analyzing differences in the socioeconomic, healthcare access, and cost of care barriers, significant differences among unmatched and matched samples were noted (Table 3). In a matched sample, with the exception of healthcare insurance, there were statistically significant differences in the proportion. A higher proportion of the TGNB group belong to the low-income group (39.5% vs. 29.8%, $p < 0.001$), were unable to work (12.5% vs. 8.6%, $p < 0.001$) and delayed care due to cost barriers (19.0% vs. 12.4%, $p < 0.001$; Table 3) compared with the cisgender group. However, for personal doctor and medical check-ups, a significantly higher proportion of the TGNB respondents reported having a medical checkup within the past year (76.6% vs. 70.3%, $p < 0.001$) and had at least one personal doctor (69.9% vs. 61.1%, $p < 0.001$; Table 3). The results of outcome analysis indicated that TGNB individuals were more likely to experience difficulty seeing (7.9% vs. 6.1%, $p = 0.008$) and making decisions (22.5% vs. 12.5%, $p < 0.001$, Table 4) as opposed

to their cisgender counterparts. A significantly higher proportion of TGNB respondents reported difficulty walking (20.6% vs. 14.3%, $p < 0.001$), dressing (7.6 vs. 4.3%, $p < 0.001$), and doing things alone (15.4% vs. 8.3%, $p < 0.001$, Table 4). As compared with the cisgender group, the TGNB group reported having poor physical (18.4% vs. 13.9%, $p < 0.001$) and mental health outcomes (24.9% vs. 14.1%, $p < 0.001$), which persisted for at least 14 days (Table 4).

After adjusting for the healthcare access factors, cost-of-care barriers, income, and education, having difficulties making decisions, walking, dressing, and doing things alone were associated with the TGNB group. Compared with the cisgender group, the odds of having difficulties making decisions were 1.94 times higher (95% CI: 1.67–2.27) and odds of difficulties walking were 1.38 times higher (95% CI: 1.19, 1.59, Table 5) among the TGNB group. Additionally, the TGNB group had 59.8% higher odds (aOR 1.598, 95% CI: 1.256, 2.034) of experiencing difficulties dressing and 83.3% higher odds (aOR 1.833, 95% CI: 1.533, 2.191, Table 5) having difficulties doing things alone.

Table 3. Socioeconomic, health-specific and cost-of-care barriers among unmatched and matched samples.

Outcome	Unmatched			Matched		
	TGNB N (Weighted %)	Cisgender N (Weighted%)	<i>p</i> Value	TGNB N (%)	Cisgender N (%)	<i>p</i> Value
Income in dollars						
Up to 25 K	926 (43.7)	138,849 (26.3)	<0.001	884 (39.5)	655 (29.8)	<0.001
25–50 K	573 (19.7)	13,4271 (23.2)		549 (24.6)	533(24.2)	
50–75 K	311 (12.5)	89,208 (15.1)		304 (13.6)	338 (15.4)	
75+ K	515 (24.1)	192,787 (35.4)		499 (22.3)	675 (30.7)	
Employment status						
In labor force or working	1350 (49.2)	326,136 (56.8)	<0.001	1301 (48.4)	1385 (51.5)	<0.001
Out of labor force or unemployed	1091 (36.9)	280,908 (36.0)		1049 (39.1)	1072 (39.9)	
Unable to work	353 (13.9)	48,893 (7.2)		337 (12.5)	230 (8.6)	
Healthcare insurance						
Yes	2434 (82.2)	605,968 (87.2)	<0.001	2317 (86.7)	2332 (87.4)	0.5
No	389 (17.8)	54,374 (12.8)		354 (13.3)	337 (12.6)	
Personal doctor						
Yes, only one	1967 (67.3)	500,447 (70.6)	<0.001	1876 (69.9)	1638 (61.1)	<0.001
Yes, more than one	237 (7.4)	54,825 (7.6)		224 (8.3)	240 (8.9)	
No	619 (25.3)	10,469 (21.8)		584 (21.8)	805 (30.0)	
Delayed care due to cost						
Yes	547 (21.9)	67,990 (13.3)	<0.001	511 (19.0)	334 (12.4)	<0.001
No	2277 (78.1)	592,187 (86.7)		2173 (81.0)	2352 (87.6)	
Medical check up						
Yes, within past year	2161 (72.6)	521,770 (74.6)	<0.001	2057 (76.6)	1889 (70.3)	<0.001

Yes, within past 2 years	259 (11.6)	62,672 (11.0)	244 (9.1)	319 (11.9)
Within past 5 years	171 (6.1)	33,406 (6.5)	163 (6.1)	204 (7.6)
5 or more years	176 (7.9)	31,738 (5.8)	168 (6.3)	208 (7.7)
No	57 (1.8)	10,423 (2.0)	52 (1.9)	67 (2.5)

TGNB = transgender, non-binary.

Table 4. Burden of physical and mental disabilities among unmatched and matched samples.

Outcome	Unmatched			Matched		
	TGNB N (%)	Cisgender N (%)	<i>p</i> Value	TGNB N (%)	Cisgender N (%)	<i>p</i> Value
Difficulty hearing						
Yes	302 (9.8)	58,784 (6.6)	<0.001	286 (10.7)	247 (9.3)	0.07
No	2493 (90.2)	596,657 (93.4)		2375 (89.3)	2411 (90.7)	
Difficulty seeing						
Yes	219 (7.8)	34,750 (5.1)	<0.001	211 (7.9)	161 (6.1)	0.008
No	2572 (92.2)	620,253 (94.9)		2448 (92.1)	2492 (93.9)	
Difficulty making decisions						
Yes	629 (27.3)	69,714 (11.4)	<0.001	598 (22.5)	323 (12.2)	<0.001
No	2158 (72.7)	584,543 (88.6)		2058 (77.5)	2328 (87.8)	
Difficulty walking						
Yes	576 (17.4)	115,267 (14.5)	<0.001	546 (20.6)	379 (14.3)	<0.001
No	2207 (82.6)	538,729 (85.5)		2108 (79.4)	2270 (85.7)	
Difficulty dressing						
Yes	217 (9.6)	29,252 (4.1)	<0.001	202 (7.6)	114 (4.3)	<0.001
No	2567 (90.4)	624,558 (95.6)		2452 (92.4)	2531 (95.7)	
Difficulty doing things alone						
Yes	427 (20.0)	50,627 (7.2)	<0.001	409 (15.4)	218 (8.3)	<0.001
No	2352 (80.0)	602,566 (92.8)		2240 (84.6)	2423 (91.7)	
General health						
Excellent	106,025 (17.2)	399 (16.9)	<0.001	380 (14.2)	388 (14.5)	<0.001
Very Good	216,242 (31.4)	787 (25.7)		754 (28.1)	825 (30.8)	
Good	209,887 (32.4)	891 (30.3)		844 (31.5)	896 (33.4)	
Fair	509 (17.6)	91,996 (14.1)		490 (18.3)	426 (15.9)	
Poor	232 (9.4)	35,467 (4.9)		213 (7.9)	146 (5.4)	
Poor physical health						
0 days	1411 (50.2)	402,618 (63.6)	<0.001	1345 (51.8)	1523 (58.0)	<0.001
1–13 days	812 (32.4)	152,847 (23.8)		772 (29.7)	738 (28.1)	
≤14 days	505 (17.4)	91,189 (12.6)		478 (18.4)	365 (13.9)	
Poor mental health						
0 days	1408 (44.9)	433,007 (64.0)	<0.001	1351 (51.5)	1577 (59.8)	<0.001
1–13 days	651 (25.0)	141,039 (23.3)		617 (23.5)	687 (26.1)	
≤14 days	694 (30.1)	75,359 (12.7)		654 (24.9)	373 (14.1)	

TGNB = transgender, non-binary.

Table 5. Adjusted and unadjusted odds ratio for disabilities among matched samples.

Variable	Unadjusted Odds Ratio	95% CI		<i>p</i> Value	AOR	95% CI		<i>p</i> Value
		LCL	UCL			LCL	UCL	
Difficulty hearing								
TGNB	1.175	0.982	1.406	0.07	1.066	0.888	1.280	0.5
Cisgender	REF	-	-	-	-	-	-	-
Difficulty seeing								
TGNB	1.334	1.079	1.650	0.008	1.236	0.995	1.534	0.05 *
Cisgender	REF	-	-	-	-	-	-	-
Difficulty making decisions								
TGNB	2.094	1.807	2.428	<0.001	1.948	1.673	2.269	<0.001
Cisgender	REF	-	-	-	-	-	-	-
Difficulty walking								
TGNB	1.551	1.344	1.791	<0.001	1.381	1.192	1.599	<0.001
Cisgender	REF	-	-	-	-	-	-	-
Difficulty dressing								
TGNB	1.829	1.444	2.316	<0.001	1.598	1.256	2.034	<0.001
Cisgender	REF	-	-	-	-	-	-	-
Difficulty doing things alone								
TGNB	2.029	1.705	2.415	<0.001	1.833	1.533	2.191	<0.001
Cisgender	REF	-	-	-	-	-	-	-

TGNB = transgender, non-binary. *p* values less than 0.05 are statistically significant; Adjusted odds ratios (AOR) were obtained after controlling for healthcare access factors, income, and employment; LCL-Lower Confidence Limit; UCL-Upper Confidence Limit. * Marginally Significant.

4. Discussion

Pre- and post-matching, TGNB people experience disparities in two important factors that are determinants of health (i.e., income and employment) when compared with cisgender people. This is consistent with other studies that did not use matched samples [5,6]. TGNB people tend to have lower rates of employment and make less money than their cisgender peers, possibly due to stigma at the interpersonal and structural levels. While TGNB people may have more protections under the law than their lesbian, gay, and bisexual peers due to the Title VII federal sex discrimination law, they still experience unacceptably high rates of workplace discrimination. A recent survey of TGNB people found that 44% had experienced discrimination in employment and hiring in the prior 12 months [47–49]. Research has also shown that sexual and gender minority adults experience discrimination in hiring practices and compensation, leading to higher unemployment and lower income despite high rates of higher education [50,51].

Access to healthcare is important for diagnosing, preventing, and treating disabilities. However, TGNB people face discrimination in healthcare that results in a multitude of barriers to accessing care, which include a lack of health insurance or insurance coverage for gender affirming care, financial barriers, transphobic healthcare providers, verbal or physical abuse and maltreatment by healthcare providers, and healthcare providers' lack of knowledge about TGNB care [52–54]. Prior to matching in this sample, TGNB people in this study were less likely to report having a personal doctor or having had a medical check-up in the past year than cisgender people. However, after matching, we saw these results switch, with TGNB people being more likely to have a personal doctor (70%) or a medical checkup in the past year (77%). This is similar to results from the U.S. Transgender Survey [54]. This study demonstrates that matched results are particularly important as they provide more accurate estimates after accounting for the imbalance among covariates, which might otherwise have contributed to the

hypothesized effects in the unmatched findings. Knowing the barriers to healthcare experienced by TGNB, these findings from the matched analysis are encouraging, and may be due to the healthcare needs of TGNB people, especially those who are on gender-affirming hormones or receiving care from a transgender provider.

In the matched analysis, TGNB people were more likely to report all disabilities (except difficulty hearing) and 14 days or more of poor physical and mental health. After controlling for healthcare access factors, income, and employment, TGNB people were found to have greater disability disparities including difficulties making decisions, walking, dressing, and doing things alone. Previous research using an unmatched sample also found that the TGNB people were more likely to have mobility, cognitive, and independent living disabilities compared with cisgender people, which this study confirms [5]. TGNB people in this study were almost two times more likely to have difficulties making decisions or doing things alone. This may indicate a need for help that might not be readily available for TGNB people. The largest survey of TGNB people in the U.S. found that they were more likely to be single and never married compared to the U.S. population [3]. Our findings are similar with regard to marital status. Additionally, TGNB people are more likely to experience family rejection and homelessness and are less likely to have children [3]. People with disabilities often need additional assistance with activities of daily living and usually this assistance comes from family members and friends [3,55,56]. TGNB people who have disabilities may rely on their friends for support and help rather than their family of origin. Indeed, older sexual and gender minorities are more likely to rely on friends for care than do their heterosexual, cisgender peers [55,57]. Research has found that, within the sexual and gender minority community, friends who provide informal care have less access to social support, which leads to greater stress and depressive symptoms [58]. Our finding of increased risk of disability highlights the need for support services that are culturally sensitive to those with disabilities who identify as TGNB and their caregivers.

A disability may make healthcare access issues more profound for TGNB people. According to Healthy People 2020, 76.8% of adults with disabilities experience barriers that hinder their use of available healthcare and wellness services [59]. These barriers include structural barriers (e.g., lack of accessible medical equipment), financial barriers (e.g., lack of health insurance, medical equipment that is too expensive), and personal/cultural barriers (e.g., misconceptions about people with disabilities, maltreatment from healthcare providers) [19,60,61]. Healthcare barriers persist despite having two federal laws (Americans with Disabilities Act and section 504 of the Rehabilitation Act), which require healthcare providers to grant their patients with disabilities full and equal access to healthcare services and facilities [62–64]. Having a disability and identifying as TGNB may present a double barrier to healthcare. As mentioned above, TGNB people experience unique barriers to healthcare compared with the cisgender population. Previous research has found that TGNB people with disabilities are two times more likely to experience financial barriers to health care [57]. Additionally, financial barriers to health care, lifetime victimization, internalized stigma, and lack of social support are predictors for disability among older sexual minority adults [57]. Policies at the federal, state, and institutional levels are not only needed to address structural stigma, but need to be implemented and enforced to address healthcare barriers experienced by TGNB people and people with disabilities.

TGNB people with disabilities are likely to experience multiple minority stress due to their multiple minority identities, that of being TGNB and that of having a disability. Both groups are at an increased risk of discrimination [65]. For example, stigma and incorrect stereotypes lead some healthcare providers to see people with a disability as asexual, and, in turn, they do not offer certain services such as sexually transmitted disease testing [58,59]. On the other hand, healthcare providers may have limited knowledge of gender-affirming care. Together, these two situations would impede healthcare for a TGNB person with a disability. Individually, TGNB stigma and the stigma

of disability impact depression, psychological distress, quality of life, and quality of healthcare [49]. Together, these two forms of stigma may have a profound impact on the health, well-being, and healthcare of TGNB people with disabilities. However, because of the paucity of research at the intersection of disability and TGNB identity, little is known about the health burden of this form of multiple minority stress. More research is needed at this intersection to understand the health impacts.

Strengths and Limitations

To our knowledge, this is the first study that utilized propensity-matched analysis to investigate the burden of disabilities among TGNB people. Transgender and cisgender participants were appropriately matched on demographic characteristics to assess the true differences across groups. Like other studies, this study is not without limitations. First, the cross-sectional design of this study does not allow us to infer causality. Moreover, BRFSS utilizes a repeated cross-sectional design; it is likely that some participants were sampled across the years, which might have introduced some bias in the estimates. However, authors expect minimal effect as estimates were generated on a matched subset. Second, due to inconsistency in the method of collecting sexual orientation information, matching did not account for the sexual orientation. Third, this study may be subject to information bias, including reporting bias and recall bias. Lastly, although groups of transgender and cisgender participants were matched on demographic characteristics, a residual confounding may still exist due to differences within the subgroups.

5. Conclusions

TGNB people were more likely to report disabilities and poorer mental and physical health than cisgender people. The minority stress theory would posit that these disparities are, at least in part, due to the excess stigma that TGNB people experience at the structural/social, interpersonal, and individual levels. TGNB people with disabilities may experience multiple minority stress and experience both the stigma of disability and of being a gender minority. These stigma may impact other health outcomes as well as access to healthcare. However, more research is needed to examine the intersection of disability and gender minority status. Additionally, interventions (culturally competent care) and enforceable policies are needed to address the unique barriers to healthcare experienced by people with disabilities and TGNB people.

Author Contributions: Conceptualization, J.R.P.; methodology, J.R.P. and K.B.; software, K.B.; validation, J.R.P. and K.B.; formal analysis, K.B.; investigation, K.B.; resources, J.R.P.; data curation, J.R.P. and K.B.; writing—original draft preparation, J.R.P. and K.B.; writing—review and editing, J.R.P. and K.B.; visualization, J.R.P. and K.B.; supervision, J.R.P.; project administration, K.B. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: This study was deemed excluded from the ethical review by the University of Nevada, Las Vegas Institutional Review Board, because it involves secondary data analysis of publicly available, deidentified data with no direct involvement of the human subjects.

Informed Consent Statement: Not applicable.

Data Availability Statement: Data are available in a publicly accessible repository that does not issue DOIs. Publicly available datasets were analyzed in this study. These data can be found here: https://www.cdc.gov/brfss/data_documentation/index.htm, accessed on 24 June 2020.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. American Psychological Association Guidelines for psychological practice with transgender and gender nonconforming people. *Am. Psychol.* **2015**, *70*, 832–864, <https://doi.org/10.1037/a0039906>.
2. The Williams Institute. How Many Adults Identify as Transgender in the United States. 2016. Available online: <https://williamsinstitute.law.ucla.edu/wp-content/uploads/Trans-Adults-US-Aug-2016.pdf> (accessed on 20 August 2021).
3. James, S.E.; Herman, J.L.; Rankin, S.; Keisling, M.; Mottet, L.; Anafi, M. *The Report of the 2015 U.S. Transgender Survey*; National Agency for Transgender Equality: Washington, DC, USA, 2016.
4. Nolan, I.T.; Kuhner, C.J.; Dy, G.W. Demographic and temporal trends in transgender identities and gender confirming surgery. *Transl. Androl. Urol.* **2019**, *8*, 184–190, <https://doi.org/10.21037/tau.2019.04.09>.
5. Downing, J.M.; Przedworski, J.M. Health of Transgender Adults in the U.S., 2014–2016. *Am. J. Prev. Med.* **2018**, *55*, 336–344, <https://doi.org/10.1016/j.amepre.2018.04.045>.
6. Crissman, H.P.; Berger, M.B.; Graham, L.F.; Dalton, V.K. Transgender Demographics: A Household Probability Sample of US Adults, 2014. *Am. J. Public Health* **2017**, *107*, 213–215, <https://doi.org/10.2105/ajph.2016.303571>.
7. Baker, K.E. Findings from the Behavioral Risk Factor Surveillance System on Health-Related Quality of Life Among US Transgender Adults, 2014–2017. *JAMA Intern. Med.* **2019**, *179*, 1141–1144, <https://doi.org/10.1001/jamainternmed.2018.7931>.
8. Newfield, E.; Hart, S.; Dibble, S.; Kohler, L. Female-to-male transgender quality of life. *Qual. Life Res.* **2006**, *15*, 1447–1457, <https://doi.org/10.1007/s11136-006-0002-3>.
9. Streed, C.G.; McCarthy, E.P.; Haas, J.S. Self-Reported Physical and Mental Health of Gender Nonconforming Transgender Adults in the United States. *LGBT Health* **2018**, *5*, 443–448, <https://doi.org/10.1089/lgbt.2017.0275>.
10. Pharr, J.R.; Batra, K. Propensity Score Analysis Assessing the Burden of Non-Communicable Diseases among the Transgender Population in the United States Using the Behavioral Risk Factor Surveillance System (2017–2019). *Healthcare* **2021**, *9*, 696.
11. Meyer, I.H.; Brown, T.N.T.; Herman, J.L.; Reisner, S.L.; Bockting, W.O. Demographic Characteristics and Health Status of Transgender Adults in Select US Regions: Behavioral Risk Factor Surveillance System, 2014. *Am. J. Public Health* **2017**, *107*, 582–589, <https://doi.org/10.2105/ajph.2016.303648>.
12. Meyer, I.H. Minority Stress and Mental Health in Gay Men. *J. Health Soc. Behav.* **1995**, *36*, 38–56, <https://doi.org/10.2307/2137286>.
13. Meyer, I.H. Prejudice, social stress, and mental health in lesbian, gay, and bisexual populations: Conceptual issues and research evidence. *Psychol. Bull.* **2003**, *129*, 674–697, <https://doi.org/10.1037/0033-2909.129.5.674>.
14. Hatzenbuehler, M.L.; Pachankis, J.E. Stigma and Minority Stress as Social Determinants of Health Among Lesbian, Gay, Bisexual, and Transgender Youth. *Pediatr. Clin. N. Am.* **2016**, *63*, 985–997, <https://doi.org/10.1016/j.pcl.2016.07.003>.
15. Hatzenbuehler, M.L. Structural stigma: Research evidence and implications for psychological science. *Am. Psychol.* **2016**, *71*, 742–751, <https://doi.org/10.1037/amp0000068>.
16. Testa, R.J.; Michaels, M.S.; Bliss, W.; Rogers, M.L.; Balsam, K.F.; Joiner, T. Suicidal ideation in transgender people: Gender minority stress and interpersonal theory factors. *J. Abnorm. Psychol.* **2017**, *126*, 125–136, <https://doi.org/10.1037/abn0000234>.
17. Testa, R.J.; Habarth, J.; Peta, J.; Balsam, K.; Bockting, W. Development of the Gender Minority Stress and Resilience Measure. *Psychol. Sex. Orientat. Gend. Divers.* **2015**, *2*, 65–77, <https://doi.org/10.1037/sgd0000081>.
18. Connelly, P.; Freely, E.M.; Perry, C.; Ewan, J.; Touyz, R.M.; Currie, G.; Delles, C. Gender-Affirming Hormone Therapy, Vascular Health and Cardiovascular Disease in Transgender Adults. *Hypertension* **2019**, *74*, 1266–1274, <https://doi.org/10.1161/hypertensionaha.119.13080>.
19. Pharr, J.R.; James, T.; Yeung, Y.-L. Accessibility and accommodations for patients with mobility disabilities in a large healthcare system: How are we doing? *Disabil. Health J.* **2019**, *12*, 679–684, <https://doi.org/10.1016/j.dhjo.2019.03.008>.
20. Alzahrani, T.; Nguyen, T.; Ryan, A.; Dwairy, A.; McCaffrey, J.; Yunus, R.; Forgione, J.; Krepp, J.; Nagy, C.; Mazhari, R.; et al. Cardiovascular Disease Risk Factors and Myocardial Infarction in the Transgender Population. *Circ. Cardiovasc. Qual. Outcomes* **2019**, *12*, e005597, <https://doi.org/10.1161/circoutcomes.119.005597>.
21. Morales-Estrella, J.L.; Boyle, M.; Zein, J.G. Transgender status is associated with higher risk of lifetime asthma. In: *A34. Asthma Clinical Studies I*; American Thoracic Society: New York, NY, USA, 2018; p. A1371.
22. Armstrong, B.K.; Cust, A.E. Sun exposure and skin cancer, and the puzzle of cutaneous melanoma: A perspective on fears et al. mathematical models of age and ultraviolet effects on the incidence of skin cancer among whites in the United States. *American journal of epidemiology* **1977**, *105*, 420–427. *Cancer Epidemiol.* **2017**, *48*, 147–156.
23. Silverberg, M.J.; Nash, R.; Becerra-Culqui, T.A.; Cromwell, L.; Getahun, D.; Hunkeler, E.; Lash, T.L.; Millman, A.; Quinn, V.P.; Robinson, B.; et al. Cohort study of cancer risk among insured transgender people. *Ann. Epidemiol.* **2017**, *27*, 499–501, <https://doi.org/10.1016/j.annepidem.2017.07.007>.
24. Fredriksen-Goldsen, K.I.; Cook-Daniels, M.L.; Kim, H.-J.; Erosheva, E.A.; Emlet, C.; Hoy-Ellis, P.C.P.; Goldsen, B.J.; Muraco, A. Physical and Mental Health of Transgender Older Adults: An At-Risk and Underserved Population. *Gerontology* **2013**, *54*, 488–500, <https://doi.org/10.1093/geront/gnt021>.
25. U.S. Equal Employment Opportunity Commission. The Americans with Disabilities Act Amendments Acts of 2008. Available online: <https://www.eeoc.gov/statutes/ada-amendments-act-2008> (accessed on 10 August 2021).
26. Okoro, C.A.; Hollis, N.D.; Cyrus, A.C.; Griffin-Blake, S. Prevalence of Disabilities and Health Care Access by Disability Status and Type Among Adults—United States, 2016. *MMWR. Morb. Mortal. Wkly. Rep.* **2018**, *67*, 882–887, <https://doi.org/10.15585/mmwr.mm6732a3>.

27. Ward, M.J. *Chapter 4A Personal Perspective on Historical Views of Disability*; American Public Health Association: Washington, DC, USA, 2009; <https://doi.org/10.2105/9780875531915ch04>.
28. Buljevac, M.; Majdak, M.; Leutar, Z. The stigma of disability: Croatian experiences. *Disabil. Rehabil.* **2011**, *34*, 725–732, <https://doi.org/10.3109/09638288.2011.616570>.
29. Fredriksen-Goldsen, K.I.; Kim, H.J.; Shui, C.; Bryan, A.E. Chronic health conditions and key health indicators among lesbian, gay, and bisexual older US adults, 2013–2014. *Am. J. Public Health* **2017**, *107*, 1332–1338.
30. Fredriksen-Goldsen, K.I.; Kim, H.-J.; Barkan, S.E.; Muraco, A.; Hoy-Ellis, C.P. Health Disparities Among Lesbian, Gay, and Bisexual Older Adults: Results from a Population-Based Study. *Am. J. Public Health* **2013**, *103*, 1802–1809, <https://doi.org/10.2105/ajph.2012.301110>.
31. Eliason, M.J.; Martinson, M.; Carabez, R.M. Disability Among Sexual Minority Women: Descriptive Data from an Invisible Population. *LGBT Health* **2015**, *2*, 113–120, <https://doi.org/10.1089/lgbt.2014.0091>.
32. Centers for Disease Control and Prevention. Behavioral Risk Factor Surveillance System (BRFSS) Overview 2017. 2018. Available online: https://www.cdc.gov/brfss/annual_data/2017/pdf/overview-2017-508.pdf (accessed on 21 August 2021).
33. Centers for Disease Control and Prevention: CDC—BRFSS—2019 BRFSS Modules Used by Category. 2020. Available online: <https://www.cdc.gov/brfss/questionnaires/modules/category2019.htm> (accessed on 5 June 2021).
34. Marcelo, G. Bioaerosol Size Effect in COVID-19 Transmission. 2020. Available online: <https://www.preprints.org/manuscript/202004.0093/v1/download> (accessed on 21 August 2021).
35. Center for Disease Control and Prevention. Behavioral Risk Factor Surveillance System Complex Sampling Weighting and Preparing 2017 BRFSS Module Data for Analysis. 2018. Available online: https://www.cdc.gov/brfss/annual_data/2017/pdf/Complex-Smple-Weights-Prep-Module-Data-Analysis-2017-508.pdf (accessed on 2 January 2021).
36. Centers for Disease Control and Prevention. Behavioral Risk Factor Surveillance System: 2017 Codebook Report. Available online: https://www.cdc.gov/brfss/annual_data/2017/pdf/codebook17_llcp-508.pdf (accessed on June 2021).
37. Centers for Disease Control and Prevention. Behavioral Risk Factor Surveillance System: 2018 Codebook Report. Available online: https://www.cdc.gov/brfss/annual_data/2018/pdf/codebook18_llcp-v2-508.pdf (accessed on 7 June 2021).
38. Centers for Disease Control and Prevention. Behavioral Risk Factor Surveillance System: 2019 Codebook Report. Available online: https://www.cdc.gov/brfss/annual_data/2019/pdf/codebook19_llcp-v2-508.HTML (accessed on 7 June 2021).
39. Montana Department of Public Health & Human Services. *Interpreting and Reporting BRFSS Data*; Montana Department of Public Health & Human Services: Missoula, MT, USA, 2015.
40. North Carolina Department of Health and Human Services. *Technical Notes 2012*; North Carolina Department of Health and Human Services: Raleigh, NC, USA, 2012.
41. Alaska Department of Health and Social Services. *Complete Health Indicator Report of Mental Health—Frequent Mental Distress—Adults (18+)*; Alaska Department of Health and Social Services: Juneau, AK, USA, 2018.
42. Austin, P.C. Optimal caliper widths for propensity-score matching when estimating differences in means and differences in proportions in observational studies. *Pharm. Stat.* **2011**, *10*, 150–161, <https://doi.org/10.1002/pst.433>.
43. Sekhon, J.S.; Grieve, R.D. A matching method for improving covariate balance in cost-effectiveness analyses. *Health Econ.* **2011**, *21*, 695–714, <https://doi.org/10.1002/hec.1748>.
44. Greifer, N. *cobalt: Covariate Balance Tables and Plots, version 4.3.1*; R package; Comprehensive R Archive Network (CRAN): 2021. Available online: <https://cran.r-project.org/web/packages/cobalt/vignettes/cobalt.html> (accessed on 21 August 2021).
45. Zhang, Z.; written on behalf of AME Big-Data Clinical Trial Collaborative Group; Kim, H.J.; Lonjon, G.; Zhu, Y. Balance diagnostics after propensity score matching. *Ann. Transl. Med.* **2019**, *7*, 16, <https://doi.org/10.21037/atm.2018.12.10>.
46. Venables, W.N.; Ripley, B.D. Package MASS. Available online: <http://www.r-project.org> (accessed on 17 October 2012).
47. Center for American Progress. The State of the LGBTQ Community in 2020. Available online: <https://www.americanprogress.org/issues/lgbtq-rights/reports/2020/10/06/491052/state-lgbtq-community-2020/#Ca=10> (accessed on 14 August 2021).
48. Transgender Law Center. Available online: <https://transgenderlawcenter.org/resources/employment/eeocfa> (accessed on 21 August 2021).
49. Sawyer, K.; Thoroughgood, C.; Webster, J. Queering the Gender Binary: Understanding Transgender Workplace Experiences. In *Sexual Orientation and Transgender Issues in Organizations*; Springer: Cham, Switzerland, 2016; pp. 21–42, https://doi.org/10.1007/978-3-319-29623-4_2.
50. Tilcsik, A. Pride and Prejudice: Employment Discrimination against Openly Gay Men in the United States. *Am. J. Sociol.* **2011**, *117*, 586–626, <https://doi.org/10.1086/661653>.
51. Badgett, M.V. Bias in the workplace: Consistent evidence of sexual orientation and gender identity discrimination 1998–2008. *Chi.-Kent L. Rev.* **2009**, *84*, 559.
52. Kcomt, L. Profound health-care discrimination experienced by transgender people: rapid systematic review. *Soc. Work. Health Care* **2019**, *58*, 201–219, <https://doi.org/10.1080/00981389.2018.1532941>.
53. Roberts, T.K.; Fantz, C.R. Barriers to quality health care for the transgender population. *Clin. Biochem.* **2014**, *47*, 983–987, <https://doi.org/10.1016/j.clinbiochem.2014.02.009>.
54. Kachen, A.; Pharr J.R. Health care access and utilization by transgender populations: A United States transgender survey study. *Transgender Health* **2020**, *5*, 141–148.

55. Canadian Mental Health Association. Mental Health for All. Available online: <https://cmhahkpr.ca/wp-content/uploads/2018/02/yadegarfarand-and-meinhold-bergmann.2014.family-rejection-social-isolation-and-loneliness-as-predictors....pdf> (accessed on 13 August 2021).
56. Thompson, L. *Long-Term Care: Support for Family Caregivers*; Georgetown University, Health Policy Institute: Washington, DC, USA, 2004.
57. Fredriksen-Goldsen, K.I.; Emlet, C.; Kim, H.-J.; Muraco, A.; Erosheva, E.A.; Goldsen, J.; Hoy-Ellis, C.P. The Physical and Mental Health of Lesbian, Gay Male, and Bisexual (LGB) Older Adults: The Role of Key Health Indicators and Risk and Protective Factors. *Gerontol.* **2012**, *53*, 664–675, <https://doi.org/10.1093/geront/gns123>.
58. Shiu, C.; Muraco, A.; Fredriksen-Goldsen, K. Invisible Care: Friend and Partner Care Among Older Lesbian, Gay, Bisexual, and Transgender (LGBT) Adults. *J. Soc. Soc. Work. Res.* **2016**, *7*, 527–546, <https://doi.org/10.1086/687325>.
59. U.S. Department of Health and Human Services. Healthy People 2020. Disability and Secondary Conditions. 2010. Available online: <http://www.healthypeople.gov/HP2020/Objectives/ViewObjective.aspx?Id=457&TopicArea=Disability+and+Secondary+Conditions&Objective=DSC+HP2020%e2%80%91914&TopicAreaId=17> accessed on 15 August 2021).
60. Pharr, J. Accessible medical equipment for patients with disabilities in primary care clinics: Why is it lacking? *Disabil. Health J.* **2013**, *6*, 124–132, <https://doi.org/10.1016/j.dhjo.2012.11.002>.
61. Drainoni, M.L.; Lee-Hood, E.; Tobias, C.; Bachman, S.S.; Andrew, J.; Maisels, L. Cross-disability experiences of barriers to health-care access: Consumer perspectives. *J. Disabil. Policy Stud.* **2006**, *17*, 101–115.
62. U.S. Department of Justice (USDJ), U.S. Department of Health and Human Rights (DHHR). Access to Medical Care for Individuals with Mobility Disabilities. 2010. Available online: http://www.ada.gov/medcare_mobility_ta/medcare_ta.pdf (Accessed on 10 August 2021).
63. Brodwin, M.G. Frederick, P.C. Sexuality and societal beliefs regarding persons living with disabilities. *J. Rehabil.* **2010**, *76*, 37.
64. Ganle, J.K.; Baatiema, L.; Quansah, R.; Danso-Appiah, A. Barriers facing persons with disability in accessing sexual and reproductive health services in sub-Saharan Africa: A systematic review. *PLoS ONE* **2020**, *15*, e0238585, <https://doi.org/10.1371/journal.pone.0238585>.
65. Meyer, I.H. Resilience in the study of minority stress and health of sexual and gender minorities. *Psychol. Sex. Orientat. Gend. Divers.* **2015**, *2*, 209–213, <https://doi.org/10.1037/sgd0000132>.