

SUPPLEMENTARY MATERIALS

Biomarkers of response to immunotherapy in poorly differentiated sinonasal carcinomas

Eva Villanueva-Fernández¹, Mario A. Hermsen², Laura Suárez-Fernández², Blanca Vivanco³, Alessandro Franchi⁴, Rocío García-Marín², Virginia N. Cabal², Helena Codina-Martínez², Sara Lucila Lorenzo-Guerra², José L. Llorente¹, Fernando López¹.

Table S1. PMS2, MLH1, MSH2 and MSH6 staining pattern in 5 cases designated as MSI positive.

	PMS2	MLH1	MSH2	MSH6
SNEC patient 1	negative	reduced	positive	positive
SNEC patient 2	negative	negative	negative	negative
SNUC patient 1	negative	reduced	reduced	reduced
SNUC patient 2	negative	negative	reduced	reduced
HG-non-ITAC	reduced	reduced	positive	positive

Figure S1. Representative images of SMARCB1-deficient carcinoma (**A–D**), SMARCA4-deficient carcinoma (**E–H**) and NUT carcinoma (**I–L**) showing hematoxylin and eosin staining (**A,E,I**), SmarCB1 expression (**B**), SmarCA4 expression (**F**) and NUT1 expression (**J**), as well as CD8+ TILs (**C,G,K**) and PD-L1 (**D,H,L**). All images ×200 magnification.

