

Comment

Unreliable Findings Due to Miscalculations and Errors. Comment on Nally et al. The Effectiveness of School-Based Interventions on Obesity-Related Behaviours in Primary School Children: A Systematic Review and Meta-Analysis of Randomised Controlled Trials. *Children* 2021, 8, 489

Keara Ginell ^{1,2}, Beate Henschel ², Lilian Golzarri-Arroyo ², Yasaman Jamshidi-Naeini ²,
Andrew W. Brown ^{3,4,5} and David B. Allison ^{2,*}

¹ Department of Cognitive Science, Vassar College, Poughkeepsie, NY 12604, USA

² Department of Epidemiology and Biostatistics, Indiana University School of Public Health-Bloomington, Bloomington, IN 47405, USA

³ Department of Applied Health Science, Indiana University School of Public Health-Bloomington, Bloomington, IN 47405, USA

⁴ Arkansas Children's Research Institute, Little Rock, AR 72202, USA

⁵ Department of Biostatistics, University of Arkansas for Medical Sciences, Little Rock, AR 72205, USA

* Correspondence: allison@iu.edu



Citation: Ginell, K.; Henschel, B.; Golzarri-Arroyo, L.; Jamshidi-Naeini, Y.; Brown, A.W.; Allison, D.B. Unreliable Findings Due to Miscalculations and Errors. Comment on Nally et al. The Effectiveness of School-Based Interventions on Obesity-Related Behaviours in Primary School Children: A Systematic Review and Meta-Analysis of Randomised Controlled Trials. *Children* 2021, 8, 489. *Children* 2023, 10, 1660. <https://doi.org/10.3390/children10101660>

Academic Editor: George Antonogeorgos

Received: 28 September 2022

Revised: 18 August 2023

Accepted: 20 September 2023

Published: 7 October 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Nally et al. [1] conducted a meta-analysis to examine the effectiveness of school-based interventions in changing BMI among primary school children and reported significant differences in the change in BMI and BMI z-score between intervention and control conditions in favor of the intervention. However, the conduct of the meta-analysis as reported in Figures 3 and 4 is not consistent with the reported methods and involves invalid input data and some undisclosed analytical approaches.

First, the results in Figures 3 and 4 do not match the methods and figure captions. The captions of Figures 3 and 4 read “Forest plot for standardised mean difference of change in BMI (kg/m²) between intervention and control groups [...]” and “Forest plot for standardised mean difference of change in BMI z-score between intervention and control groups [...]” [1], respectively. Upon reviewing Adab et al. [2], Anderson et al. [3], Angelopoulos et al. [4], and Tarro et al. [5], it appears that Figure 3 does not show *standardized* mean differences of *change scores*, but rather *raw* mean differences of *post-intervention* values. The data shown in Figure 4 are post-intervention values and not change scores.

Second, of the 48 studies in Nally et al.'s [1] qualitative analysis, they report excluding 10 studies from the meta-analysis for not reporting the variance in data and other reasons. The study by Rausch Herscovici et al. [6], which has the largest contribution (35.8%) to the weighted average (Figure 3), did not report a measure of variation to calculate between-group effect sizes using standard accepted calculations. Nonetheless, it was included in the meta-analysis. Even if the study by Rausch Herscovici et al. [6] was to be included, it reported findings that were not statistically significant. Yet, Nally et al. [1] concluded statistically significant intervention effects in BMI [−1.00 (−1.15, −0.85)] and BMI z-scores [−2.93 (−3.23, −2.63)] for that study (Figures 3 and 4). Methods for including studies with incomplete information are available (e.g., imputing or estimating variance, and other approaches [7]), but no such methods were disclosed by Nally et al. [1], and Nally et al.'s [1] approach led to estimates that conflict with Rausch Herscovici et al.'s [6] conclusion about statistical significance.

Third, the data used for the study by Ford et al. [8] in Figures 3 and 4 do not match the report in the original study. Values reported in Figure 3 are implausibly small (i.e., mean of 1.085 and 1.04 for the intervention and control, respectively) to be post-intervention BMI

values as reported for all other studies in Figure 3, and they do not match the change from baseline BMI values reported by Ford et al. [8]. We infer that Nally et al. [1] incorrectly used change in body mass (kg) values instead of BMI in Figure 3's analyses and change in BMI values instead of BMI z-scores in Figure 4's analyses. Ford et al. [8] did not report BMI z-scores; rather, they reported the z-scores for the change in BMI that would be used in a z-test to obtain a *p*-value, whereas BMI z-score is an anthropometric measure in children. Thus, the Ford et al. [8] study does not report the outcome of interest to be included in the meta-analysis in Figure 4. However, we can keep the Ford et al. [8] study in Figure 3 with corrected values for the mean difference of 0.00 (95% CI: −0.35, 0.35).

After removing Rausch Herscovici et al. [6] and replacing the values for Ford et al. [8] for Figure 3, we re-analyzed the data using a fixed-effects model (consistent with Nally et al.'s [1] methods) and show no evidence indicating significant effects for the mean difference in BMI between intervention and control groups (−0.04 kg/m²; 95% CI = −0.15, 0.06; *I*² = 53.8%, Table 1). This contrasts with Nally et al.'s [1] conclusion that BMI and BMI z-score were significantly reduced in the intervention group. We note that given that the included studies have interventions that are expected to introduce design and exposure heterogeneity, combined with evidence of moderate to substantial statistical heterogeneity, an appropriate analysis is a random-effects model. Thus, we also re-analyzed the data using a random-effects model, which resulted in similar conclusions about intervention efficacy to our fixed-effects model (Table 1). Likewise, for Figure 4, after removing Rausch Herscovici et al. [6] and Ford et al. [8], the overall standardized mean difference in BMI z-score between intervention and control groups was no longer statistically significantly different in both the fixed- and random-effects models but with a lower heterogeneity compared to Nally et al. [1] (Table 1).

Table 1. Results from re-analysis after corrected data extraction and including random-effects results.

	Figure 3: Mean Differences in BMI (95% CI)	Figure 4: Standardized Mean Differences in BMI z-scores (95% CI)
Reported in Nally et al. [1]	<i>I</i> ² = 86%	<i>I</i> ² = 96%
- Fixed-effects model	−0.39 (−0.47, −0.30)	−0.05 (−0.08, −0.02)
Re-analysis ¹	<i>I</i> ² = 53.8%	<i>I</i> ² = 59.3%
- Fixed-effects model	−0.04 (−0.15, 0.06)	−0.02 (−0.05, 0.01)
- Random-effects model	−0.07 (−0.24, 0.11)	−0.02 (−0.07, 0.03)

¹ Re-analysis for Figure 3 excluded the Rausch Herscovici et al. [6] study and used a mean difference of 0.00 (95% CI: −0.35, 0.35) for Ford et al. [8]. Re-analysis for Figure 4 excluded both the Rausch Herscovici et al. [6] and Ford et al. [8] studies.

The Committee on Publication Ethics' Retraction Guidelines state that retraction should be considered if there is "clear evidence that the findings are unreliable . . . as a result of major error (e.g., miscalculation or experimental error)" [9]. Although we focus herein on examples of studies that substantially affect the results, we noted other inconsistencies in reporting of the original studies' data. We further note that there are appropriate ways to include some studies that were excluded by Nally et al. [1] because of the incomplete reporting of effects by the original investigators. The findings are therefore unreliable because of miscalculation and error, and we therefore respectfully conclude that the meta-analysis by Nally et al. [1] should be retracted.

Author Contributions: Conceptualization, K.G. and D.B.A.; data curation, K.G.; formal analysis, K.G.; validation, B.H. and L.G.-A.; writing—original draft preparation, K.G.; writing—review and editing, B.H., L.G.-A., Y.J.-N., A.W.B. and D.B.A.; supervision, D.B.A. All authors have read and agreed to the published version of the manuscript.

Funding: This work was funded in part by the National Institutes of Health (NIH): R25HL124208 and R25DK099080, and the Gordon and Betty Moore Foundation. The opinions expressed are those of the authors and not necessarily of the NIH or any other organization.

Data Availability Statement: Data from our re-analysis are readily available at <https://doi.org/10.17605/OSF.IO/6Q57B> (accessed on 14 July 2022).

Conflicts of Interest: The authors collectively declare the following disclosures. In the 36 months prior to original submission: Personal payments or promises for same, donations to a foundation on their behalf, or travel expenses: Alkermes, Inc.; Alliance for Potato Research and Education; Amin Talati Wasserman for KSF Acquisition Corp (Glanbia); Big Sky Health, Inc.; Clark Hill PLC; Northarvest Bean Growers Association; International Food Information Council; Kaleido Biosciences; Law Offices of Ronald Marron; Medpace/Gelesis; Novo Nordisk Fonden; Purchaser Business Group on Health; Soy Nutrition Institute Global; and Sports Research Corp. Family member employed by Reckitt Benckiser. Indiana University, which employs or employed D.B.A, B.H., L.G.-A., A.W.B., and Y.J.-N., and the Indiana University Foundation has received funds or donations to support research or educational activities from Alliance for Potato Research and Education; American Egg Board; Arnold Ventures; Eli Lilly and Company; Haas Avocado Board; Mars, Inc.; National Cattlemen’s Beef Association; Reckitt Benckiser Group PLC; Soleno Therapeutics; WW (formerly Weight Watchers); Center for Open Science; Gordon and Betty Moore Foundation; Indiana CTSI; Sloan Foundation; and numerous other for-profit and non-profit organizations to support the work of the School of Public Health and the university more broadly. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

References

1. Nally, S.; Carlin, A.; Blackburn, N.E.; Baird, J.S.; Salmon, J.; Murphy, M.H.; Gallagher, A.M. The effectiveness of school-based interventions on obesity-related behaviours in primary school children: A systematic review and meta-analysis of randomised controlled trials. *Children* **2021**, *8*, 489. [\[CrossRef\]](#) [\[PubMed\]](#)
2. Adab, P.; Pallan, M.J.; Lancashire, E.R.; Hemming, K.; Frew, E.; Barrett, T.; Bhopal, R.; E Cade, J.; Canaway, A.; Clarke, J.; et al. Faculty Opinions recommendation of Effectiveness of a childhood obesity prevention programme delivered through schools, targeting 6 and 7 year olds: Cluster randomised controlled trial (WAVES study). *BMJ* **2018**, *360*, k211. [\[CrossRef\]](#) [\[PubMed\]](#)
3. Anderson, E.L.; Howe, L.D.; Kipping, R.R.; Campbell, R.; Jago, R.; Noble, S.M.; Wells, S.; Chittleborough, C.; Peters, T.; A Lawlor, D. Long-term effects of the Active for Life Year 5 (AFLY5) school-based cluster-randomised controlled trial. *BMJ Open* **2016**, *6*, e010957. [\[CrossRef\]](#) [\[PubMed\]](#)
4. Angelopoulos, P.D.; Milonitis, H.J.; Grammatikaki, E.; Moschonis, G.; Manios, Y. Changes in BMI and blood pressure after a school based intervention: The CHILDREN study. *Eur. J. Public Health* **2009**, *19*, 319–325. [\[CrossRef\]](#) [\[PubMed\]](#)
5. Tarro, L.; Llauro, E.; Albaladejo, R.; Morina, D.; Arija, V.; Solà, R.; Giral, M. A primary-school-based study to reduce the prevalence of childhood obesity—The EdAl (Educació en Alimentació) study: A randomized controlled trial. *Trials* **2014**, *15*, 58. [\[CrossRef\]](#) [\[PubMed\]](#)
6. Rausch Herscovici, C.; Kovalskys, I.; De Gregorio, M.J. Gender differences and a school-based obesity prevention program in Argentina: A randomized trial. *Rev. Panam. Salud Pública* **2013**, *34*, 75–82. [\[PubMed\]](#)
7. Higgins, J.P.T.; Li, T.; Deeks, J.J. (Eds.) Chapter 6: Choosing effect measures and computing estimates of effect. In *Cochrane Handbook for Systematic Reviews of Interventions Version 6.2 (Updated February 2021)*; Cochrane: Oxford, UK, 2021; Available online: www.training.cochrane.org/handbook (accessed on 14 July 2022).
8. Ford, P.A.; Perkins, G.; Swaine, I. Effects of a 15-week accumulated brisk walking programme on the body composition of primary school children. *J. Sports Sci.* **2012**, *31*, 114–122. [\[CrossRef\]](#) [\[PubMed\]](#)
9. Barbour, V.; Kleinert, S.; Wager, E.; Yentis, S. *Guidelines for Retracting Articles*; Committee on Publication Ethics: Harlestone, UK, 2009; Volume 10.

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.