

Correction

Correction: Supramani et al. Performance of Biomass and Exopolysaccharide Production from the Medicinal Mushroom *Ganoderma lucidum* in a New Fabricated Air-L-Shaped Bioreactor (ALSB). *Processes* 2023, 11, 670

Sugenendran Supramani ¹, Nur Ardiyana Rejab ², Zul Ilham ^{3,4} , Rahayu Ahmad ^{5,*}, Pau-Loke Show ^{6,7,8,9} , Mohamad Faizal Ibrahim ¹⁰  and Wan Abd Al Qadr Imad Wan-Mohtar ^{1,3,11,*} 

- ¹ Functional Omics and Bioprocess Development Laboratory, Institute of Biological Sciences, Faculty of Science, University Malaya, Kuala Lumpur 50603, Malaysia
- ² Programme of Genetics, Institute of Biological Sciences, Faculty of Science, University Malaya, Kuala Lumpur 50603, Malaysia
- ³ Bioresources and Bioprocessing Research Group, Institute of Biological Sciences, Faculty of Science, University Malaya, Serdang 50603, Malaysia; ilham@um.edu.my
- ⁴ Environmental Science and Management Program, Institute of Biological Sciences, Faculty of Science, University Malaya, Kuala Lumpur 50603, Malaysia
- ⁵ Halal Action Laboratory, Kolej GENIUS Insan, University Sains Islam Malaysia, Nilai 71800, Malaysia
- ⁶ Department of Chemical Engineering, Khalifa University, Shakhboub Bin Sultan St-Zone 1, Abu Dhabi P.O. Box 127788, United Arab Emirates
- ⁷ Zhejiang Provincial Key Laboratory for Subtropical Water Environment and Marine Biological Resources Protection, Wenzhou University, Wenzhou 325035, China
- ⁸ Department of Chemical and Environmental Engineering, Faculty of Science and Engineering, University of Nottingham Malaysia, Semenyih 43500, Malaysia
- ⁹ Department of Sustainable Engineering, Saveetha School of Engineering, SIMATS, Chennai 602105, India
- ¹⁰ Department of Bioprocess Technology, Faculty of Biotechnology and Biomolecular Sciences, University Putra Malaysia, Serdang 43400, Malaysia
- ¹¹ Bioscience Research Institute, Technical University Shannon Midlands Midwest, N37 HD68 Athlone, Ireland
- * Correspondence: rahayu@usim.edu.my (R.A.); qadyr@um.edu.my (W.A.A.Q.I.W.-M.)



Received: 8 April 2025
Accepted: 9 April 2025
Published: 23 April 2025

Citation: Supramani, S.; Rejab, N.A.; Ilham, Z.; Ahmad, R.; Show, P.-L.; Ibrahim, M.F.; Wan-Mohtar, W.A.A.Q.I. Correction: Supramani et al. Performance of Biomass and Exopolysaccharide Production from the Medicinal Mushroom *Ganoderma lucidum* in a New Fabricated Air-L-Shaped Bioreactor (ALSB). *Processes* 2023, 11, 670. *Processes* 2025, 13, 1280. <https://doi.org/10.3390/pr13051280>

Copyright: © 2025 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Error in Table

In the original publication [1], there was a mistake in the caption for Table 2, “Kinetic parameters of *G. lucidum* in shake flask compared to ALSB”, as published. The caption was misleading. The corrected Table 2 Caption, “Total biomass and EPS yield of *Ganoderma lucidum* in both shake flask and ALSB”, appears below.

Table 2. Total biomass and EPS yield of *Ganoderma lucidum* in both shake flask and ALSB.

Mushroom Cultivation Method	Biomass Concentration (g/L)	Biomass Productivity (g/L/day)	EPS Concentration (g/L)	EPS Productivity (g/L/day)	Specific Production Rate of EPS (lg/g/L/day)
Shake Flask	12.31 ± 0.07	0.820 ± 0.02	8.22 ± 0.07	0.548 ± 0.02	0.045 ± 0.02
ALSB	33.96 ± 0.03	2.264 ± 0.03	15.52 ± 0.05	1.035 ± 0.03	0.305 ± 0.03

Text Correction

A correction has been made to Section 3.1.

As shown in Figure 2, the standard growth curve of the ALSB is evidently higher than that of the shake flask and follows the typical fungal fermentation growth curve (exponential phase from day 5 and transition phase from day 10) [18]. The highest biomass concentration in the shake flask was 2.6 g/L on day 9 compared to 5.7 g/L for the ALSB on

day 11. Meanwhile, the highest EPS in the shake flask was 1.68 g/L on day 11 compared to ALSB (3.8 g/L) on day 11. In Table 2, the total biomass and EPS yield of *Ganoderma lucidum* in both the shake flask and the ALSB are shown. The total biomass yield for the 15 days was 12.31 g/L in the shake flask and 33.96 g/L in the ALSB (2.8-fold higher in the ALSB).

The authors state that the scientific conclusions are unaffected. This correction was approved by the Academic Editor. The original publication has also been updated.

Reference

1. Supramani, S.; Rejab, N.A.; Ilham, Z.; Ahmad, R.; Show, P.-L.; Ibrahim, M.F.; Wan-Mohtar, W.A.A.Q.I. Performance of Biomass and Exopolysaccharide Production from the Medicinal Mushroom *Ganoderma lucidum* in a New Fabricated Air-L-Shaped Bioreactor (ALSB). *Processes* **2023**, *11*, 670. [[CrossRef](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.