

Article

Imbalance-Based Irregularity Molecular Descriptors of Nanostar Dendrimers

Zafar Hussain ¹, Mobeen Munir ², Shazia Rafique ³, Tayyab Hussnain ³, Haseeb Ahmad ², Young Chel Kwun ^{4,*} and Shin Min Kang ^{5,6}

¹ Department of Mathematics and Statistics, The University of Lahore, Lahore 54500, Pakistan

² Department of Mathematics, Division of Science and Technology, University of Education, Lahore 54000, Pakistan

³ Center for Excellence in Molecular Biology, Punjab University Lahore, Lahore 53700, Pakistan

⁴ Department of Mathematics, Dong-A University, Busan 49315, Korea

⁵ Department of Mathematics, Gyeongsang National University, Jinju 52828, Korea

⁶ Center for General Education, China Medical University, Taichung 40402, Taiwan

* Correspondence: yckwun@dau.ac.kr

Received: 27 June 2019; Accepted: 29 July 2019; Published: 6 August 2019



Abstract: Dendrimers are branched organic macromolecules with successive layers of branch units surrounding a central core. The molecular topology and the irregularity of their structure plays a central role in determining structural properties like enthalpy and entropy. Irregularity indices which are based on the imbalance of edges are determined for the molecular graphs associated with some general classes of dendrimers. We also provide graphical analysis of these indices for the above said classes of dendrimers.

Keywords: nanostar dendrimer; irregularity measure; complexity of structure; $NS_1[p]$; $NS_2[p]$; $NS_3[p]$

1. Introduction

Algebra, topology, geometry and combinatorics are the main branches of mathematics which are employed to study the symmetries and irregularities of the structures of different substances. Dendrimers have consistently attracted the attention of both chemists as well as pure mathematicians because of the complexities of the underlying molecular graphs. Dendrimers are highly branched, star-shaped macromolecules with nanometer-scale dimensions. Dendrimers are constituted by main parts: A central core, an internal part called ‘branch’, and an exterior surface with functional surface groups. The varied combination of these components yields products of different shapes and sizes with shielded interior cores that are ideal candidates for applications in both biological and materials sciences. Some recent applications include drug delivery, gene transfection, catalysis, energy harvesting, photo activity, molecular weight and size determination, rheology modification, and nanoscale science and technology [1–3].

Graphs can be used to study theoretical and computational aspects of dendrimers. Recently this approach has proved remarkable in relating properties of substances with involved structural parameters [4–7]. Topological indices are used here as major ingredients [7–14]. Some nanotubes, modified electrodes, chemical sensors, micro- and macro-capsule, and colored glasses can be designed using nanostar dendrimers. The structure of polymer molecules in a plane depends on the adjacency of their units. For detailed insight, see [1–3,14–18] and references therein. Figure 1 shows the spatial arrangements of $NS_1[1]$, $NS_1[2]$ polypropylenimine octaamin dendrimers in plane. The recursive nature of these dendrimers is evident from this figure. Graph theoretic models of these dendrimers can potentially be used in fractals.




Figure 1. $NS_1[1]$ and $NS_1[2]$ polypropylenimine octaamin dendrimers.

In Figure 1, G_1 shows the structure of polypropylenimine octaamin dendrimers $NS_1[p]$ when $p = 1$, and G_2 represents the structure of $NS_1[p]$ when $p = 2$.

The next object will be polypropylenimine octaamin dendrimer $NS_2[p]$. Figure 2 is a graph theoretical representation for this dendrimer.




Figure 2. $NS_2[p]$ polypropylenimine octaamin dendrimers.

The third object of interest is the $NS_3[p]$, also known as polymer dendrimer. Figure 3 shows the molecular structure of this dendrimer.




Figure 3. $NS_3[p]$ polymer de ndrimer.

The above three families have been used a lot recently for their theoretical properties. De et al. in [14], computed the F-index of nanostar dendrimers, Siddiqui et al. computed the Zagreb indices for different nanostar dendrimers in [15], and Madanshekaf computed the Randi index for some different classes of nanostar dendrimers in [16,17]. Munir, et al. computed M-polynomial and related indices of these nanostar dendrimers in [18], titania nanotubes in [19], polyhez nanotubes in [20] and circulant graphs in [21].

In the current article, we are interested in imbalance-based irregularity indices of the above discussed families of three dendrimers. We use techniques from combinatorics and graph theory to avoid the use of quantum mechanics, as has been done recently in most of the cases, see [7–14]. Important tools which are used for this purpose are structural and functional polynomials. These tools use structural parameters as inputs and the outputs are the key information that is used to determine properties of the material under discussion. Certain properties of matters like standard enthalpy, toxicity, entropy as well as reactivity and biological mechanics are theoretically based on these tools [4–6]. Estrada related the atom bond connectivity index with energies of the branched alkanes in [9]. Some applications of indices in pharmacy are given in [10] and in Quantitative structure activity analysis in [11].

2. Preliminaries and Notations

In this part we lay out some basic material and notations which will be used throughout the article. All graphs will be connected. We fix the symbol G for a simple connected graph, V for the set of vertices of G , E for the set of edges, du and dv are the degrees of vertices u and v , respectively. Topological index is an invariant of the graph that preserves the structural aspects of the graph. A degree based topological index is based on the end degrees of edges. A graph is said to be regular if every vertex of the graph has the same degree. A topological invariant is called irregularity index if the index vanishes for a regular graph and is non-zero for a non-regular graph. Regular graphs have been investigated a lot, particularly in mathematics. Their applications in chemical graph theory came to be known after the discovery of nanotubes and fullerenes. Paul Erdos emphasized the study of irregular graphs for the first time in history in [22]. At the Second Krakow Conference on Graph Theory (1994), Erdos officially

posed an open problem about determination of the extreme size of highly irregular graphs of given order [23]. Since then, the irregular graphs and the degree of irregularity have become the basic open problem of graph theory. A graph in which each vertex has a different degree than other vertices is known as a perfect graph. Authors in [24], proved that no graph is perfect. The graphs lying in between are called quasi-perfect graphs in which each, except two vertices, have different degrees [25]. A simplified way of expressing the irregularity is the irregularity index. Irregularity indices have been studied recently in a novel way [26,27]. The first such irregularity index was introduced in [28]. Most of these indices used the concept of imbalance of an edge defined as $imball_{uv} = |d_u - d_v|$, [25–27]. The Albertson index, AL(G), was defined by Alberston in [29] as $AL(G) = \sum_{UV \in E} |d_u - d_v|$. In this index, the imbalance of edges is computed. The irregularity index IRL(G) and IRLU(G) is introduced by Vukicevic and Gasparov, [30] as $IRL(G) = \sum_{UV \in E} |lnd_u - lnd_v|$, and $IRLU(G) = \sum_{UV \in E} \frac{|d_u - d_v|}{\min(d_u, d_v)}$. Recently, Abdoo et al. introduced the new term “total irregularity measure of a graph G”, which is defined as [31–33] $IRR_t(G) = \frac{1}{2} \sum_{UV \in E} |d_u - d_v|$. Recently, Gutman et al. introduced the $\sigma(G)$ irregularity index of the graph G, which is described as $\sigma(G) = \sum_{UV \in E} (d_u - d_v)^2$ in [34]. The Randic index itself is directly related to an irregularity measure, which is described as $IRA(G) = \sum_{UV \in E} \left(\frac{1}{d_u} - \frac{1}{d_v} \right)^2$ in [35]. Further irregularity indices of similar nature can be traced in [34] in detail. These indices are given as $IRDIF(G) = \sum_{UV \in E} \left| \frac{d_u}{d_v} - \frac{d_v}{d_u} \right|$, $IRLF(G) = \sum_{UV \in E} \frac{|d_u - d_v|}{\sqrt{(d_u d_v)}}$, $LA(G) = 2 \sum_{UV \in E} \frac{|d_u - d_v|}{(d_u + d_v)}$, $IRD1 = \sum_{UV \in E} \ln \{1 + |d_u - d_v|\}$, $IRGA(G) = \sum_{UV \in E} \ln \frac{d_u + d_v}{2 \sqrt{(d_u d_v)}}$, and $IRB(G) = \sum_{UV \in E} \left(\frac{1}{d_u} - \frac{1}{d_v} \right)^2$. Futher details are given in [28–46]. Recently authors computed irregularity indices of a nanotube [47]. Recently Gao et al. computed irregularity measures of some dendrimer structures in [48] and molecular structures in [49]. Actually, the authors computed only four irregularity measures for some classes of dendrimers in [48]. These structures are used as long infinite chain macromolecules in chemistry and related areas. Hussain et al. computed these irregularity measures for some classes of benzenoid systems in [50].

3. Main Results

In this part, we give our main theoretical results.

Theorem 1. Let $NS_1[p]$ be the polypropylenimine octaamin dendrimers, then the irregularity indices of $NS_1[p]$ are:

1. $IRDIF(NS_1[p]) = 1.5(2^{p+1}) + 22.3372^p - 22.334$
2. $IRR(NS_1[p]) = 2^{p+1} + 22(2^p) - 22$
3. $IRL(NS_1[p]) = 0.69314718(2^{p+1}) + 10.070961(2^p) - 10.070961$
4. $IRLU(NS_1[p]) = 2^{p+1} + 15(2^p) - 15$
5. $IRLF(NS_1[p]) = 0.7071068(2^{p+1}) + 10.334278(2^p) - 10.334278$
6. $\sigma(NS_1[p]) = 2^{p+1} + 30(2^p) - 30$
7. $IRLA(NS_1[p]) = 0.6667(2^p) + 9.6(2^p) - 9.6$
8. $IRD1 = 0.69314(2^{p+1}) + 14.098(2^p) - 14.098$
9. $IRA(NS_1[p]) = 0.085786432(2^{p+1}) + 0.950245633(2^p) - 0.9502456337$
10. $IRGA(NS_1[p]) = 0.06036(2^{p+1}) + 0.8610954(2^p) - 0.8610954$
11. $IRB(NS_1[p]) = 0.171528753(2^{p+1}) + 3.557593539(2^p) - 3.557593539$
12. $IRR_t(NS_1[p]) = 12(2^p) - 11$

Proof. In order to prove the above theorem we have to consider Figure 1.

We can see that the edges $NS_1[p]$ admit the following partition in Table 1. □

Table 1. Edge partition of $NS_1[p]$ polypropylenimine octaamin dendrimers.

Edge Type (d_u, d_v)	Number of Edges
(1,2)	2^{p+1}
(1,3)	$4(2^p - 1)$
(2,2)	$\{12 \times 2^p - 11\}$
(2,3)	$14(2^p - 1)$

Now, using above Table 1 and definitions we have:

1. $IRDIF(G) = \sum_{UV \in E} \left| \frac{d_u}{d_v} - \frac{d_v}{d_u} \right|$

$$IRDIF(NS_1[p], x, y) = 2^{p+1} \left| \frac{2}{1} - \frac{1}{2} \right| + 4(2^p - 1) \left| \frac{3}{1} - \frac{1}{3} \right| + \{12 \times 2^p - 11\} \left| \frac{2}{2} - \frac{2}{2} \right| + 14(2^p - 1) \left| \frac{3}{2} - \frac{2}{3} \right|$$

$$= 2^{p+1} \left| \frac{2}{1} - \frac{1}{2} \right| + 4(2^p - 1) \left| \frac{3}{1} - \frac{1}{3} \right| + 14(2^p - 1) \left| \frac{3}{2} - \frac{2}{3} \right|.$$
2. $IRR(G) = \sum_{UV \in E} imb(e)$

$$IRR(NS_1[p], x, y) = 2^{p+1} |2 - 1| + 4(2^p - 1) |3 - 1| + \{12 \times 2^p - 11\} |2 - 2| + 14(2^p - 1)$$

$$= 2^{p+1} + 4(2^p - 1) |2| + 14(2^p - 1).$$
3. $IRL(G) = \sum_{UV \in E} |lnd_u - lnd_v|$

$$IRL(NS_1[p], x, y) = 2^{p+1} |ln2 - ln1| + 4(2^p - 1) |ln3 - ln1| + \{12 \times 2^p - 11\} |ln2 - ln2| + 14(2^p - 1) |ln3 - ln2|$$

$$= 2^{p+1} |ln2| + 4(2^p - 1) |ln3| + 14(2^p - 1) |ln\frac{3}{2}|.$$
4. $IRLU(G) = \sum_{UV \in E} \frac{|d_u - d_v|}{\min(d_u, d_v)}$

$$IRLU(NS_1[p], x, y) = 2^{p+1} \frac{|2-1|}{1} + 4(2^p - 1) \frac{|3-1|}{1} + \{12 \times 2^p - 11\} \frac{|2-2|}{2} + 14(2^p - 1) \frac{|3-2|}{2}$$

$$= 2^{p+1} + 8(2^p - 1) + 7(2^p - 1).$$
5. $IRLF(G) = \sum_{UV \in E} \frac{|d_u - d_v|}{\sqrt{(d_u d_v)}}$

$$IRLF(NS_1[p], x, y) = 2^{p+1} \frac{|2-1|}{\sqrt{2}} + 4(2^p - 1) \frac{|3-1|}{\sqrt{3}} + \{12 \times 2^p - 11\} \frac{|2-2|}{\sqrt{4}} + 14(2^p - 1) \frac{|3-2|}{\sqrt{6}}$$

$$= \frac{2^{p+1}}{\sqrt{2}} + \frac{8(2^p - 1)}{\sqrt{3}} + \frac{14(2^p - 1)}{\sqrt{6}}.$$
6. $\sigma(G) = \sum_{UV \in E} (d_u - d_v)^2$

$$\sigma(NS_1[p], x, y) = 2^{p+1} (2 - 1)^2 + 4(2^p - 1) (3 - 1)^2 + \{12 \times 2^p - 11\} (2 - 2)^2 + 14(2^p - 1) (3 - 2)^2$$

$$= 2^{p+1} + 16(2^p - 1) + 14(2^p - 1).$$
7. $IRLA(G) = 2 \sum_{UV \in E} \frac{|d_u - d_v|}{(d_u + d_v)}$

$$IRLA(NS_1[p], x, y) = 2 \left[2^{p+1} \frac{|2-1|}{(2+1)} + 4(2^p - 1) \frac{|3-1|}{(3+1)} + \{12 \times 2^p - 11\} \frac{|2-2|}{(2+2)} + 14(2^p - 1) \frac{|3-2|}{(3+2)} \right]$$

$$= 2 \left[2^{p+1} \frac{|2-1|}{(2+1)} + 4(2^p - 1) \frac{|3-1|}{(3+1)} + 14(2^p - 1) \frac{|3-2|}{(3+2)} \right].$$
8. $IRD1 = \sum_{UV \in E} \ln\{1 + |d_v - d_u|\}$

$$IRD1(NS_1[n], x, y) = 2^{p+1} \ln\{1 + |2 - 1|\} + 4(2^p - 1) \ln\{1 + |3 - 1|\} + \{12 \times 2^p - 11\} \ln\{1 + |2 - 2|\} + 14(2^p - 1) \ln\{1 + |3 - 2|\}$$

$$= 2^{p+1} \ln\{1 + |2 - 1|\} + 4(2^p - 1) \ln\{1 + |3 - 1|\} + 14(2^p - 1) \ln\{1 + |3 - 2|\}.$$

$$9. \quad IRA(G) = \sum_{UV \in E} \left(d_u^{\frac{-1}{2}} - d_v^{\frac{-1}{2}} \right)^2$$

$$\begin{aligned} IRA(NS_1[p], x, y) &= 2^{p+1} \left(\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{1}} \right)^2 + 4(2^p - 1) \left(\frac{1}{\sqrt{3}} - \frac{1}{\sqrt{1}} \right)^2 + \{12 \times 2^p - 11\} \left(\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}} \right)^2 + 14(2^p - 1) \left(\frac{1}{\sqrt{3}} - \frac{1}{\sqrt{2}} \right)^2 \\ &= 2^{p+1} \left(\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{1}} \right)^2 + 4(2^p - 1) \left(\frac{1}{\sqrt{3}} - \frac{1}{\sqrt{1}} \right)^2 + 14(2^p - 1) \left(\frac{1}{\sqrt{3}} - \frac{1}{\sqrt{2}} \right)^2. \end{aligned}$$

$$10. \quad We have IRGA(G) = \sum_{UV \in E} \ln \frac{d_u + d_v}{2 \sqrt{(d_u d_v)}}$$

$$\begin{aligned} IRGA(NS_1[p], x, y) &= 2^{p+1} \ln \frac{|2+1|}{2\sqrt{2}} + 4(2^p - 1) \ln \frac{|3+1|}{2\sqrt{3}} + \{12 \times 2^p - 11\} \ln \frac{|2+2|}{2\sqrt{4}} + 14(2^p - 1) \ln \frac{|3+2|}{2\sqrt{6}} \\ &= 2^{p+1} \ln \frac{|2+1|}{2\sqrt{2}} + 4(2^p - 1) \ln \frac{|3+1|}{2\sqrt{3}} + 14(2^p - 1) \ln \frac{|3+2|}{2\sqrt{6}}. \end{aligned}$$

$$11. \quad IRB(G) = \sum_{UV \in E} \left(d_u^{\frac{1}{2}} - d_v^{\frac{1}{2}} \right)^2$$

$$\begin{aligned} IRB(NS_1[p], x, y) &= 2^{p+1} (\sqrt{2} - \sqrt{1})^2 + 4(2^p - 1) (\sqrt{3} - \sqrt{1})^2 + \{12 \times 2^p - 11\} (\sqrt{2} - \sqrt{2})^2 + 14(2^p - 1) (\sqrt{3} - \sqrt{2})^2 \\ &= 2^{p+1} (\sqrt{2} - \sqrt{1})^2 + 4(2^p - 1) (\sqrt{3} - \sqrt{1})^2 + 14(2^p - 1) (\sqrt{3} - \sqrt{2})^2. \end{aligned}$$

$$12. \quad IRR_t(G) = \frac{1}{2} \sum_{UV \in E} |d_u - d_v|$$

$$\begin{aligned} IRR_t(NS_1[p], x, y) &= \frac{1}{2} \left[2^{p+1} |2 - 1| + 4(2^p - 1) |3 - 1| + \{12 \times 2^p - 11\} |2 - 2| + 14(2^p - 1) |3 - 2| \right] \\ &= \frac{1}{2} [2^{p+1} + 4(2^p - 1) |2| + 14(2^p - 1)]. \end{aligned}$$

The following Table 2 shows the values of these irregularity indices for some test values of parameter p .

Table 2. Irregularity indices for $NS_1[p]$ polypropylenimine octaamin dendrimers.

Irregularity Indices	$p = 1$	$p = 2$	$p = 3$	$p = 4$	$p = 5$
$IRDIF(G) = \sum_{UV \in E} \left \frac{d_u}{d_v} - \frac{d_v}{d_u} \right $	28.340	79.014	180.335	383.005	788.344
$AL(G) = \sum_{UV \in E} d_u - d_v $	26	74	170	362	746
$IRLU(G) = \sum_{UV \in E} \frac{ d_u - d_v }{\min(d_u, d_v)}$	19	53	121	257	529
$IRLU(G) = \sum_{UV \in E} \frac{ d_u - d_v }{\sqrt{(d_u d_v)}}$	13.1627	36.6596	83.6536	177.6415	365.6174
$IRF(G) = \sum_{UV \in E} (d_u - d_v)^2$	34	98	226	482	994
$IRLA(G) = 2 \sum_{UV \in E} \frac{ d_u - d_v }{(d_u + d_v)}$	12.268	34.136	77.872	165.344	340.288
$IRD1 = \sum_{UV \in E} \ln \{1 + d_v - d_u \}$	16.870	47.839	109.776	233.650	481.398
$IRA(G) = \sum_{UV \in E} \left(d_u^{\frac{-1}{2}} - d_v^{\frac{-1}{2}} \right)^2$	1.2934	3.5370	8.0243	16.9988	34.9479
$IRGA(G) = \sum_{UV \in E} \ln \frac{d_u + d_v}{2 \sqrt{(d_u d_v)}}$	1.1025	3.0661	6.9934	14.8480	30.5571
$IRB(G) = \sum_{UV \in E} \left(d_u^{\frac{1}{2}} - d_v^{\frac{1}{2}} \right)^2$	4.2437	12.0450	27.6476	58.8528	121.2632
$IRR_t(G) = \frac{1}{2} \sum_{UV \in E} d_u - d_v $	13	37	85	181	373

Now we proceed to irregularity indices of $NS_2[p]$.

Theorem 2. Let $NS_2[p]$ be the nanostar polypropylenimine octaamin dendrimers, then the irregularity indices of $NS_2[p]$ are:

1. $IRDIF(NS_2[p]) = 1.5(2^{p+1}) + 5(2^p) - 5$
2. $IRR(NS_2[p]) = 2^{p+1} + 6(2^p) - 6$
3. $IRL(NS_2[p]) = 0.6931471(2^{p+1}) + 2.4328(2^p) - 2.4328$

4. $IRLU(NS_2[p]) = 2^{p+1} + 3(2^p) - 3$
5. $IRLF(NS_2[p]) = 0.707106(2^{p+1}) + 2.4494897(2^p) - 2.4494897$
6. $\sigma(NS_2[p]) = 2^{p+1} + 6(2^p) - 6$
7. $IRLA(NS_2[p]) = 0.6667(2^{p+1}) + 2.4(2^p) - 2.4$
8. $IRD1 = 0.6931471806(2^{p+1}) + 4.15888302(2^p) - 4.15888302$
9. $IRA(NS_2[p]) = 0.08578644(2^{p+1}) + 0.1010205144(2^p) - 0.1010205144$
10. $IRGA(NS_2[p]) = 0.0588915178(2^{p+1}) + 0.1224659836(2^p) - 0.1224659836$
11. $IRB(NS_2[p]) = 0.171578(2^{p+1}) + 0.606123086(2^p) - 0.606123086$
12. $IRR_t(NS_2[p]) = 4(2^p) - 3$

Proof. In order to prove the above theorem, we have to consider Figure 2. We can see that the edges of $NS_2[p]$ admit the following partition in Table 3. \square

Table 3. Edge partition of nanostar polypropylenimine octaamin dendrimers.

Edge Type (d_u, d_v)	Number of Edges
(1,2)	2^{p+1}
(2,2)	$\{8 \times 2^p - 5\}$
(2,3)	$6(2^p - 1)$

We can see that the edges of $NS_2[p]$ admit the following partition in Table 3.

Now using above Table 3 and the above definitions, we have:

1. $IRDIF(G) = \sum_{UV \in E} \left| \frac{d_u}{d_v} - \frac{d_v}{d_u} \right|$

$$IRDIF(NS_2[p], x, y) = 2^{p+1} \left| \frac{\frac{2}{1}}{\frac{1}{2}} - \frac{\frac{1}{2}}{\frac{2}{1}} \right| + \{8 \times 2^p - 5\} \left| \frac{\frac{2}{1}}{\frac{2}{2}} - \frac{\frac{2}{2}}{\frac{1}{1}} \right| + 6(2^p - 1) \left| \frac{\frac{3}{2}}{\frac{2}{3}} - \frac{\frac{2}{3}}{\frac{3}{2}} \right|$$

$$= 2^{p+1} \left| \frac{2}{1} - \frac{1}{2} \right| + 6(2^p - 1) \left| \frac{3}{2} - \frac{2}{3} \right|.$$
2. $IRR(G) = \sum_{UV \in E} imb(e) \therefore imb(e) = |d_u - d_v|$

$$IRR(NS_2[p], x, y) = 2^{p+1} |2 - 1| + \{8 \times 2^p - 5\} |2 - 2| + 6(2^p - 1) |3 - 2|$$

$$= 2^{p+1} + 6(2^p - 1).$$
3. $IRL(G) = \sum_{UV \in E} |lnd_u - lnd_v|$

$$IRL(NS_2[p], x, y) = 2^{p+1} |ln2 - ln1| + \{8 \times 2^p - 5\} |ln2 - ln2| + 6(2^p - 1) |ln3 - ln2|$$

$$= 2^{p+1} |ln2| + 6(2^p - 1) |ln\frac{3}{2}|.$$
4. $IRLU(G) = \sum_{UV \in E} \frac{|d_u - d_v|}{\min(d_u d_v)}$

$$IRLU(NS_2[p], x, y) = 2^{p+1} \frac{|2-1|}{\sqrt{2}} + \{8 \times 2^p - 5\} \frac{|2-2|}{\sqrt{4}} + 6(2^p - 1) \frac{|3-2|}{\sqrt{6}}$$

$$= 2^{p+1} + 3(2^p - 1).$$
5. $IRLF(G) = \sum_{UV \in E} \frac{|d_u - d_v|}{\sqrt{(d_u d_v)}}$

$$IRLU(NS_2[p], x, y) = 2^{p+1} \frac{|2-1|}{\sqrt{2}} + \{8 \times 2^p - 5\} \frac{|2-2|}{\sqrt{4}} + 6(2^p - 1) \frac{|3-2|}{\sqrt{6}}$$

$$= \frac{2^{p+1}}{\sqrt{2}} + \frac{6(2^p - 1)}{\sqrt{6}}.$$

$$6. \quad \sigma(G) = \sum_{UV \in E} (d_u - d_v)^2$$

$$\begin{aligned} \sigma(NS_2[p], x, y) &= 2^{p+1}(2-1)^2 + \{8 \times 2^p - 5\}(2-2)^2 + 6(2^p-1)(3-2)^2 \\ &= 2^{p+1}(2-1)^2 + 6(2^p-1)(3-2)^2. \end{aligned}$$

$$7. \quad IRLA(G) = 2 \sum_{UV \in E} \frac{|d_u - d_v|}{(d_u + d_v)}$$

$$\begin{aligned} IRLA(NS_2[p], x, y) &= 2 \left[2^{p+1} \frac{|2-1|}{(2+1)} + \{8 \times 2^p - 5\} \frac{|2-2|}{(2+2)} + 6(2^p-1) \frac{|3-2|}{(3+2)} \right] \\ &= 2 \left[\frac{2^{p+1}}{3} + \frac{6(2^p-1)}{5} \right]. \end{aligned}$$

$$8. \quad IRD1 = \sum_{UV \in E} \ln\{1 + |d_v - d_u|\}$$

$$\begin{aligned} IRD1(NS_2[p], x, y) &= 2^{p+1} \ln\{1 + |2-1|\} + \{8 \times 2^p - 5\} \ln\{1 + |2-2|\} + 6(2^p-1) \ln\{1 + |3-2|\} \\ &= 2^{p+1} \ln 2 + 6(2^p-1) \ln 2. \end{aligned}$$

$$9. \quad IRA(G) = \sum_{UV \in E} \left(d_u^{\frac{-1}{2}} - d_v^{\frac{-1}{2}} \right)^2$$

$$\begin{aligned} IRA(NS_2[p], x, y) &= 2^{p+1} \left(\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{1}} \right)^2 + \{8 \times 2^p - 5\} \left(\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}} \right)^2 + 6(2^p-1) \left(\frac{1}{\sqrt{3}} - \frac{1}{\sqrt{2}} \right)^2 \\ &= 2^{p+1} \left(\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{1}} \right)^2 + 6(2^p-1) \left(\frac{1}{\sqrt{3}} - \frac{1}{\sqrt{2}} \right)^2. \end{aligned}$$

$$10. \quad IRGA(G) = \sum_{UV \in E} \ln \frac{d_u + d_v}{2 \sqrt{(d_u d_v)}}$$

$$\begin{aligned} IRGA(NS_2[p], x, y) &= 2^{p+1} \ln \frac{|2+1|}{2\sqrt{2}} + \{8 \times 2^p - 5\} \ln \frac{|2+2|}{2\sqrt{4}} + 6(2^p-1) \ln \frac{|3+2|}{2\sqrt{6}} \\ &= 2^{p+1} \ln \frac{3}{2\sqrt{2}} + 6(2^p-1) \ln \frac{5}{2\sqrt{6}}. \end{aligned}$$

$$11. \quad IRB(G) = \sum_{UV \in E} \left(d_u^{\frac{1}{2}} - d_v^{\frac{1}{2}} \right)^2$$

$$\begin{aligned} IRB(NS_2[p], x, y) &= 2^{p+1} (\sqrt{2} - \sqrt{1})^2 + \{8 \times 2^p - 5\} (\sqrt{2} - \sqrt{2})^2 + 6(2^p-1) (\sqrt{3} - \sqrt{2})^2 \\ &= 2^{p+1} (\sqrt{2} - \sqrt{1})^2 + 14(2^p-1) (\sqrt{3} - \sqrt{2})^2. \end{aligned}$$

$$12. \quad IRR_t(G) = \frac{1}{2} \sum_{UV \in E} |d_u - d_v|$$

$$\begin{aligned} IRR_t(NS_2[p], x, y) &= \frac{1}{2} \left[2^{p+1} |2-1| + \{8 \times 2^p - 5\} |2-2| + 6(2^p-1) |3-2| \right] \\ &= \frac{1}{2} [2^{p+1} + 6(2^p-1)]. \end{aligned}$$

Table 4 represents some calculated values of irregularity indices of $NS_2[p]$ for some test values of p .

Table 4. Irregularity indices for $NS_2[p]$ polypropylenimine octaamin dendrimers.

Irregularity indices	$p = 1$	$p = 2$	$p = 3$	$p = 4$	$p = 5$
$IRDIF(G) = \sum_{UV \in E} \left \frac{d_u}{d_v} - \frac{d_v}{d_u} \right $	11	27	59	123	251
$AL(G) = \sum_{UV \in E} d_u - d_v $	10	26	58	122	250
$IRL(G) = \sum_{UV \in E} ln d_u - ln d_v $	5.2054	12.844	28.1199	58.6727	119.7782
$IRLU(G) = \sum_{UV \in E} \frac{ d_u - d_v }{\min(d_u, d_v)}$	7	17	37	77	157
$IRLU(G) = \sum_{UV \in E} \frac{ d_u - d_v }{\sqrt{(d_u d_v)}}$	5.2779	13.0053	28.4601	59.3697	121.1889
$IRF(G) = \sum_{UV \in E} (d_u - d_v)^2$	10	26	58	122	250
$IRLA(G) = 2 \sum_{UV \in E} \frac{ d_u - d_v }{(d_u + d_v)}$	5.0668	12.5336	27.4672	57.3344	117.0688
$IRD1 = \sum_{UV \in E} \ln \{1 + d_u - d_v \}$	6.9315	18.0218	40.2025	84.5639	173.2867
$IRA(G) = \sum_{UV \in E} \left(\frac{d_u^{-\frac{1}{2}}}{d_v^{\frac{1}{2}}} - \frac{d_v^{-\frac{1}{2}}}{d_u^{\frac{1}{2}}} \right)^2$	0.4442	0.9894	2.0797	4.2605	8.62197
$IRGA(G) = \sum_{UV \in E} \ln \frac{d_u + d_v}{2 \sqrt{(d_u d_v)}}$	0.3580	0.8385	1.7995	3.7215	7.5655
$IRB(G) = \sum_{UV \in E} \left(d_u^{\frac{1}{2}} - d_v^{\frac{1}{2}} \right)^2$	1.2924	3.1909	6.9881	14.5823	29.7708
$IRR_t(G) = \frac{1}{2} \sum_{UV \in E} d_u - d_v $	5	13	29	61	125

Our next aim is to determine the of irregularity indices of polymer dendrimers.

Theorem 3. Let $NS_3[p]$ be polymer dendrimer then the irregularity indices of $NS_3[p]$ are:

1. $IRDIF(NS_3[p]) = 4.5(2^p) + 54.999978(2^{p-1}) - 14.999994$
2. $IRR(NS_3[p]) = 3(2^p) + 66(2^{p-1}) - 18$
3. $IRL(NS_3[p]) = 2.079441(2^p) + 26.76069(2^{p-1}) - 7.29837$
4. $IRLU(NS_3[p]) = 3(2^p) + 33(2^{p-1}) - 9$
5. $IRLF(NS_3[p]) = 2.12132(2^p) + 26.944384(2^{p-1}) - 7.348468$
6. $\sigma(NS_3[p]) = 3(2^p) + 66(2^{p-1}) - 18$
7. $IRLA(NS_3[p]) = 2(2^p) + 26.4(2^{p-1}) - 7.2$
8. $IRD1 = 2.079441(2^p) + 45.747702(2^{p-1}) - 12.476646$
9. $IRA(NS_3[p]) = 1.5(2^p) + 1.111242(2^{p-1}) - 0.303066$
10. $IRGA(NS_3[p]) = 0.176673(2^p) + 1.347126(2^{p-1}) - 0.367398$
11. $IRB(NS_3[p]) = 0.514719(2^p) + 6.667386(2^{p-1}) - 1.8378$
12. $IRR_t(NS_3[p]) = 1.5(2^p) + 33(2^{p-1}) - 9$

Proof. In order to prove the above theorem we have to consider Figure 3. \square

We can see that the edges of $NS_3[p]$ admit the following partition in Table 5.

Table 5. Edge partition of polymer dendrimer.

Edges Type (d_u, d_v)	Number of Edges
(1,2)	3.2^p
(2,2)	$54(2^{p-1}) - 24$
(2,3)	$66(2^{p-1} - 1) + 48$
(3,3)	3.2^{p+1}

Now using above Table 5 and the above definitions we have:

$$1. \quad IRDIF(G) = \sum_{UV \in E} \left| \frac{d_u}{d_v} - \frac{d_v}{d_u} \right|$$

$$\begin{aligned} IRDIF(NS_3[p], x, y) &= 3.2^p \left| \frac{2}{1} - \frac{1}{2} \right| + 54(2^{p-1}) - 24 \left| \frac{2}{2} - \frac{2}{2} \right| + 66(2^{p-1} - 1) + 48 \left| \frac{3}{2} - \frac{2}{3} \right| + 3.2^{p+1} \left| \frac{3}{3} - \frac{3}{3} \right| \\ &= 3.2^p \left| \frac{2}{1} - \frac{1}{2} \right| + 66(2^{p-1} - 1) + 48 \left| \frac{3}{2} - \frac{2}{3} \right|. \end{aligned}$$

$$2. \quad IRR(G) = \sum_{UV \in E} imb(e) \therefore imb(e) = |d_u - d_v|$$

$$\begin{aligned} IRR(NS_3[p], x, y) &= 3.2^p |2 - 1| + 54(2^{p-1}) - 24 |2 - 2| + 66(2^{p-1} - 1) + 48 |3 - 2| + 3.2^{p+1} |3 - 3| \\ &= 3.2^p + 66(2^{p-1} - 1) + 48. \end{aligned}$$

$$3. \quad IRL(G) = \sum_{UV \in E} |lnd_u - lnd_v|$$

$$\begin{aligned} IRL(NS_3[p], x, y) &= 3.2^p |ln2 - ln1| + 54(2^{p-1}) - 24 |ln2 - ln2| + 66(2^{p-1} - 1) + 48 |ln3 - ln2| + 3.2^{p+1} |ln3 - ln3| \\ &= 3.2^p |ln2 - ln1| + 66(2^{p-1} - 1) + 48 |ln3 - ln2|. \end{aligned}$$

$$4. \quad IRLU(G) = \sum_{UV \in E} \frac{|d_u - d_v|}{\min(d_u, d_v)}$$

$$\begin{aligned} IRLU(NS_3[p], x, y) &= 3.2^p \frac{|2-1|}{1} + 54(2^{p-1}) - 24 \frac{|2-2|}{2} + 66(2^{p-1} - 1) + 48 \frac{|3-2|}{2} + 3.2^{p+1} \frac{|3-3|}{3} \\ &= 3.2^p + \frac{66(2^{p-1}-1)+48}{2}. \end{aligned}$$

$$5. \quad IRLF(G) = \sum_{UV \in E} \frac{|d_u - d_v|}{\sqrt{(d_u d_v)}}$$

$$\begin{aligned} IRLF(NS_3[p], x, y) &= 3.2^p \frac{|2-1|}{\sqrt{2}} + 54(2^{p-1}) - 24 \frac{|2-2|}{\sqrt{4}} + 66(2^{p-1} - 1) + 48 \frac{|3-2|}{\sqrt{6}} + 3.2^{p+1} \frac{|3-3|}{\sqrt{9}} \\ &= \frac{3.2^p}{\sqrt{2}} + \frac{66(2^{p-1}-1)+48}{\sqrt{6}}. \end{aligned}$$

$$6. \quad \sigma(G) = \sum_{UV \in E} (d_u - d_v)^2$$

$$\begin{aligned} \sigma(NS_3[p], x, y) &= 3.2^p (2-1)^2 + 54(2^{p-1}) - 24(2-2)^2 + 66(2^{p-1}-1) + 48(3-2)^2 + 3.2^{p+1}(3-3)^2 \\ &= 3.2^p + 66(2^{p-1}-1) + 48. \end{aligned}$$

$$7. \quad IRLA(G) = 2 \sum_{UV \in E} \frac{|d_u - d_v|}{(d_u + d_v)}$$

$$\begin{aligned} IRLA(NS_3[p], x, y) &= 2 \left[3.2^p \frac{|2-1|}{(2+1)} + 54(2^{p-1}) - 24 \frac{|2-2|}{(2+2)} + 66(2^{p-1}-1) + 48 \frac{|3-2|}{(2+2)} + 3.2^{p+1} \frac{|3-3|}{(3+3)} \right] \\ &= 2 \left[\frac{3.2^p}{(3)} + 66(2^{p-1}-1) + \frac{48}{(5)} \right]. \end{aligned}$$

$$8. \quad IRD1 = \sum_{UV \in E} \ln \{1 + |d_v - d_u|\}$$

$$\begin{aligned} IRD1(NS_3[n], x, y) &= 3.2^p \ln \{1 + |2 - 1|\} + 54(2^{p-1}) - 24 \ln \{1 + |2 - 2|\} + 66(2^{p-1} - 1) + 48 \ln \{1 + |3 - 2|\} + 3.2^{p+1} \ln \{1 + |3 - 3|\} \\ &= 3.2^p \ln 2 + (66(2^{p-1} - 1) + 48) \ln 2. \end{aligned}$$

$$9. \quad IRA(G) = \sum_{UV \in E} \left(d_u^{\frac{-1}{2}} - d_v^{\frac{-1}{2}} \right)^2$$

$$\begin{aligned} IRA(NS_3[p], x, y) &= 3.2^p \left(\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{1}} \right)^2 + 54(2^{p-1}) - 24 \left(\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}} \right)^2 + 66(2^{p-1} - 1) + 48 \left(\frac{1}{\sqrt{3}} - \frac{1}{\sqrt{2}} \right)^2 + 3.2^{p+1} \left(\frac{1}{\sqrt{3}} - \frac{1}{\sqrt{3}} \right)^2 \\ &= 3.2^p \left(\frac{1}{\sqrt{2}} \right)^2 + 66(2^{p-1} - 1) + 48 \left(\frac{1}{\sqrt{3}} - \frac{1}{\sqrt{2}} \right)^2. \end{aligned}$$

$$10. \quad IRGA(G) = \sum_{UV \in E} \ln \frac{d_u + d_v}{2 \sqrt{(d_u d_v)}}$$

$$\begin{aligned} IRGA(NS_3[p], x, y) &= 3.2^p \ln \frac{|2+1|}{2\sqrt{2}} + 54(2^{p-1}) - 24 \ln \frac{|2+2|}{2\sqrt{3}} + 66(2^{p-1} - 1) + 48 \ln \frac{|3+2|}{2\sqrt{4}} + 3.2^{p+1} \ln \frac{|3+3|}{2\sqrt{6}} \\ &= 3.2^p \ln \frac{|3|}{2\sqrt{2}} + (66(2^{p-1} - 1) + 48) \ln \frac{|5|}{2\sqrt{6}}. \end{aligned}$$

$$11. \quad IRB(G) = \sum_{UV \in E} \left(d_u^{\frac{1}{2}} - d_v^{\frac{1}{2}} \right)^2$$

$$\begin{aligned} IRB(NS_3[p], x, y) &= 3.2^p (\sqrt{2} - \sqrt{1})^2 + 54(2^{p-1}) - 24(\sqrt{2} - \sqrt{2})^2 + 66(2^{p-1} - 1) + 48(\sqrt{3} - \sqrt{2})^2 + 3.2^{p+1} (\sqrt{3} - \sqrt{3})^2 \\ &= 3.2^p (\sqrt{2} - \sqrt{1})^2 + (66(2^{p-1} - 1) + 48)(\sqrt{3} - \sqrt{2})^2. \end{aligned}$$

$$12. \quad IRR_t(G) = \frac{1}{2} \sum_{UV \in E} |d_u - d_v|$$

$$\begin{aligned} IRR_t(NS_3[p], x, y) &= \frac{1}{2} [3.2^p |2-1| + 54(2^{p-1}) - 24|2-2| + 66(2^{p-1} - 1) + 48|3-2| + 3.2^{p+1}|3-3|] \\ &= \frac{1}{2} [3.2^p + (66(2^{p-1} - 1) + 48)]. \end{aligned}$$

The following Table 6 represents some calculated values of irregularity indices of $NS_3[p]$ for some test values of p .

Table 6. Irregularity indices for $NS_3[p]$ polymer dendrimers.

Irregularity Indices	$p = 1$	$p = 2$	$p = 3$	$p = 4$	$p = 5$
$IRDIF(G) = \sum_{UV \in E} \left \frac{d_u}{d_v} - \frac{d_v}{d_u} \right $	48.9999	112.9999	240.9999	496.9998	1008.99
$AL(G) = \sum_{UV \in E} d_u - d_v $	54	126	270	558	1134
$IRL(G) = \sum_{UV \in E} lnd_u - lnd_v $	23.6212	54.541	116.379	240.058	487.414
$IRLU(G) = \sum_{UV \in E} \frac{ d_u - d_v }{\min(d_u, d_v)}$	30	69	147	303	615
$IRLU(G) = \sum_{UV \in E} \frac{ d_u - d_v }{\sqrt{(d_u d_v)}}$	23.8385	55.025	117.399	242.147	491.644
$IRF(G) = \sum_{UV \in E} (d_u - d_v)^2$	54	126	270	558	1134
$IRLA(G) = 2 \sum_{UV \in E} \frac{ d_u - d_v }{(d_u + d_v)}$	23.2	53.6	114.4	236	479.2
$IRD1 = \sum_{UV \in E} \ln \{1 + d_v - d_u \}$	37.4299	87.3365	187.1496	386.7760	786.0286
$IRA(G) = \sum_{UV \in E} \left(d_u^{\frac{-1}{2}} - d_v^{\frac{-1}{2}} \right)^2$	3.8082	7.9194	16.1419	32.5868	65.4768
$IRGA(G) = \sum_{UV \in E} \ln \frac{d_u + d_v}{2 \sqrt{(d_u d_v)}}$	1.3331	3.0335	6.4345	13.236	26.84015
$IRB(G) = \sum_{UV \in E} \left(d_u^{\frac{1}{2}} - d_v^{\frac{1}{2}} \right)^2$	5.8590	13.5558	28.9495	59.7368	121.3113
$IRR_t(G) = \frac{1}{2} \sum_{UV \in E} d_u - d_v $	27	63	135	279	567

4. Graphical Analysis, Discussions and Conclusions

In this part we give our comparative analysis of some of the irregularity indices of the above discussed dendrimers and the dependences of the irregularity indices on the parameter of the structures. Figures 4–7 contain three graphs of irregularity indices. The horizontal axis is used for step size p and the vertical axis shows the value of irregularity index. In the graphs, the red color shows the irregularity of $NS_1[p]$, the blue color shows the irregularity of $NS_2[p]$ and the green color shows the irregularity of the $NS_3[p]$. In each graph, three different colored curves are depicted which shows the behavior of the irregularity indices with increase in p .




Figure 4. Graph of $IRDIF(G)$.




Figure 5. Graph of $IRLU(G)$.




Figure 6. Graph of $IRA(G)$.




Figure 7. Graph of $IRB(G)$.

From above graph it seems obvious that irregularities have a slight increase with an increase in the step size p for the range $p \leq 12$. But after $p \geq 14$, these irregularity indices drastically increase with increase in p . So $NS_3[p]$ is the most irregular and asymmetric structure as far as most of the irregularity indices are concerned. Nanostar dendrimers are relatively less irregular, and $NS_1[p]$ are the most regular dendrimers. This trend is not restricted to only irregularity index $IRDIF$. Most of the irregularity indices behave pretty similarly as shown in the following figures. All other figures show the trends which can easily be understood in the Figures 5–7.

All above graphs (Figures 4–7) indicate that $NS_3[p]$ is highly asymmetric as far as all irregularity indices are concerned, whereas $NS_1[p]$ is less asymmetric, and $NS_2[p]$ is the most regular structure with respect to all indices. In IRB, $NS_1[p]$ and $NS_3[p]$ show the same irregularity behavior. These facts typically relate geometry and topology of the structure of these dendrimers and can be used in modelling purposes.

We foresee that our results could play an important role in determining properties of these dendrimers such as enthalpy, toxicity, resistance and entropy. Similar research has been done in [29], where authors discussed some properties of alkane isomers.

Author Contributions: Z.H. and M.M. gave the idea; H.A., T.H. and S.R. wrote the article; S.M.K. and Y.C.K. edited and verified the results.

Funding: This work was supported by the Dong-A University research fund.

Acknowledgments: We are greatly thankful to reviewers for their comments which have helped bring this article to its current form.

Conflicts of Interest: Authors declare no conflict of interests.

References

1. Hodge, P. Polymer science branches out. *Nature* **1993**, *362*, 18–19. [[CrossRef](#)]
2. Fischer, M.; Vögtle, F. Dendrimers: From design to applications—A progress report. *Angew. Chem. Int. Ed.* **1999**, *38*, 884–905. [[CrossRef](#)]
3. Wiener, E.C.; Auteri, F.P.; Chen, J.W.; Brechbiel, M.W.; Gansow, O.A.; Schneider, D.S.; Belford, R.L.; Clarkson, R.B.; Lauterbur, P.C. Molecular dynamics of ion-chelate complexes attached to dendrimers. *J. Am. Chem. Soc.* **1996**, *118*, 7774–7782. [[CrossRef](#)]
4. Rucker, G.; Rucker, C. On topological indices, boiling points, and cycloalkanes. *J. Chem. Inf. Comput. Sci.* **1999**, *39*, 788–802. [[CrossRef](#)]
5. Klavžar, S.; Gutman, I. A Comparison of the Schultz molecular topological index with the Wiener index. *J. Chem. Inf. Comput. Sci.* **1996**, *36*, 1001–1003. [[CrossRef](#)]
6. Gutman, I.; Polansky, O.E. *Mathematical Concepts in Organic Chemistry*; Springer: New York, NY, USA, 1986.
7. Randic, M. On the characterization of molecular branching. *J. Am. Chem. Soc.* **1975**, *97*, 6609–6615. [[CrossRef](#)]
8. Wiener, H. Structural determination of paraffin boiling points. *J. Am. Chem. Soc.* **1947**, *69*, 17–20. [[CrossRef](#)] [[PubMed](#)]
9. Estrada, E. Atomic bond connectivity and the energetic of branched alkanes. *Chem. Phys. Lett.* **2008**, *463*, 422–425. [[CrossRef](#)]
10. Kier, L.; Hall, L. *Molecular Connectivity in Chemistry and Drug Research*; Academic Press: New York, NY, USA, 1976.
11. Kier, L.B.; Hall, L.H. *Molecular Connectivity in Structure Activity Analysis*; Wiley: New York, NY, USA, 1986.
12. Deng, H.; Yang, J.; Xia, F. A general modeling of some vertex-degree based topological indices in benzenoid systems and phenylenes. *Comput. Math. Appl.* **2011**, *61*, 3017–3023. [[CrossRef](#)]
13. Zhang, H.; Zhang, F. The Clar covering polynomial of hexagonal systems. *Discret. Appl. Math.* **1996**, *69*, 147–167. [[CrossRef](#)]
14. De, N.; Nayeem, S.M.A. Computing the F-index of nanostar dendrimers. *Pac. Sci. A Nat. Sci. Eng.* **2016**, *18*, 14–21. [[CrossRef](#)]
15. Siddiqui, M.K.; Imran, M.; Ahmad, A. On Zagreb indices, Zagreb polynomials of some nanostar dendrimers. *Appl. Math. Comput.* **2016**, *280*, 132–139. [[CrossRef](#)]
16. Madanshekaf, A.; Ghaneei, M. Computing two topological indices of nanostars dendrimer, Optoelectron. *Adv. Mater. Rapid Commun.* **2010**, *4*, 1849–1851.
17. Madanshekaf, A.; Moradi, M. The first geometric-arithmetic index of some nanostar dendrimers. *Iran. J. Math. Chem.* **2014**, *5*, 1–6.
18. Munir, M.; Nazeer, W.; Rafique, S.; Kang, S.M. M-polynomial and related topological indices of Nanostar dendrimers. *Symmetry* **2016**, *8*, 97. [[CrossRef](#)]
19. Munir, M.; Nazeer, W.; Rafique, S.; Nizami, A.R.; Kang, S.M. M-polynomial and degree-based topological indices of Titania nanotubes. *Symmetry* **2016**, *8*, 117. [[CrossRef](#)]

20. Munir, M.; Nazeer, W.; Rafique, S.; Kang, S.M. M-Polynomial and Degree-Based Topological Indices of Polyhex Nanotubes. *Symmetry* **2016**, *8*, 149. [[CrossRef](#)]
21. Munir, M.; Nazeer, W.; Shahzadi, S.; Kang, S.M. Some invariants of circulant graphs. *Symmetry* **2016**, *8*, 134. [[CrossRef](#)]
22. Chartrand, G.; Erdos, P.; Oellermann, O. How to define an irregular graph. *Coll. Math. J.* **1988**, *19*, 36–42. [[CrossRef](#)]
23. Majcher, Z.; Michael, J. Highly irregular graphs with extreme numbers of edges. *Discret. Math.* **1997**, *164*, 237–242. [[CrossRef](#)]
24. Behzad, M.; Chartrand, G. No graph is perfect. *Am. Math. Mon.* **1967**, *74*, 962–963. [[CrossRef](#)]
25. Horoldagva, B.; Buyantogtokh, L.; Dorjsembe, S.; Gutman, I. Maximum size of maximally irregular graphs. *MATCH Commun. Math. Comput. Chem.* **2016**, *76*, 81–98.
26. Liu, F.; Zhang, Z.; Meng, J. The size of maximally irregular graphs and maximally irregular triangle-free graphs. *Graphs Comb.* **2014**, *30*, 699–705. [[CrossRef](#)]
27. Collatz, L.; Sinogowitz, U. Spektren endlicher Graphen. In *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg*; Springer: Berlin, Germany, 1957; Volume 21, pp. 63–77.
28. Bell, F.K. A note on the irregularity of graphs. *Linear Algebra Appl.* **1992**, *161*, 45–54. [[CrossRef](#)]
29. Albertson, M. The irregularity of a graph. *Ars Comb.* **1997**, *46*, 219–225.
30. Vukičević, D.; Graovac, A. Valence connectivities versus Randić, Zagreb and modified Zagreb index: A linear algorithm to check discriminative properties of indices in acyclic molecular graphs. *Croat. Chem. Acta* **2004**, *77*, 501–508.
31. Abdo, H.; Brandt, S.; Dimitrov, D. The total irregularity of a graph. *Discret. Math. Theor. Comput. Sci.* **2014**, *16*, 201–206.
32. Abdo, H.; Dimitrov, D. The total irregularity of graphs under graph operations. *Miskolc Math. Notes* **2014**, *15*, 3–17. [[CrossRef](#)]
33. Abdo, H.; Dimitrov, D. The irregularity of graphs under graph operations. *Discuss. Math. Graph Theory* **2014**, *34*, 263–278. [[CrossRef](#)]
34. Gutman, I. Topological Indices and Irregularity Measures. *J. Bull.* **2018**, *8*, 469–475.
35. Reti, T.; Sharfdini, R.; Dregelyi-Kiss, A.; Hagobin, H. Graph irregularity indices used as molecular descriptors in QSPR studies. *MATCH Commun. Math. Comput. Chem.* **2018**, *79*, 509–524.
36. Hu, Y.; Li, X.; Shi, Y.; Xu, T.; Gutman, I. On molecular graphs with smallest and greatest zeroth-Corder general randic index. *MATCH Commun. Math. Comput. Chem.* **2005**, *54*, 425–434.
37. Caporossi, G.; Gutman, I.; Hansen, P.; Pavlovic, L. Graphs with maximum connectivity index. *Comput. Biol. Chem.* **2003**, *27*, 85–90. [[CrossRef](#)]
38. Li, X.; Gutman, I. Mathematical aspects of Randic, type molecular structure descriptors. In *Mathematical Chemistry Monographs*; University of Kragujevac and Faculty of Science: Kragujevac, Serbia, 2006.
39. Das, K.; Gutman, I. Some properties of the second Zagreb Index. *MATCH Commun. Math. Comput. Chem.* **2004**, *52*, 103–112.
40. Trinajstić, N.N.; Ikolic, S.; Milicevic, A.; Gutman, I. On Zagreb indices. *Kemija Industriji* **2010**, *59*, 577–589.
41. Milicevic, A.; Nikolic, S.; Trinajstić, N. On reformulated Zagreb indices. *Mol. Divers.* **2004**, *8*, 393–399. [[CrossRef](#)] [[PubMed](#)]
42. Gupta, C.K.; Lokesha, V.; Shwetha, S.B.; Ranjini, P.S. On the symmetric division DEG index of graph. *Southeast Asian Bull. Math.* **2016**, *40*, 59–80.
43. Balaban, A.T. Highly discriminating distance based numerical descriptor. *Chem. Phys. Lett.* **1982**, *89*, 399–404. [[CrossRef](#)]
44. Furtula, B.; Graovac, A.; Vukičević, D. Augmented Zagreb index. *J. Math. Chem.* **2010**, *48*, 370–380. [[CrossRef](#)]
45. Das, K.C. Atom-bond connectivity index of graphs. *Discret. Appl. Math.* **2010**, *158*, 1181–1188. [[CrossRef](#)]
46. Estrada, E.; Torres, L.; Rodríguez, L.; Gutman, I. An atom-bond connectivity index: Modeling the enthalpy of formation of alkanes. *Indian J. Chem.* **1998**, *37A*, 849–855.
47. Iqbal, Z.; Aslam, A.; Ishaq, M.; Aamir, M. Characteristic study of irregularity measures of some Nanotubes. *Can. J. Phys.* **2019**. [[CrossRef](#)]
48. Gao, W.; Aamir, M.; Iqbal, Z.; Ishaq, M.; Aslam, A. On Irregularity Measures of Some Dendrimers Structures. *Mathematics* **2019**, *7*, 271. [[CrossRef](#)]
49. Gao, W.; Abdo, H.; Dimitrov, D. On the irregularity of some molecular structures. *Can. J. Chem.* **2017**, *95*, 174–183.

50. Hussain, Z.; Rafique, S.; Munir, M.; Athar, M.; Chaudhary, M.; Ahmad, H.; Min Kang, S. Irregularity Molecular Descriptors of Hourglass, Jagged-Rectangle, and Triangular Benzenoid Systems. *Processes* **2019**, *7*, 413. [[CrossRef](#)]



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).