

Supplementary Materials

Hydrothermal Functionalization of Oil Palm Empty Fruit Bunch Biochar for Wastewater Treatment Purposes

Izzudin Ibrahim ¹, Toshiki Tsubota ², Mohd Ali Hassan ³ and Yoshito Andou ^{1,4,*}

¹ Department of Biological Functions Engineering, Graduate School of Life Science and Systems Engineering, Kyushu Institute of Technology, 2-4 Hibikino, Wakamatsu, Kitakyushu, Fukuoka 808-0196, Japan; ibrahim.izzudin516@mail.kyutech.jp

² Department of Applied Chemistry, Faculty of Engineering, Kyushu Institute of Technology, 1-1 Sensuicho, Tobata, Kitakyushu, Fukuoka 804-8550, Japan; tsubota@che.kyutech.ac.jp

³ Department of Bioprocess Technology, Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia; alihass@upm.edu.my

⁴ Collaborative Research Centre for Green Materials on Environmental Technology, Kyushu Institute of Technology, 2-4 Hibikino, Wakamatsu, Kitakyushu, Fukuoka 808-0196, Japan

* Correspondence: yando@life.kyutech.ac.jp







Figure S1. Langmuir isotherm model linear fitting curve for MB adsorption on EFB-FBC ($C = 125, 150, 175$ and 250 mg/L, contact time = 24 h, temperature = 30 °C, dosage = 2 g/L)



(a)



(b)



(c)

Figure S2. Methylene blue kinetic adsorption data for EFB-FBC fitted for (a) pseudo-first order (b) pseudo-second order and (c) intraparticle diffusion kinetic models.