

Supplementary materials

Development and Validation of UPLC-MS/MS Method for Quantitative Analysis of 5-Fluorouracil in Aqueous Humor of Rabbits

Mohd Abul Kalam ^{1,†}, Sulaiman S. Alhudaithi ^{1,†}, Adel Ali Alhowyan ¹, Muzaffar Iqbal ²,
Mohammad Raish ¹, Abdullah K. Alshememry ¹, Musaed Alkholief ¹, Aliyah A. Almomen ²
and Aws Alshamsan ^{1,*}

- ¹ Department of Pharmaceutics, College of Pharmacy, King Saud University, Riyadh 11451, Saudi Arabia; makalam@ksu.edu.sa (M.A.K.); salhudaithi@ksu.edu.sa (S.S.A.); adel-ali@ksu.edu.sa (A.A.A.); mraish@ksu.edu.sa (M.R.); aalshememry@ksu.edu.sa (A.K.A.); malkholief@ksu.edu.sa (M.A.)
- ² Department of Pharmaceutical Chemistry, College of Pharmacy, King Saud University, Riyadh 11451, Saudi Arabia; muziqbal@ksu.edu.sa (M.I.); alalmomen@ksu.edu.sa (A.A.A.)
- * Correspondence: aalshamsan@ksu.edu.sa
- † These authors contributed equally to this work.




Figure S1a. Representative LLOQ chromatogram of 5-FU and IS in AqH in sample prepared by protein precipitation method.




Figure S1b. Representative LLOQ chromatogram of 5-FU and IS in AqH in sample prepared by liquid-liquid extraction method.




Figure S2. The comparative extraction recovery (%) and matrix effects (%) data of different extracting solvents obtained during sample preparation optimization.