

Supplementary

Isolation and Identification of Non-Conjugated Linoleic Acid from Processed *Panax ginseng* Using LC-MS/MS and ¹H-NMR

Tae-Kyung Kim ¹, Changsuk Lee ¹, Taek-Hee Nam ¹, Yong-Ki Seo ², Kyeong-Soo Kim ², Kimoon Kang ², Seok-Hun Yun ¹, Seok-Hwan Yoon ¹ and Jaeho Jeong ^{1,*}

- ¹ Bio Research Institutes, CJ CheilJedang, Suwon 16495, Korea; tk.kim5@cj.net (T.-K.K.); changsuk.lee@cj.net (C.L.); th.nam@cj.net (T.-H.N.); sh.yun@cj.net (S.-H.Y.); seokhwan.yoon@cj.net (S.-H.Y.)
- ² Food Research Institutes, CJ CheilJedang, Suwon 16495, Korea; yk.seo@cj.net (Y.-K.S.); ks.kim30@cj.net (K.-S.K.); kimoon.kang@cj.net (K.K.)
- * Correspondence: jaeho.jeong@cj.net; biodory@cj.net

Citation: Kim, T.-K.; Lee, C.; Nam, T.-H.; Seo, Y.-K.; Kim, K.-S.; Kang, K.; Yun, S.-H.; Yoon, S.-H.; Jeong, J. Isolation and Identification of Non-Conjugated Linoleic Acid from Processed *Panax ginseng* Using LC-MS/MS and ¹H-NMR. *Separations* **2021**, *8*, 208. <https://doi.org/10.3390/separations8110208>

Academic Editor: Gavino Sanna

Received: 15 October 2021

Accepted: 29 October 2021

Published: 4 November 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).




Figure S1. ^1H NMR spectra overlay of the commercial linoleic acid (A) and a representative fractionated sample (peak at 61 minutes) of black ginseng (B). ^1H NMR spectrum of linoleic acid (Cas.No. 60-33-3, MilliporeSigma, St. Louis, MO, USA) was exactly same with fractionated sample. The magnified spectrum of olefinic hydrogen is indicated by the dotted rectangle.