

Article

A Simple Microextraction Method for Toxic Industrial Dyes Using a Fatty-Acid Solvent Mixture

Danielle P. Arcon and Francisco C. Franco, Jr. *

Chemistry Department, De La Salle University, 2401 Taft Avenue, Manila 0922, Philippines;
danielle_arcon@dlsu.edu.ph

* Correspondence: francisco.franco@dlsu.edu.ph

Table S1. Summary of the ¹H NMR chemical shifts.

	δ (ppm)				
	H _E	H _D	H _C	H _B	H _A
C12	11.84	2.13	1.46	1.21	0.82
C6	1.51	2.09	1.44	1.19	0.77
C12:C6	11.88	2.10	1.46	1.21	0.79
C12:C6 with water	9.44	2.18	1.49	1.18	0.79

Citation: Arcon, D.; Franco Jr., F. A Simple Microextraction Method for Toxic Industrial Dyes Using a Fatty Acid Solvent Mixture. *Separations* **2021**, *8*, 135. <https://doi.org/10.3390/separations8090135>

Academic Editor: Marcello Locatelli

Received: 2 August 2021

Accepted: 19 August 2021

Published: 26 August 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).