

Review

Effect of Nature of Substituents on Coordination Properties of Mono- and Disubstituted Derivatives of Boron Cluster Anions $[B_nH_n]^{2-}$ ($n = 10, 12$) and Carboranes with exo-Polyhedral B–X Bonds ($X = N, O, S, \text{Hal}$)

Evgenii Yu. Matveev ¹, Varvara V. Avdeeva ^{2,*}, Konstantin Yu. Zhizhin ², Elena A. Malinina ² and Nikolay T. Kuznetsov ²

¹ Institute of Fine Chemical Technologies Named after M. V. Lomonosov, MIREA—Russian Technological University, Vernadskogo pr. 86, 119571 Moscow, Russia

² Kurnakov Institute of General and Inorganic Chemistry, Russian Academy of Sciences, Leninskii pr. 31, 119991 Moscow, Russia

* Correspondence: avdeeva.varvara@mail.ru

Citation: Matveev, E.Y.; Avdeeva, V.V.; Zhizhin, K.Y.; Malinina, E.A.; Kuznetsov, N.T. Effect of Nature of Substituents on Coordination Properties of Mono- and Disubstituted Derivatives of Boron Cluster Anions $[B_nH_n]^{2-}$ ($n = 10, 12$) and Carboranes with exo-Polyhedral B–X Bonds ($X = N, O, S, \text{Hal}$). *Inorganics* **2022**, *10*, 238. <https://doi.org/10.3390/inorganics10120238>

Academic Editor: Marina Yu. Stogniy

Received: 7 November 2022

Accepted: 1 December 2022

Published: 2 December 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: This review systematizes data on the coordination ability of mono- and disubstituted derivatives of boron cluster anions and carboranes in complexation with transition metals. Boron clusters anions $[B_nH_n]^{2-}$, monocarborane anions $[CB_nH_{n-1}]^-$, and dicarboranes $[C_2B_nH_{n-2}]$ (with non-functionalized carbon atoms) ($n = 10, 12$) containing the B–X exo-polyhedral bonds ($X = N, O, S, \text{Hal}$) are discussed. Synthesis and structural features of complexes known to date are described. The effect of complexing metal and substituent attached to the boron cage on the composition and structures of the final complexes is analyzed. It has been established that substituted derivatives of boron cluster anions and carboranes can act as both ligands and counterions. A complexing agent can coordinate substituted derivatives of the boron cluster anions due to three-center two-electron 3c2e MHB bonds, by the substituent functional groups, or a mixed type of coordination can be realized, through the BH groups of the boron cage and the substituent. As for B-substituted carboranes, complexes with coordinated substituents or salts with non-coordinated carborane derivatives have been isolated; compounds with MHB bonding are not characteristic of carboranes.

Keywords: decahydro-*closo*-decaborate anion; dodecahydro-*closo*-dodecaborate anion; complexation

1. Introduction

Boron has a particularly rich chemistry; various structures can be found for elemental boron [1–3], boric acids and borates [4–6], metal and non-metal borides [7–9], boranes and carboranes [10–18]. In the number of hydrides that it forms, boron is second only to carbon. Boron cluster anions $[B_nH_n]^{2-}$ ($n = 10, 12$) and carboranes (Figure 1) are fascinating objects with versatile chemistry. Owing to their tendency to participate in reactions of substitution of exo-polyhedral hydrogen atoms, they allow one to design new cage systems differing in geometry and electronic structure.

The application fields of boron clusters are traditionally explained by their high energy intensity [19]. Moreover, it was proposed to use them in boron neutron capture therapy of tumor tissues because of the high neutron absorption capacity of boron atoms [20–22]. Boron clusters can be used to manufacture heat-resistant polymers and materials with neutron-protective properties [23–26], and contrast agents for MRI diagnostics [27]. Metal complexes with boron clusters are applied as heavy metal extractants, as coordination polymers, etc. [28–32]. Recently, complexes containing boron clusters have been proposed

to prepare metal borides [33–36]. The physiological properties of boron clusters and their applications in medicine are discussed in recent reviews [37–41].




Figure 1. Structures of *closo*-decaborates anions, 1-carba-*closo*-borates anions and 1,2-dicarba-*closo*-boranes.

To date, the coordination ability of boron clusters $[B_nH_n]^{2-}$ ($n = 10, 12$), perchlorinated clusters $[B_{10}Cl_{10}]^{2-}$, and dimeric clusters $[B_{20}H_{18}]^{2-}$ has been studied in sufficient detail and analyzed in recent reviews [42–51]. It is concluded that boron cluster anions are soft bases according to Pearson; therefore, their participation in complexation reactions as ligands is observed in complexation with metals that act as Pearson’s soft acids (Cu(I), Ag(I), and Pb(II)). In this case, boron cluster anions allow synthesizing of a great number of mono-nuclear, binuclear, polymeric complexes with boron polyhedra coordinated by the metal atom via vertices (BH group), edges (HBBH group) or faces (BBB) of the boron cage. When interacting with metals which act as hard acids (Fe(III), Co(III)), boron cluster anions act as reducing agents and reduce the oxidation state of metals to M(II) or M(0). Classical 3d metals in the 2+ oxidation state are acids of the intermediate group (Zn(II), Ni(II), Cu(II), etc.) and usually afford salts consisting of cationic metal complexes $[ML_x]^{2+}$ with neutral ligands L (organic, inorganic or solvent molecules) and boron cluster anions as counterions. In this case, numerous non-covalent interactions can be found in the structures of compounds, including hydrogen and dihydrogen bonds between BH groups of the boron clusters and organic cations, ligand molecules or solvents.

Boron cluster anions tend to participate in reactions of substitution of exo-polyhedral hydrogen atoms to form substituted derivatives with various functional groups [52–56]. The resulting substituted boron clusters can also participate in complexation as ligands. However, the currently known complexes are scarcely studied, prepared in different systems, in the presence of various metals and ligands, which makes it difficult to analyze and compare the reactivity of substituted derivatives of boron cluster anions as ligands.

Carboranes differ from the boron clusters in their charge: monocarboranes $[CB_9H_{10}]^-$ and $[CB_{11}H_{12}]^-$ are monoanions, whereas dicarboranes $[C_2B_8H_{10}]$ and $[C_2B_{10}H_{12}]$ are neutral compounds. The boron atoms in carboranes have a lower negative charge as compared to boron clusters; therefore, their ability to act as a donor of electronic density reduces. In

this case, there are the following ways to form complexes with carboranes: (i) the formation of a B–Hg bond during electrophilic mercurization, which proceeds to the position furthest from the carbon atoms (actually, formation of a C–Hg bond is also possible) (see review [57] and references thereof); (ii) B–H activation with the formation of a B–M bond in the presence of a substituent with a donor atom in the *ortho* position acting as a ligand (the substituent is usually attached to the carbon atom); therefore, activation of neighboring positions is observed (see reviews [58,59] and recent articles [60,61]); (iii) coordination by a substituent, as a rule, attached to the carbon atom of carborane cage (see, for example, review [62]).

The chemistry of carboranes is realized as a rule via carbon atoms of the carborane cage which provides wide opportunities to vary functional groups and metal atoms bonded to them. Here, we wanted to compare the coordination ability of the boron cage in substituted boron clusters and B-substituted carboranes without C-functionalization.

In this work, we have systematized the data on complexes of mono- and disubstituted derivatives of boron cluster anions $[B_nH_n]^{2-}$, monocarborane anions $[CB_nH_{n-1}]^-$, and dicarboranes $[C_2B_nH_{n-2}]$ (with non-functionalized carbon atoms) ($n = 10, 12$) containing exo-polyhedral B–X bonds ($X = N, O, S, \text{Hal}$) in order to compare the coordination ability of substituted boron cages and determine directions for further systematic research.

2. Metal Complexes with Substituted Derivatives of Boron Cluster Anions

It is known that unsubstituted boron clusters make it possible to prepare complexes of soft acids with coordinated boron hydride ligands [42,43] (first of all, copper(I), silver(I), lead(II)). Heteroleptic silver(I) complexes with inner-sphere boron cluster anions are the most studied group of complexes. This is probably due to the correspondence between the softness of silver(I) acid and softness of the boron cluster bases, as compared to other metals.

The introduction of a substituent containing electron-donor groups makes it possible to obtain complexes in which the coordination of the boron cage is realized via the functional groups of the substituent; thus, those metals that are too hard to coordinate boron clusters are able to coordinate substituted derivatives through a substituent being introduced. Moreover, the substituent can decrease the total charge of the boron cluster, thus decreasing its coordination ability in complexation.

2.1. Halogen Atoms

In the literature, there are a great number of complexes containing perhalogenated *closo*-borate anions $[B_n\text{Hal}_n]^{2-}$ ($n = 10, 12$; $\text{Hal} = \text{F}, \text{Cl}, \text{Br}, \text{I}$). These anions are interesting because they are weak coordinating ligands; they form various complexes with silver atoms as complexing agents. Complexes with these anions are discussed in detail in a number of reviews [45–47] and are beyond this work. Here, we would discuss only mono- and disubstituted derivatives containing halogen atoms.

When the *closo*-dodecaborate anion was allowed to react with hydrogen halides (HCl, HBr or HI) in dichloroethane, **monochloro- and dichlorosubstituted *closo*-dodecaborate derivatives** were isolated [63]. The source of the chlorine atoms is the solvent used, whereas halohydrogens act as electrophilic initiators. The structure of Bipy-containing tris-chelate **nickel(II)** complex $[\text{Ni}(\text{Bipy})_3][\text{B}_{12}\text{H}_{10.668}\text{Cl}_{1.332}]$ was determined by single-crystal X-ray diffraction. It was found that the compound consists of tris-chelate cationic complex $[\text{Ni}(\text{Bipy})_3]^{2+}$, chlorine-substituted anions $[\text{B}_{12}\text{H}_{11}\text{Cl}]^{2-}$ or $[\text{B}_{12}\text{H}_{10}\text{Cl}_2]^{2-}$, and crystallization solvent molecules. In the crystal, mono- and disubstituted derivatives present a ratio of 2:1. Thus, it is clear that Cl has no effect on the coordination ability of the boron cage.

Silver(I) complex $[\text{Ag}(\text{CH}_3\text{CN})_3]_2[\text{Ag}_2[2\text{-B}_{10}\text{H}_9\text{F}]_2]$ [64] with the **monofluoro-substituted derivative of the *closo*-decaborate anion** was isolated by the reaction between tetraphenylphosphonium 2-fluoro-*closo*-decaborate $[2\text{-B}_{10}\text{H}_9\text{F}]^{2-}$ and silver trifluoroacetate. The structure of this complex (Figure 2) is built of anions $[\text{Ag}_2[2\text{-B}_{10}\text{H}_9\text{F}]_2]^{2-}$ ($\text{Ag} \cdots \text{H} 2.02(5) \text{--} 2.36(5) \text{ \AA}$) linked by cations $[\text{Ag}(\text{CH}_3\text{CN})_3]^+$ ($\text{Ag} \cdots \text{H} 2.36 \text{ \AA}$) forming double chains.




Figure 2. Structure of polymeric silver(I) complex $[\text{Ag}(\text{CH}_3\text{CN})_3]_2[\text{Ag}_2[2\text{-B}_{10}\text{H}_9\text{F}]_2]_n$ (F is disordered into two positions).

Silver(I) complex $[\text{Ag}_2(\text{Ph}_3\text{P})_4[\text{B}_{12}\text{H}_{11}\text{Cl}]]$ [65] with triphenylphosphine Ph_3P was also obtained for the **monochloro-substituted derivative of the *closo*-dodecaborate anion**. The binuclear complex was isolated when the $[\text{B}_{10}\text{H}_{11}\text{Cl}]^{2-}$ anions reacted with $[\text{Ag}(\text{Ph}_3\text{P})_3\text{NO}_3]$ in CH_3CN or DMF. The compound is a centrosymmetric complex in which the silver atoms coordinate the *closo*-dodecaborate anion along opposite edges (Ag–B 2.779 and 2.790 Å) that are as far as possible from the chlorine atom introduced (Figure 3).




Figure 3. Structure of binuclear silver(I) complex $[\text{Ag}_2(\text{Ph}_3\text{P})_4[\text{B}_{12}\text{H}_{11}\text{Cl}]]$.

Two positional isomers of the monoanions $[\text{Ag}[\text{B}_{10}\text{H}_{10}](\text{PPh}_3)_2]^-$ (with unsubstituted *closo*-decaborate anion) and $[\text{Ag}[\text{B}_{10}\text{H}_9\text{Cl}](\text{PPh}_3)_2]^-$ (**with monochlorosubstituted *closo*-decaborate anion**) were found co-crystallized in complexes $[\text{Ag}(\text{PPh}_3)_4][\text{Ag}[\text{B}_{10}\text{H}_{9.14}\text{Cl}_{0.86}](\text{PPh}_3)_2]$ and $[\text{Ag}(\text{PPh}_3)_4][(\text{PPh}_3)_2\text{Ag}[\text{B}_{10}\text{H}_{9.5}\text{Cl}_{0.5}]]$ [66] with equatorial and apical coordination of the boron cage, which were obtained from DMF and acetonitrile, respectively (Figure 4). It should be noted that in both compounds, monochlorosubstituted derivatives are coordinated by edges opposite to the chlorine atom introduced.

The data present show that one (or two) halogen atoms have no effect on the coordination ability of the boron cage. For both the $[\text{B}_{10}\text{H}_{10}]^{2-}$ and $[\text{B}_{12}\text{H}_{12}]^{2-}$ anions, nickel(II) resulted in complex salts, whereas silver(I) afforded complexes with boron clusters coordinated by edges located as far as possible from the substituent introduced.




Figure 4. Structures of isomers of mononuclear complex $[\text{Ag}[2\text{-B}_{10}\text{H}_9\text{Cl}](\text{PPh}_3)_2]^-$ with (a) apical and (b) equatorial coordination of the boron cluster.

2.2. Hydroxy Substituent

When the *closo*-decaborate anion reacts with sulfolane in the presence of *p*-toluenesulfonic acid followed by alkaline hydrolysis of the resulting product, salts of the **2-hydroxy-*closo*-decaborate anion** $[2\text{-B}_{10}\text{H}_9\text{OH}]^{2-}$ are isolated [56]. The formation of tris-chelate **nickel(II)** complex $[\text{NiL}_3][\text{B}_{10}\text{H}_9\text{OH}]$ ($\text{L} = 2,2'$ -bipyridyl (Bipy), 1,10-phenanthroline (Phen), 2,2'-bipyridylamine (BPA), 1,2-diaminobenzene (DAB)) [67] with the $[2\text{-B}_{10}\text{H}_9\text{OH}]^{2-}$ anion as a counterion were observed in the nickel(II) complexation reactions in the presence of organic ligands L . Structures of solvates $[\text{Ni}(\text{Phen})_3][\text{B}_{10}\text{H}_9\text{OH}] \cdot 0.75\text{CH}_3\text{CN} \cdot 0.5\text{H}_2\text{O}$ (Figure 5a) and $[\text{Ni}(\text{Phen})_3][\text{B}_{10}\text{H}_9\text{OH}] \cdot 2\text{CH}_3\text{CN} \cdot 0.67\text{DMF}$ were isolated and studied by X-ray diffraction.




Figure 5. Structures of (a) $[\text{Ni}(\text{Phen})_3][\text{B}_{10}\text{H}_9\text{OH}] \cdot 0.75\text{CH}_3\text{CN} \cdot 0.5\text{H}_2\text{O}$ and (b) $[\text{Cu}_2(\text{bipy})_4(\mu\text{-CO}_3)][2\text{-B}_{10}\text{H}_9\text{OH}] \cdot 2\text{DMSO} \cdot \text{H}_2\text{O}$.

The synthesis and structure of **copper(II)** complex with the $[2\text{-B}_{10}\text{H}_9\text{OH}]^{2-}$ anion was reported [68]. It was found that the $[2\text{-B}_{10}\text{H}_9\text{OH}]^{2-}$ anion can be prepared in situ, starting from an unsubstituted $[\text{B}_{10}\text{H}_{10}]^{2-}$ boron cluster. Long-term heating of copper(II) complex $[\text{Cu}_2(\text{bipy})_4(\mu\text{-CO}_3)][\text{B}_{10}\text{H}_{10}]$ in DMSO led to partial substitution of one hydrogen atom by the OH group to form copper(II) complex $[\text{Cu}_2(\text{bipy})_4(\mu\text{-CO}_3)][2\text{-B}_{10}\text{H}_{9.83}\text{OH}_{0.17}] \cdot 2\text{DMSO} \cdot \text{H}_2\text{O}$ (Figure 5b). In this compound, the cationic part is the same as in the starting copper(II) complex, while the anionic part contains the unsubstituted *closo*-decaborate anion and its monohydroxy-substituted derivative cocrystallized in one crystal in the ratio 0.17:0.83.

Titanium complex with the monohydroxy-substituted *closo*-dodecaborate anion $[B_{12}H_{11}OH]^{2-}$ and the cyclopentadienyl ligand was prepared and characterized [69] (Figure 6a). The complex was isolated when $[TiCpCl_2]$ was allowed to react with $(Ph_3MeP)_2[B_{12}H_{11}OH]$. As it was found, titanium coordinates the oxygen atom of the 2-hydroxy-*closo*-decaborate anion; the distances Ti–O and B–O are 1.711(9) Å and 1.45(2) Å, respectively.




Figure 6. Structure of (a) titanium complex $[Ph_3(CH_3)P]_2[CpTiCl_2[B_{12}H_{11}OH]]$ and (b) lead(II) complex $[Pb(Bipy)(DMF)[2-B_{10}H_9OH]] \cdot DMF$.

In **lead(II) complex** with the monohydroxy-substituted *closo*-decaborate derivative $[Pb(Bipy)(DMF)[2-B_{10}H_9OH]] \cdot DMF$ [70] (Figure 6b), Pb(II) coordinates one Bipy molecule (Pb–N 2.520(6), 2.583(7) Å), the hydroxyl substituent (Pb–O 2.285(6) Å), and one DMF molecule (Pb–O 2.504(7) Å). The coordination environment of lead(II) is completed by two BH groups (Pb–H 2.944, 3.286 Å); the corresponding distances allow one to conclude that the boron polyhedron participates in coordination.

For the OH substituent, it can be concluded that copper(II) and nickel(II) are able to coordinate a boron cluster neither by the BH group nor by the substituent. At the same time, titanium(II) coordinates the functional group in the corresponding complexes, whereas lead(II) exhibits combination coordination via the OH substituent and BH groups of the boron cage.

2.3. Ammonium Substituents

As a rule, the presence of substituents reduces the coordination ability of the boron ligand. This is especially true for substituents that decrease the total charge of the boron cluster. Thus, the reaction of salts of the *closo*-decaborate anion with hydroxylamine sulfonic acid leads to the formation of the **ammonium-substituted derivative** $[2-B_{10}H_9NH_3]^-$ [70]. Silver(I) complexation with triphenylphosphine and the substituted anion afforded solvate $[Ag(PPh_3)_4][B_{10}H_9NH_3] \cdot 2DMF$ with the $[B_{10}H_9NH_3]^-$ derivative acting as a counterion [71]. According to the single-crystal X-ray diffraction data, the compound consists of silver(I) cationic complexes $[Ag(PPh_3)_4]^+$ and the ammonium-substituted anions as counterions.

Lead(II) complexes with the $[2-B_{12}H_{11}NEt_3]^-$ anion and ligands Bipy, BPA were described [72]. Complexes $\{PbL_2[2-B_{12}H_{11}NEt_3]_2\}$ were obtained in lead(II) complexation in the presence of ligands L. The complexes were characterized by IR spectroscopy as well as elemental analysis. The authors concluded that lead(II) coordinates the boron cage via the 3c2e PBHB bonds.

When $[NH_4Et_3]_2[B_{12}H_{12}]$ was allowed to react with $[RuCl_2(PPh_3)_3]$, **ruthenium(II) complex** $[(PPh_3)_2ClRu[B_{12}H_{11}(NEt_3)]]$ with the **singly charged $[B_{12}H_{11}NEt_3]^-$** was isolated [73] as

solvate $[(PPh_3)_2ClRu[B_{12}H_{11}(NEt_3)]] \cdot CH_2Cl_2$. The boron cage anion demonstrates facial coordination (Ru–B 2.268–2.485 Å) (Figure 7), whereas the substituent remains non-coordinated.




Figure 7. Structure of ruthenium(II) complex $[(PPh_3)_2ClRu[B_{12}H_{11}(NEt_3)]] \cdot CH_2Cl_2$ (solvent molecules are omitted).

We can conclude that this ruthenium(II) complex is the first example of complexes where the metal atom is able to coordinate singly charged anion $[2-B_{12}H_{11}NEt_3]^-$ with lower coordination ability as compared to unsubstituted boron clusters. Lead(II) can be expected to coordinate the boron cage (the conclusions are based on IR spectral data), whereas silver(I) affords complexes with an inner-sphere position of the mono-charged substituted derivatives.

2.4. Amino Group

Salt $Na_2[B_{12}H_{11}NH_2]$ salt can be obtained by the reaction of $(Bu_3NH)_2[B_{12}H_{11}NH_3]$ with sodium hydride in THF [74]. When the **amino-closo-dodecaborate ion** reacted with **nickel(II)** complex $[Ni(THF)_2Br_2]$, compound $[Na_6(THF)_{15}][Ni[B_{12}H_{11}NH_2]]_4 \cdot THF$ was isolated. Nickel coordinates four N atoms from the substituents and is in a square planar environment (Ni–N 1.924(4), 1.931(4) Å).

When $Na_2[B_{12}H_{11}NH_2]$ was allowed to react with $[Au(Ph_3P)Cl]$, neutral **gold(I)** complex bis((triphenylphosphine)-(1-amino-closo-dodecaborate(11))-gold) $[Au(PPh_3)[NH_2-B_{12}H_{11}]]$ [74] was isolated. The Au–N bonds are 2.076(8) and 2.067(9) Å.

An effective method for preparation of complexes starting from the singly charged **ammonio-substituted derivative** $[B_{12}H_{11}NH_3]^+$ is the deprotonation of the monoanion with a strong base to form the **amino-substituted dianion** $[B_{12}H_{11}NH_2]^{2-}$, which can participate in complexation reactions. Thus, **ruthenium(II)** complexes $Bu_3MeN[Ru(PPh_3)_2Cl[B_{12}H_{11}NH_2]] \cdot CH_2Cl_2$ and $Bu_4N[Ru(dppb)Cl[B_{12}H_{11}NH_2]] \cdot CH_2Cl_2$ (dppb = bis(diphenylphosphine)butane) (Figure 8b) [75] were isolated when the $[B_{12}H_{11}NH_2]^{2-}$ reacted with $[Ru(Ph_3P)_3Cl_2]$ or $[Ru(dppb)(Ph_3P)Cl_2]$, respectively. In both complexes, the complexing agent coordinates the boron cage by the N atom of the substituent and two BH groups (Ru–B 2.323(3), 2.591(3) Å in the first one and 2.350(2), 2.490(3) Å in the second; Ru–N 2.186(2) and 2.2177(19) Å, respectively).

In addition, it was found [75] that sodium salt $Na[Ru(PPh_3)_2Cl[B_{12}H_{11}NH_2]]$ reacts with carbon monoxide affording ruthenium(II) complex $[Ru(PPh_3)_2CO[B_{12}H_{11}NH_2]]$. In this compound, the metal atom coordinates the substituted derivative of the boron cluster by the nitrogen atom of the substituent and two BH groups (Ru–N, 2.211(4) Å, Ru–B 2.418(5), 2.442(5) Å).

Rhodium(I) complex $MePPh_3[Rh(PPh_3)_2[B_{12}H_{11}NH_2]]$ with the **amino-substituted derivative** $[B_{12}H_{11}NH_2]^{2-}$ is known [74], which was isolated when $(MePPh_3)Na[B_{12}H_{11}NH_2]$ was allowed to react with $[Rh(PPh_3)_3Cl]$. The boron cage is coordinated by the rhodium atom through the N atom of the substituent and the BH group (Rh–N 2.146(6) Å, Rh–B 2.592(8) Å, Rh–H 1.919 Å).

Thus, ruthenium(II) and rhodium(I) are metals that could form a combined coordination mode of NH_2 -substituted derivatives: via 3c2e MHB bonds with the boron cage and functional groups of the substituent.




Figure 8. Structures of (a) $[\text{Au}(\text{PPh}_3)[\text{NH}_2\text{-B}_{12}\text{H}_{11}]]$ and (b) $\text{Bu}_4\text{N}[\text{Ru}(\text{dppb})\text{Cl}[\text{B}_{12}\text{H}_{11}\text{NH}_2]] \cdot \text{CH}_2\text{Cl}_2$ (solvent molecules and cation are omitted).

2.5. Sulfonium Group

The reaction of the *closo*-decaborate anion with DMSO in the presence of HCl leads to the formation of the singly charged **dimethylsulfonium derivative** $[\text{1-B}_{10}\text{H}_9\text{SMe}_2]^-$ [76]. **Lead(II)** complexes $[\text{PbL}_2[\text{1-B}_{10}\text{H}_9\text{SMe}_2]_2]$ ($\text{L} = \text{Bipy}, \text{BPA}$) [72] were obtained when aqueous solutions of salts with $[\text{1-B}_{10}\text{H}_9\text{SMe}_2]^-$ were allowed to react with $\text{Pb}(\text{NO}_3)_2$ and organic ligand L . It was determined by X-ray diffraction that in complex $[\text{Pb}(\text{Bipy})_2[\text{1-B}_{10}\text{H}_9\text{SMe}_2]_2]$ (Figure 9) lead(II) coordinates two Bipy and two boron cages by the equatorial and apical faces, both anions being coordinated by edges connecting two equatorial belts of the B_{10} polyhedron (Pb-B 3.24–3.55 Å).




Figure 9. Structure of lead(II) complex $[\text{Pb}(\text{Bipy})_2[\text{1-B}_{10}\text{H}_9\text{SMe}_2]_2]$. Hydrogen atoms are omitted.

The interaction of the 2-sulfanyl derivative of the *closo*-decaborate anion $(\text{Bu}_4\text{N})_2[\text{2-B}_{10}\text{H}_9\text{SH}]$ with phthalimide in the presence of cesium carbonate leads to the **sulfonium derivative of the *closo*-decaborate anion** $[\text{2-B}_{10}\text{H}_9\text{S}(\text{CH}_2\text{N}(\text{CO})_2\text{C}_6\text{H}_4)_2]^-$. The introduction of this compound into the **silver(I)** complexation makes it possible to obtain a complex with 2-[bis(N-phthalimidomethyl)sulfonio]-*closo*-decaborate as a counterion $[\text{Ag}(\text{PPh}_3)_4][\text{2-B}_{10}\text{H}_9\text{S}(\text{CH}_2\text{N}(\text{CO})_2\text{C}_6\text{H}_4)_2]$ [77]. According to the single-crystal X-ray diffraction data, the compound contains cationic complexes $[\text{Ag}(\text{PPh}_3)_4]^+$ and the derivative of the boron cluster as counterions.

Lead(II) and silver(I) complexes with $[2-B_{10}H_9S(CH_2C(O)NH_2)_2]^-$ were briefly discussed [78]. The compounds were identified by IR spectroscopy and elemental analysis. It was concluded that silver(I) compound $[Ag_2(bipy)_2[2-B_{10}H_9S(CH_2C(O)NH_2)_2]NO_3]$ contains boron clusters coordinated by complexing agents via the AgHB bonds, whereas in $[Pb(bipy)_2[2-B_{10}H_9S(CH_2C(O)NH_2)_2]]$ there are no 3c2e PbHB bonds in the IR spectrum, indicating that the complexing agent coordinates the boron cage via the substituent.

Thus, lead(II) is the second metal (after ruthenium(II)) that is able to coordinate singly-charged substituted derivatives of the boron clusters. Silver(I) seems to be harder than lead(II) and ruthenium(II) because it gives complexes with an inner-sphere position of the monocharged derivatives.

2.6. Sulfanyl Group

A number of **lead(II) and silver(I)** complexes with **sulfanyl-substituted *closo*-decaborate derivative** $[2-B_{10}H_9SH]^{2-}$ were reported [78]. Based on the data of IR spectroscopy and elemental analysis, it was concluded that $[Ag_2(bipy)_2[2-B_{10}H_9SH]]$ contains boron clusters coordinated by complexing agents via the AgHB bonds; the $[2-B_{10}H_9SH]^{2-}$ anion is involved in coordination by lead(II) in complexes $[Pb[2-B_{10}H_9SH]]$ and $[Pb(bipy)_2[2-B_{10}H_9SH]]$ via 3c2e PbHB bonds and the substituent.

The thiol derivative $[B_{12}H_{11}SH]^{2-}$ is formed by the interaction of the $[B_{12}H_{12}]^{2-}$ anion with thiourea under electrochemical oxidation conditions followed by hydrolysis of the resulting compound [79]. The SH-substituted derivative $[B_{12}H_{11}SH]^{2-}$ was used in **ruthenium(III)** complexation; complex $[Ru(SB_{12}H_{11})(NH_3)_5] \cdot 2H_2O$ (Figure 10) was isolated when the salt of the substituted boron cluster was allowed to react with $[RuCl(NH_3)_5]Cl_2$. The metal atom coordinates five NH_3 groups and the substituent via the S atom [80] (S–B 1.878 Å, Ru–S 2.240 Å).




Figure 10. Structure of ruthenium(III) complex $[Ru(SB_{12}H_{11})(NH_3)_5] \cdot 2H_2O$ (water molecules are omitted).

2.7. Oxonium Substituents

The introduction of cyclic ether molecules as an exo-polyhedral substituent into the boron cluster results in reduction of the total charge of the derivatives of the *closo*-decaborate anion. Thus, the reaction of salts of the undecahydrodecaborate anion $[B_{10}H_{11}]^-$ with 1,4-dioxane and tetrahydropyran gives derivatives of the *closo*-decaborate anion containing the organic molecules as substituents in position B(2) of the polyhedron [65,81]. These derivatives react with $[Ag(PPh_3)_3]NO_3$ to form the corresponding silver(I) complexes $[Ag(PPh_3)_4][2-B_{10}H_9O(CH_2)_5] \cdot CH_2Cl_2$ and $[Ag(PPh_3)_4][2-B_{10}H_9O(CH_2)_4O]$. In both complexes, silver coordinates four molecules of Ph_3P forming cationic complex $[Ag(PPh_3)_4]^+$ [65], while the substituted derivative plays the role of a counterion. Both compounds are additional evidence that the silver(I) atom tends to form salts with monocharged derivatives of the boron clusters instead of forming complexes with them.

2.8. Opening of the Cyclic Substituent

Activation of the coordination ability of the oxonium derivatives of the boron clusters can be realized by opening of the cyclic ether substituent. In this case, the resulting derivative acquires the double negative charge and all the O atoms present in the structure of the resulting derivative potentially can be coordinated by the metal atom.

Particularly, reactions of the 1,4-dioxane derivative of the $[B_{10}H_{10}]^{2-}$ anion with ethylenediamine in ethanol lead to the formation of the derivative of the *closo*-decaborate anion **2- $B_{10}H_9O(CH_2)_2O(CH_2)_2NH(CH_2)_2NH_2$] $^{2-}$ with pendant ethylenediamine (en) group** separated from the boron cluster by an alkoxy spacer. This anion was used in nickel(II) complexation [82]. In the compounds obtained, the metal atom coordinates only the substituent introduced into the boron cluster, whereas all BH groups remain uncoordinated. Complexes $[Ni(en)[2-B_{10}H_9O(CH_2)_2O(CH_2)_2NH(CH_2)_2NH_2]] \cdot H_2O$ and $[Ni(H_2O)(en)_3[2-B_{10}H_9O(CH_2)_2O(CH_2)_2NH(CH_2)_2NH_2]]$ were isolated. The complexes were obtained by heterophasic (using $NiCO_3-Ni(OH)_2$) and homophasic reactions, respectively. In the first complex, the water molecule is located in the outer sphere, whereas nickel(II) coordinates two O atoms and two N atoms of the substituent chain. In the second, nickel(II) coordinates two N atoms and one O atom of the substituent, two N atoms of ethylenediamine, and one O atom of coordinated water. The complexes are hydrated isomers resulting from the changed denticity of the substituted boron ligand (Figure 11).




Figure 11. Schematic representations of the structures of nickel(II) complexes $[Ni(H_2O)(en)_3[2-B_{10}H_9O(CH_2)_2O(CH_2)_2NH(CH_2)_2NH_2]]$ (left) and $[Ni(en)[2-B_{10}H_9O(CH_2)_2O(CH_2)_2NH(CH_2)_2NH_2]] \cdot H_2O$ (right).

The opening of the cyclic substituent in the $[2-B_{10}H_9O(CH_2)_2O]^-$ anion using the ethylate ion leads to the formation of the **2- $B_{10}H_9O(CH_2CH_2)_2OEt$] $^{2-}$ derivative with the pendant ethoxy group** separated from the boron cluster by an alkoxy spacer. Lead(II) complex $[Pb(Bipy)[2-B_{10}H_9O(CH_2CH_2)_2OEt]]_2 \cdot 0.5DMF$ was obtained in lead complexation in the presence of Bipy [83] (Figure 12). In the crystal, Pb(II) coordinates two N atoms of Bipy (Pb–N 2.470(4), 2.483(5) Å) and three O atoms of the alkoxy spacer of the substituent (Pb–O 2.497(4)–2.859(4) Å). In addition, Pb(II) coordinates the apical BH group of one boron cage and the apical edge of the other (Pb–B 3.128(6) and 3.288(7) Å, Pb–H 2.70(5) and 2.73(6) Å).

It was found that a monosubstituted derivative of the *closo*-decaborate with 1,4-dioxane $[B_{10}H_9O(CH_2)_4O]^-$ reacts with polyhydric alcohols (ethylene glycol, glycerol, triethanolamine), giving **derivatives with pendant hydroxy groups** **$[B_{10}H_9OCH_2CH_2OCH_2CH_2OR]^{2-}$** (R = CH_2OH , $CH(OH)CH_2OH$, $CH_2N(CH_2CH_2OH)_2$) [84]. On the basis of these compounds, **gadolinium(III) complexes** $Gd_2[B_{10}H_9OCH_2CH_2OCH_2CH_2OR]_3$ were obtained when reacting with gadolinium(III) carbonate. The final compounds were characterized by NMR and IR spectroscopies as well as mass-spectrometry [84]. According to the data obtained, it seems that the *closo*-decaborate derivatives are coordinated by the substituents.




Figure 12. Structure of lead(II) complex $[\text{Pb}(\text{Bipy})][2\text{-B}_{10}\text{H}_9\text{O}(\text{CH}_2\text{CH}_2)_2\text{OEt}]_2 \cdot 0.5\text{DMF}$.

Thus, it is obvious that lead(II) coordinates the discussed derivatives of the boron clusters with the B–O bonds via the 3c2e PbHB bonds and functional groups of the substituents, whereas nickel(II) and gadolinium(III) are too hard to coordinate the boron cage and are able to coordinate the substituents.

2.9. S-thiocyanato Substituents

The reaction between *closo*-dodecaborate and dirodane in dichloromethane afforded the **S-thiocyanato-derivative of the *closo*-decaborate anion** $[\text{B}_{12}\text{H}_{11}\text{SCN}]^{2-}$ [81]; its interaction with trinuclear mercury complex $(o\text{-C}_6\text{F}_4\text{Hg})_3$ yielded half-sandwich and sandwich mercury complexes $\{(o\text{-C}_6\text{F}_4\text{Hg})_3[\text{B}_{12}\text{H}_{11}\text{SCN}]\}^{2-}$ and $\{[(o\text{-C}_6\text{F}_4\text{Hg})_3]_2[\text{B}_{12}\text{H}_{11}\text{SCN}]\}^{2-}$ [85,86]. The latter complex (Figure 13) has the structure of a wedge-shaped sandwich with $[\text{B}_{12}\text{H}_{11}\text{SCN}]^{2-}$ located between two $(o\text{-C}_6\text{F}_4\text{Hg})_3$ molecules (Hg–H 2.56–3.18 Å, Hg–B 3.317(10)–3.546(11) Å).




Figure 13. Structure of mercury complex in $(\text{Bu}_4\text{N})_2\{[(o\text{-C}_6\text{F}_4\text{Hg})_3]_2[\text{B}_{12}\text{H}_{11}\text{SCN}]\}$.

2.10. Diazo Substituents

The reaction of triethylammonium *closo*-decaborate with 2,4,6-tribromophenyldiazonium tetrafluoroborate in acetonitrile leads to the formation of triethylammonium 1-diazo-*closo*-decaborate $(\text{Et}_3\text{NH})[1\text{-B}_{10}\text{H}_9\text{N}_2]$. When it reacted with copper(I) chloride, copper(I) complex $[\text{Et}_3\text{NH}][\text{Cu}[1\text{-B}_{10}\text{H}_9\text{N}_2]_2]$ with a **singly charged diazo-substituted derivative** $[1\text{-B}_{10}\text{H}_9\text{N}_2]^-$ was isolated [87] containing a linear B–N≡N group. This complex is built of copper(I) anionic complex $[\text{Cu}[1\text{-B}_{10}\text{H}_9\text{N}_2]_2]^-$ (Figure 14a), in which copper(I) coordinates two $[1\text{-B}_{10}\text{H}_9(\text{N}_2)]^-$ monoanions along the apical edge (Cu–B 2.184(9), 2.168(8) Å; Cu–H 1.96, 1.99 Å), opposite to the introduced N≡N substituent.




Figure 14. (a) Anionic copper(I) complex in $[Et_3NH][Cu[1-B_{10}H_9N_2]_2]$ and (b) molecular ruthenium(II) complex $[RuH_2[N_2B_{10}H_8SMe_2](Ph_3P)_3] \cdot 3C_6H_6$.

Ruthenium complexation starting from $[RuH_2(N_2)(PPh_3)_3]$ with the **neutral disubstituted derivative 1,10-(dimethylsulfonio)diazo-*closo*-decaborane** $[N_2B_{10}H_8SMe_2]$ gave ruthenium(II) complex $[RuH_2[N_2B_{10}H_8SMe_2](Ph_3P)_3] \cdot 3C_6H_6$ [88]. In this complex, the $Ru-N \equiv N-B$ group is linear according to X-ray diffraction (Figure 14b) ($Ru-N$ 1.889 Å; $Ru-H$ 1.53(7), 1.74(7); $B-N$ 1.498 Å).

Thus, copper(I) coordinates the diaza-substituted monocharged derivative forming 3c2e CuHB bonds, whereas ruthenium(II) coordinates the neutral derivative with additional dimethylsulfonium substitution by the diaza-group. The absence of RuHB bonds can be explained by two hydride atoms bonded with the metal atom and the neutral charge of the boron cage.

2.11. Cyano-Substituents

The **1,10-dicyano-*closo*-decaborate ion** $[1,10-B_{10}H_8(CN)_2]^{2-}$ was obtained by heating the $[1,10-B_{10}H_8(CONH_2)_2]^{2-}$ derivative, which in turn is formed upon sequential processing of the *closo*-decaborate anion with nitric acid followed by reduction with sodium borohydride to obtain a neutral compound $[1,10-B_{10}H_8(CO)_2]$ and its further interaction with ammonia [89–91]. The disubstituted derivative $[1,10-B_{10}H_8(CN)_2]^{2-}$ was used in iron(III) complexation in the presence of cyclopentadienyl and phosphine ligands, which resulted in a binuclear iron(III) complex with linear $Fe-N \equiv C-B$ bond of the composition $[(Cp)(dppe)Fe]_2[1,10-B_{10}H_8(NC)_2] \cdot H_2O$ [91]. In the resulting complex, the disubstituted derivative of the boron cluster bridges two metal atoms (Figure 15). In this compound, the derivative is coordinated only by the functional groups of the substituent ($Fe-N$ 1.9102(17) Å, $N-C$ 1.152(3) Å).





Figure 15. Structure of iron(III) complex $[(Cp)(dppe)Fe]_2[1,10-B_{10}H_8(NC)_2] \cdot H_2O$ with linear $M-N \equiv C-B$ groups. Solvent molecules are omitted.

The obtained complex is the first iron(III) complex with boron clusters. Usually, boron clusters reduce metal(III) to metal(II) in the course of complexation (as was observed for cobalt and iron); the presence of the substituent and shielding the metal by the cyclopentadienyl and phosphine ligands resulted in the preparation of iron(III) complex.

2.12. Azaheterocycles as Substituents


The ability of the *closo*-decaborate anion to participate in reactions of the substitution of hydrogen atoms in the course of copper(II) complexation was mentioned above for the OH-substituted derivative. Another example of substitution reactions accompanying the complexation is the reaction of copper(I) complex $[\text{Cu}_2[\text{B}_{10}\text{H}_{10}]]$ with 2,2'-bipyridylamine BPA [92,93] which resulted in redox reaction and afforded copper(II) complex $[\text{Cu}(\text{BPA})_2(\text{NCCH}_3)_2][2\text{-B}_{10}\text{H}_9\text{BPA}]_2 \cdot 2\text{H}_2\text{O}$ (Scheme 1).



Scheme 1. Preparation of complex $[\text{Cu}(\text{BPA})_2(\text{NCCH}_3)_2][2\text{-B}_{10}\text{H}_9\text{BPA}]_2$.

The BPA molecule is bonded to equatorial position B(2) of the boron cage and plays the role of a substituent. The **monosubstituted N-dipyridylamine derivative** $[2\text{-B}_{10}\text{H}_9\text{BPA}]^-$ acts as a counter ion for the mixed-ligand mononuclear cationic Cu(II) complex $[\text{Cu}(\text{BPA})_2(\text{NCCH}_3)_2]^{2+}$ [92]. Copper coordinates two chelating BPA ligands, and acetonitrile complete the coordination sphere of the metal to a distorted octahedron (4 + 2).

The **monosubstituted N-bipyridyl derivative** $[\text{B}_{10}\text{H}_9\text{Bipy}]^-$ was found to form in situ in copper(I) complexation with the unsubstituted *closo*-decaborate anion [93]. In this reaction (Scheme 2), the process of substitution of an exo-polyhedral hydrogen atom for the ligand molecule is observed, which accompanies the copper(I) complexation.



Scheme 2. Preparation of complex $[\text{Cu}(\text{CH}_3\text{CN})_2[\text{B}_{10}\text{H}_9\text{Bipy}]]$.


The substituted derivative $[2\text{-B}_{10}\text{H}_9\text{Bipy}]^-$ in the complex is coordinated by the metal atom via nitrogen atoms (Cu–N 2.028(2) Å) and apical BH group (Cu–B(H) 2.601(3), Cu–H(B) 1.85(3) Å). In addition, two acetonitrile molecules are involved in copper(I) coordination (Cu–N 1.995(3), 1.976(3) Å).

Note that these reactions are of particular interest because the $[2\text{-B}_{10}\text{H}_9\text{L}]^{2-}$ derivatives with azaheterocyclic ligands L cannot be obtained in the course of the acid-catalyzed nucleophilic substitution [94,95]; under acidic conditions, organic ligands (Bipy, Phen, BPA) are protonated to form salts $[\text{LH}]_2[\text{B}_{10}\text{H}_{10}]$ or $[\text{LH}_2][\text{B}_{10}\text{H}_{10}]$ [96,97].

The data obtained indicate that copper(II) cannot coordinate the monocharged substituted derivatives whereas copper(I) is able to form complexes with the inner-sphere boron cluster. The possibility of the Bipy molecule to bend across the linker C–C bond allows it to be coordinated by copper(I) and simultaneously to be attached to the boron cage. Note that similar copper(I) complex with Phen ligand cannot be formed because of the rigidity of the ligand.

2.13. Carboxy Groups as Substituents

During the reactions of cobalt(II) complexation in DMF, the solvent can act as a reagent to give substituted derivatives of the boron clusters. Monosubstituted **formoxy derivative** $[2\text{-B}_{10}\text{H}_9\text{OC}(\text{H})\text{O}]^{2-}$ containing the exo-polyhedral B–O bond was isolated when $(\text{Et}_3\text{NH})_2[\text{B}_{10}\text{H}_{10}]$ was allowed to react with CoCl_2 in DMF on heating (Scheme 3). The resulting boron cluster anion contains a formic acid residue $-\text{OC}(\text{H})\text{O}$ as a substituent. The addition of a threefold excess of Phen to the reaction mixture afforded cobalt(II) complex $[\text{Co}(\text{Phen})_3][2\text{-B}_{10}\text{H}_9\text{OC}(\text{H})\text{O}]\cdot 3\text{DMF}$ [98].



Scheme 3. Preparation of complex $[\text{Co}(\text{Phen})_3][2\text{-B}_{10}\text{H}_9\text{OC}(\text{H})\text{O}]$.

Monosubstituted and disubstituted derivatives of the B_{10} polyhedron with acetoxo groups as exo-polyhedral substituents can be obtained by reacting salts of the $[\text{B}_{10}\text{H}_{10}]^{2-}$ anion with acetic acid varying synthesis conditions [99,100]. These derivatives were also found to act as ligands in lead(II) complexation with Bipy [101]. The target compounds were obtained when the substituted derivative was allowed to react with solid $\text{Pb}(\text{NO}_3)_2$ in organic solvent; some part of lead(II) nitrate dissolved in the reaction mixture; after filtration of the unsolved $\text{Pb}(\text{NO}_3)_2$, a Bipy solution in the same solvent was added, giving a yellow color to the resulting mixture. As a result of lead(II) complexation, the target complexes precipitated.

Lead(II) complex $(\text{Ph}_4\text{P})[\text{Pb}(\text{Bipy})[2\text{-B}_{10}\text{H}_9\text{OC}(\text{O})\text{CH}_3]_2]$ [100] and $[\text{Pb}(\text{Bipy})_2[2,7(8)\text{-B}_{10}\text{H}_8(\text{OC}(\text{O})\text{CH}_3)_2]]$ [99] with **monosubstituted** $[2\text{-B}_{10}\text{H}_9\text{OC}(\text{O})\text{CH}_3]^{2-}$ and **disubstituted** $[2,7(8)\text{-B}_{10}\text{H}_8(\text{OC}(\text{O})\text{CH}_3)_2]^{2-}$ **acetoxo derivatives** are known (Figure 16). In both compounds, Pb(II) coordinates one or two Bipy molecules (Pb–N 2.551(3)–2.581(3) Å), one or two oxygen atoms of the carboxylate groups of the substituent (Pb–O 2.749(3)–2.760(3) Å), and BH groups of the boron cage forming PbHB bonds (Pb–B 2.989(5)–3.263(8) Å; Pb–H 2.60(4)–2.93(4) Å).





Figure 16. Structures of lead(II) complexes (a) $(\text{Ph}_4\text{P})[\text{Pb}(\text{Bipy})[2\text{-B}_{10}\text{H}_9\text{OC}(\text{O})\text{CH}_3]_2]$ (cation is omitted) and (b) $[\text{Pb}(\text{Bipy})_2[2,8\text{-B}_{10}\text{H}_8(\text{OC}(\text{O})\text{CH}_3)_2]]$.

The **monocharged disubstituted *closo*-decaborate derivative with the bidentate acetate group** $\text{Cat}[2,6(9)\text{-B}_{10}\text{H}_8\text{>}(\text{O})_2\text{CCH}_3]$ ($\text{Cat} = \text{Ph}_4\text{P}^+, \text{Ph}_4\text{As}^+$) can be used to prepare a compound with two different substituents (Scheme 4). In the course of lead(II) complexation, it undergoes partial hydrolysis, the acetate group remains monodentately bound to the boron cluster in the B(2) position, while the OH group was found to act as a substituent in the B(6) position.



Scheme 4. Preparation of monoanion $[2,6(9)\text{-B}_{10}\text{H}_8\text{>}(\text{O})_2\text{CCH}_3]^- = [\text{An}]^-$ as salt $\text{Cat}[\text{An}]$ or complex $[\text{Pb}(\text{Bipy})_2[\text{An}]]_2$.

The resulting **derivative with two different substituents $-\text{OH}$ and $-\text{OC}(\text{O})\text{CH}_3$** was isolated as lead(II) complex $[\text{Pb}(\text{Bipy})_2[2,6(9)\text{-B}_{10}\text{H}_8(\text{OC}(\text{O})\text{CH}_3)(\text{OH})]]_2 \cdot 3\text{H}_2\text{O}$ [102]. The lead(II) atom coordinates two Bipy ($\text{Pb}(1)\text{-N}$ 2.510(6), 2.610(7) Å), the OH group ($\text{Pb}(1)\text{-O}$ 2.85(3), 2.95(3) Å), and the BH group of the boron cage ($\text{Pb}(1)\text{-H}$ 2.58, 2.66 Å) (Figure 17). In this case, the acetoxy substituent remains non-coordinated. It can be concluded that despite both the $-\text{OH}$ and $-\text{OC}(\text{O})\text{CH}_3$ groups being potentially active in lead(II) coordination, the hydroxyl group is more favorable for coordination, which can be explained by steric factors, as the acetoxy group create some steric hindrances for lead(II) atoms.





Figure 17. Structure of lead(II) complex $[\text{Pb}(\text{Bipy})_2][2,6(9)\text{-B}_{10}\text{H}_8(\text{OC}(\text{O})\text{CH}_3)(\text{OH})]_2$.

The nucleophilic addition of diethylaminomalonate to the acetonitrile derivative of the *closo*-decaborate anion and the following alkaline hydrolysis of ester groups gives the aminomalononic acid-based product $(\text{NBu}_4)[2\text{-B}_{10}\text{H}_9\text{NHC}(\text{CH}_3)\text{NHCH}(\text{COOH})_2]$ [103]. The complexation reaction between the **derivative of the *closo*-decaborate anion with aminomalononic acid** and hafnium(IV) butoxide and hafnium(IV) diethylamide afforded **hafnium(IV) complex** $(\text{NBu}_4)_2[[2\text{-B}_{10}\text{H}_9\text{NHC}(\text{CH}_3)\text{NHCH}(\text{COO})_2]_2\text{Hf}]$ (Scheme 5), which was characterized by IR spectroscopy, mass-spectrometry, and elemental analysis [103].



Scheme 5. Preparation of complex $(\text{NBu}_4)_2[[2\text{-B}_{10}\text{H}_9\text{NHC}(\text{CH}_3)\text{NHCH}(\text{COO})_2]_2\text{Hf}]_2$.

Thus, in cobalt(II) complexes with carboxy substituents, boron anions act as counterions; lead(II) affords a number of complexes with combined coordination (MHB + substituent). In hafnium(IV) complexes, the substituted derivative is assumed to be coordinated by the metal via the functional group of the substituent introduced into the boron cage.

2.14. Amide Groups as Substituents

Rhodium(III) complex $[\text{Rh}(\text{Me}_5\text{Cp})[\text{B}_{12}\text{H}_{11}\text{NHC}(\text{O})\text{NMe}_2]] \cdot \text{CH}_3\text{CN}$ with a cyclopentadienyl ligand and **3,3-dimethylureido-*closo*-dodecaborate anion** $[\text{B}_{12}\text{H}_{11}\text{NHC}(\text{O})\text{NMe}_2]^{2-}$ was reported [104]. This substituted derivative was prepared by successive treatment of $\text{Cs}[\text{B}_{12}\text{H}_{11}\text{NH}_3]$ with NaH and DMF. The complex was prepared by the reaction of $[\text{B}_{12}\text{H}_{11}\text{NHC}(\text{O})\text{NMe}_2]^{2-}$ with $[\text{Rh}(\text{Me}_5\text{Cp})(\text{CH}_3\text{CN})_3][\text{SbF}_6]_2$ in acetonitrile. The metal coordinates the boron ligand through the O atom of the substituent (Rh–O 2.085(2) Å) and two BH groups of the boron cage (Rh–H 1.951, 1.974 Å; Rh–B 2.428(4), 2.436(4) Å) (Figure 18a).

Later, another rhodium(III) complex with the **monosubstituted benzamido-*closo*-dodecaborate anion** $[\text{B}_{12}\text{H}_{11}\text{NHC}(\text{O})\text{Ph}]^{2-}$ was synthesized and isolated $[\text{Rh}(\text{Me}_5\text{Cp})[\text{B}_{12}\text{H}_{11}\text{NHC}(\text{O})\text{Ph}]] \cdot \text{CH}_3\text{CN}$ [105]. The $[\text{B}_{12}\text{H}_{11}\text{NHC}(\text{O})\text{Ph}]^{2-}$ anion was obtained by acylation of the ammonio-*closo*-dodecaborate anion $[\text{B}_{12}\text{H}_{11}\text{NH}_3]^-$ with benzene chloride [106]. This rhodium complex was prepared by reacting tetrabutylammonium salt of the $[\text{B}_{12}\text{H}_{11}\text{NHC}(\text{O})\text{Ph}]^{2-}$ anion with rhodium complex $[\text{Rh}(\text{Me}_5\text{Cp})(\text{CH}_3\text{CN})_3][\text{SbF}_6]_2$. The resulting complex also demonstrates the combined coordination of the boron cluster via two BH groups (Rh–H 1.954, 1.957 Å; Rh–B 2.427(4), 2.431(4) Å) and the O atom of the substituent (Rh–O 2.095(2) Å) (Figure 18b).




Figure 18. Structure of rhodium(III) complexes (a) $[\text{Rh}(\text{Me}_5\text{Cp})[\text{B}_{12}\text{H}_{11}\text{NHC}(\text{O})\text{NMe}_2]] \cdot \text{CH}_3\text{CN}$ and (b) $[\text{Rh}(\text{Me}_5\text{Cp})[\text{B}_{12}\text{H}_{11}\text{NHC}(\text{O})\text{Ph}]] \cdot \text{CH}_3\text{CN}$.

2.15. Phthalocyanine Derivatives as Substituents

A number of compounds based on boron clusters containing phthalocyanine derivatives are known. Aluminum(III) and cobalt(II) complexes with **tetrakis(methylamino-*closo*-dodecaborato)phthalocyanines** and **octakis(methylamino-*closo*-dodecaborato)phthalocyanines** were isolated [107]. To prepare the final compounds, anion $[\text{B}_{12}\text{H}_{11}\text{NH}_3]^-$ was reduced to $[\text{B}_{12}\text{H}_{11}\text{NH}_2]^{2-}$ with sodium hydride, and halogen-containing phthalocyanines were introduced into the resulting solution. The corresponding sodium salts were prepared (Figure 19). These derivatives of the *closo*-dodecaborate anion contain **the B–N *exo*-polyhedral bond**. The synthesized sodium and cesium salts are the first water-soluble phthalocyanines based on the *closo*-dodecaborate anion, almost unlimitedly soluble in water.




Figure 19. Structures of cobalt(II) and aluminum(III) complexes with the *closo*-dodecaborate derivatives of phthalocyanine containing B–N bonds ($\text{M} = \text{Co}$ or AlOH).

Zinc(II) and cobalt(II) complexes with phthalocyanine derivatives containing pendant *closo*-dodecaborate anions with *exo*-polyhedral B–O and B–S groups were isolated (Figure 20) [108–110]. The possibility to prepare sodium salts of the compounds under discussion soluble in water is very important for their potential application in boron-neutron capture therapy.




Figure 20. Structures of (a) zinc(II) and (b) cobalt(II) complexes with the *closo*-dodecaborate derivatives of phthalocyanine containing B-S and B-O bonds.

In addition, a zinc(II) complex with the *closo*-dodecaborate derivatives of phthalocyanine based on eight 1,4-dioxane derivatives of the *closo*-decaborate anion was isolated (Figure 21) in order to estimate its ability to act as a boron delivery agent for boron neutron capture therapy [111]. The compound was prepared by cyclotetramerization of 4-(3,5-dimethoxyphenoxy)phthalonitrile in the presence of zinc(II) acetate. The boronated phthalocyanine was found to accumulate in A549 human lung adenocarcinoma cells. The maximal cytoplasmic concentration was achieved at an extracellular concentration of $32 \pm 3 \mu\text{M}$. The compound was found to deliver 2.4×10^7 boron atoms per cell.




Figure 21. Structure of zinc(II) complex with the *closo*-dodecaborate derivatives of phthalocyanine containing B-O bonds.

3. Metal Complexes with B-Substituted Derivatives of Carboranes

As indicated above, carboranes have a versatile chemistry involving functionalization of carbon atoms of their cage. In complexation reactions, there are a great number of complexes with derivatives containing C-X exo-polyhedral bonds. Thus, carboranylphosphine ligands (with C-P exo-polyhedral bonds) are generalized in a recent review [62]; chalcogenocarboranes with C-X bonds (X = S, Se, Te) were summarized [112–114]; carboxy [115–121], carbene [122], and acetylene [123] derivatives with the C-C exo-polyhedral bonds were discussed in the corresponding reviews. The carboranyl-based N,O-donor compounds functionalized by the carbon atom of the carborane cage have been thoroughly studied [123–129].

Here, we want to discuss the effect of the substituent attached to the boron atom on the coordination ability of the carborane cage without involving the carbon atoms in the functionalization.

3.1. Derivatives with B–Hal Bonds

Actually, there are a great number of perhalogenated carboranes and complexes based on them. The obtained compounds are weakly coordinating ligands and are studied in detail. Here, we wanted to discuss partially halogenated carboranes as well, because there are few representatives of complexes containing mono- and dihalogenated carboranes.

Monofluoro-substituted derivative of monocarborane $[\text{CB}_{11}\text{H}_{11}\text{F}]^-$ was prepared by fluorination of $\text{Cs}[\text{CB}_{11}\text{H}_{11}\text{F}]^-$ with anhydrous HF [130]. In **silver(I)** complex $[\text{Ag}(\text{C}_6\text{H}_6)_2][12\text{-CB}_{11}\text{H}_{11}\text{F}]$ (Figure 22a), there is no interaction between the silver ion and the fluorine atom: the carborane cage is coordinated by 3c2e BHAg bonds (Ag-H 2.181 Å, Ag-B 2.762 Å). In the related silver(I) complex with **monobrominated derivative** $[\text{CB}_{11}\text{H}_{11}\text{Br}]^-$, on the contrary, there is a strong Ag–Br interaction with the Ag–Br distance of 2.642(1) Å [131].




Figure 22. Structures of (a) $[\text{Ag}(\text{C}_6\text{H}_6)_2][12\text{-CB}_{11}\text{H}_{11}\text{F}]$ and (b) $[\text{Ag}(\text{C}_6\text{H}_6)_2][12\text{-CB}_{11}\text{H}_{11}\text{Br}]$.

Disubstituted derivative of monocarborane $[\text{CB}_9\text{H}_8\text{F}_2]^-$ was used in **silver(I)** complexation, which results in silver(I) complex $[\text{Ag}(\text{C}_6\text{H}_6)_2][6,8\text{-CB}_9\text{H}_8\text{F}_2]$ [132] (Figure 23). It was found that the compound is a molecular silver(I) complex with two coordinated benzene molecules and carborane anion coordinated by two AgHB bonds (Ag-H 2.09, 2.10 Å). The coordination is similar to that observed for monofluorinated carborane in complex $[\text{Ag}(\text{C}_6\text{H}_6)_2][12\text{-CB}_{11}\text{H}_{11}\text{F}]$.




Figure 23. Structure of silver(I) complex $[\text{Ag}(\text{C}_6\text{H}_6)_2][6,8\text{-CB}_9\text{H}_8\text{F}_2]$.

The reaction between molybdenum complex with cyclopentadienyl ligand $[\text{Cp}(\text{CO})_3\text{MoI}]$ and silver salt $[\text{Ag}[\text{CB}_{11}\text{H}_{11}\text{Br}]]$ initially results in an intermediate dimeric **molybdenum-silver** complex $[\text{MoCp}(\text{CO})_3\text{Ag}(\text{CB}_{11}\text{H}_{11}\text{Br})]_2$ (Figure 24a), which has a central $\{\text{AgI}\}_2$ core appended by two carborane anions [133]. The carborane anions are coordinated by the complexing metal via the bromine atom (the Ag–Br bond is 2.6456(8) Å). Prolonged reaction results in elimination of AgI to form molybdenum complex $[\text{MoCp}(\text{CO})_3(\text{CB}_{11}\text{H}_{11}\text{Br})]$ (Figure 24b) with the Mo–Br bond equal to 2.6759(2) Å.




Figure 24. Structures of (a) $[\text{MoCp}(\text{CO})_3\text{IAg}(\text{CB}_{11}\text{H}_{11}\text{Br})]_2$ and (b) $[\text{MoCp}(\text{CO})_3(\text{CB}_{11}\text{H}_{11}\text{Br})]$.

Three isostructural polymeric **silver complexes** with **hexahalogenocarboranes** $[\text{CB}_{11}\text{H}_6\text{Hal}_6]^-$ were isolated [134]: complex $[\text{Ag}(\text{CB}_{11}\text{H}_6\text{Cl}_6)(p\text{-Me}_2\text{C}_6\text{H}_4)]_n$ with coordinated solvent molecules, solvent-free $[\text{Ag}(\text{CB}_{11}\text{H}_6\text{Br}_6)]_n$ (Figure 25a), and $[\text{Ag}(\text{CB}_{11}\text{H}_6\text{I}_6)]_n \cdot 0.5\text{C}_6\text{H}_6$ (Figure 25b) with non-coordinated solvent. The compounds were obtained from *p*-xylene, toluene, and benzene, respectively. In all three compounds, silver atoms coordinate two or three halogen substituents, whereas BH groups around the carbon atom remain uncoordinated. The Ag–Hal bonds are 2.640(1)–2.926(1) Å, av. 2.862(2) Å, and 2.777(4)–3.306(5) Å for Cl, Br, and I, respectively.




Figure 25. Structures of (a) $[\text{Ag}(\text{CB}_{11}\text{H}_6\text{Br}_6)]_n$ and (b) $[(\text{CB}_{11}\text{H}_6\text{Br}_6)\text{PtMe}_3]$.

Platinum complex with **hexabromocarborane** $[\text{CB}_{11}\text{H}_6\text{Br}_6]^-$ was isolated as $[(\text{CB}_{11}\text{H}_6\text{Br}_6)\text{PtMe}_3]$ when the corresponding cesium salt of the carborane reacted with $[\text{Me}_3\text{Pt}(\text{OTf})_4]$ [135]. In the final complex, the complexing agent coordinates the carborane anion by three bromine atoms forming MHB bonds; the Pt–Br bond lengths are 2.7129(17)–2.7279(18) Å.

It is interesting to discuss the position of **silver(I)** atoms in **mixed halocarboranes containing both Cl and Br** substituents. Compounds $[1\text{-H-CB}_{11}\text{Y}_5\text{X}_6]^-$ (X, Y = Cl, Br, I) were prepared in high yield when $[\text{Me}_3\text{NH}][1\text{-H-CB}_{11}\text{H}_5\text{X}_6]$ (X = Cl, Br, I) was treated with proper halogenating reagents at 180–220 °C in a sealed tube [136]. Interestingly, mixed halocarboranes in silver(I) complexes are coordinated from the side opposite to the carbon atom of the carborane cage (Figure 26). Thus, carborane $[1\text{-H-CB}_{11}\text{Br}_5\text{Cl}_6]^-$ in molecular complex $[(\text{solv})_2\text{Ag}[1\text{-H-CB}_{11}\text{Br}_5\text{Cl}_6] \cdot \text{solv}]$ (solv = mesitylene) is coordinated by two chlorine atoms (Ag–Cl 2.986(2) and 2.889(2) Å), whereas $[1\text{-H-CB}_{11}\text{Cl}_5\text{Br}_6]^-$ in related polymeric

complex $[(\text{solv})_2\text{Ag}[1\text{-H-CB}_{11}\text{Cl}_5\text{Br}_6]\cdot\text{solv}]$ is coordinated by two bromine atoms (Ag–Br 2.750(2)–2.873(1) Å).




Figure 26. Structures of (a) $[(\text{solv})_2\text{Ag}[1\text{-H-CB}_{11}\text{Br}_5\text{Cl}_6]\cdot\text{solv}]$ and (b) $[(\text{solv})_2\text{Ag}[1\text{-H-CB}_{11}\text{Cl}_5\text{Br}_6]\cdot\text{solv}]$ (solv = mesitylene).

Iridium(III) hydridophosphine complexes $[\text{IrL}_2\text{H}_2(\text{anion})]$ with $\text{L} = \text{PPh}_3$ or PMe_2Ph and **hexahalogenocarborane anions** $[1\text{-CB}_{11}\text{H}_6\text{Cl}_6]^-$ and $[1\text{-CB}_{11}\text{H}_6\text{I}_6]^-$ were prepared by hydrogenation of cyclooctadiene precursor complexes [137]. In the complexes, the carborane cage is coordinated by two halogen bonds, whereas the BH groups near the *ortho*-position of the carbon atom are uncoordinated. In the structure of $[\text{Ir}(\text{PPh}_3)_2\text{H}_2(1\text{-CB}_{11}\text{H}_6\text{Cl}_6)]$ (Figure 27a), the Ir–Cl bonds are 2.680(1) and 2.655(1) Å.

Palladium(II) complex with uncoordinated **monochlorosubstituted carborane anion** $[\text{CB}_{11}\text{H}_{11}\text{Cl}]^-$ was prepared when $[\text{Pd}(\text{dppe})_2]\text{Cl}_2$ reacted with $[\text{Ag}[\text{CB}_{11}\text{H}_{12}]]$ in CH_2Cl_2 [138]. The monosubstituted derivative was isolated as complex $[\text{Pd}(\text{dppe})_2][\text{CB}_{11}\text{H}_{11}\text{Cl}]_2\cdot 3\text{CH}_2\text{Cl}_2$ in low yield as a result of the complexation reaction affording compound $[\text{Pd}(\text{dppe})][\text{CB}_{11}\text{H}_{12}][\text{CB}_{11}\text{H}_{12}]$ as the main product. In the by-product, the monochlorosubstituted carborane anion acts as a counterion (Figure 27b).





Figure 27. Structures of (a) $[\text{Ir}(\text{PPh}_3)_2\text{H}_2(1\text{-CB}_{11}\text{H}_6\text{Cl}_6)]$ and (b) $[\text{Pd}(\text{dppe})_2][\text{CB}_{11}\text{H}_{11}\text{Cl}]_2\cdot 3\text{CH}_2\text{Cl}_2$.

Hexachloro- or hexabromocarboranes $[\text{CB}_{11}\text{H}_6\text{Cl}_6]^-$ or $[\text{CB}_{11}\text{H}_6\text{Br}_6]^-$ were used in **rhodium(II)** complexation with diphenylphosphine ligands present below (Scheme 6) [139,140], which were synthesized by reacting $[\text{RhCl}(\text{nbd})]_2$ with cesium or sodium salts $[\text{CB}_{11}\text{H}_6\text{Cl}_6]^-$ or $[\text{CB}_{11}\text{H}_6\text{Br}_6]^-$ in methanol at room temperature.



Scheme 6. Structures of ligands in complexes $[\text{RhL}][\text{CB}_{11}\text{H}_6\text{Hal}_6]$ (Hal = Cl, Br).

In all cases, rhodium(II) complexes $[\text{RhL}][\text{CB}_{11}\text{H}_6\text{Cl}_6]$ or $[\text{RhL}][\text{CB}_{11}\text{H}_6\text{Br}_6]$ with carboranes as counter ions were isolated.

When **ruthenium(I)** complex $[\text{CpRu}(\text{NO})(\text{CH}_3)_2]$ was allowed to stand in the presence of an excess of carborane-based protonating agent $[(\text{C}_2\text{H}_5\text{OC}_2\text{H}_5)_2\text{H}][\text{CB}_{11}\text{H}_6\text{Br}_6]$ in acetonitrile, complex $[\text{Ru}(\text{CH}_3\text{CN})_6][\text{CB}_{11}\text{H}_6\text{Br}_6]$ was isolated [141]. In the obtained complex, ruthenium(I) coordinates acetonitrile molecules, whereas hexabromocarborane $[\text{CB}_{11}\text{H}_6\text{Br}_6]^-$ acts as a counterion.

3.2. Derivatives with B–S Bonds

Rhodium(II) complex $[\text{cis-Rh}(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{S-9-}closo\text{-1,7-C}_2\text{B}_{10}\text{H}_{11})_2]\text{Cl}$ with 2-((2-(di-phenylphosphaneyl)ethyl)thio)-substituent was synthesized by the reaction between $[\text{Rh}(\text{coe})\text{Cl}_2]$ and $[\text{9-(Ph}_2\text{PCH}_2\text{CH}_2\text{S-}closo\text{-1,7-C}_2\text{B}_{10}\text{H}_{11})]$ in dichloromethane at room temperature [(coe) = cyclooctene] (Figure 28a). In this complex, the Rh atom coordinates the sulfur derivative of carborane via the substituent; the Rh–S bonds are 2.3541(16) and 2.3592(16) Å [142].




Figure 28. Structures of (a) $[\text{Rh}(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{S-9-1,7-C}_2\text{B}_{10}\text{H}_{11})_2]\text{Cl}$ and (b) $[\text{cis-Rh}(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{S-9-1,7-C}_2\text{B}_{10}\text{H}_{11})_2][\text{Rh}(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{S-1-CB}_{11}\text{H}_{11})_2]$.

Complex $\text{NMe}_4[\text{cis-Rh}(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{S-1-CB}_{11}\text{H}_{11})_2]$ was synthesized when $[\text{Rh}(\text{coe})\text{Cl}_2]$ was allowed to react with $\text{NMe}_4[1-(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{S-CB}_{11}\text{H}_{11})]$ in dichloromethane at room temperature [142]. When complex $[\text{cis-Rh}(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{S-9-1,7-C}_2\text{B}_{10}\text{H}_{11})_2]\text{Cl}$ reacted with $\text{NMe}_4[\text{cis-Rh}(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{S-1-CB}_{11}\text{H}_{11})_2]$ in methanol, compound $[\text{cis-Rh}(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{S-9-1,7-C}_2\text{B}_{10}\text{H}_{11})_2][\text{cis-Rh}(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{S-1-CB}_{11}\text{H}_{11})_2]$ was isolated, which contains a complex cation and a complex anion. Note that in the complex anion, monocarborane is functionalized via the carbon atom; in the complex cation, the neutral dicarborane with the exo-polyhedral B–S bond is coordinated via the substituent. The Ru–S bond falls in the range 2.3585(11)–2.3612(12) Å.

Thioethyldiphenylphosphineplatinum(II) complexes based on *ortho*-carboranes were isolated [143]. Complex $[\text{cis-Pt}(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{S-9-}closo\text{-1,7-C}_2\text{B}_{10}\text{H}_{11})_2]\text{Cl}_2$ (Figure 29a) was synthesized by the reaction between $[\text{Pt}(\text{cod})\text{Cl}_2]$ and $[\text{9-(Ph}_2\text{PCH}_2\text{CH}_2\text{S-}closo\text{-1,7-C}_2\text{B}_{10}\text{H}_{11})]$.

C₂B₁₀H₁₁] in deuterated dichloromethane. The *ortho*-dicarborane is coordinated by the substituent; the Pt–S bond is 2.2739(16) Å. In the structurally related complex with *meta*-dicarborane (Figure 29b), the Pt–S 2.2719(7) Å.

Complex [cis-Pt(Ph₂PCH₂CH₂S-{9-*closo*-1,7-C₂B₁₀H₁₁})₂](BF₄)₂ (Figure 29c) was synthesized via the reaction between AgBF₄ and [cis-Pt(Ph₂PCH₂CH₂S-{9-*closo*-1,7-C₂B₁₀H₁₁})₂]Cl or [cis-Pt(Ph₂PCH₂CH₂S-{9-*closo*-1,7-C₂B₁₀H₁₁})₂]Cl₂ in deuterated dichloromethane. The platinum(II) atom coordinates the carborane derivatives by the substituent; the Pt–S bond is 2.379 and 2.377 Å.




Figure 29. Structures of (a) [cis-Pt(Ph₂PCH₂CH₂S-{9-1,7-C₂B₁₀H₁₁})Cl₂], (b) [cis-Pt(Ph₂PCH₂CH₂S-{9-1,2-C₂B₁₀H₁₁})Cl₂], and (c) [cis-Pt(Ph₂PCH₂CH₂S-{9-1,7-C₂B₁₀H₁₁})₂](BF₄)₂ (BF₄[−] anions are omitted).

First **gold(I)** complex with triphenylphosphine ligand [Au₂(Ph₃P)₂S{9,12-S₂C₂B₁₀H₁₀}] was isolated [144] (Figure 30a). In the compound, a six-member ring is formed involving the BB edge, two sulfur atoms of the substituents and two metal atoms; the Au–S bonds are 2.3171(16), 2.3161(9) Å; the Au–Au bond is 2.9937(2) Å.

Copper(I) complex [Cu-S-9-*closo*-1,7-C₂B₁₀H₁₁]₄ was synthesized by mechanochemical treatment of copper(I) *meta*-carborane-9-thiolate [Cu-S-9-*closo*-1,7-C₂B₁₀H₁₁] (Figure 30b). In the final complex, (μ-1,7-dicarb-*closo*-dodecaborane(11)-9-thiolato)-tetra-copper(I), four *meta*-carborane derivatives are coordinated by four metal atoms; the Cu–S bond falls in the range 2.163–2.176 Å [145].




Figure 30. Structures of (a) gold(I) complex $[\text{Au}_2(\text{Ph}_3\text{P})_2\text{S}\{9,12\text{-S}_2\text{C}_2\text{B}_{10}\text{H}_{10}\}]$ and (b) copper(I) complex $[\text{Cu-S-9-closo-1,7-C}_2\text{B}_{10}\text{H}_{11}]_4$.

A number of **rhodium(III)** complexes with **9,12-dithiolato-1,2-dicarborane** was isolated [146,147]. Half-sandwich complex $[\text{Cp}^*\text{Rh}\{9,12\text{-S}_2\text{C}_2\text{B}_{10}\text{H}_{10}\}]$ was obtained by reacting *o*-carborane-9,12-dithiol with $[\text{Cp}^*\text{RhCl}_2]_2$ in the presence of a base [147]. Its structure was determined by X-ray diffraction (Figure 31b). Various complexes can be obtained based on this compound; for example, $[\text{Cp}^*\text{Rh}_2(\text{Ph}_3\text{P})_2\{9,12\text{-S}_2\text{C}_2(\text{B}_{10}\text{H}_{10})\}]\text{PF}_6$ was isolated in the reaction with $\text{Rh}(\text{Ph}_3\text{P})_3\text{Cl}$ in the presence of ammonium hexafluorophosphate [146].




Figure 31. Structure of (a) rhodium complex $[\text{Cp}^*\text{Rh}\{9,12\text{-S}_2\text{C}_2(\text{B}_{10}\text{H}_{10})\}]$ and (b) iridium complex $[\text{Cp}^*\text{Ir}\{9,10\text{-S}_2\text{C}_2(\text{B}_{10}\text{H}_{10})\}]$.

Similar **iridium(III)** complex $[\text{Cp}^*\text{Ir}\{9,10\text{-S}_2\text{C}_2\text{B}_{10}\text{H}_{10}\}]$ (Figure 31b) was found to react with R_3P ($\text{R} = \text{Me}, \text{Et}, \text{Ph}, 4\text{-F-C}_6\text{H}_4, 4\text{-OMe-C}_6\text{H}_4$) at room temperature to form a series of phosphine complexes $[\text{Cp}^*\text{Ir}(\text{R}_3\text{P})\{9,10\text{-S}_2\text{C}_2(\text{B}_{10}\text{H}_{10})\}]$ [148–150], which were characterized by X-ray diffraction. The compounds are built in a similar manner (see Figure 32a).

It was found that $[\text{Cp}^*\text{Ir}\{9,10\text{-S}_2\text{C}_2\text{B}_{10}\text{H}_{10}\}]$ reacts with phosphine ligands L ($\text{L} = \text{Ph}_3\text{P}, \text{Me}_2\text{PhP}, \text{MePh}_2\text{P}, \text{Ph}_2\text{PCH}_2\text{PPh}_2$) in dichloromethane at room temperature to give complexes $[\text{Cp}^*\text{Ir}(\text{L})\{9,10\text{-S}_2\text{C}_2\text{B}_{10}\text{H}_{10}\}]$. At the same time, the reaction of $[\text{Cp}^*\text{Ir}\{9,10\text{-S}_2\text{C}_2\text{B}_{10}\text{H}_{10}\}]$ with dppe leads to the formation of a corresponding dimeric complex $[\text{Cp}^*\text{Ir}(\text{Ph}_2\text{PCH}_2)\{9,10\text{-S}_2\text{C}_2\text{B}_{10}\text{H}_{10}\}]_2$ (Figure 32b), where $\text{CH}\cdots\text{HB}$, $\text{CH}\cdots\text{B}$, $\text{CH}\cdots\text{S}$, $\text{CH}\cdots\text{HC}$, and $\text{BH}\cdots\pi$ intermolecular interactions are observed [151].

Similar **cobalt(III)** complex with cyclopentadienyl ligand $[\text{Cp}^*\text{Co}\{9,12\text{-S}_2\text{C}_2\text{B}_{10}\text{H}_{10}\}]$ was isolated and structurally characterized [152]. This complex is built similarly to the previously discussed rhodium(III) and iridium(III) complexes (see Figure 31). The Co-B bonds are 1.174(1) and 2.1735(10) Å. A series of boron-fused 1,4-dithiin compounds were prepared by the reactions of the boron-substituted half-sandwich complex $[\text{Cp}^*\text{Co}\{9,12\text{-S}_2\text{C}_2\text{B}_{10}\text{H}_{10}\}]$ with alkynes.




Figure 32. Structures of iridium complexes (a) $[\text{Cp}^*\text{Ir}((4\text{-F-C}_6\text{H}_4)_3\text{P})\{9,10\text{-S}_2\text{C}_2\text{B}_{10}\text{H}_{10}\}]$ and (b) $[\text{Cp}^*\text{Ir}(\text{Ph}_2\text{PCH}_2)\{9,10\text{-S}_2\text{C}_2\text{B}_{10}\text{H}_{10}\}]_2$.

The neutral **tricobalt(II)** complex $[(\text{Cp})_3\text{Co}_3\{9,10,12\text{-S}_3\text{C}_2\text{B}_{10}\text{H}_9\}]$ (Figure 33a) was obtained by the interaction of trisubstituted *o*-carborane-9,10,12-trithiol with $[\text{CpCo}(\text{CO})_2]$ and three equivalents of FcPF_6 ($\text{Fc} = (\text{C}_5\text{H}_5)_2\text{Fe}$) in the presence of triethylamine in dichloromethane [153]. The addition of an excess of ferrocenium hexafluorophosphate causes a redox reaction to form salt $[(\text{Cp})_3\text{Co}_3\{9,10,12\text{-S}_3\text{C}_2\text{B}_{10}\text{H}_9\}]\text{PF}_6$ (Figure 33b).




Figure 33. Structures of (a) $[(\text{Cp})_3\text{Co}_3\{9,10,12\text{-S}_3\text{C}_2\text{B}_{10}\text{H}_9\}]$ and salt (b) $[(\text{Cp})_3\text{Co}_3\{9,10,12\text{-S}_3\text{C}_2\text{B}_{10}\text{H}_9\}]\text{PF}_6$.

3.3. Derivatives with B–N Bonds

From the Cambridge Structural Database, there are only three examples of carboranes with exo-polyhedral B–N bonds. The authors [154] synthesized new **B-carboranyl phosphine-iminophosphorane ligands** $[3\text{-(N=PPh}_2\text{CH}_2\text{PPh}_2\text{)-1,2-B}_{10}\text{C}_2\text{H}_{11}]$ with the carboranyl group directly attached to the iminophosphorane nitrogen atom through the B(3) boron atom; the obtained derivative was used in **palladium(II)** complexation with $[\text{PdCl}_2(\text{PhCN})_2]$ to give complex $[\text{PdCl}_2(\text{Ph}_2\text{PCH}_2\text{PPh}_2\text{CN})_2\text{B}_{10}\text{C}_2\text{H}_{11}]$ (Figure 34). In the final compound, palladium coordinates N and P atoms of the substituent with the Pd–N bond 2.103(4) Å and Pd–P bond 2.2258(17) Å.




Figure 34. Structure of palladium(II) complex $[\text{PdCl}_2(\text{Ph}_2\text{PCH}_2\text{PPh}_2\text{CN})_2\text{B}_{10}\text{C}_2\text{H}_{11}]$.

Rhenium(I) complexes with **3-isocyano-1,2-dicarba-*closo*-dodecaborane** were isolated when 3-isocyanoderivative of 1,2-carborane was allowed to react with $[\text{NEt}_4][\text{Re}(\text{CO})_3\text{Br}_3]$ and $[\text{Re}(\text{CO})_3(\text{solvent})_3][\text{PF}_6]$ to form final compounds $[\text{Re}(\text{CO})_3\text{L}_3][\text{PF}_6]$ and $[\text{Re}(\text{CO})_3\text{L}_2\text{Br}]$ ($\text{L} = 3\text{-CN-1,2-B}_{10}\text{C}_2\text{H}_{10}$) (Figure 35). In both compounds, the complexing agent coordinates carborane derivatives via the C atom of the CN substituent; the Re–C bond falls in the range 1.958(5)–2.075(2) Å [155].




Figure 35. Structures of complexes (a) $[\text{Re}(\text{CO})_3(3\text{-CN-1,2-B}_{10}\text{C}_2\text{H}_{10})_3][\text{PF}_6]$ (anion is omitted) and (b) $[\text{Re}(\text{CO})_3(3\text{-CN-1,2-B}_{10}\text{C}_2\text{H}_{10})_2\text{Br}]$.

Among the structures closest to the systems discussed we can note the carborane-fused triazole radical anion $[1,2-(3\text{-Ph-1-CH}_3\text{-3-N}_3\text{)-1-CB}_{11}\text{Cl}_{10}]^-$, formed when 1,2-(3-phenyl-3-triazene)-decachloro-1-carba-*closo*-dodecaborate reacts with methyl phthalate [156]. When it is treated with **bis(cyclopentadienyl)cobalt**, a redox reaction occurs with the formation of compound $[\text{Co}(\text{Cp})_2][1,2-(3\text{-Ph-1-CH}_3\text{-3-N}_3\text{)-1-CB}_{11}\text{Cl}_{10}]$ (Figure 36a). In this compound, the monocarborane anion is functionalized simultaneously via the carborane atom of the cage and adjacent BH group; thus, the obtained derivative contains the B–N and C–N exo-polyhedral bonds. The metal atom is coordinated by two cyclopentadienyl ligands, whereas the carborane derivative acts as a counterion. The authors declare that it is an interesting example of a relatively stable radical anion that can be used to obtain functional materials.

Another representative of carborane-derivatives containing the B–N bond is (*t*-butylamino-1,7,9-tricarba-*nido*-undecaborato)-(9-dimethylamine-7,8-dicarba-*nido*-undecaborato)-iron [157] (Figure 36b). This compound is a metallocarborane containing a tricarborane derivative functionalized by the carbon atom and dicarborane cage with the B–N exo-polyhedral bond.




Figure 36. Structures of (a) $[\text{Co}(\text{Cp})_2][1,2-(3\text{-Ph-1-CH}_3\text{-3-N}_3)\text{-1-CB}_{11}\text{Cl}_{10}]$ and (b) $(1\text{-BuNH-1,7,9-C}_3\text{B}_8\text{H}_{10})\text{Fe}(9\text{-N}(\text{CH}_3)_3\text{-7,8-C}_2\text{B}_9\text{H}_{10})$.

3.4. Derivatives with B–O Bonds

The complexes of this type have been isolated only for perhalogenated carboranes. Mono-triflyloxy-substituted carborane can be halogenated to form decachloro derivatives with the exopolyhedral B–OTf bond. The use of $[\text{HCB}_{11}\text{Cl}_{10}\text{OTf}]^-$ in palladium(II) complexation demonstrates that this weakly coordinating anion can act as a counterion in palladium complexes $[(\text{POCOP})\text{Pd}(\text{C}_6\text{D}_5\text{Br})][\text{HCB}_{11}\text{Cl}_{10}\text{OTf}]$ (Figure 37a) and binuclear $[(\text{POCOP})\text{Pd-Cl-Pd}(\text{POCOP})][\text{HCB}_{11}\text{Cl}]$ (POCOP is P,P-1,3-phenylene bis(P,P-diphenylphosphinite) [158]. In structurally related complex $\{[(\text{POCOP})\text{Pd}][\text{HCB}_{11}\text{Cl}_{10}\text{OTf}]\}$ (Figure 37b), palladium coordinated the triflyloxy-substituted carborane via the O atom of the substituent; the B–O bond is 2.2076(15) Å [159].




Figure 37. Structures of palladium complexes (a) $[(\text{POCOP})\text{Pd}(\text{C}_6\text{D}_5\text{Br})][\text{HCB}_{11}\text{Cl}_{10}\text{OTf}]$ and (b) $\{[(\text{POCOP})\text{Pd}][\text{HCB}_{11}\text{Cl}_{10}\text{OTf}]\}$.

4. Conclusions

Complexes with substituted derivatives of boron cluster anions $[\text{B}_n\text{H}_n]^{2-}$, monocarboranes $[\text{CB}_n\text{H}_{n-1}]^-$ and dicarboranes $[\text{C}_2\text{B}_n\text{H}_{n-2}]$ ($n = 10, 12$) isolated to date are listed in Tables 1 and 2. Analyzing the data shown, it can be concluded that the following types of complexes with substituted derivatives of *closo*-borate anions and carboranes can be isolated:

(a) metal complexes with substituted derivatives as counterions

These compounds are built of a cationic metal complex, whereas boron clusters are not coordinated by the metal atom. Note that in these compounds, specific non-bonding interactions $B-H\cdots H-X$ ($X = C, O, N$) are usually observed between the BH group of boron clusters and ligands, organic cations, or solvent molecules;

(b) metal complexes with coordinated substituted derivatives

These compounds contain derivatives of boron cluster anions or carboranes as coordinated ligands. Owing to the electronic and geometrical structure of boron clusters and carboranes and their chemical behavior, the following types of metal bonding with the boron cage are observed: (i) coordination via the 3c2e MHB interactions, in which the metal, boron, and hydrogen are involved in coordination; (ii) coordination of functional groups of the substituent introduced into the borane or carborane cage, while the BH groups remain non-coordinated by the metal atom; (iii) combined coordination, where the boron cluster is coordinated by 3c2e MHB bonds and the functional groups of the substituent.

The presence of 3c2e MHB bonds in the synthesized compounds is clearly manifested in the IR spectra by the appearance of absorption bands $\nu(BH)_{MHB}$ in the range 2400–2100 cm^{-1} , which correspond to the stretching vibrations of coordinated BH groups $\nu(BH)_{MHB}$; at the same time, the $\nu(BH)$ band of “non-coordinated” BH bonds is observed near 2500 cm^{-1} .

In the case of coordination of a substituted derivative due to the functional groups of the substituent, absorption bands characteristic of the corresponding groups appear in the IR spectra of complexes, but these bands are often split into several components and shift towards higher wavenumbers. These characteristic changes indicate the involvement of exo-polyhedral functional groups in coordination by metals.

Analyzing data shown in Tables 1 and 2, it is clear that substituted derivatives of the boron cluster anions give all four types of compounds, whereas compounds with carborane derivatives acting as counterions or coordinated via a substituent are generally formed (Figure 38). Several examples of compounds with MHB bonds or combined coordination were isolated for carborane derivatives, indicating that these types of compounds are not characteristic for carboranes.




Figure 38. Coordination of the substituted derivatives of the boron clusters and carboranes.

In addition, it is clear that metals involved in the complexation of boranes and carboranes are different. Particularly, metals acting as Pearson's soft acids (Cu(I), Ag, Pb, Ru) form complexes with boron clusters coordinated via three-center two-electron MHB bonds. If the substituent reduces the total charge of the system (NH_3 , thionium, oxonium groups), the coordination ability of the obtained derivatives decreases; thus, silver complexes with non-coordinated boron cluster anions are formed as end products, while Ru and Pb are still able to coordinate the singly-charged anions.

It can be concluded that lead and ruthenium demonstrate the greatest affinity for boron cluster anions: it is possible to obtain complexes of these metals with the substituted derivatives coordinated by MHB bonds, through the functional group of the substituent, and combined coordination can also be realized. Note that salts of these metals with non-coordinated boron ligands have not been obtained.

For carboranes, it can be seen that the series of metals forming compounds with coordinated derivatives of carboranes are platinum group metals (Ru, Rh, Pd, Ir, Pt). Only three representatives of substituted derivatives of carboranes were used in silver(I) complexation, whereas no lead(II) complexes have been isolated. It seems that this field of chemistry should be studied intensively and new compounds with MHB bonds could be prepared.

As for substituents introduced into the boron cluster, it is clear from Table 1 that the most studied are substituted derivatives of the *closo*-decaborate and *closo*-dodecaborate anions containing the chlorine atom, hydroxy group or acetoxy group. For Hal atoms, it is clear that the corresponding derivatives are coordinated (if at all) by forming MHB bonds from the side opposite to the positions of substituents introduced into the boron cage. The derivatives containing the OH and acetoxy groups are coordinated by lead forming combined coordination MHB + substituent. Some examples of lead and ruthenium complexes with S-substituted derivatives have been isolated with combined coordination of the boron cage (MHB + substituent).

An analysis of the obtained compounds shows that the most interesting combined variant of the coordination of substituted derivatives of boron cluster anions (MHB + substituent) can be expected with the introduction of substituents that do not reduce the total

charge of the boron cluster, using soft acid metals according to Pearson, and using functional groups that correspond in hardness/softness to the metal.

The majority of complexes with B-substituted carboranes contain the B–Hal or B–S exo-polyhedral bonds. Compounds with other substituents are extremely rare. Analyzing Table 2, it can be concluded that carboranes have lower coordination ability; they form compounds with non-coordinated carboranes (salts) or complexes with coordination of the substituent (Figure 38). Thus, it is clear that the carborane cage should be functionalized to act as inner-sphere ligands owing to a substituent being introduced.

Table 1. Coordination modes found for mono- and disubstituted derivatives of boron cluster anions $[B_nH_n]^{2-}$ ($n = 10, 12$) in complexes. Metals are shown in bold and are highlighted using different colors in order to compare the composition of complexes with the same metal but various types of coordination.

Boron Cluster Anion as a Counterion	Coordination with the Formation of 3c2e MHB Bonds	Coordination by a Substituent	Combined Coordination: MHB + Substituent
[Ni(Bipy)₃][B₁₀H₉OH] [Ni(Phen)₃][B₁₀H₉OH] [Ni(Bipy)₃][B₁₂H₁₁Cl] [Co(Phen)₃][2-B₁₀H₉OC(H)O] [Cu^{II}(BPA)₂(NCCH₃)₂][2-B₁₀H₉BPA]₂ [Cu^{II}₂(bipy)₄(μ-CO₃)][2-B₁₀H₉OH] [Ag(PPh₃)₄][B₁₀H₉NH₃] [Ag(PPh₃)₄][2-B₁₀H₉S(CH₂N(CO)₂C₆H₄)₂] [Ag(PPh₃)₄][2-B₁₀H₉O(CH₂)₅] [Ag(PPh₃)₄][2-B₁₀H₉O(CH₂)₄O]	<i>Et</i> ₃ NH[Cu^I [1-B ₁₀ H ₉ N ₂] ₂] [Ag(CH₃CN)₃]₂[Ag₂[2-B₁₀H₉F]₂]_n [Ag(PPh₃)₄][(PPh₃)₂Ag[B₁₀H₉Cl]] [Ag₂(Ph₃P)₄[B₁₂H₁₁Cl]] [Ag₂(Bipy)₂[2-B₁₀H₉SH]] [Ag₂(Bipy)₂[2-B₁₀H₉S(CH₂C(O)NH₂)]NO₃ (Bu ₄ N)[[(<i>o</i> -C ₆ F ₄ Hg) ₃] ₂ [B ₁₂ H ₁₁ SCN]] (Bu ₄ N) ₂ [(<i>o</i> -C ₆ F ₄ Hg) ₃][B ₁₂ H ₁₁ SCN]] [(PPh ₃) ₂ Cl Ru [B ₁₂ H ₁₁ (NEt ₃)] [Pb[2-B₁₀H₉SH]] [Pb(Bipy)₂[2-B₁₀H₉SH]] [Pb(Bipy)₂[1-B₁₀H₉SMe₂]₂]	[Ph ₃ MeP)] ₂ [Cp Ti Cl ₂ [B ₁₂ H ₁₁ OH] [Ni(en)[2-B₁₀H₉O(CH₂)₂O(CH₂)₂NH(CH₂)₂NH₂]] [Na ₆ (THF) ₁₅][Ni [B ₁₂ H ₁₁ NH ₂]] ₄ [Au(PPh₃)[NH₂-B₁₂H₁₁]] [RuH₂[N₂B₁₀H₈SMe₂](Ph₃P)₃] [Ru[SB₁₂H₁₁](NH₃)₅] [(Cp)(dppe) Fe] ₂ [1,10-B ₁₀ H ₈ (NC) ₂]·H ₂ O phthalocyanine Al(III) , Co(II) , Zn(II) complexes Gd(III) and Hf(IV) complexes	[Cu^I (NCCH ₃) ₂ [2-B ₁₀ H ₉ Bipy]] [Ag₂(Ph₃P)₄[B₁₀H₉C(O)OCH₃]] (Ph ₄ P) ₂ [Pb (Bipy)[2-B ₁₀ H ₉ OC(O)CH ₃] ₂] [Pb(Bipy)[2-B₁₀H₉O(CH₂CH₂)₂OEt]] [Pb(Bipy)(DMF)[2-B₁₀H₉OH]] [Pb(Bipy)(2-B₁₀H₉(OCH₂CH₂)₂OEt)] [Pb(Bipy)(DMF)[B₁₀H₉OH] [Pb(Bipy)₂[2,6(9)-B₁₀H₈(OC(O)CH₃)(OH)]]₂ [Pb(Bipy)₂[2,7(8)-B₁₀H₈(OC(O)CH₃)₂]] Bu ₃ MeN[Ru (PPh ₃) ₂ Cl[B ₁₂ H ₁₁ NH ₂]] Bu ₄ N[Ru (dppb)Cl[B ₁₂ H ₁₁ NH ₂]] [Ru(PPh₃)₂CO[B₁₂H₁₁NH₂]] MePPh ₃ [Rh (PPh ₃) ₂ [B ₁₂ H ₁₁ NH ₂]] [Rh(Me₅Cp)[B₁₂H₁₁NHC(O)NMe₂]] [Rh(Me₅Cp)[B₁₂H₁₁NHC(O)Ph]]

Table 2. Coordination modes found for mono- and disubstituted carboranes in complexes. Metals are shown in bold and are highlighted using different colors in order to compare the composition of complexes with the same metal but various types of coordination.

Boron Cluster Anion as a Counterion	Coordination with the Formation of 3c2e MHB Bonds	Coordination by a Substituent	Combined Coordination: MHB + Substituent
[Pd(dppe)₂][CB₁₁H₁₁Cl]₂·3CH₂Cl₂ [(POCOP)Pd(C₆D₅Br)][HCB₁₁Cl₁₀OTf] [Ru(CH₃CN)₆][CB₁₁H₆Br₆] [RhL][CB₁₁H₆Cl₆] [RhL][CB₁₁H₆Br₆] (L = diphenylphosphine ligands) [Co(Cp)₂][1,2-(3-Ph-1-CH₃-3-N₃)-1-CB₁₁Cl₁₀]	[Ag(C₆H₆)₂[12-CB₁₁H₁₁F]] [Ag(C₆H₆)₂][6,8-CB₉H₈F₂]	[Ag(C₆H₆)[12-CB₁₁H₁₁Br]] [MoCp(CO)₃(CB₁₁H₁₁Br)] [Ir(PPh₃)₂H₂(1-CB₁₁H₆Cl₆)] [Cp*Ir{9,10-S₂C₂(B₁₀H₁₀)}] [Cp*Ir(R₃P){9,10-S₂C₂(B₁₀H₁₀)}] [Cp*Co{9,12-S₂C₂B₁₀H₁₀}] [Cp*Rh{9,12-S₂C₂B₁₀H₁₀}] [<i>cis</i>-Rh(Ph₂PCH₂CH₂S-{9-1,7-C₂B₁₀H₁₁})₂]Cl [<i>cis</i>-Pt(Ph₂PCH₂CH₂S-{1-CB₁₁H₁₁})₂] [<i>cis</i>-Pt(Ph₂PCH₂CH₂S-{9-<i>closo</i>-1,7-C₂B₁₀H₁₁})Cl₂] [<i>cis</i>-Pt(Ph₂PCH₂CH₂S-{9-<i>closo</i>-1,7-C₂B₁₀H₁₁})₂](BF₄)₂ [Au₂(Ph₃P)₂S{9,12-S₂C₂B₁₀H₁₀}] [Cu-S-9-<i>closo</i>-1,7-C₂B₁₀H₁₁]₄ [PdCl₂(Ph₂PCH₂PPh₂CN)₉B₁₀C₂H₁₁] [Re(CO)₃L₂Br] (L = 3-CN-1,2-B₁₀C₂H₁₀) {[(POCOP)Pd][HCB₁₁Cl₁₀OTf]}	[MoCp(CO)₃IAg(CB₁₁H₁₁Br)]₂

Author Contributions: Conceptualization, E.A.M. and K.Y.Z.; methodology, V.V.A.; validation, E.Y.M.; writing—original draft preparation, E.Y.M.; writing—review and editing, V.V.A.; visualization, E.Y.M.; supervision, N.T.K.; project administration, E.A.M. All authors have read and agreed to the published version of the manuscript.

Funding: The work was carried out within the framework of the State Assignment of the Kurnakov Institute of General and Inorganic Chemistry, Russian Academy of Sciences in the field of fundamental scientific research.

Acknowledgments: The authors express their endless gratitude to Igor B. Sivaev (A. N. Nesmeyanov Institute of Organoelement Chemistry, Russian Academy of Sciences) for his comments that helped to improve the quality of the review.

Conflicts of Interest: The authors declare that they have no conflict of interest.

References

1. Hoard, J.L. Structure and Polymorphism in Elemental Boron. In *Borax to Boranes*; American Chemical Society: Washington, DC, USA, 1961; Volume 32, pp. 42–52. <https://doi.org/10.1021/ba-1961-0032.ch006>.
2. Jemmisab, E.D.; Prasadab, D.L.V.K. Icosahedral B₁₂, macropolyhedral boranes, β -rhombohedral boron and boron-rich solids. *J. Solid State Chem.* **2006**, *179*, 2768–2774. <https://doi.org/10.1016/j.jssc.2005.11.041>.
3. Jemmis, E.D.; Balakrishnarajan, M.M. Polyhedral Boranes and Elemental Boron: Direct Structural Relations and Diverse Electronic Requirements. *J. Am. Chem. Soc.* **2001**, *123*, 18, 4324–4330. <https://doi.org/10.1021/ja0026962>.
4. Topnikova, A.P.; Belokoneva, E.L. The structure and classification of complex borates. *Russ. Chem. Rev.* **2019**, *88*, 204. <https://doi.org/10.1070/rcr4835>.
5. Zhou, Y.; Fang, C.; Fang, Y.; Zhu, F. Polyborates in aqueous borate solution: A Raman and DFT theory investigation. *Spectrochim. Acta A Mol. Biomol. Spectrosc.* **2011**, *83*, 82–87. <https://doi.org/10.1016/j.saa.2011.07.081>.
6. Kozerozhets, I.V.; Avdeeva, V.V.; Buzanov, G.A.; Semenov, E.A.; Ioni, Y.V.; Gubin, S.P. A New Approach for the Synthesis of Powder Zinc Oxide and Zinc Borates with Desired Properties. *Inorganics* **2022**, *10*, 212. <https://doi.org/10.3390/inorganics10110212>.
7. Jiang, Y.; Lu, Y. Designing transition-metal-boride-based electrocatalysts for applications in electrochemical water splitting. *Nanoscale* **2020**, *12*, 9327–9351. <https://doi.org/10.1039/d0nr01279c>.
8. Tanaka, T. Crystal Structures of Boron-Rich Rare-Earth Borides, NIMS Digital Library. 2010. Available online: <https://library.nims.go.jp> (accessed on 10 October 2022).
9. Gunda, H.; Klebanoff, L.E.; Sharma, P.A.; Varma, A.K.; Dolia, V.; Jasuja, K.; Stavila, V. Progress, challenges, and opportunities in the synthesis, characterization, and application of metal-boride-derived two-dimensional nanostructures. *ACS Mater. Lett.* **2021**, *3*, 535–556. <https://doi.org/10.1021/acsmaterialslett.1c00086>.
10. *Boron Science: New Technologies and Applications*; Hosmane, N.S., Ed.; CRC Press: Boca Raton, FL, USA, 2012.
11. Grimes, R.N. *Carboranes*; Academic Press: London, UK, 2016. <https://doi.org/10.1016/B978-0-12-801894-1.09989-3>.
12. Murphy, N.; McCarthy, E.; Dwyer, R.; Farràs, P. Boron clusters as breast cancer therapeutics. *J. Inorg. Biochem.* **2021**, *218*, 111412. <https://doi.org/10.1016/j.jinorgbio.2021.111412>.
13. Sivaev, I.B.; Bregadze, V.V. Polyhedral boranes for medical applications: Current status and perspectives. *Eur. J. Inorg. Chem.* **2009**, *2009*, 1433–1450. <https://doi.org/10.1002/ejic.200900003>.
14. *Handbook of Boron Science: With Applications In Organometallics, Catalysis, Materials And Medicine*; Hosmane, N.S., Eagling, R.D., Eds.; World Scientific: Singapore, 2018; Volume 1–4. <https://doi.org/10.1142/q0130>.
15. Boustani, I. Boron Clusters. In *Molecular Modelling and Synthesis of Nanomaterials*; Springer: Cham, Switzerland, 2020; pp. 113–254. https://doi.org/10.1007/978-3-030-32726-2_4.
16. Alexandrova, A.N.; Boldyrev, A.I.; Zhai, H.J.; Wang, L.S. All-boron aromatic clusters as potential new inorganic ligands and building blocks in chemistry. *Coord. Chem. Rev.* **2006**, *250*, 2811–2866. <https://doi.org/10.1039/D2QI00890D>.
17. Kaniowski, D.; Ebenryter-Olbinska, K.; Kulik, K.; Janczak, S.; Maciaszek, A.; Bednarska-Szczepaniak, K.; Lesnikowski, Z. Boron clusters as a platform for new materials: Composites of nucleic acids and oligofunctionalized carboranes (C₂B₁₀H₁₂) and their assembly into functional nanoparticles. *Nanoscale* **2020**, *12*, 103–114. <https://doi.org/10.1039/C9NR06550D>.
18. Abakumov, G.A.; Piskunov, A.V.; Cherkasov, V.K.; Fedushkin, I.L.; Ananikov, V.P.; Eremin, D.B.; Gordeev, E.G.; Beletskaya, I.P.; Averin, A.D.; Bochakarev, M.N.; et al. Organoelement chemistry: Promising growth areas and challenges. *Russ. Chem. Rev.* **2018**, *87*, 393. <https://doi.org/10.1070/RCR4795>.
19. Zhang, Z.; Zhao, Z.; Wang, B.; Zhang, J. Boron based hypergolic ionic liquids: A review. *Green Energy Environ.* **2021**, *6*, 794–822. <https://doi.org/10.1016/j.gee.2020.12.002>.
20. Corrias, A.; Ennas, G.; Musinu, A.; Marongiu, G.; Paschina, G. Amorphous transition metal-boron ultrafine particles prepared by chemical methods. *Chem. Mater.* **1993**, *5*, 1722–1726. <https://doi.org/10.1021/cm00036a009>.
21. Ali, F.; Hosmane, N.S.; Zhu, Y. Boron chemistry for medical applications. *Molecules* **2020**, *25*, 828. <https://doi.org/10.3390/molecules25040828>.

22. Hawthorne, M. F.; Pushechnikov, A. Polyhedral borane derivatives: Unique and versatile structural motifs. *Pure Appl. Chem.* **2012**, *84*, 2279–2288. <http://dx.doi.org/10.1351/PAC-CON-12-02-11>
23. Knoth Jr., W. H.; Michel, R. H. Polymers of Polyhedral Boron Compounds. U.S. Patent 3354121A, 21 November 1967.
24. Hey-Hawkins, E.; Teixidor, C.V. Boron-based compounds. In *Potential and Emerging Applications in Medicine*; Wiley: Chichester, UK, 2018. <https://doi.org/10.1002/9781119275602>.
25. Plešek, J. Potential applications of the boron cluster compounds. *Chem. Rev.* **1992**, *92*, 269–278. <https://doi.org/10.1021/cr00010a005>.
26. Sivaev, I.B.; Bregadze, V.I. Borane, carborane and metallacarborane anions for stabilization of transient and highly reactive intermediates. In *Handbook of Boron Science: with Applications in Organometallics, Catalysis, Materials and Medicine*; World Scientific: Singapore, 2018; Volume 1, p. 147. https://doi.org/10.1142/9781786344649_0004.
27. Goswami, L.N.; Ma, L.; Chakravarty, S.; Cai, Q.; Jalisatgi, S.S.; Hawthorne, M.F. Discrete nanomolecular polyhedral borane scaffold supporting multiple gadolinium (III) complexes as a high performance MRI contrast agent. *Inorg. Chem.* **2013**, *52*, 1694–1700. <https://doi.org/10.1021/ic3017613>.
28. Sivaev, I.B.; Bregadze, V.I.; Kuznetsov, N.T. Derivatives of the *closo*-dodecaborate anion and their application in medicine. *Russ. Chem. Bull.* **2002**, *51*, 1362–1374. <https://doi.org/10.1023/A:1020942418765>.
29. Teixidor, F.; Viñas, C.; Demonceau, A.; Nuñez, R. Boron clusters: Do they receive the deserved interest? *Pure Appl. Chem.* **2003**, *75*, 1305–1313. <https://doi.org/10.1351/pac200375091305>.
30. Lesnikowski, Z.J. Boron units as pharmacophores—new applications and opportunities of boron cluster chemistry. *Collect. Czechoslov. Chem. Commun.* **2007**, *72*, 1646–1658. <https://doi.org/10.1135/cccc20071646>.
31. Hawthorne, M.F. Biochemical applications of boron cluster chemistry. *Pure Appl. Chem.* **1991**, *63*, 327–334. <https://doi.org/10.1351/pac199163030327>.
32. Axtell, J.C.; Saleh, L.M.; Qian, E.A.; Wixtrom, A.I.; Spokoyny, A.M. Synthesis and applications of perfunctionalized boron clusters. *Inorg. Chem.* **2018**, *57*, 2333–2350. <https://doi.org/10.1021/acs.inorgchem.7b02912>.
33. Avdeeva, V.V.; Polyakova, I.N.; Vologzhanina, A.V.; Goeva, L.V.; Buzanov, G.A.; Generalova, N.B.; Malinina, E.A.; Zhizhin, K.Y.; Kuznetsov, N.T. $[\text{Co}(\text{solv})_6][\text{B}_{10}\text{H}_{10}](\text{solv} = \text{DMF and DMSO})$ for low-temperature synthesis of borides. *Russ. J. Inorg. Chem.* **2016**, *61*, 1125–1134. <https://doi.org/10.1134/S0036023616090023>.
34. Malinina, E.A.; Goeva, L.V.; Buzanov, G.A.; Retivov, V.M.; Avdeeva, V.V.; Kuznetsov, N.T. Synthesis and Thermal Reduction of Complexes $[\text{NiL}_n][\text{B}_{10}\text{H}_{10}](\text{L} = \text{DMF, H}_2\text{O, } n = 6; \text{L} = \text{N}_2\text{H}_4, n = 3)$: Formation of Solid Solutions $\text{Ni}_3\text{C}_{1-x}\text{B}_x$. *Russ. J. Inorg. Chem.* **2020**, *65*, 126–132. <https://doi.org/10.1134/S0036023620010118>.
35. Malinina, E.A.; Goeva, L.V.; Buzanov, G.A.; Avdeeva, V.V.; Efimov, N.N.; Kozerozhets, I.V.; Kuznetsov, N.T. Synthesis and Physicochemical Properties of Binary Cobalt (II) Borides. Thermal Reduction of Precursor Complexes $[\text{CoL}_n][\text{B}_{10}\text{H}_{10}](\text{L} = \text{H}_2\text{O, } n = 6; \text{N}_2\text{H}_4, n = 3)$. *Russ. J. Inorg. Chem.* **2019**, *64*, 1325–1334. <https://doi.org/10.1134/S0036023619110123>.
36. Malinina, E.A.; Goeva, L.V.; Buzanov, G.A.; Avdeeva, V.V.; Efimov, N.N.; Kuznetsov, N.T. A New Method for Synthesis of Binary Borides with Desired Properties. *Dokl. Chem.* **2019**, *487*, 180–183. <https://doi.org/10.1134/S0012500819070061>.
37. Fink, K.; Uchman, M. Boron cluster compounds as new chemical leads for antimicrobial therapy. *Coord. Chem. Rev.* **2021**, *431*, 213684. <https://doi.org/10.1016/j.ccr.2020.213684>.
38. Caminade, A.M. Inorganic dendrimers: Recent advances for catalysis, nanomaterials, and nanomedicine. *Chem. Soc. Rev.* **2016**, *45*, 5174–5186. <https://doi.org/10.1039/C6CS00159A>.
39. Barba-Bon, A.; Salluce, G.; Lostalé-Seijo, I.; Assaf, K.; Hennig, A.; Montenegro, J.; Nau, W.M. Boron clusters as broadband membrane carriers. *Nature* **2002**, *603*, 637–642. <https://doi.org/10.1038/s41586-022-04413-w>.
40. Gabel, D. Boron clusters in medicinal chemistry: Perspectives and problems. *Pure Appl. Chem.* **2015**, *87*, 173–179. <https://doi.org/10.1515/pac-2014-1007>.
41. Avdeeva, V.V.; Garaev, T.M.; Malinina, E.A.; Zhizhin, K.Y.; Kuznetsov, N.T. Physiologically Active Compounds Based on Membranotropic Cage Carriers—Derivatives of Adamantane and Polyhedral Boron Clusters. *Russ. J. Inorg. Chem.* **2022**, *67*, 28–47. <https://doi.org/10.1134/S0036023622010028>.
42. Avdeeva, V.V.; Malinina, E.A.; Kuznetsov, N.T. Boron cluster anions and their derivatives in complexation reactions. *Coord. Chem. Rev.* **2022**, *469*, 214636. <https://doi.org/10.1016/j.ccr.2022.214636>.
43. Avdeeva, V.V.; Malinina, E.A.; Sivaev, I.B.; Bregadze, V.I.; Kuznetsov, N.T. Silver and copper complexes with *closo*-polyhedral borane, carborane and metallacarborane anions: Synthesis and X-ray structure. *Crystals* **2016**, *6*, 60. <https://doi.org/10.3390/cryst6050060>.
44. Avdeeva, V.V.; Malinina, E.A.; Kuznetsov, N.T. Coordination chemistry of iron triad metals with organic N-donor ligands and boron cluster anions $[\text{B}_{10}\text{H}_{10}]^{2-}$, $[\text{B}_{12}\text{H}_{12}]^{2-}$, and $[\text{B}_{10}\text{Cl}_{10}]^{2-}$: Complexation and accompanying processes. *Russ. J. Inorg. Chem.* **2017**, *62*, 1673–1702. <https://doi.org/10.1134/S0036023617130022>.
45. Avdeeva, V.V.; Malinina, E.A.; Zhizhin, K.Y.; Kuznetsov, N.T. Salts and Complexes Containing the Decachloro-*closo*-Decaborate Anion. *Russ. J. Coord. Chem.* **2021**, *47*, 519–545. <https://doi.org/10.1134/S1070328421080017>.
46. Kravchenko, E.A.; Gippius, A.A.; Kuznetsov, N.T. Noncovalent Interactions in Compounds Based on Perchlorinated Boron Cluster as Monitored by ^{35}Cl NQR. *Russ. J. Inorg. Chem.* **2020**, *65*, 546–566. <https://doi.org/10.1134/S0036023620040105>.
47. Knapp, C. Weakly coordinating anions: Halogenated borates and dodecaborates. In *Comprehensive Inorg. Chem. II*, 2nd ed.; Reedijk, J., Poeppelmeier, K., Eds.; Elsevier: Amsterdam, The Netherlands, 2013; pp. 651–679. <https://doi.org/10.1016/B978-0-08-097774-4.00125-X>.

48. Avdeeva, V.V.; Malinina, E.A.; Zhizhin, K.Y.; Bernhardt, E.; Kuznetsov, N.T. Structural Diversity of Dimer Clusters Based on the Octadecahydro-Eicosaborate Anion. *J. Struct. Chem.* **2019**, *60*, 692–712. <https://doi.org/10.1134/S0022476619050020>.
49. Avdeeva, V.V.; Korolenko, S.E.; Malinina, E.A.; Kuznetsov, N.T. Solvent Molecules as Ligands in Coordination Compounds of Metals with Boron Cluster Anions and Their Derivatives (A Review). *Russ. J. Gen. Chem.* **2022**, *92*, 393–417. <https://doi.org/10.1134/S1070363222030070>.
50. Avdeeva, V.V.; Malinina, E.A.; Kuznetsov, N.T. Isomerism in Salts and Complexes with Boron Cluster Anions $[B_{10}H_{10}]^{2-}$ and $[B_{20}H_{18}]^{2-}$. *Russ. J. Inorg. Chem.* **2020**, *65*, 335–358. <https://doi.org/10.1134/S003602362003002X>.
51. Avdeeva, V.V.; Malinina, E.A.; Zhizhin, K.Y.; Kuznetsov, N.T. Structural Diversity of Cationic Copper (II) Complexes with Neutral Nitrogen-Containing Organic Ligands in Compounds with Boron Cluster Anions and Their Derivatives. *Russ. J. Inorg. Chem.* **2020**, *65*, 514–534. <https://doi.org/10.1134/S0036023620040026>.
52. Zhizhin, K.Y.; Zhdanov, A.P.; Kuznetsov, N.T. Derivatives of *closo*-decaborate anion $[B_{10}H_{10}]^{2-}$ with exo-polyhedral substituents. *Russ. J. Inorg. Chem.* **2010**, *55*, 2089–2127. <https://doi.org/10.1134/S0036023610140019>.
53. Stogniy, M.Y.; Erokhina, S.A.; Sivaev, I.B.; Bregadze, V.I. Nitrilium derivatives of polyhedral boron compounds (boranes, carboranes, metallocarboranes): Synthesis and reactivity. *Phosphorus Sulfur Silicon Relat. Elem.* **2019**, *194*, 983–988. <https://doi.org/10.1080/10426507.2019.1631312>.
54. Sivaev, I.B.; Prikaznov, A.V.; Naoufal, D. Fifty years of the *closo*-decaborate anion chemistry. *Collect. Czechoslov. Chem. Commun.* **2010**, *75*, 1149–1199. <https://doi.org/10.1135/cccc2010054>.
55. Hansen, B.R.; Paskevicius, M.; Li, H.W.; Akiba, E.; Jensen, T.R. Metal boranes: Progress and applications. *Coord. Chem. Rev.* **2016**, *323*, 60–70. <https://doi.org/10.1016/j.ccr.2015.12.003>.
56. Sivaev, I.B.; Bregadze, V.I.; Sjöberg, S. Chemistry of *closo*-dodecaborate anion $[B_{12}H_{12}]^{2-}$: A review. *Collect. Czechoslov. Chem. Commun.* **2002**, *67*, 679–727. <https://doi.org/10.1135/cccc20020679>.
57. Sivaev, I.B.; Stogniy, M.Y. Mercury derivatives of polyhedral boranes, carboranes, and metallocarboranes. *Russ. Chem. Bull.* **2019**, *68*, 217–253. <https://doi.org/10.1007/s11172-019-2379-5>.
58. Sivaev, I.B. Functional group directed B–H activation of polyhedral boron hydrides by transition metal complexes. *Russ. J. Inorg. Chem.* **2021**, *66*, 1289–1342. <https://doi.org/10.1134/S0036023621090151>.
59. Au, Y.K.; Xie, Z. Recent advances in transition metal-catalyzed selective BH functionalization of *o*-carboranes. *Bull. Chem. Soc. Jpn.* **2021**, *94*, 879–899. <https://doi.org/10.1246/bcsj.20200366>.
60. Hamdaoui, M.; Liu, F.; Cornaton, Y.; Lu, X.; Shi, X.; Zhang, H.; Liu, J.; Spingler, B.; Djukic, J.-P.; Duttwyler, S. An Iridium-Stabilized Borenum Intermediate. *J. Am. Chem. Soc.* **2022**, *144*, 18359–18374. <https://doi.org/10.1021/jacs.2c06298>.
61. Tsang, M.Y.; Vinas, C.; Teixidor, F.; Planas, J.G.; Conde, N.; SanMartin, R.; Herrero, M.T.; Domínguez, E.; Lledós, A.; Vidossich, P.; et al. Synthesis, structure, and catalytic applications for ortho- and meta-carboranyl based NBN pincer-Pd complexes. *Inorg. Chem.* **2014**, *53*, 9284–9295. <https://doi.org/10.1021/ic5013999>.
62. Sivaev, I.B.; Stogniy, M.Y.; Bregadze, V.I. Transition metal complexes with carboranylphosphine ligands. *Coord. Chem. Rev.* **2021**, *436*, 213795. <https://doi.org/10.1016/j.ccr.2021.213795>.
63. Drozdova, V.V.; Malinina, E.A.; Polyakova, I.N.; Razgonyaeva, G.A.; Zhizhin, K.Y.; Kuznetsov, N.T. Reactions of the *closo*-dodecaborate anion $B_{12}H_{12}^{2-}$ with hydrogen halides in dichloroethane. *Russ. J. Inorg. Chem.* **2007**, *52*, 52–57. <https://doi.org/10.1134/S003602360701010X>.
64. Ivanov, S.V.; Ivanova, S.M.; Miller, S.M.; Anderson, O.P.; Solntsev, K.A.; Strauss, S.H. Fluorination of $B_{10}H_{10}^{2-}$ with an *N*-Fluoro Reagent. A New Way To Transform B–H Bonds into B–F Bonds. *Inorg. Chem.* **1996**, *35*, 6914–6915. <https://doi.org/10.1021/ic961043c>.
65. Kubasov, A.S.; Matveev, E.Y.; Klyukin, I.N.; Nichugovskiy, A.I.; Zhizhin, K.Y.; Kuznetsov, N.T. Silver(I) complexes with substituted derivatives of the boron cluster anions as ligands. *Inorg. Chim. Acta* **2020**, *510*, 119749. <https://doi.org/10.1016/j.ica.2020.119749>.
66. Avdeeva, V.V.; Polyakova, I.N.; Vologzhanina, A.V.; Malinina, E.A.; Zhizhin, K.Y.; Kuznetsov, N.T. Positional isomers of mononuclear silver(I) anionic complex $[Ag(Ph_3P)_2[B_{10}H_{10-x}Cl_x]]^-$ ($x = 0$ or 1) with apically and equatorially coordinated decahydrido-*closo*-decaborate and 2-chlorononahydrido-*closo*-decaborate ligands. *Polyhedron* **2017**, *123*, 396–403. <https://doi.org/10.1016/j.poly.2016.12.009>.
67. Matveev, E.Y.; Novikov, I.V.; Kubasov, A.S.; Retivov, V.M.; Zhizhin, K.Y.; Kuznetsov, N.T. Nickel (II) Complexes with Azaheterocyclic Ligands and 2-Hydroxy-*closo*-Decaborate Anion $[2-B_{10}H_9OH]^{2-}$. *Russ. J. Inorg. Chem.* **2021**, *66*, 187–192. <https://doi.org/10.1134/S0036023621020121>.
68. Malinina, E.A.; Korolenko, S.E.; Zhdanov, A.P.; Avdeeva, V.V.; Privalov, V.I.; Kuznetsov, N.T. Metal-Promoted Exopolyhedral Substitution of Terminal Hydrogen Atoms in the *closo*-Decaborate Anion $[B_{10}H_{10}]^{2-}$ in the Presence of Copper (II): Formation of the Substituted Derivative $[2-B_{10}H_9OH]^{2-}$. *J. Clust. Sci.* **2021**, *32*, 755–763. <https://doi.org/10.1007/s10876-020-01840-5>.
69. Peymann, T.; Knobler, C.B.; Hawthorne, M.F. A Study of the Sequential Acid-Catalyzed Hydroxylation of Dodecahydro-*closo*-dodecaborate ($2-$). *Inorg. Chem.* **2000**, *39*, 1163–1170. <https://doi.org/10.1021/ic991105+>.
70. Hertler, W.R.; Raasch, M.S. Chemistry of Boranes. XIV. Amination of $B_{10}H_{10}^{2-}$ and $B_{12}H_{12}^{2-}$ with Hydroxylamine-O-sulfonic Acid. *J. Am. Chem. Soc.* **1964**, *86*, 3661–3668. <https://doi.org/10.1021/ja01072a014>.
71. Zhdanov, A.P.; Voinova, V.V.; Klyukin, I.N.; Buzanov, G.A.; Grigor'ev, M.S.; Zhizhin, K.Y.; Kuznetsov, N.T. Complex $[Ag(PPh_3)_4][2-B_{10}H_9NH_3 \cdot 2DMF]$: Synthesis and Structure. *Russ. J. Coord. Chem.* **2019**, *45*, 563–568. <https://doi.org/10.1134/S1070328419080098>.

72. Orlova, A.M.; Sivaev, I.B.; Lagun, V.L.; Katser, S.B.; Solntsev, K.A.; Kuznetsov, N.T. Synthesis and structure of PbBipy-2-(1-B₁₀H₉S(CH₃)₂)₂. *Koord. Khim.* **1993**, *19*, 116–121.
73. Elrington, M.; Greenwood, N.N.; Kennedy, J.D.; Thornton-Pett, M. Ruthenium(II) complexes of closo-dodecaboranyl anions and the molecular structure of the pileo thirteen-vertex ruthenaborane [(PPh₃)₂ClRuB₁₂H₁₁(NEt₃)]. *J. Chem. Soc. Dalton Trans.* **1987**, 451–456. <https://doi.org/10.1039/DT9870000451>.
74. Kirchmann, M.; Wesemann, L. η^1 and η^2 Coordination of 1-amino-closo-dodecaborate. *Dalton Trans.* **2008**, 2144–2148. <https://doi.org/10.1039/B718569C>.
75. Kirchmann, M.; Wesemann, L. Amino-closo-dodecaborate—A new ligand in coordination chemistry. *Dalton Trans.* **2008**, 444–446. <https://doi.org/10.1039/b715305h>.
76. Knoth, W.H.; Hertler, W.R.; Muetterties, E.L. Chemistry of Boranes. XXIII. B₁₀H₉S(CH₃)₂- and B₁₀H₈[S(CH₃)₂]₂. *Inorg. Chem.* **1965**, *4*, 280–287. <https://doi.org/10.1021/ic50025a004>.
77. Golubev, A.V.; Kubasov, A.S.; Turyshchev, E.S.; Bykov, A.Y.; Zhizhin, K.Y.; Kuznetsov, N.T. Perbrominated Sulfonium-Substituted closo-Decaborates with exo-Polyhedral Amino Groups [2-B₁₀Br₉S((CH₂)_nNH₂)₂][−] (*n* = 1–3). *Russ. J. Inorg. Chem.* **2020**, *65*, 1333–1342. <https://doi.org/10.1134/S0036023620090041>.
78. Malinina, E.A.; Korolenko, S.E.; Goeva, L.V.; Kuznetsov, N.T. Synthesis and Structure of New Water-Soluble Ag(I) and Pb(II) Complexes with Sulfonyl-Substituted Derivatives of the closo-Decaborate Anion. *Dokl. Chem.* **2018**, *483*, 297–300. <https://doi.org/10.1134/S0012500818120042>.
79. Brattsev, V.A.; Morris, J.H. *Advances in Neutron Capture Therapy*; Larsson, B., Crawford, J., Weinreich, R., Eds.; Elsevier: Amsterdam, The Netherlands, 1997; Volume 2, p. 51.
80. Broomhead, J.A.; Lynch, M.J. The use of the borocaptate anion as a ligand. Synthesis and X-ray crystal structure of pentaammine (1-thiolato-closo-undecahydrododecaborane) ruthenium(III) dihydrate. *Inorg. Chim. Acta* **1995**, *240*, 19–23. <https://doi.org/10.1016/0020-169304645-3>.
81. Zhizhin, K.Y.; Mustaysta, V.N.; Malinina, E.A.; Votnova, N.A.; Matveev, E.Y.; Goeva, L.V.; Kuznetsov, N.T. Interaction of closo-decaborate anion B₁₀H₁₀^{2−} with cyclic ethers. *Russ. J. Inorg. Chem.* **2004**, *49*, 180–189.
82. Kubasov, A.S.; Matveev, E.Y.; Retivov, V.M.; Akimov, S.S.; Razgonyaeva, G.A.; Polyakova, I.N.; Votnova, N.A.; Zhizhin, K.Y.; Kuznetsov, N.T. Nickel(II) complexes with nitrogen-containing derivatives of the closo-decaborate anion. *Russ. Chem. Bull.* **2014**, *63*, 187. <https://doi.org/10.1007/s11172-014-0412-2>.
83. Zhizhin, K.Y.; Mustyatsa, V.N.; Malinina, E.A.; Matveev, E.Y.; Goeva, L.V.; Polyakova, I.N.; Kuznetsov, N.T. Nucleophilic cleavage of cyclic substituents in derivatives of the closo-decaborate anion. *Russ. J. Inorg. Chem.* **2005**, *50*, 243.
84. Akimov, S.S.; Matveev, E.Y.; Kubasov, A.S.; Razgonyaeva, G.A.; Zhizhin, K.Y.; Kuznetsov, N.T. Polydentate ligands based on closo-decaborate anion for the synthesis of gadolinium(III) complexes. *Russ. Chem. Bull.* **2013**, *62*, 1417–1421. <https://doi.org/10.1007/s11172-013-0204-0>.
85. Sivaev, I.B.; Katser, S.B.; Solntsev, K.A.; Kuznetsov, N.T. Some peculiarities of the chemistry of the B₁₂H₁₁SCN^{2−} anion: Crystal structure of (thiocyanato)undecahydro-closo-dodecaborates. *Russ. J. Inorg. Chem.* **1995**, *40*, 779.
86. Tikhonova, I.A.; Shubina, E.S.; Dolgushin, F.M.; Tugashov, K.I.; Teplitskaya, L.N.; Filin, A.M.; Sivaev, I.B.; Petrovskii, P.V.; Furin, G.G.; Bregadze, V.I.; et al. Crown compounds for anions. Sandwich complex of cyclic trimeric perfluoro-*o*-phenylenemercury with [B₁₂H₁₁SCN]^{2−} anion. *Russ. Chem. Bull.* **2003**, *52*, 594–600. <https://doi.org/10.1023/A:1023994320029>.
87. Ng, L.L.; Ng, B.K.; Shelly, K.; Knobler, C.B.; Hawthorne, M.F. Structural analysis of [Et₃NH][Cu(1-B₁₀H₉N₂)₂] and [Et₃NH][1-B₁₀H₉N₂]. *Inorg. Chem.* **1991**, *30*, 4278–4280. <https://doi.org/10.1021/ic00022a034>.
88. Schramm, K.D.; Ibers, J.A. A linear RN₂ transition metal linkage. The structure of Ru(H₂)(N₂)(B₁₀H₈)(SMe₂)(PPh₃)₃. *Inorg. Chem.* **1977**, *16*, 3287–3293. <https://doi.org/10.1021/ic50178a060>.
89. Jacob, L.; Rzeszotarska, E.; Pietrzak, A.; Young Jr, V.G.; Kaszyński, P. Synthesis, Structural Analysis, and Functional Group Interconversion in the [closo-B₁₀H₈-1, 10-X₂]^{2−} (X = CN, [OCRNMe₂]⁺, OCOR, and [OH₂]⁺) Derivatives. *Eur. J. Inorg. Chem.* **2020**, *2020*, 3083–3093. <https://doi.org/10.1002/eqic.202000456>.
90. Hertler, W.R.; Knoth, W.H.; Muetterties, E.L. Chemistry of Boranes. XXII. Polyhedral Rearrangement of Derivatives of B₁₀H₁₀^{2−} and B₁₂H₁₂^{2−}. *J. Am. Chem. Soc.* **1964**, *86*, 5434–5439. <https://doi.org/10.1021/ja01078a015>.
91. Guschlbauer, J.; Shaughnessy, K.H.; Pietrzak, A.; Chung, M.C.; Sponsler, M.B.; Kaszyński, P. [closo-B₁₀H₈-1, 10-(CN)₂]^{2−} as a Conduit of Electronic Effects: Comparative Studies of Fe··Fe Communication in [(η^5 -Cp)(dppe)Fe]₂[μ_2 -(NC-X-CN)]^{*n*+} (*n* = 0, 2). *Organometallics* **2021**, *40*, 2504–2515. <https://doi.org/10.1021/acs.organomet.1c00269>.
92. Dziova, A.E.; Avdeeva, V.V.; Polyakova, I.N.; Belousova, O.N.; Malinina, E.A.; Kuznetsov, N.T. exo-Polyhedral substitution in B₁₀H₁₀^{2−} anion induced by redox reactions in the Cu(I)-B₁₀H₁₀^{2−}-L system (L = bipy, bpa). *Dokl. Chem.* **2011**, *440*, 253–256. <https://doi.org/10.1134/S0012500811090035>.
93. Avdeeva, V.V.; Dziova, A.E.; Polyakova, I.N.; Goeva, L.V.; Malinina, E.A.; Kuznetsov, N.T. Redox, complexation, and substitution reactions in [Cu₂B₁₀H₁₀]-2, 2'-bipyridylamine-CH₃CN system. *Russ. J. Inorg. Chem.* **2013**, *58*, 657–663. <https://doi.org/10.1134/S003602361306003X>.
94. Mahfouz, N.; Abi Ghaida, F.; El Hajj, Z.; Diab, M.; Floquet, S.; Mehdi, A.; Naoufal, D. Recent Achievements on Functionalization within closo-Decahydrodecaborate [B₁₀H₁₀]^{2−} Clusters. *ChemistrySelect* **2022**, *7*, e202200770 <https://doi.org/10.1002/slct.202200770>.
95. Retivov, V.M.; Matveev, E.Y.; Lisovskiy, M.V.; Razgonyaeva, G.A.; Ochertyanova, L.I.; Zhizhin, K.Y.; Kuznetsov, N.T. Nucleophilic substitution in closo-decaborate [B₁₀H₁₀]^{2−} in the presence of carbocations. *Russ. Chem. Bull.* **2010**, *59*, 550–555. <https://doi.org/10.1007/s11172-010-0123-2>.

96. Avdeeva, V.V.; Vologzhanina, A.V.; Malinina, E.A.; Kuznetsov, N.T. Dihydrogen Bonds in Salts of Boron Cluster Anions $[B_nH_n]^{2-}$ with Protonated Heterocyclic Organic Bases. *Crystals* **2019**, *9*, 330. <https://doi.org/10.3390/cryst9070330>.
97. Goeva, L.V.; Zhuchkova, A.F.; Malinina, E.A.; Korolenko, S.E.; Avdeeva, V.V.; Kuznetsov, N.T. Radiation-Chemical Transformations of 7-NH₃-4-CH₃-Coumarin Decahydro-*closo*-Decaborate as a Potential Inhibitor of Free Radicals. *Russ. J. Inorg. Chem.* **2022**, *67*, 1144–1150. <https://doi.org/10.1134/S0036023622080149>.
98. Avdeeva, V.V.; Polyakova, I.N.; Churakov, A.V.; Vologzhanina, A.V.; Malinina, E.A.; Zhizhin, K.Y.; Kuznetsov, N.T. Complexation and exopolyhedral substitution of the terminal hydrogen atoms in the decahydro-*closo*-decaborate anion in the presence of cobalt(II). *Polyhedron* **2019**, *162*, 65–70. <https://doi.org/10.1016/j.poly.2019.01.051>.
99. Zhizhin, K.Y.; Vovk, O.O.; Malinina, E.A.; Mustyatsa, V.N.; Goeva, L.V.; Polyakova, I.N.; Kuznetsov, N.T. Interaction between a Decahydro-*closo*-Decaborate (2-) Anion and Aliphatic Carboxylic Acids. *Russ. J. Coord. Chem.* **2001**, *27*, 613–619. <https://doi.org/10.1023/A:1017937102647>.
100. Zhizhin, K.Y.; Malinina, E.A.; Polyakova, I.N.; Lisovskij, M.V.; Kuznetsov, N.T. Acid-catalyzed nucleophilic substitution in the *closo*-decaborate $B_{10}H_{10}^{2-}$ anion. *Russ. J. Inorg. Chem.* **2002**, *47*, 1168.
101. Malinina, E.A.; Goeva, L.V.; Kuznetsov, N.T. (2009). Multicenter interactions in lead (II) coordination compounds with $B_nH_n^{2-} = 6, 10, 12$ cluster anions and their derivatives. *Russ. J. Inorg. Chem.* **2009**, *54*, 417–424. <https://doi.org/10.1134/S0036023609030152>.
102. Avdeeva, V.V.; Polyakova, I.N.; Goeva, L.V.; Malinina, E.A.; Kuznetsov, N.T. $[2,6(9)-B_{10}H_8(O)_2CCH_3]^-$ and $[2,7(8)-B_{10}H_8(OC(O)CH_3)_2]^{2-}$ derivatives in synthesis of position isomers of the $[B_{10}H_8(OC(O)CH_3)(OH)]^{2-}$ anion with the 2,6(9)- and 2,7(8)-arrangement of functional groups. *Russ. J. Inorg. Chem.* **2014**, *59*, 1247–1258. <https://doi.org/10.1134/S0036023614110047>.
103. Voinova, V.V.; Klyukin, I.N.; Zhdanov, A.P.; Grigor'ev, M.S.; Zhizhin, K.Y.; Kuznetsov, N.T. Synthesis of New Boron-Containing Ligands and Their Hafnium (IV) Complexes. *Russ. J. Inorg. Chem.* **2020**, *65*, 839–845. <https://doi.org/10.1134/S0036023620060261>.
104. Zhang, Y.; Sun, Y.; Lin, F.; Liu, J.; Duttwyler, S. Rhodium (III)-Catalyzed Alkenylation–Annulation of *closo*-Dodecaborate Anions through Double B–H Activation at Room Temperature. *Angew. Chem.* **2016**, *128*, 15838–15843. <https://doi.org/10.1002/anie.201607867>.
105. Zhang, Y.; Wang, T.; Wang, L.; Sun, Y.; Lin, F.; Liu, J.; Duttwyler, S. RhIII-Catalyzed Functionalization of *closo*-Dodecaborates by Selective B–H Activation: Bypassing Competitive C–H Activation. *Chem. A Eur. J.* **2018**, *24*, 15812–15817. <https://doi.org/10.1002/chem.201803455>.
106. Sun, Y.; Zhang, J.; Zhang, Y.; Liu, J.; van der Veen, S.; Duttwyler, S. The *closo*-Dodecaborate Dianion Fused with Oxazoles Provides 3D Diboraheterocycles with Selective Antimicrobial Activity. *Chem. A Eur. J.* **2018**, *24*, 10364–10371. <https://doi.org/10.1002/chem.201801602>.
107. Zhizhin, K.Y.; Malinina, E.A.; Goeva, L.V.; Chernyavskaya, A.S.; Ivanov, S.V.; Luk'yanets, E.A.; Kuznetsov, N.T. New boron-containing water-soluble phthalocyanine. *Dokl. Akad. Nauk Ross. Akad. Nauk* **1997**, *357*, 206–208.
108. Fabris, C.; Jori, G.; Giuntini, F.; Roncucci, G. Photosensitizing properties of a boronated phthalocyanine: Studies at the molecular and cellular level. *J. Photochem. Photobiol. B Biol.* **2001**, *64*, 1–7. <https://doi.org/10.1016/S1011-134400184-1>.
109. Sivaev, I.B.; Bregadze, V.I.; Gül, A.; Mironov, A.F. Boron-containing phthalocyanines and porphyrans. *Macrocyclic Chem.* **2012**, *5*, 292–301. <https://doi.org/10.6060/mhc2012.121211s>.
110. Semioshkin, A.; Tsaryova, O.; Zhidkova, O.; Bregadze, V.; Wöhrle, D. Reactions of oxonium derivatives of $(B_{12}H_{12})^{2-}$ with phenols, and synthesis and photochemical properties of a phthalocyanine containing four $(B_{12}H_{12})^{2-}$ groups. *J. Porphyr. Phthalocyanines* **2006**, *10*, 1293–1300. <https://doi.org/10.1142/S1088424606000661>.
111. Patel, H.; Takagaki, M.; Bode, B.P.; Snajdr, I.; Patel, D.; Sharman, C.; Bux, M.; Bux, S.; Kotara, M.; Hosmane, N.S. Carborane-appended saccharides: Prime candidates for boron neutron capture therapy (BNCT) clinical trials. *Biochem. Biophys. J. Neutron Ther. Cancer Treat.* **2013**, *1*, 15–21.
112. Jin, G.X. Advances in the chemistry of organometallic complexes with 1, 2-dichalcogenolato-o-carborane ligands. *Coord. Chem. Rev.* **2004**, *248*, 587–602. <https://doi.org/10.1016/j.ccr.2004.01.002>.
113. Liu, S.; Han, Y.F.; Jin, G.X. Formation of direct metal–metal bonds from 16-electron “pseudo-aromatic” half-sandwich complexes $Cp^*M[E_2C_2(B_{10}H_{10})]$. *Chem. Soc. Rev.* **2007**, *36*, 1543–1560. <https://doi.org/10.1039/B701869j>.
114. Jain, L.; Jain, V.K.; Kushwah, N.; Pal, M.K.; Wadawale, A.P.; Bregadze, V.I.; Glazun, S.A. Chalcogenocarboranes: A family of multifaceted sterically demanding ligands. *Coord. Chem. Rev.* **2014**, *258*, 72–118. <https://doi.org/10.1016/j.ccr.2013.09.003>.
115. Fontanet, M.; Rodríguez, M.; Fontrodona, X.; Romero, I.; Viñas, C.; Teixidor, F. Intramolecular hydrogen bonding stabilizes the nuclearity of complexes. A comparative study based on the H-carborane and Me-carborane framework. *Dalton Trans.* **2015**, *44*, 10399–10409. <https://doi.org/10.1039/C5DT00305A>.
116. Li, Z.; Choquesillo-Lazarte, D.; Fraile, J.; Viñas, C.; Teixidor, F.; Planas, J.G. Rational design of carborane-based Cu 2-paddle wheel coordination polymers for increased hydrolytic stability. *Dalton Trans.* **2022**, *51*, 1137–1143. <https://doi.org/10.1039/D1DT04065K>.
117. Kennedy, R.D.; Krungleviciute, V.; Clingerman, D.J.; Mondloch, J.E.; Peng, Y.; Wilmer, C.E.; Mirkin, C.A. Carborane-based metal–organic framework with high methane and hydrogen storage capacities. *Chem. Mater.* **2013**, *25*, 3539–3543. <https://doi.org/10.1021/cm4020942>.
118. Choi, S.; Lee, H.E.; Ryu, C.H.; Lee, J.; Lee, J.; Yoon, M.; Kim, Y.; Park, M.H.; Lee, K.M.; Kim, M. Synthesis of o-carborane-functionalized metal–organic frameworks through ligand exchanges for aggregation-induced emission in the solid state. *Chem. Commun.* **2019**, *55*, 11844–11847. <https://doi.org/10.1039/C9CC06386B>.

119. Bae, Y.S.; Farha, O.K.; Spokoyny, A.M.; Mirkin, C.A.; Hupp, J.T.; Snurr, R.Q. Carborane-based metal–organic frameworks as highly selective sorbents for CO₂ over methane. *Chem. Commun.* **2008**, 4135–4137. <https://doi.org/10.1039/B805785K>.
120. Spokoyny, A.M.; Farha, O.K.; Mulfort, K.L.; Hupp, J.T.; Mirkin, C.A. Porosity tuning of carborane-based metal–organic frameworks (MOFs) via coordination chemistry and ligand design. *Inorg. Chim. Acta* **2010**, *364*, 266–271. <https://doi.org/10.1016/j.ica.2010.08.007>.
121. Li, Z.; Núñez, R.; Light, M.E.; Ruiz, E.; Teixidor, F.; Viñas, C.; Ruiz-Molina, D.; Roscini, C.; Planas, J.G. Water-Stable Carborane-Based Eu³⁺/Tb³⁺ Metal–Organic Frameworks for Tunable Time-Dependent Emission Color and Their Application in Anticounterfeiting Bar-Coding. *Chem. Mater.* **2022**, *34*, 4795–4808. <https://doi.org/10.1021/acs.chemmater.2c00323>.
122. Fisher, S.P.; McArthur, S.G.; Tej, V.; Lee, S.E.; Chan, A.L.; Banda, I.; Gregory, A.; Berkley, K.; Tsay, C.; Rheingold, A.L.; et al. Strongly Coordinating Ligands To Form Weakly Coordinating Yet Functional Organometallic Anions. *J. Am. Chem. Soc.* **2019**, *142*, 251–256. <https://doi.org/10.1021/jacs.9b10234>.
123. Wang, Q.-Y.; Wang, J.; Wang, S.; Wang, Z.-Y.; Cao, M.; He, C.-L.; Yang, J.-Q.; Zang, S.-Q.; Mak, T.C.W. *J. Am. Chem. Soc.* **2020**, *142*, 12010–12014. <https://doi.org/10.1021/jacs.0c04638>.
124. Planas, J.G.; Teixidor, F.; Viñas, C. N, O-Type carborane-based materials. *Crystals* **2016**, *6*, 50. <https://doi.org/10.3390/cryst6050050>.
125. Di Salvo, F.; Tsang, M.Y.; Teixidor, F.; Viñas, C.; Planas, J.G.; Crassous, J.; Vanthuyne, N.; Aliaga-Alcalde, N.; Ruiz, E.; Coquerel, G.; et al. A Racemic and Enantiopure Unsymmetric Diiron(III) Complex with a Chiral o-Carborane-Based Pyridylalcohol Ligand: Combined Chiroptical, Magnetic, and Nonlinear Optical Properties. *Chem. A Eur. J.* **2014**, *20*, 1081–1090. <https://doi.org/10.1002/chem.201303037>.
126. Tsang, M.Y.; Rodríguez-Hermida, S.; Stylianou, K.C.; Tan, F.; Negi, D.; Teixidor, F.; Viñas, C.; Choquesillo-Lazarte, D.; Verdugo-Escamilla, C.; Guerrero, M.; et al. Carborane bis-pyridylalcohols as linkers for coordination polymers: Synthesis, crystal structures, and guest-framework dependent mechanical properties. *Cryst. Growth Des.* **2017**, *17*, 846–857. <https://doi.org/10.1021/acs.cgd.6b01682>.
127. Tsang, M.Y.; Di Salvo, F.; Teixidor, F.; Vinas, C.; Planas, J.G.; Choquesillo-Lazarte, D.; Vanthuyne, N. Is molecular chirality connected to supramolecular chirality? The particular case of chiral 2-pyridyl alcohols. *Cryst. Growth Des.* **2015**, *15*, 935–945. <https://doi.org/10.1021/cg501748x>.
128. Tsang, M.Y.; Viñas, C.; Teixidor, F.; Choquesillo-Lazarte, D.; Giner Planas, J. Crystalline Inclusion Compounds of a Palladacyclic Tetraol Host Featuring o-Carborane Units. *Eur. J. Inorg. Chem.* **2017**, *2017*, 4589–4598. <https://doi.org/10.1002/ejic.201700535>.
129. Tsang, M.Y.; Teixidor, F.; Vinas, C.; Choquesillo-Lazarte, D.; Aliaga-Alcalde, N.; Planas, J.G. Synthesis, structures and properties of iron(III) complexes with (o-carboranyl) bis-(2-hydroxymethyl) pyridine: Racemic versus meso. *Inorg. Chim. Acta* **2016**, *448*, 97–103. <https://doi.org/10.1016/j.ica.2016.04.028>.
130. Ivanov, S.V.; Lupinetti, A. J.; Miller, S. M.; Anderson, O. P.; Solntsev, K. A.; Strauss, S. H. Regioselective Fluorination of CB₁₁H₁₂[−]. New Weakly Coordinating Anions. *Inorg. Chem.* **1995**, *34*, 6419–6420. <https://doi.org/10.1021/ic00130a003>.
131. Jelinek, T.; Baldwin, P.; Scheidt, W.R.; Reed, C.A. New weakly coordinating anions. 2. Derivatization of the carborane anion CB₁₁H₁₂[−]. *Inorg. Chem.* **1993**, *32*, 1982–1990. <https://doi.org/10.1021/ic00062a018>.
132. Ivanov, S.V.; Rockwell, J.J.; Miller, S.M.; Anderson, O.P.; Solntsev, K.A.; Strauss, S.H. Reactions of CB₉H₁₀[−] with electrophiles, including the regioselective mono- and dihalogenation of the lower belt. *Inorg. Chem.* **1996**, *35*, 7882–7891. <https://doi.org/10.1021/ic960859a>.
133. Patmore, N.J.; Ingleson, M.J.; Mahon, M.F.; Weller, A.S. Investigation of the synthesis of {Mo(η⁵-C₅H₅)(CO)₃}⁺ fragments partnered with the monoanionic carboranes [closo-CB₁₁H₁₁Br][−], [closo-CB₁₁H₆Br₆][−] and [closo-HCB₁₁Me₁₁][−] by silver saltmetathesis and hydride abstraction. *Dalton Trans.* **2003**, 2894–2904. <https://doi.org/10.1039/b303537a>.
134. Mak, T.W.; Reed, C. Structural diversity in silver salts of hexahalogenocarborane anions, Ag(CB₁₁H₆X₆) (X = Cl, Br or I). *J. Chem. Soc. Dalton Trans.* **1997**, 1213–1218. <https://doi.org/10.1039/A608336F>.
135. De Crisci, A.G.; Kleingardner, J.; Lough, A.J.; Larsen, A.; Fekl, U. A weakly coordinating anion as a tripodal “Br₃”-ligand for platinum (IV) — Structure of [(closo-CB₁₁H₆Br₆)PtMe₃]. *Can. J. Chem.* **2009**, *87*, 95–102. <https://doi.org/10.1139/V08-085>.
136. Tsang, C.W.; Yang, Q.; Sze, E.T. P.; Mak, T.C.; Chan, D.T.; Xie, Z. Weakly coordinating nature of a carborane cage bearing different halogen atoms. Synthesis and structural characterization of icosahedral mixed halocarborane anions, 1-H-CB₁₁Y₅X₆[−] (X, Y = Cl, Br, I). *Inorg. Chem.* **2000**, *39*, 5851–5858. <https://doi.org/10.1021/ic000354r>.
137. Moxham, G.L.; Douglas, T.M.; Brayshaw, S.K.; Kociok-Köhn, G.; Lowe, J.P.; Weller, A.S. The role of halogenated carborane monoanions in olefin hydrogenation catalysed by cationic iridium phosphine complexes. *Dalton Trans.* **2006**, 5492–5505. <https://doi.org/10.1039/b612049k>.
138. Lassahn, P.G.; Lozan, V.; Wu, B.; Weller, A.S.; Janiak, C. Dihalogeno (diphosphane) metal (ii) complexes (metal = Co, Ni, Pd) as pre-catalysts for the vinyl/addition polymerization of norbornene—elucidation of the activation process with B(C₆F₅)₃/AlEt₃ or Ag [closo-1-CB₁₁H₁₂] and evidence for the in situ formation of “naked” Pd²⁺ as a highly active species. *Dalton Trans.* **2003**, 4437–4450. <https://doi.org/10.1039/b302937a>.
139. Moxham, G.L.; Randell-Sly, H.E.; Brayshaw, S.K.; Woodward, R.L.; Weller, A.S.; Willis, M.C. A Second-Generation Catalyst for Intermolecular Hydroacylation of Alkenes and Alkynes Using β-S-Substituted Aldehydes: The Role of a Hemilabile P–O–P Ligand. *Angew. Chem. Int. Ed.* **2006**, *45*, 7618–7622. <https://doi.org/10.1002/anie.200603133>.

140. Moxham, G.L.; Randell-Sly, H.; Brayshaw, S.K.; Weller, A.S.; Willis, M.C. Intermolecular Alkene and Alkyne Hydroacylation with β -S-Substituted Aldehydes: Mechanistic Insight into the Role of a Hemilabile P–O–P Ligand. *Chem. A Eur. J.* **2008**, *14*, 8383–8397. <https://doi.org/10.1002/chem.200800738>.
141. Masland, J.; Diaz, J.; Eady, S.; Lobkovsky, E.; Larsen, A. Hexakis (acetonitrile- κ N) ruthenium (II) bis (hexabromocarbododecaborate) acetonitrile solvate. *Acta Crystallogr. Sect. E Struct. Rep. Online* **2010**, *66*, m325–m326. <https://doi.org/10.1107/S1600536810006252>.
142. Kennedy, R.D.; Stern, C.L.; Mirkin, C.A. Zwitterionic weak-link approach complexes based on anionic icosahedral monocarbaboranes. *Inorg. Chem.* **2013**, *52*, 14064–14071. <https://doi.org/10.1021/ic401851z>.
143. Spokoyny, A.M.; Machan, C.W.; Clingerman, D.J.; Rosen, M.S.; Wiester, M.J.; Kennedy, R.D.; Stern, C.L.; Sarjeant, A.A.; Mirkin, C.A. A coordination chemistry dichotomy for icosahedral carborane-based ligands. *Nat. Chem.* **2011**, *3*, 590–596. <https://doi.org/10.1038/nchem.1088>.
144. Jones, P.G. CSD Communication, 2016. Refcode UXITOM.
145. Yan, H.; Yang, F.; Pan, D.; Lin, Y.; Hohman, J.N.; Solis-Ibarra, D.; Li, F.H.; Dahl, J.E.P.; Carlson, R.M.K.; Tkachenko, B.A.; et al. Sterically controlled mechanochemistry under hydrostatic pressure. *Nature* **2018**, *554*, 505–510. <https://doi.org/10.1038/nature25765>.
146. Zhang, X.; Zhou, Z.; Yan, H. Metal–metal redox synergy in selective B–H activation of ortho-carborane-9,12-dithiolate. *Chem. Commun.* **2014**, *50*, 13077–13080. <https://doi.org/10.1039/C4CC06171C>.
147. Herberhold, M.; Jin, G.X.; Yan, H.; Milius, W.; Wrackmeyer, B. 16- and 18-Electron Cp*Rh complexes with 1,2-dicarba-closo-dodecaborane (12)-1, 2-dichalcogenolato ligands, as studied by multinuclear magnetic resonance. *J. Organomet. Chem.* **1999**, *587*, 252–257. [https://doi.org/10.1016/S0022-328X\(99\)00330-7](https://doi.org/10.1016/S0022-328X(99)00330-7).
148. Zhang, X.; Dai, H.; Yan, H.; Zou, W.; Cremer, D. B–H $\cdots\pi$ interaction: A new type of nonclassical hydrogen bonding. *J. Am. Chem. Soc.* **2016**, *138*, 4334–4337. <https://doi.org/10.1021/jacs.6b01249>.
149. Zou, W.; Zhang, X.; Dai, H.; Yan, H.; Cremer, D.; Kraka, E. Description of an unusual hydrogen bond between carborane and a phenyl group. *J. Organomet. Chem.* **2018**, *865*, 114–127. <https://doi.org/10.1016/j.jorganchem.2018.02.014>.
150. Dai, H.; Li, C.X.; Yu, C.; Wang, Z.; Yan, H.; Lu, C. Copper(II) catalyzed domino synthesis of quinoline derivatives from arylamines and alkynes. *Org. Chem. Front.* **2017**, *4*, 2008–2011. <https://doi.org/10.1039/C7QO00333A>.
151. Cao, H.J.; Dai, H.; Zhang, X.; Yan, H.; Lu, C. Synthesis and characterization of Cp*Ir-dithiolene-o-carborane phosphine complexes: A continuous investigation of B–H $\cdots\pi$ interaction. *Mol. Phys.* **2019**, *117*, 1287–1297. <https://doi.org/10.1080/00268976.2018.1539257>.
152. Zhang, X.; Zou, X.; Yan, H. Synthesis of Boron-Fused 1, 4-Dithiin via Cobalt-Mediated Disulfuration of Alkyne at the o-Carborane-9, 12-dithiolate Unit. *Organometallics* **2014**, *33*, 2661–2666. <https://doi.org/10.1021/om500411b>.
153. Zhang, X.; Tang, X.; Yang, J.; Li, Y.; Yan, H.; Bregadze, V.I. Boron-substituted o-carboranetrithiol to construct trimeric cobalt clusters. *Organometallics* **2013**, *32*, 2014–2018. <https://doi.org/10.1021/om400097m>.
154. Rodríguez-Rey, J.L.; Esteban-Gómez, D.; Platas-Iglesias, C.; Sousa-Pedrares, A. Electronic versus steric control in palladium complexes of carboranyl phosphine-iminophosphorane ligands. *Dalton Trans.* **2019**, *48*, 486–503. <https://doi.org/10.1039/C8DT04006K>.
155. Morel, P.; Schaffer, P.; Valliant, J.F. The synthesis and X-ray crystallographic structure determination of 3-isocyano-1, 2-dicarba-closo-dodecaborane-Re (I) complexes. *J. Organomet. Chem.* **2003**, *668*, 25–30. [https://doi.org/10.1016/S0022-328X\(02\)02094-6](https://doi.org/10.1016/S0022-328X(02)02094-6).
156. Asay, M.; Kefalidis, C.E.; Estrada, J.; Weinberger, D.S.; Wright, J.; Moore, C.E.; Lavallo, V. Isolation of a Carborane-Fused Triazole Radical Anion. *Angew. Chem. Int. Ed.* **2013**, *52*, 11560–11563. <https://doi.org/10.1002/anie.201306764>.
157. Vinogradov, M.M.; Loginov, D.A.; Starikova, Z.A.; Petrovskii, P.V.; Holub, J.; Kudinov, A.R. Nonsymmetrical iron bis (carborane) complexes (η -1-Bu⁴NH-1, 7, 9-C₃B₈H₁₀) Fe (η -9-L-7, 8-C₂B₉H₁₀)(L = SMe₂, NMe₃). *Russ. Chem. Bull.* **2010**, *59*, 2143–2146. <https://doi.org/10.1007/s11172-010-0370-2>.
158. Press, L.P.; McCulloch, B.J.; Gu, W.; Chen, C.H.; Foxman, B.M.; Ozerov, O.V. Triflyloxy-substituted carboranes as useful weakly coordinating anions. *Chem. Commun.* **2015**, *51*, 14034–14037. <https://doi.org/10.1039/C5CC05047B>.
159. Jeans, R.J.; Chan, A.P.; Murrell, A.H.; Chouman, H.A.; Rosair, G.M.; Welch, A.J. Metalation of Bis (meta-carborane). *J. Organomet. Chem.* **2021**, *950*, 121980. <https://doi.org/10.1016/j.jorganchem.2021.121980>.