

Supplementary Materials for

**Crystal Structures of DNA Intercalating Agents
Dipyrido[3,2-f:2',3'-h]quinoxaline (dpq),
(Benzo[i]dipyrido[3,2-a:2',3'c]phenazine (dppn), and
[Ir(ppy)₂(dppn)][PF₆] (Where Hppy = 2-Phenylpyridine)**

	Page
Figure S1: ¹ H NMR of dipyrido[3,2-f:2',3'-h]quinoxaline (dpq) in CDCl ₃	S2
Figure S2: ¹ H NMR of benzo[i]dipyrido[3,2-a:2',3'c]phenazine (dppn) in CDCl ₃	S2
Figure S3: ¹ H NMR [Ir(ppy) ₂ (dppn)][PF ₆] in (CD ₃) ₂ CO	S3




Figure S1. ¹H NMR of dipyrido[3,2-f:2',3'-h]quinoxaline (dpq) in CDCl₃.




Figure S2. ¹H NMR of benzo[i]dipyrido[3,2-a:2',3'-c]phenazine (dppn) in CDCl₃.




Figure S3. ^1H NMR $[\text{Ir}(\text{ppy})_2(\text{dppn})][\text{PF}_6]$ in $(\text{CD}_3)_2\text{CO}$.