

Supplementary Materials

Characterization of dextran produced by the food-related strain *Weissella cibaria* C43-11 and of the relevant *dextranucrase* gene

Palmira De Bellis ^{1,*a}, Massimo Ferrara ^{1,a}, Anna Rita Bavaro ¹, Vito Linsalata ¹, Mariaelena Di Biase ¹, Biagia Musio², Vito Gallo ², Giuseppina Mulè ¹ and Francesca Valerio ¹

¹ Institute of Sciences of Food Production (ISPA), National Research Council (CNR), Via G. Amendola 122/O, 70126 Bari, Italy

² Dipartimento di Ingegneria Civile, Ambientale, del Territorio, Edile e di Chimica (DICATECh), Politecnico di Bari, via Orabona 4, I-70125, Bari, Italy

* Correspondence: mirella.debellis@ispa.cnr.it; Tel.: +39-080-5929459; Fax: +39-080-5929374

^a These authors contributed equally to this work.

Figure S1. Typical 1D ¹H NOESY NMR spectra (D₂O, 400 MHz, 303 K) of free mono- (arabinose, galactose, glucose, fructose, and mannose) and disaccharides (sucrose and maltose), and purified samples from C43-11+S and C2-32+S.




Figure S2. Typical 1D ^1H NOESY NMR spectra (D_2O , 400 MHz, 303 K) of free mono- (arabinose, galactose, glucose, fructose, and mannose) and disaccharides (sucrose and maltose), and purified samples from C43-11+S and C2-32+S.

