Sugarcane pulp lunch-boxes produce more microparticles in acidic foods

Yi Hu[a], Chunru Mo[b], Zhiwei Wang[a], Wenwen Yu[b*], Changying Hu[a*]

[a] Packaging Engineering Institute, Jinan University, Zhuhai, Guangdong, 519070, China.

[b] Department of Food Science & Engineering, Jinan University, Huangpu West Avenue 601, Guangzhou City, 510632, Guangdong, China.

Corresponding authors at:

Changying Hu, hucy0000@sina.com

Address: Packaging Engineering Institute, Jinan University, Zhuhai, Guangdong, 519070, China.

Wenwen Yu, yuwenwen@*jnu*.edu.cn

Address: Department of Food Science & Engineering, Jinan University, Huangpu West Avenue 601, Guangzhou City, 510632, Guangdong, China.

Supplementary Information

Prepared for Submission
To *Foods*

Item	Description	Page
Figure S1	The leaking phenomenon of lunchbox when contacting with the 95% EtOH food simulant.	S3
Figure S2	Infrared spectra of 15 SCP lunchboxes.	S4



Figure S1. The leaking phenomenon of lunchbox when contacting with the 95% EtOH food simulant.



Figure S2. Infrared spectra of 15 SCP lunchboxes.