foods

MDPI

Article

Inhibitory Mechanism of Lactic Acid Bacteria Cell-Free Supernatant Against Stored Grain Molds

Can Cui, Xiaopeng Fu, Tianjie Qi, Tianci Zou, Yijun Liu, Yanfei Li, Yan Zhao and Haoxin Lv *

School of Food and Strategic Reserves, Henan University of Technology, Zhengzhou 450001, China; cc13526618922@163.com (C.C.); 13698871691@163.com (X.F.); 15003798360@163.com (T.Q.); 13915338078@163.com (T.Z.); lyj3500@163.com (Y.L.); liyanfei@haut.edu.cn (Y.L.); zhaoyanss10@126.com (Y.Z.)
* Correspondence: lvhaoxin0129@126.com

Abstract

Grain is highly vulnerable to contamination by fungi during storage, leading to reduced product quality and substantial economic losses. Lactic acid bacteria (LAB) have the potential to be used as antifungal agents; however, strains aimed at inhibiting stored grain molds remain limited, and the inhibitory mechanisms require further investigation. To solve this problem, 71 LAB strains were isolated from various samples. Among these, 17 strains exhibiting inhibitory activity against *A. flavus*, *A. niger* and *P. citrinum* were selected using a dual-layer plate assay. Based on morphological characterization and *16S rRNA* gene sequence analysis, these strains were classified as *Weissella cibaria*, *Pediococcus pentosaceus*, and *Lactiplantibacillus paraplantarum*. Further investigations involving pH adjustment, catalase, and proteinase K treatments confirmed that organic acids were the primary antifungal substances in LAB cell-free supernatant (CFS). HPLC quantification identified acetic acid, malic acid, lactic acid, phenyllactic acid and citric acid contained in the CFS. Antifungal assays verified that acetic acid and lactic acid exhibited the strongest inhibitory effects against *P. citrinum* and *A. flavus*, whereas phenyllactic acid and acetic acid demonstrated the most potent suppression against *A. niger*. These findings established a theoretical basis for the application of LAB CFS in grain storage.

Keywords: grain storage; fungal contamination; lactic acid bacteria; inhibitory mechanism; organic acid

1. Introduction

Grain constitutes a fundamental cornerstone for maintaining social stability. It has been estimated that 25–35% of grains are lost after harvest annually, among which fungi infection accounts for 9–13% [1]. Stored grains are particularly susceptible to contamination by species of *Aspergillus*, *Fusarium* and *Penicillium* [2]. These molds not only degrade the quality of grains, but also synthesize mycotoxins, posing a significant threat to human and animal health [3]. According to FAO, around 25% of global grain production is contaminated with mycotoxin each year, reaching hundreds of billions of dollars worldwide [4]. Consequently, there is an urgent need to develop effective strategies to mitigate and prevent fungal contamination during grain storage.

To mitigate the impact of fungal contamination, diverse strategies have been developed. These encompass physical approaches such as mechanical ventilation, cold plasma technology, and electron beam irradiation [5–7]. However, physical techniques are limited

Check for updates

Academic Editor: Vania Patrone

Received: 22 January 2026
Revised: 3 February 2026
Accepted: 7 February 2026
Published: 11 February 2026

Copyright: © 2026 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license.

by high equipment costs, substantial energy consumption, and insufficient technical validation. Chemical inhibition methods, including hydrogen sulfide fumigation and solid fungicides for example benzoic acid, sorbic acid, and sodium diacetate, have also been employed [8,9]. However, these chemical agents may adversely affect the sensory characteristics and nutritional quality of grains, pose negative environmental impacts, and increase the risk of cardiovascular diseases with excessive intake [10,11].

In recent years, the application of lactic acid bacteria (LAB) and their bioactive metabolites for preventing fungi contamination has gradually attracted attention due to their potent antifungal activity, favorable safety profile, cost-effectiveness, and broad-spectrum applicability. Krishnan et al. demonstrated that the cell-free supernatant (CFS) from *Weissella confusa* strains ML2 and BF2 effectively inhibited the germination of *Fusarium verticillioides* NCIM 1100 conidia, thereby preventing early-stage infection in wheat [12]. CFS from *Lactiplantibacillus pentosus* CW5 has been shown to serve a dual purpose, it effectively delays the softening and decay of kiwifruit in storage and concurrently reduces the population of postharvest pathogenic fungi [13]. The peptide mixture within the CFS of *Lactobacillus plantarum* TE10 exhibits significant antifungal activity, effectively inhibiting the growth of *Aspergillus flavus* on fresh corn and extending its shelf life [14]. Similarly, Møller et al. reported that *Levilactobacillus* spp. 3QB398 and *Levilactobacillus brevis* 2QB422 can inhibit the infection by *Aspergillus parasiticus* [15]. In another study, Cao et al. identified a strain of *Lactobacillus plantarum* AR524 that showed strong antagonistic activity against *Fusarium graminearum* [16].

The antifungal activity of LAB has been attributed to the production of bioactive compounds such as organic acids, hydrogen peroxide and proteinaceous antimicrobials. Organic acids could lower the environmental pH, disrupt fungal cell membrane integrity, and inhibit spore germination and mycelial growth [17,18]. Additionally, hydrogen peroxide exerts antifungal effects by attacking cell membranes and inducing oxidative stress [19]. While proteinaceous antimicrobials were able to inhibit fungal cell wall synthesis and interfere with key metabolic enzymes [20,21]. Although these mechanisms are well-documented in high-moisture substrates such as fruits and dairy products [22,23], their efficacy and dominance in the low-water-activity, complex environment of stored grains remain unclear.

In this study, LAB strains with antifungal activity were isolated from various environmental samples. Then the selected LAB strains were characterized morphologically and identified molecularly based on 16S rDNA sequencing. The antifungal activity of the CFS was assessed, and the primary bioactive substances were identified. Furthermore, the underlying antifungal mechanism of the LAB CFS was investigated, thereby establishing a theoretical foundation for developing biological preservatives for grain storage.

2. Materials and Methods

2.1. Materials

The LAB strains were isolated from various samples including soil, corn kernel, pickled cabbage and wheat bran samples. Reference fungal strains *A. flavus* MA (accession number of 18S rRNA gene sequence: MZ557834), *A. niger* BQM (accession number of 18S rRNA gene sequence: MT903388) and *P. citrinum* MB (accession number of 18S rRNA gene sequence: MZ557835), previously isolated and preserved in the laboratory, were employed to screen LAB strains exhibiting antifungal activity. DeMan, Rogosa, and Sharpe (MRS) and Potato dextrose agar (PDA) media were sourced from Beijing Aobo Star Biotechnology Co., Ltd. (Beijing, China). PCR Mix (with dye) was purchased from Beijing Liuhe Biotechnology Co., Ltd. (Beijing, China). Tris-acetic acid electrophoresis buffer was obtained from Leagene Biotechnology Co., Ltd. (Beijing, China). Agarose was bought from Biowest (Loire valley, France). Catalase and proteinase K were procured from Beijing Solarbio Science & Tech-

nology Co., Ltd. (Beijing, China). Lactic acid, acetic acid, propionic acid, L-3-phenyl-lactic acid, citric acid and malic acid (all HPLC grade) were acquired from Sigma-Aldrich (St. Louis, MO, USA).

2.2. Screening of LAB Strains for Antifungal Activity

2.2.1. Preliminary Screening of LAB Strains for Antifungal Activity

A. flavus MA, *A. niger* BQM and *P. citrinum* MB were plated onto PDA plates and grown at a constant temperature of 28 °C for 3 days to allow abundant spore production. Spores were collected by rinsing the PDA plates with sterile distilled water, and the resulting suspension was homogenized using a vortex mixer. The spore concentration was adjusted to 1×10^5–5×10^5 CFU/mL and stored at 4 °C prior use [24].

Around 5 g of each sample (soil, corn kernels, fermented cabbage and wheat bran) was added into 45 mL of sterile saline. The mixture was vortexed at 180 rpm for one hour. The suspension was subjected to serial diluted into five gradients: 10^{-1}, 10^{-2}, 10^{-3}, 10^{-4} and 10^{-5} using sterile saline [25]. A 100 µL aliquot of the mold spore suspension was transferred into a sterile Petri dish, and then 20 mL of MRS agar medium cooled to 50 °C was added. After the agar being solidified, 100 µL of each sample dilution was spread onto the surface. A control plate was prepared using an equivalent volume of sterile saline instead of the sample dilution. Cultivation was carried out on all plates at 37 °C with a duration of 48 h [26]. Colonies exhibiting inhibition of mold growth were selected from each plate based on morphological characteristics such as color, glossiness, size and the capacity of growth under co-culture conditions [27].

2.2.2. Evaluation of the Antifungal Activity of LAB Strains

The modified dual-culture assay was applied to assess the antifungal activity of LAB strains against *A. flavus* MA, *A. niger* BQM and *P. citrinum* MB [28]. Briefly, 10 mL of MRS agar medium was dispensed into sterile Petri dishes and allowed to solidify. Following solidification, two parallel lines spaced 3 cm apart were inoculated with LAB strains on MRS plates, which were then incubated at 37 °C for 24 h. Then, 10 mL of PDA semi-solid medium supplemented with a mold spore suspension at a concentration of 1×10^5 to 5×10^5 CFU/mL was overlaid onto the plates. After solidification of the overlay medium, the plates were incubated at 28 °C for 48 h. Plates without LAB inoculation were used as the control group. All experiments were performed in triplicate. The diameter of inhibition zone perpendicular to the LAB inoculation lines was measured using a vernier caliper.

2.3. Identification of LAB Strains with Antifungal Activity

2.3.1. Morphological Identification of LAB Isolates

Physiological and biochemical identification of LAB strains with antifungal activity was conducted in accordance with the procedures outlined in Bergey's Manual of Systematic Bacteriology. The LAB strains were streaked onto MRS solid medium and incubated at 37 °C for 48 h. Colony morphology was observed and recorded, and single colonies were selected for Gram staining [29].

2.3.2. Molecular Identification of LAB Isolates

Genomic DNA of selected LAB strains was extracted using the DNeasy Blood and Tissue kit (QIAGEN). PCR amplification of the *16S rRNA* gene was performed with the universal primers set (Eu8f, 5′-AGAGTTTGATCCTGGCTCAG-3′; Eu1492r, 5′-GGCTACCTTGTTACGACTT-3′), each at a final concentration of 0.2 µM [30]. Following amplification, the PCR products were subjected to electrophoretic separation on a 2% (*w*/*w*) agarose gel at a constant voltage of 120 V. Upon completion of electrophoresis, purified PCR products were submitted to the Beijing Genomics Institute for Sanger sequencing of

the *16S rRNA* gene. The resulting sequences were submitted to the EzBioCloud database (https://www.ezbiocloud.net/ accessed on 5 June 2023) for similarity comparison with closely related type strains [31]. The MEGA 12.1 program was utilized for conducting phylogenetic and molecular evolutionary analysis [32].

2.4. Evaluation of the Antifungal Activity of LAB CFS

The LAB strains were grown in MRS broth at 37 °C with shaking 200 rpm for 24 h. When the OD_{600} reached approximately 1, a 5% inoculum was transferred into fresh MRS broth and incubated under identical conditions for an additional 48 h. The CFS was collected by centrifugation at 6791× *g* for 10 min at 4 °C and subsequently filtered through a 0.22 μm sterile membrane. Following pH measurement, the supernatant was stored at 4 °C for further analysis [33].

The antifungal activity of LAB CFS against *A. flavus* MA, *A. niger* BQM and *P. citrinum* MB was evaluated using a 96-well plate assay. A mixture of 190 μL LAB CFS and 10 μL fungal spore suspension (final concentration: 1×10^5 to 5×10^5 CFU/mL) was added to each well, while the control group received 190 μL sterile MRS broth and 10 μL mold spore suspension [34]. The 96-well plates were incubated at 28 °C for 72 h with three replicates per group. The optical density at 600 nm (OD_{600}) was measured using a microplate reader (Powerscan HT, DS Pharma Biomedical Co., Ltd., Osaka, Japan), and the inhibitory rate was calculated to assess the antifungal effect of LAB CFS.

$$Inhibitory\ rate(\%) = \left(1 - \frac{72\text{ h }OD_{LAB} - 0\text{ h }OD_{LAB}}{72\text{ h }OD_{control} - 0\text{ h }OD_{control}}\right) \times 100\%$$

where OD_{LAB} represents the OD_{600} value of the mold spore suspension cultured in LAB CFS at 0 h or 72 h, respectively; and $OD_{control}$ indicates the OD_{600} value of the mold spore suspension cultured in sterile MRS broth at 0 h or 72 h.

2.5. Effect of pH, Catalase and Proteinase K Treatments on the Antifungal Activity of LAB CFS

2.5.1. Effect of pH Treatments on the Antifungal Activity of LAB CFS

Following the protocol described by Zhang et al. [35,36], the LAB CFS was adjusted to pH 4.0, 5.0 and 6.0, respectively. Untreated LAB CFS and sterile MRS broth were used as controls. The inhibitory effect of pH-treated LAB CFS against *A. flavus* MA, *A. niger* BQM and *P. citrinum* MB was evaluated using the 96-well plate assay [37].

2.5.2. Effect of Catalase Treatment on the Antifungal Activity of LAB CFS

First, 3% (*w*/*v*) catalase was dissolved in 0.05 mol·L^{-1} phosphate buffer (pH 7.0). The enzyme solution was then added to LAB CFS to yield a final concentration of 5 mg/mL, the mixture was then incubated in a water bath at 37 °C for 2 h. Catalase activity in LAB CFS was subsequently inactivated by heating at 100 °C for 5 min. The pH of the LAB CFS was adjusted to pH 6.0, and the antifungal activity was assessed using the 96-well plate assay [38]. Sterile MRS broth and LAB CFS (pH 6.0) without catalase treatment were used as controls, with three replicates performed for each condition [21].

2.5.3. Effect of Protease K Treatment on the Antifungal Activity of LAB CFS

Protease K was dissolved in 0.05 mol·L^{-1} phosphate buffer (pH 7.0). The enzyme preparation was added to the LAB CFS to achieve a final concentration of 1 mg/mL and then incubated at 37 °C in a water bath for 2 h. Protease K activity in the LAB CFS was subsequently inactivated by heating at 100 °C for 5 min. After treatment, the pH was adjusted to 6.0, and the antifungal activity was evaluated using the 96-well plate assay [39].

Sterile MRS broth and LAB CFS (pH 6.0) without protease K treatment were used as controls, with three replicates performed for each condition [40].

2.6. Analysis of Organic Acids in LAB CFS

The organic acids in LAB CFS were analyzed using HPLC (Waters, e2695, Milford, CT, USA) [41]. The detection wavelength was set at 210 nm. Chromatographic separation was achieved using an Agilent ZORBAX SB-C18 column (25 cm × 4.6 mm, 5 μm). The mobile phase consisted of 0.01 mol/L KH_2PO_4 (pH 2.7) and methanol at a ratio of 95:5 (*v*/*v*). The flow rate was 0.6 mL/min, and the column temperature was kept at 30 °C. The injection volume was 20 μL, and the total elution time was 20 min.

Standard solutions of each organic acid—phenyllactic acid, lactic acid, acetic acid, citric acid, malic acid and propionic acid—were prepared individually at 20 mg/mL by solubilizing the respective pure compounds in ultrapure water. Each standard solution was injected separately into the system to determine its retention time. A mixed organic acid standard solution was then prepared by combining the six individual standard solutions with 0.1% (*w*/*w*) phosphoric acid solution. In the mixed solution, standard solutions of phenyllactic acid, lactic acid, malic acid and citric acid were prepared at concentrations of 0.1, 0.5, 1, 2, 5 mg/mL, respectively, while those of acetic acid and propionic acid were 0.5, 1, 2, 5, 10 mg/mL, respectively. The mixed standard solution was filtered through a 0.22 μm sterile filter membrane into an injection vial for sample introduction and analysis. Calibration curves were constructed by plotting the peak area (y-axis) against the corresponding organic acid concentration (x-axis). The instrument-generated data were processed to construct calibration curves and determine the correlation coefficients.

LAB CFS was prepared based on the method described in [42]. In order to ensure that the organic acid concentrations fell within the detection range of the calibration curve established from the standard solutions, the LAB CFS was diluted 5 times with 0.1% (*w*/*w*) phosphoric acid solution and filtered through a 0.22 μm sterile membrane prior to instrumental analysis, then transferred into an injection vial.

2.7. Evaluation of the Antifungal Activity of Organic Acids

According to the results of organic acid analysis in LAB CFS, the antifungal activity of five organic acids—lactic acid, acetic acid, citric acid, malic acid and phenyllactic acid—was evaluated. The concentrations of the individual organic acid solutions were adjusted to 0.5, 1, 5, 10 and 20 mg/mL using sterile MRS liquid medium. After filtration through a 0.22 μm sterile membrane filter, the antifungal activity of each solution against *A. flavus* MA, *A. niger* BQM and *P. citrinum* MB was assessed using the 96-well plate method [43]. The experiment was performed in triplicate with sterile MRS liquid medium serving as the control.

2.8. Statistical Analysis

Data were primarily organized and descriptive statistics were calculated using Microsoft Excel 2019. Statistical analysis was performed using SPSS 23. Significant differences among groups were evaluated by Duncan's multiple range test at a threshold of $p < 0.05$. Figures were plotted and prepared using Origin 2018 and final compositing was done with Adobe Photoshop CS6 for improved presentation clarity. All experiments were conducted in triplicate, and data are expressed as mean ± standard deviation.

3. Results and Discussion

3.1. Screening of LAB Strains with Antifungal Activity

3.1.1. Preliminary Screening of LAB Strains with Antifungal Activity

Figure 1 illustrates that the control group's MRS agar plate presented a uniform lawn of mold colonies (Figure 1a), whereas several smooth, round and white colonies—presumed to be LAB with antifungal potential—appeared on the MRS agar plate of the experiment group (Figure 1b). According to distinct morphological characteristics such as color, size, shape and morphology, a collection of 71 strains was obtained from diverse materials, including soil, corn kernel, fermented cabbage and wheat bran samples (Table 1). The isolation strategy of this study leverages the broad natural distribution of LAB [44]. This integrated niche-sampling approach serves a dual purpose: LAB isolated directly from grain substrates may exhibit enhanced environmental adaptability for direct application in storage ecosystems, while those from highly competitive environments like soil and fermented pickles are more likely to yield robust strains, naturally selected for strong acid production or novel antifungal mechanisms [45]. The considerable biodiversity of LAB underpins their significant application potential [46]. Therefore, this study aims to systematically screen strains from these distinct sources and elucidate their inhibitory mechanisms against key storage molds, thereby establishing a theoretical and resource foundation for developing targeted, efficient bioprotective agents.




Figure 1. Screening of LAB strains with antifungal potential. (**a**) control group, mold spore suspension with sterile saline; (**b**) experiment group, mold spore suspension with sample diluent.

Table 1. LAB strains isolated from various experimental sources.

Experimental Sources	Number of Strains	LAB Strains
Rhizosphere soil of wheat	7	TA1, TA2, TA3, TA4, TA5, TA6, TA7
Rhizosphere soil of vegetables	6	TB1, TB2, TB3, TB4, TB5, TB6
Rhizosphere soil of garlic	11	TC1, TC2, TC3, TC4, TC5, TC6, TC7, TC8, TC9, TC10, TC11
Soil from flower beds	13	TD1, TD2, TD3, TD4, TD5, TD6, TD7, TD8, TD9, TD10, TD11, TD12, TD13
Corn kernel from Northeast China, variety No. 3	4	DA1, DA2, DA3, DA4
Corn kernels from Heihe city, Heilongjiang province, China	12	DB1, DB2, DB3, DB4, DB5, DB6, DB7, DB8, DB9, DB10, DB11, DB12
Corn kernel from Northeast China, variety No. 6	8	DC1, DC2, DC3, DC4, DC5, DC6, DC7, DC8
Fermented cabbage	6	P1, P2, P3, P4, P5, P6
Wheat bran	4	F1, F2, F3, F4

3.1.2. Evaluation of Antifungal Activity of LAB Strains

According to the inhibition zone diameter [47], the antifungal activity of LAB strains was divided into four levels: no inhibitory effect (−), defined as the absence of an inhibition zone; low inhibitory effect (+), with a diameter less than 20 mm; moderate inhibitory effect (++), ranging from greater than 20 mm to less than 30 mm; high inhibitory effect (+++), higher than 30 mm (Figure S1).

A modified dual-culture assay was used to screen the inhibitory potential of the 71 LAB strains against the three mold species: *A. flavus* MA, *A. niger* BQM, and *P. citrinum* MB. The results showed that 17 LAB strains demonstrated inhibitory effects against all three pathogenic molds (Table 2). As shown in Table 2, 11 LAB strains inhibited the growth of *A. flavus* MA, with strain P3 demonstrating the strongest inhibitory effect and yielding an inhibition zone diameter exceeding 30 mm. Moreover, 14 LAB strains showed inhibitory effect on *A. niger* BQM, among which, strain P3 had the highest inhibitory effect. In addition, 17 strains inhibited *P. citrinum* MB, with 12 strains, including DA1, DA2, DA4, DB2, DB5, DB6, DB11, DC5, DC6, P3, P5, P6, had strong inhibitory effect, and strain DC5 produced the largest inhibition zone. Similarly, Zhang et al. reported that *Lactococcus lactis* PLL15 and *Lactobacillus sakei* PLL12 could significantly inhibit the germination of *A. niger* spores and disrupt the integrity of the cell membrane and cell wall [48]. Concurrently, Kim et al. found that *Lactiplantibacillus plantarum* G2 and *Levilactobacillus brevis* H8 efficiently suppressed the growth of *Aspergillus* species in ham [49].

Table 2. The antifungal activity of LAB strains against *A. flavus* MA, *A. niger* BQM and *P. citrinum* MB.

LAB Strains	*A. flavus* MA		*A. niger* BQM		*Penicillium citrinum* MB	
	Inhibition Zone Diameter Value (mm)	Antifungal Activity	Inhibition Zone Diameter Value (mm)	Antifungal Activity	Inhibition Zone Diameter Value (mm)	Antifungal Activity
DA1	0	−	0	−	30.86	+++
DA2	0	−	30.91	+++	38.11	+++
DA4	0	−	28.17	++	32.23	+++
DB1	15.22	+	0	−	29.39	++
DB2	12.23	+	22.12	++	30.37	+++
DB3	13.86	+	21.53	++	27.31	++
DB5	0	−	39.43	+++	31.68	+++
DB6	14.55	+	26.65	++	38.04	+++
DB7	13.37	+	29.87	++	29.92	++
DB8	0	−	0	−	29.81	++
DB11	13.75	+	37.87	+++	30.55	+++
DB12	19.52	+	35.81	+++	29.01	++
DB5	18.75	+	23.05	++	46.53	+++
DB6	19.39	+	25.07	++	30.53	+++
P3	31.85	+++	40.63	+++	42.37	+++
P5	21.41	++	29.38	++	38.41	+++
P6	0	−	26.62	++	31.69	+++

Note: −, no inhibitory effect; +, low inhibitory effect; ++, moderate inhibitory effect; +++, high inhibitory effect.

3.2. Analysis of LAB Strains with Antifungal Activity

3.2.1. Morphological Identification

Table 3 summarizes the key morphological traits of the antifungal LAB strains. Colonies were circular, smooth, white or cream-white, and either semi-transparent or non-transparent and all isolates were Gram-stain-positive. These characteristics were consistent with the established morphological profile of typical LAB strains. As a defined group of Gram-positive, non-sporulating bacteria, they are usually in the form of cocci or rods [50].

Table 3. Morphological characteristics of LAB strains with antifungal activity.

LAB Strains	Morphology	Color	Transparency	Gram-Stain
DA1	Circular, smooth	White	Semitransparent	+
DA2	Circular, smooth	White	Semitransparent	+
DA4	Circular, smooth	White	Semitransparent	+
DB1	Circular, smooth	Cream-white	Non-transparent	+
DB2	Circular, smooth	Cream-white	Non-transparent	+
DB3	Circular, smooth	Cream-white	Semitransparent	+
DB5	Circular, smooth	Cream-white	Non-transparent	+
DB6	Circular, smooth	Cream-white	Non-transparent	+
DB7	Circular, smooth	Cream-white	Non-transparent	+
DB8	Circular, smooth	Cream-white	Non-transparent	+
DB11	Circular, smooth	Cream-white	Non-transparent	+
DB12	Circular, smooth	Cream-white	Non-transparent	+
DC5	Circular, smooth	White	Semitransparent	+
DC6	Circular, smooth	White	Semitransparent	+
P3	Circular, smooth	Cream-white	Non-transparent	+
P5	Circular, smooth	Cream-white	Non-transparent	+
P6	Circular, smooth	Cream-white	Non-transparent	+

Note: +, positive.

3.2.2. Molecular Identification

Table 4 presents the results of molecular identification for the LAB strains using 16S rDNA sequencing. It is indicated that 17 LAB strains were divided into 3 genera and 3 species. The 16S rDNA sequences of strains DA1, DA2, DA4, DB3, DC5 and DC6 were most closely related to that of the reference strain *Weissella cibaria* KACC 11862 ^{T}, with sequence identity exceeding 99%. Based on both molecular and morphological characteristics, these six strains were identified as *Weissella cibaria*. Strains DB1, DB2, DB5, DB6, DB7, DB8, DB11, DB12 and P6 showed the closest phylogenetic relationship to *Pediococcus pentosaceus* DSM 20336 ^{T}, with sequence similarities also higher than 99%. Integrating molecular data with morphological features, these nine strains were identified as *Pediococcus pentosaceus*. The 16S rDNA sequences of strains P3 and P5 were closest to that of *Lactiplantibacillus paraplantarum* DSM 10667 ^{T}, with sequence similarities of 99.59% and 99.25%, respectively. According to their morphological characteristics, both strains were identified as *Lactiplantibacillus paraplantarum*. Vasundaradevi et al. demonstrated that organic acids produced by *Lactobacillus paraplantarum* significantly suppress the growth of uropathogens [51]. Furthermore, Wang et al. demonstrated that applying the CFS of *Weissella cibaria* effectively delays quality deterioration in sweet corn [52]. Separately, *Pediococcus pentosaceus* has also been employed in litchi postharvest preservation to mitigate postharvest losses [53]. Collectively, these studies confirm that *Weissella cibaria*, *Pediococcus pentosaceus*, and *Lactiplantibacillus paraplantarum* exhibit antifungal properties. However, their potential in controlling major molds specifically associated with grain spoilage during storage remains underexplored. Therefore, this study aimed to elucidate the antifungal mechanisms of these LAB strains against key storage-borne pathogenic fungi.

3.3. Analysis of the Antifungal Activity of LAB CFS

Table 5 presents a comparative summary of the antifungal activity exhibited by the CFS of 17 LAB strains against *A. flavus* MA, *A. niger* BQM and *P. citrinum* MB. Among these strains, the CFS of strains DA2, DA4, DB2, DB11, DB12, DC5, P3 and P5 exhibited inhibitory rates exceeding 99%. These results align with precious report that *Lactobacillus plantarum* strains were used as biocontrol agents against *A. niger*, *A. flavus*, *P. chrysogenum* in wheat and fresh maize seeds; *Pediococcus pentosaceous* strain was utilized to control the

growth of *P. expansum* in pear, grape and plum [54]; and *W. cibaria* strains have previously been employed to prevent the growth of foodborne pathogens [55]. Moreover, molecular identification analysis revealed that these eight high-performing strains belong to three distinct genera and three species, highlighting their taxonomic diversity and providing a sound rationale for selecting them for mechanistic investigation.

Table 4. Molecular identification of LAB strains based on 16S rDNA sequencing.

LAB Strains	Reference Strains	16S rDNA Sequence Identity/%
DA1	*W. cibaria* KACC 11862 ^{T}	99.19
DA2	*W. cibaria* KACC 11862 ^{T}	100.00
DA4	*W. cibaria* KACC 11862 ^{T}	99.85
DB3	*W. cibaria* KACC 11862 ^{T}	99.26
DC5	*W. cibaria* KACC 11862 ^{T}	99.92
DC6	*W. cibaria* KACC 11862 ^{T}	100.00
DB1	*P. pentosaceus* DSM 20336 ^{T}	99.26
DB2	*P. pentosaceus* DSM 20336 ^{T}	99.39
DB5	*P. pentosaceus* DSM 20336 ^{T}	99.12
DB6	*P. pentosaceus* DSM 20336 ^{T}	99.32
DB7	*P. pentosaceus* DSM 20336 ^{T}	99.53
DB8	*P. pentosaceus* DSM 20336 ^{T}	100.00
DB11	*P. pentosaceus* DSM 20336 ^{T}	99.33
DB12	*P. pentosaceus* DSM 20336 ^{T}	100.00
P6	*P. pentosaceus* DSM 20336 ^{T}	99.39
P3	*L. paraplantarum* DSM 10667 ^{T}	99.59
P5	*L. paraplantarum* DSM 10667 ^{T}	99.25

Note: *W. cibaria*, *Weissella cibaria*; *P. pentosaceus*, *Pediococcus pentosaceus*; *L. paraplantarum*, *Lactiplantibacillus paraplantarum*. T: Type strain.

Table 5. Antifungal activity of LAB CFS against *A. flavus* MA, *A. niger* BQM and *P. citrinum* MB.

LAB Strains	Inhibitory Rate/%		
	***A. flavus* MA**	***A. niger* BQM**	***P. citrinum* MB**
DA1	95.08	99.05	95.96
DA2	99.29	99.35	99.53
DA4	99.13	99.50	99.55
DB1	98.22	97.88	91.41
DB2	99.49	99.11	99.08
DB3	97.52	99.23	91.41
DB5	97.69	95.23	93.03
DB6	97.59	98.54	92.22
DB7	98.89	98.90	98.78
DB8	98.58	98.63	98.30
DB11	99.20	99.12	99.88
DB12	99.05	99.41	99.36
DC5	98.94	99.36	99.65
DC6	98.09	95.16	98.88
P3	99.43	99.88	99.81
P5	99.31	99.50	99.10
P6	97.94	97.64	97.75

3.4. Analysis of the Effect of pH, Catalase and Proteinase K Treatments on the Antifungal Activity of LAB CFS

3.4.1. Analysis of the Effect of pH Treatments on the Antifungal Activity of LAB CFS

The inhibitory activity of the CFS from 8 LAB strains at different pH values against *A. flavus* MA, *A. niger* BQM and *P. citrinum* MB is displayed in Figure 2. A marked dependence on pH was observed for the inhibitory activity of the LAB CFS. At pH 4, the inhibitory rates against all three molds remained comparable to those of the control, with no significant difference ($p > 0.05$), maintaining at approximately 99%. However, when pH value was increased to 5 and 6, the inhibitory rate of LAB CFS was markedly reduced ($p < 0.05$). At pH 6, the inhibitory rate ranged from 12.42% to 39.28%, indicating a substantial decrease ($p < 0.05$). Similar results were reported by Szczerbiec et al., who demonstrated that *Lactobacillus* spp. strains exhibited significant inhibitory activity against *Proteus mirabilis*; however, the antifungal activity of the CFS was entirely eliminated at pH 6.0 [56]. In an acidic environment, organic acids could disrupt transmembrane proton gradients and inhibit essential metabolic enzymes, causing cell death or growth arrest; when the environmental pH rises, organic acids cannot easily cross the lipid bilayer, thereby losing their antifungal efficacy [57]. Additionally, LAB CFS may contain other metabolites which possess antifungal activity or exhibit synergistic effects with organic acids.




Figure 2. Inhibitory rate of LAB CFS with different pH values against *A. flavus* MA (**a**), *A. niger* BQM (**b**) and *P. citrinum* MB (**c**). Different small letters denote significant differences ($p < 0.05$) within the same LAB strain across processing methods.

3.4.2. Analysis of the Effect of Catalase Treatment on the Antifungal Activity of LAB CFS

Figure 3 illustrates the inhibitory activity against *A. flavus* MA, *A. niger* BQM, and *P. citrinum* MB exhibited by LAB CFS following catalase treatment. Compared to untreated LAB CFS, no significant change in inhibitory activity against the three molds was found following catalase treatment ($p > 0.05$), indicating that hydrogen peroxide was not secreted by the eight LAB strains. The study by Ma et al. also indicated that the CFS from *Pediococcus acidilactici* JY03, *Weissella paramesenteroides* JT13 and *Lactobacillus sucicola* JT03 remained antifungal activity following catalase treatment, thus indicating that hydrogen peroxide was not included among its antifungal substances [58].




Figure 3. Inhibitory rate of LAB CFS treated with catalase against *A. flavus* MA (**a**), *A. niger* BQM (**b**) and *P. citrinum* MB (**c**).

3.4.3. Analysis of the Effect of Protease K Treatment on the Antifungal Activity of LAB CFS

The inhibitory activity of LAB CFS treated with protease K against *A. flavus* MA, *A. niger* BQM and *P. citrinum* MB is demonstrated in Figure 4. Protease K treatment of the LAB CFS showed no significant difference in inhibitory rate relative to the untreated control against all three molds ($p > 0.05$). Therefore, it was presumed that the antifungal substances in LAB CFS are not proteinaceous. Similarly, it has been demonstrated that the antifungal activity of the CFS from *Lactiplantibacillus plantarum* G2 showed no significant change after protease K treatment, confirming that the antifungal substances in this CFS are not proteins [49].




Figure 4. Inhibitory rate of LAB CFS treated with protease K against *A. flavus* MA (**a**), *A. niger* BQM (**b**) and *P. citrinum* MB (**c**).

3.5. Analysis of Organic Acid Composition of LAB CFS

3.5.1. HPLC Analysis of Organic Acids

Previous studies have demonstrated that organic acids including phenyllactic acid, lactic acid, acetic acid, citric acid, malic acid and propionic acid were produced by LAB [26,59]. Therefore, HPLC calibration curves (Figure 5) for these six kinds of organic acid standards were constructed to investigate the content of each organic acid in LAB CFS. As can be seen from Figure 5, the retention times of the organic acid standards were as follows: phenyllactic acid at 5.075 min, lactic acid at 6.158 min, acetic acid at 6.766 min, citric acid at 7.133 min, malic acid at 8.081 min and propionic acid at 15.183 min. The calibration curves and correlation coefficient (R^2) of the organic acid standards are summarized in Table 6. The results demonstrate that the R^2 values ranged from 0.9981 to 0.9999, confirming an excellent linear correlation between analyte concentration and chromatographic response. Under the selected chromatographic conditions, all six organic acids exhibited well-defined peaks and were effectively separated. Therefore, the developed calibration curves are suitable for the quantitative determination of organic acids in LAB CFS.




Figure 5. HPLC calibration curve of organic acid standards: (1) phenyllactic acid; (2) lactic acid; (3) acetic acid; (4) citric acid; (5) malic acid; (6) propionic acid.

Table 6. Regression equation and correlation coefficient of organic acid standards.

Organic Acid Standards	Retention Time (min)	Regression Equation	Correlation Coefficient (R^2)
Phenyllactic acid	5.075	$y = 1 \times 10^6x + 199{,}099$	0.9987
Lactic acid	6.158	$y = 2 \times 10^6x + 22{,}119$	0.9999
Acetic acid	6.766	$y = 969{,}221x + 27{,}340$	0.9999
Citric acid	7.133	$y = 2 \times 10^6x + 19{,}146$	0.9999
Malic acid	8.081	$y = 948{,}650x + 3952.9$	0.9981
Propionic acid	15.183	$y = 1 \times 10^6x + 67{,}504$	0.9999

3.5.2. Quantitative Determination of Organic Acid Content in LAB CFS

As can be seen from Table 7, five organic acids including phenyllactic acid, lactic acid, acetic acid, citric acid and malic acid were detected in the CFS of DA2, DA4, DB2, DB11, DB12, DC5 and P3, while only four organic acids such as phenyllactic acid, lactic acid, acetic acid and malic acid were found in P5 CFS. Among the CFS of the eight selected LAB strains, acetic acid and lactic acid were the main organic acids, followed by malic acid, citric acid and phenyllactic acid. *L. paraplantarum* P3 and P5 secreted the highest concentrations of lactic acid, with corresponding levels of 17.84 mg/mL and 17.82 mg/mL, respectively. The highest acetic acid contents were observed in the CFS of *W. cibaria* DC5 and DA4, at 11.73 mg/mL and 10.33 mg/mL, respectively. According to the study by Garnier et al., the primary organic acids synthesized by *Lactobacillus* CIRM-BIA1952 were identified as lactic acid and acetic acid [60]. Similarly, a study by Vasundaradevi et al. quantitatively analyzed the organic acid profile in the CFS of *Lactiplantibacillus plantarum* MYSVCF3, revealing citric acid as the predominant metabolite, which also demonstrated the strongest inhibitory efficacy against *Fusarium oxysporum* [51]. To identify the key antifungal contributor, a further investigation was conducted to evaluate the individual activity of each identified organic acid against the target molds.

Table 7. Organic acid content in LAB CFS.

LAB Strains	Organic Acid Content (mg/mL)					
	Phenyllactic Acid	**Lactic Acid**	**Acetic Acid**	**Citric Acid**	**Malic Acid**	**Propionic Acid**
Weissella cibaria DA2	1.70 ± 0.23 ^{d}	13.53 ± 0.13 ^{b}	8.69 ± 0.45 ^{bc}	1.00 ± 0.06 ^{bc}	5.81 ± 0.22 ^{b}	ND
Weissella cibaria DA4	0.91 ± 0.04 ^{e}	13.51 ± 0.08 ^{b}	10.33 ± 0.12 ^{ab}	4.32 ± 0.05 ^{a}	5.61 ± 0.27 ^{b}	ND
Pediococcus pentosaceus DB2	0.31 ± 0.17 ^{f}	17.65 ± 0.71 ^{a}	7.52 ± 0.91 ^{c}	0.97 ± 0.01 ^{c}	2.32 ± 0.85 ^{c}	ND
Pediococcus pentosaceus DB11	0.55 ± 0.02 ^{f}	12.14 ± 0.11 ^{b}	8.64 ± 0.13 ^{bc}	1.04 ± 0.02 ^{b}	6.89 ± 0.35 ^{ab}	ND
Pediococcus pentosaceus DB12	2.44 ± 0.16 ^{c}	17.36 ± 0.03 ^{a}	7.76 ± 0.06 ^{c}	1.04 ± 0.01 ^{b}	2.61 ± 0.04 ^{c}	ND
Weissella cibaria DC5	0.86 ± 0.03 ^{e}	16.26 ± 2.3 ^{a}	11.73 ± 2.73 ^{a}	0.98 ± 0.02 ^{c}	8.10 ± 1.91 ^{a}	ND
Lactiplantibacillus paraplantarum P3	4.51 ± 0.35 ^{a}	17.84 ± 0.08 ^{a}	8.93 ± 0.02 ^{bc}	0.97 ± 0.01 ^{c}	2.63 ± 0.05 ^{c}	ND
Lactiplantibacillus paraplantarum P5	3.92 ± 0.08 ^{b}	17.82 ± 0.03 ^{a}	8.48 ± 0.06 ^{bc}	ND	2.44 ± 0.06 ^{c}	ND

Note: Data are presented as mean ± standard deviation from three independent experiments. Significant differences ($p < 0.05$) within the same row are denoted by different lowercase letters (a–f). ND, not detected.

3.6. Analysis of the Antifungal Activity of Organic Acids

The inhibitory activities of five organic acids were evaluated against *A. flavus* MA, *A. niger* BQM and *P. citrinum* MB at concentrations of 0.5, 1, 5, 10 and 20 mg/mL. As shown in Figure 6, acetic acid and lactic acid exerted higher efficacy against *A. flavus* MA and *P. citrinum* MB than the other three organic acids ($p < 0.05$). While acetic acid and phenyllactic acid displayed the highest inhibitory efficacy against *A. niger* BQM ($p < 0.05$). At a concentration of 20 mg/mL, the inhibitory rate of all five organic acids against the three molds reached approximately 98%~99%. It was indicated that acetic acid and lactic acid may be the primary antifungal components in CFS against storage molds. Lactic and acetic acids, as small monobasic acids, penetrate the fungal membrane due to their lipid solubility, causing intracellular acidification and metabolic disruption that ultimately inhibits growth. This mechanism is consistent with studies confirming hyphal damage and loss of membrane integrity [61,62]. Concurrently, phenyllactic acid demonstrated notable specificity against *A. niger*. Beyond acidification, evidence suggests it may interfere with specific fungal enzyme systems and signal transduction pathways [63]. Importantly, at a concentration of 10 mg/mL, the inhibitory rate of any single organic acid remained lower than that observed for the CFS of strains such as *Pediococcus pentosaceus* DB11, *Weissella cibaria* DA2, *Weissella cibaria* DA4. It was inferred that the remarkable antifungal activity of the CFS is attributable not to a single compound, but to a synergy between various organic acids. These compounds presumably act on complementary fungal targets, producing a stronger and broader-spectrum inhibitory effect than any individual acid [34,64]. Based on these findings, a conceptual framework for a multi-target biocontrol strategy can be proposed. This antifungal agent employs acetic and lactic acids as a broad-spectrum bases, supplemented with phenyllactic acid for the specific targeting of *A. niger.* The efficacy and practicality of this approach warrant further validation through simulated grain models or pilot-scale storage trials.




Figure 6. Antifungal activity of organic acids against *A. flavus* MA (**a**), *A. niger* BQM (**b**) and *P. citrinum* MB (**c**).

4. Conclusions

In this study, a total of 71 LAB strains originated from distinct materials such as soil, corn kernels, fermented cabbage and wheat bran. Among these, 17 LAB strains demonstrated antifungal activity against three harmful molds such as *A. flavus* MA, *A. niger* BQM and *P. citrinum* MB. They were classified as three genera and species, *W. cibaria*, *P. pentosaceus* and *L. paraplantarum*, based on morphological and molecular identification. Of the 17 LAB strains, the CFS of 8 LAB strains inhibited the growth of all three molds by over 99%. Their antifungal activity was significantly reduced following pH treatment, while it was not affected by catalase or proteinase K treatment, indicating that the antifungal substances in LAB CFS were mainly organic acids. In addition, HPLC analysis revealed that the LAB CFS consists of five kinds of organic acids: malic acid, acetic acid, lactic acid, phenyllactic acid and citric acid. Among them, acetic acid and lactic acid showed strongest antifungal activity against *A. flavus* MA and *P. citrinum* MB, while acetic acid and phenyllactic acid demonstrated the highest inhibitory efficacy against *A. niger* BQM. This research reveals the antifungal mechanism of LAB strains against dominant fungal strains, providing a theoretical foundation for their application in mold prevention and postharvest grain quality preservation. Further research is planned to assess the application of the LAB CFS in grain storage, thereby laying the groundwork for its development as a practical antifungal agent.

Supplementary Materials: The following supporting information can be downloaded at https://www.mdpi.com/article/10.3390/foods15040655/s1. Figure S1: Evaluation of the antifungal activity of LAB strains. (a) no inhibitory effect; (b) low inhibitory effect; (c) moderate inhibitory effect; (d) high inhibitory effect.

Author Contributions: Conceptualization, H.L.; Methodology, C.C.; Software, X.F. and T.Z.; Validation, T.Q. and Y.L. (Yijun Liu); Formal analysis, C.C.; Investigation, X.F. and T.Z.; Resources, Y.Z.; Data curation, T.Q. and Y.L. (Yijun Liu); Writing—original draft preparation, C.C.; Writing—review and editing, C.C. and H.L.; Visualization, C.C. and H.L.; Supervision, Y.L. (Yanfei Li); Project administration, Y.L. (Yanfei Li); Funding acquisition, H.L. and Y.Z. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by Department of Science and Technology of Henan Province, grant number 242103810072; China Scholarship Council, grant number 202508410212; Henan University of Technology, grant number 21421239; the Ministry of Science and Technology of the People's Republic of China, grant number 2021YFD2100604-2.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The original contributions presented in the study are included within the article; further inquiries can be directed to the corresponding author.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Ercan, G.; Adesemoye, A.O.; Yuen, G.Y.; Everhart, S.; Campbell, J.F.; Peterson, J.A. In vitro and in planta testing of microbial agents for dual biological control of granary weevil and storage fungi on stored wheat grain. *Biol. Control* **2025**, *207*, 105812. [CrossRef]
2. Atnafu, B.; Chala, A.; Lemessa, F.; Bayissa, W.; Mohammed, A.; Gerbaba, C.A. Effectiveness of storage facilities in managing toxigenic fungi and mycotoxins associated with stored maize (*Zea mays* L.) in Ethiopia. *J. Stored Prod. Res.* **2025**, *111*, 102561. [CrossRef]
3. Pinton, P.; Terciolo, C.; Payros, D.; Oswald, I.P. Mycotoxins hazard: The European view. *Curr. Opin. Food Sci.* **2025**, *63*, 101306. [CrossRef]
4. Eskola, M.; Kos, G.; Elliott, C.T.; Hajšlová, J.; Mayar, S.; Krska, R. Worldwide contamination of food-crops with mycotoxins: Validity of the widely cited 'FAO estimate' of 25%. *Crit. Rev. Food Sci. Nutr.* **2020**, *60*, 2773–2789. [CrossRef]
5. Zhang, B.; Guo, X.; Yao, Z.; Li, J.; Zhang, M.; Xu, X. Numerical simulation and energy consumption analysis of ventilation patterns in grain silo. *J. Stored Prod. Res* **2024**, *109*, 102469. [CrossRef]
6. Zhao, Y.; Zhang, M.; Liu, Y.; Liu, Z.; Li, X.; Jia, F. A transcriptomic investigation into the intrinsic and extrinsic effects of cold plasma generated from air and nitrogen on *Pseudomonas aeruginosa*. *Food Res. Int.* **2025**, *205*, 115974. [CrossRef]
7. Sui, X.; Lu, J.; Gao, S.; Chen, Z.; Liu, H.; Li, Y.; Astruc, D.; Liu, X. Electron beam irradiation technology: A new frontier in environmental pollution control. *Water Res.* **2025**, *286*, 124240. [CrossRef]
8. Dijksterhuis, J.; Kleinhoven, P.; van Kuijk, S.; Wolters, A.H.; Bruinenberg, P.G. Synergistic antifungal effects of the preservative ammonium propionate and medium chain fatty acids against dormant and germinating conidia, germ tubes and hyphae of *Aspergillus chevalieri*, a feed spoilage fungus. *Int. J. Food Microbiol.* **2024**, *422*, 110802. [CrossRef]
9. Zhu, X.; Hui, S.; Huang, H.; Liu, R.; Wang, S.; Huang, C. Antimicrobial mechanism of chlorine dioxide and its impacts on postharvest management in horticultural produce: A review. *Postharvest Biol. Technol.* **2024**, *213*, 112921. [CrossRef]
10. Ismail, I.E.; Farag, M.R.; Alagawany, M.; Mahmoud, H.K.; Reda, F.M. Efficacy of some feed additives to attenuate the hepato-renal damage induced by aflatoxin B1 in rabbits. *J. Anim. Physiol. Anim. Nutr.* **2020**, *104*, 1343–1350. [CrossRef]
11. Dong, H.; Xu, Y.; Zhang, Q.; Li, H.; Chen, L. Activity and safety evaluation of natural preservatives. *Food Res. Int.* **2024**, *190*, 114548. [CrossRef]
12. Krishnan, S.V.; Anaswara, P.A.; Nampoothiri, K.M.; Kovács, S.; Adácsi, C.; Miklós, I.; Király, S.; Pócsi, I.; Pusztahelyi, T. Unveiling the Perspective on *Weissella confusa* as a Promising Biocontrol Agent Against Fusaria. *Microorganisms* **2025**, *13*, 666. [CrossRef]
13. Hao, Y.; Pang, H.; Lv, H.; Wu, G.; Zhang, J.; Luo, X.; Zhang, M.; Duan, Y.; Hou, Y.; Wang, Y.; et al. Treatment of kiwifruit with *Lactiplantibacillus pentosus* CW5 cell-free supernatant maintains postharvest quality and alters the surface microbiome. *Postharvest Biol. Technol.* **2025**, *224*, 113478. [CrossRef]

14. Muhialdin, B.J.; Algboory, H.L.; Kadum, H.; Mohammed, N.K.; Saari, N.; Hassan, Z.; Hussin, A.S.M. Antifungal activity determination for the peptides generated by *Lactobacillus plantarum* TE10 against *Aspergillus flavus* in maize seeds. *Food Control* **2020**, *109*, 106898. [CrossRef]
15. Møller, C.O.D.A.; Freire, L.; Rosim, R.E.; Margalho, L.P.; Balthazar, C.F.; Franco, L.T.; Anderson, D.S.S.; Carlos, H.C.; Fergal, P.R.; Oliveira, C.A.F.D. Effect of lactic acid bacteria strains on the growth and aflatoxin production potential of *Aspergillus parasiticus*, and their ability to bind aflatoxin B1, ochratoxin A, and zearalenone in vitro. *Front. Microbiol.* **2021**, *12*, 655386. [CrossRef]
16. Cao, H.; Meng, D.; Zhang, W.; Ye, T.; Yuan, M.; Yu, J.; Wu, X.; Li, Y.; Yin, F.; Fu, C.; et al. Growth inhibition of *Fusarium graminearum* and deoxynivalenol detoxification by lactic acid bacteria and their application in sourdough bread. *Int. J. Food Sci. Technol.* **2021**, *56*, 2304–2314. [CrossRef]
17. Liang, N.; Zhao, Z.; Curtis, J.M.; Gänzle, M.G. Antifungal cultures and metabolites of lactic acid bacteria for use in dairy fermentations. *Int. J. Food Microbiol.* **2022**, *383*, 109938. [CrossRef]
18. Moumnassi, S.; Gharsallaoui, A.; Brahmi, M.; Bentouhami, N.E.; Yahyaoui, M.I.; Taibi, M.; Bouchra, E.G.; Reda, B.; Emilie, D.; Asehraou, A. Carbon source modulates the antifungal and antibiofilm activities of *Lactiplantibacillus plantarum* S61 supernatants against *Rhodotorula glutinis* UMP22. *Curr. Res. Food Sci.* **2025**, *12*, 101272. [CrossRef]
19. Gajendran, V.P.; Rajamani, S. Recent advancements in harnessing lactic acid bacterial metabolites for fruits and vegetables preservation. *Probiotics Antimicrob. Proteins* **2025**, *17*, 2673–2689. [CrossRef]
20. Karbowiak, M.; Wójcicki, M.; Hyun, J.E.; Szymański, P.; Niu, Y.D.; Zielińska, D. Novel antimicrobial compounds from fermented food-derived *Lacticaseibacillus paracasei* B1 and *Lactiplantibacillus plantarum* O24 strains: Genomic and proteomic analysis. *LWT-Food Sci. Technol.* **2025**, *234*, 118597. [CrossRef]
21. Li, G.; He, X.; Li, J.; Jian, Y.; Zhu, L.; Wang, S.; Zeng, W.; Zhang, T.; Lu, H.; Wang, L. Assessment of genomic and antifungal properties of *Lactococcus garvieae* ZB15 isolated from Zhenba bacon. *Front. Microbiol.* **2025**, *16*, 1610971. [CrossRef]
22. Fernandes, K.F.D.; de Oliveira, K.Á.R.; Cirilo, Í.F.; da Costa Lima, M.; Câmara, M.P.S.; de Souza, E.L. A novel coating formulated with sodium alginate loaded with a mixed lactic acid bacteria culture to control anthracnose in guava and mango. *Food Control* **2025**, *175*, 111341. [CrossRef]
23. Elsaadany, K.; El-Sayed, A.I.; Awad, S. Identification, Safety Assessment, and Antimicrobial Characteristics of Cocci Lactic Acid Bacteria Isolated from Traditional Egyptian Dairy Products. *Foods* **2024**, *13*, 3059. [CrossRef]
24. Zhao, X.; Wang, N.; Lu, Y.; Li, Y.; Zhao, T.; Xu, J.; Liu, B.; Shao, K.; Wang, Z.; Yuan, Y.; et al. Effects of cold plasma on the growth and aflatoxin production of *Aspergillus flavus*. *Food Biosci.* **2024**, *61*, 104552. [CrossRef]
25. Dudaš, T.; Celano, G.; Budakov, D.; Cotugno, P.; Grahovac, M.; Incerti, O.; Stojšin, V.; Angelis, M.D.; Ippolito, A.; Sanzani, S.M. Efficacy and mode of action of the biocontrol yeast *Meyerozyma caribbica* BBJ against blue mold and patulin production by *Penicillium expansum*. *Int. J. Food Microbiol.* **2025**, *444*, 111442. [CrossRef]
26. Chen, H.; Ju, H.; Wang, Y.; Du, G.; Yan, X.; Cui, Y.; Yuan, Y.; Yue, T. Antifungal activity and mode of action of lactic acid bacteria isolated from kefir against *Penicillium expansum*. *Food Control* **2021**, *130*, 108274. [CrossRef]
27. Tang, P.; Cao, C.; Cai, Y.; Zhang, Z.; Liu, Z.; Pan, F.; Wang, F.; Zhu, R. Inactivation of Botrytis cinerea by pulsed light and its corresponding mechanism. *Postharvest Biol. Technol.* **2025**, *227*, 113607. [CrossRef]
28. Liu, A.; Xu, R.; Zhang, S.; Wang, Y.; Hu, B.; Ao, X.; Li, Q.; Li, J.; Hu, K.; Yang, Y.; et al. Antifungal Mechanisms and Application of Lactic Acid Bacteria in Bakery Products: A Review. *Front Microbiol.* **2022**, *13*, 924398. [CrossRef] [PubMed]
29. Parappilly, S.J.; Idicula, D.V.; Chandran, A.; Mathil Radhakrishnan, K.; George, S.M. Antifungal activity of human gut lactic acid bacteria against aflatoxigenic *Aspergillus flavus* MTCC 2798 and their potential application as food biopreservative. *J. Food Saf.* **2021**, *41*, e12942. [CrossRef]
30. Chen, J.; Pang, H.; Wang, L.; Ma, C.; Wu, G.; Liu, Y.; Guan, Y.; Zhang, M.; Qin, G.; Tan, Z. Bacteriocin-Producing Lactic Acid Bacteria Strains with Antimicrobial Activity Screened from Bamei Pig Feces. *Foods* **2022**, *11*, 709. [CrossRef] [PubMed]
31. Gao, W.; Hernandez-Garcia, L.; Castagnini, J.M.; Pallares, N.; Barba, F.J.; Culebras, P.V.M. Identification, safety evaluation and probiotic potential of lactic acid bacteria isolated from wheat. *Int. J. Food Microbiol.* **2025**, *444*, 111436. [CrossRef] [PubMed]
32. Chalita, M.; Kim, Y.O.; Park, S.; Oh, H.S.; Cho, J.H.; Moon, J.; Baek, N.; Moon, C.; Lee, K.; Yang, J.; et al. EzBioCloud: A genome-driven database and platform for microbiome identification and discovery. *Int. J. Syst. Evol. Microbiol.* **2024**, *74*, 006421. [CrossRef]
33. Kumar, S.; Stecher, G.; Suleski, M.; Sanderford, M.; Sharma, S.; Tamura, K. MEGA12: Molecular evolutionary genetics analysis version 12 for adaptive and green computing. *Mol. Biol. Evol.* **2024**, *41*, msae263. [CrossRef]
34. Li, L.; Yang, B.; Yang, S.; Tian, X.; Gao, Y.; Song, J.; Wang, H.; Li, T.; Xing, F. Inhibitory effects of *Lactobacillus brevis* on *Aspergillus westerdijkiae* and antifungal compounds identification. *Postharvest Biol. Technol.* **2024**, *214*, 112980. [CrossRef]
35. García-Viñola, V.; Ezenarro, J.; Reguant, C.; Rozès, N.; Ferreira, M.M. Interaction effects of fumaric acid, pH and ethanol on the growth of lactic and acetic acid bacteria in planktonic and biofilm states. *Food Microbiol.* **2025**, *131*, 104808. [CrossRef]
36. Zhang, H.; HuangFu, H.; Wang, X.; Zhao, S.; Liu, Y.; Lv, H.; Qin, G.; Tan, Z. Antibacterial activity of lactic acid producing *Leuconostoc mesenteroides* QZ1178 against pathogenic *Gallibacterium anatis*. *Front. Vet. Sci.* **2021**, *8*, 630294. [CrossRef]

37. Yu, X.; He, H.; Wang, J.; Lin, L.; Wang, F. *Lactiplantibacillus pentosus* YXJ 1401 combats multidrug-resistant *Salmonella* Enteritidis: Antibacterial activity, dairy preservation, and metabolomics. *J. Dairy Sci.* **2025**, *108*, 11977–11995. [CrossRef] [PubMed]
38. Son, N.Y.; Moon, G.S.; Turner, M.S.; Yuk, H.G. Anti-listeria activity of newly isolated lactic acid bacteria from fermented foods and their application to smoked salmon (Coho). *Int. J. Food Microbiol.* **2025**, *434*, 111148. [CrossRef]
39. Hu, Y.; Wang, G.; Liu, Y.; Zhang, R.; Zhang, Z.; Li, F.; Liu, Z.; Li, X.; Xing, F. Antifungal mechanism of cell-free fermentation supernatant of *Lacticaseibacillus paracasei* L54 against *Aspergillus westerdijkiae* and its potential application with sorbic acid in grape preservation. *Postharvest Biol. Tec.* **2026**, *231*, 113958. [CrossRef]
40. Santarelli, G.; Rosato, R.; Cicchinelli, M.; Iavarone, F.; Urbani, A.; Sanguinetti, M.; Delogu, G.; De Maio, F. The activity of cell-free supernatant of *Lactobacillus crispatus* M247: A promising treatment against vaginal infections. *Front. Cell. Infect. Microbiol.* **2025**, *15*, 1586442. [CrossRef]
41. Sabri, M.; El Handi, K.; Calvano, C.D.; Bianco, M.; De Stradis, A.; Valentini, F.; Elbeaino, T. *Leuconostoc mesenteroides* strain MS4-derived bacteriocins: A potent antimicrobial arsenal for controlling *Xylella fastidiosa* infection. *Microbiol. Res.* **2025**, *293*, 128071. [CrossRef]
42. Cui, C.; Lv, H.; Liu, Y.; Huo, S.; Liu, S.; Zhao, Y. Degradation of deoxynivalenol by a mixed bacteria with lanthanum chloride as a growth factor. *Food Control* **2024**, *160*, 110335. [CrossRef]
43. Xia, C.; Zhou, Q.; Meng, X.; Chen, Z.; Wu, S.; Li, P.; Gu, Q. Unveiling the mechanism of *Latilactobacillus sakei* LZ217 cell-free supernatant against *Helicobacter pylori* through metabolomics and network pharmacology. *Food Biosci.* **2025**, *64*, 105925. [CrossRef]
44. Huang, J.; Yang, G.; Chen, K.; Du, M.; Zalán, Z.; Hegyi, F.; Kan, J. Anti-fungal effects of lactic acid bacteria from pickles on the growth and sterigmatocystin production of *Aspergillus versicolor*. *Int. J. Food Microbiol.* **2024**, *422*, 110809. [CrossRef] [PubMed]
45. Jaffar, N.S.; Jawan, R.; Chong, K.P. The potential of lactic acid bacteria in mediating the control of plant diseases and plant growth stimulation in crop production-A mini review. *Front. Plant Sci.* **2023**, *13*, 1047945. [CrossRef]
46. Nasrollahzadeh, A.; Mokhtari, S.; Khomeiri, M.; Saris, P.E. Antifungal preservation of food by lactic acid bacteria. *Foods* **2022**, *11*, 395. [CrossRef] [PubMed]
47. Ogunremi, O.R.; Ganz, G.; Leischtfeld, S.F.; Schwenninger, S.M. MALDI-TOF MS profiling and antifungal activity of lactic acid bacteria from *kunu aya*, a tiger nut traditional beverage of Nigeria. *Food Biosci.* **2024**, *61*, 104581. [CrossRef]
48. Zhang, J.; Han, Z.; Huang, M.; Li, X.; Li, Q.; Song, Z.; Shi, J. Antifungal activity of *Lactobacillus sakei* PLL12 and *Lactococcus lactis* PLL15 against *Aspergillus niger* and their application in bread preservation. *Food Control* **2025**, *182*, 111883. [CrossRef]
49. Kim, J.H.; Lee, E.S.; Kim, B.M.; Oh, M.H. Antifungal lactic acid bacteria as a biopreservative against fungal contamination in dry-cured ham. *LWT-Food Sci Technol.* **2025**, *230*, 118233. [CrossRef]
50. Aguirre-Garcia, Y.L.; Nery-Flores, S.D.; Campos-Muzquiz, L.G.; Flores-Gallegos, A.C.; Palomo-Ligas, L.; Ascacio-Valdés, J.A.; Sepúlveda-Torres, L.; Rodríguez-Herrera, R. Lactic acid fermentation in the food industry and bio-preservation of food. *Fermentation* **2024**, *10*, 168. [CrossRef]
51. Vasundaradevi, R.; Sarvajith, M.; Divyashree, S.; Deepa, N.; Achar, P.N.; Sreenivasa, M.Y. Tropical fruit-derived *Lactiplantibacillus* as potential probiotic and antifungal agents against *Fusarium oxysporum*. *Sci. Rep.* **2025**, *15*, 2144. [CrossRef] [PubMed]
52. Wang, Z.; Jia, S.; Xue, Y.; Wu, W.; Zhao, Y.; Li, Y.; Lv, H. *Weissella cibaria* DA2 cell-free supernatant improves the quality of sweet corn kernels during post-harvest storage. *Postharvest Biol. Technol.* **2024**, *215*, 113021. [CrossRef]
53. Shen, F.; Cheng, X.; He, C.; Jiang, T.; Wu, J.; Fan, Y.; Zhang, J.; Tian, J.; Zhao, Y. Exogenous inoculation of *Pediococcus pentosaceus* SSC12 positively regulates anthocyanin biosynthesis to alleviate browning of postharvest litchi (*Litchi chinensis* Sonn.). *LWT-Food Sci. Technol.* **2024**, *207*, 116644. [CrossRef]
54. Raman, J.; Kim, J.S.; Choi, K.R.; Eun, H.; Yang, D.; Ko, Y.J.; Kim, S.J. Application of lactic acid bacteria (LAB) in sustainable agriculture: Advantages and limitations. *Int. J. Mol. Sci.* **2022**, *23*, 7784. [CrossRef]
55. Teixeira, C.G.; Belguesmia, Y.; da Silva Rodrigues, R.; Lucau-Danila, A.; Nero, L.A.; de Carvalho, A.F.; Drider, D. Assessment of safety and in situ antibacterial activity of *Weissella cibaria* strains isolated from dairy farms in Minas Gerais State, Brazil, for their food application. *Braz. J. Microbiol.* **2024**, *55*, 699–710. [CrossRef]
56. Szczerbiec, D.; Piechocka, J.; Głowacki, R.; Torzewska, A. Organic acids secreted by *Lactobacillus* spp. isolated from urine and their antimicrobial activity against uropathogenic *Proteus mirabilis*. *Molecules* **2022**, *27*, 5557. [CrossRef]
57. Peh, E.; Kittler, S.; Reich, F.; Kehrenberg, C. Antimicrobial activity of organic acids against Campylobacter spp. and development of combinations—A synergistic effect. *PLoS ONE* **2020**, *15*, e0239312. [CrossRef] [PubMed]
58. Ma, J.; Hong, Y.; Deng, L.; Yi, L.; Zeng, K. Screening and characterization of lactic acid bacteria with antifungal activity against *Penicillium digitatum* on citrus. *Biol. Control* **2019**, *138*, 104044. [CrossRef]
59. Zhao, S.; Hao, X.; Yang, F.; Wang, Y.; Fan, X.; Wang, Y. Antifungal activity of *Lactobacillus plantarum* ZZUA493 and its application to extend the shelf life of Chinese steamed buns. *Foods* **2022**, *11*, 195. [CrossRef]
60. Garnier, L.; Penland, M.; Thierry, A.; Maillard, M.B.; Jardin, J.; Coton, M.; Salas, M.; Coton, E.; Valence, F.; Mounier, J. Antifungal activity of fermented dairy ingredients: Identification of antifungal compounds. *Int. J. Food Microbiol.* **2020**, *322*, 108574. [CrossRef] [PubMed]

61. Debonne, E.; Van Schoors, F.; Maene, P.; Van Bockstaele, F.; Vermeir, P.; Verwaeren, J.; Eeckhout, M.; Devlieghere, F. Comparison of the antifungal effect of undissociated lactic and acetic acid in sourdough bread and in chemically acidified wheat bread. *Int. J. Food Microbiol.* **2020**, *321*, 108551. [CrossRef] [PubMed]
62. Liu, P.; Yang, Z.; Jin, C.; Li, T.; Jin, F.; Lee, H.; Lee, C.; Jin, L. Antifungal activity of metabolites of *Weissella cibaria* KM14 isolated from traditional Korean food kimchi against three spoilage fungi. *LWT-Food Sci. Technol.* **2024**, *205*, 116545. [CrossRef]
63. Fan, W.; Li, B.; Du, N.; Hui, T.; Cao, Y.; Li, X.; Ren, H. Energy metabolism as the target of 3-phenyllactic acid against *Rhizopus oryzae*. *Int. J. Food Microbiol.* **2022**, *369*, 109606. [CrossRef] [PubMed]
64. Mani-López, E.; Arrioja-Bretón, D.; López-Malo, A. The impacts of antimicrobial and antifungal activity of cell-free supernatants from lactic acid bacteria in vitro and foods. *Compr. Rev. Food Sci. Food Saf.* **2022**, *21*, 604–641. [CrossRef]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.