

Wild Italian *Prunus spinosa* L. Fruit Exerts *In Vitro* Antimicrobial Activity and Protects Against *In Vitro* and *In Vivo* Oxidative Stress

Luisa Pozzo ^{1,*}, Rossella Russo ¹, Stefania Frassinetti ¹, Francesco Vizzarri ^{2,3}, Július Árvay ⁴, Andrea Vornoli ^{1,5}, Donato Casamassima ², Marisa Palazzo ², Clara Maria Della Croce ¹ and Vincenzo Longo ¹

¹ Institute of Agricultural Biology and Biotechnology (IBBA), National Research Council, 56124 Pisa, Italy; rossella.wilbur@gmail.com (R.R.); stefania.frassinetti@ibba.cnr.it (S.F.); vornolia@ramazzini.it (A.V.); clara.dellacroce@ibba.cnr.it (C.M.D.C.); v.longo@ibba.cnr.it (V.L.)

² Department of Agricultural, Environmental and Food Sciences, University of Molise, 86100 Campobasso, Italy; francesco.vizzarri@uniba.it (F.V.); casamassima.d@unimol.it (D.C.); m.palazzo@unimol.it (M.P.)

³ Department of Agricultural and Environmental Sciences, University of Bari Aldo Moro, 70126 Bari, Italy

⁴ Department of Chemistry, Faculty of Biotechnology and Food Sciences, Slovak University of Agriculture, 94976 Nitra, Slovakia; julius.arvay@uniag.sk

⁵ Cesare Maltoni Cancer Research Center, Ramazzini Institute, Bentivoglio, 40138 Bologna, Italy

* Correspondence: luisa.pozzo@ibba.cnr.it; Tel.: +39-0503152702

Received: 8 November 2019; Accepted: 17 December 2019; Published: 19 December 2019

Abstract: Polyphenol-rich foods could have a pivotal function in the prevention of oxidative stress-based pathologies and antibacterial action. The purpose of this study was to investigate the *in vitro* antimicrobial activity, as well as the *in vitro* and *in vivo* antioxidant capacities of wild *Prunus spinosa* L. fruit (PSF) from the southeast regions of Italy. The total phenolic content (TPC) was quantified, and the single polyphenols were analyzed by HPLC-DAD, showing high rutin and 4-hydroxybenzoic acid levels, followed by gallic and trans-sinapic acids. PSF extract demonstrated antimicrobial activity against some potentially pathogenic Gram-negative and Gram-positive bacteria. Besides, we investigated the cellular antioxidant activity (CAA) and the hemolysis inhibition of PSF extract on human erythrocytes, evidencing both a good antioxidant power and a marked hemolysis inhibition. Furthermore, an *in vivo* experiment with oxidative stress-induced rats treated with a high-fat diet (HFD) and a low dose of streptozotocin (STZ) demonstrated that PSF has a dose-dependent antioxidant capacity both in liver and in brain. In conclusion, the wild Italian *Prunus spinosa* L. fruit could be considered a potentially useful material for both nutraceutical and food industries because of its antioxidant and antimicrobial effects.

Keywords: wild Italian *Prunus spinosa* L. fruit; blackthorn; phenolic compounds; antimicrobial; antioxidant

1. Introduction

Considering that traditional foods are increasingly believed healthy and wholesome, food manufacturers are developing new food products returning to natural products and traditional recipes that will be attractive to the widest potential consumers [1,2]

Blackthorn (*Prunus spinosa* L.), which belongs to the *Rosaceae* family, is a perennial plant originally growing in temperate continental climate of the northern hemisphere, particularly widespread in the Mediterranean countries and in the southeast regions of Italy. It is used for treatment of many diseases due to its diuretic, spasmolytic, antimicrobial, and antioxidant activities

[3]. Moreover, *Prunus spinosa* L. fruit (PSF) is used for the production of various traditional jams and beverages such as juice, wine, tea, and distillates in food industry [4]. It contains substantial quantities of phenolic antioxidants, including, in particular, flavonols, phenolic acids, and coumarin derivatives [5].

Epidemiological investigations demonstrated that diets rich in plant polyphenols protect against diabetes, osteoporosis, cardiovascular, and neurodegenerative diseases [6]. Dietary compounds and specific polyphenol-rich foods could have a pivotal function in the prevention of diseases associated with oxidative stress by increasing the circulation of antioxidant compounds and neutralizing the reactive oxygen species, due to their number and position of hydroxyl groups [7]. Protein nitration, lipid peroxidation, chronic inflammation, and oxidative damage to DNA may be prevented by polyphenols, which results in vasodilatory, vasoprotective, anti-atherogenic, antithrombotic, and anti-apoptotic effects, as free radical scavengers, metal chelators, inhibitors of pro-inflammatory enzymes, and modifiers of cell signaling pathways [8].

Recently, new alternatives have become desirable, and plants metabolites have been screened for antimicrobial agents for treatment of infectious diseases, due to the development of antibiotic resistance by pathogenic bacteria [9]. For instance, in several studies, dietary polyphenols have been reported to exert an antibacterial activity [10].

In the present study, the potential biological activities of wild blackthorn fruit from southeast regions of Italy were investigated. Considering the PSF as a potential natural source of phenolic compounds, this work was designed to study its *in vitro* antimicrobial, antioxidant, and antihemolytic activities. Moreover, for the first time, the *in vivo* protective effect and antioxidant capacity of PSF in high-fat diet (HFD) and streptozotocin (STZ)-induced oxidative stress has been studied.

2. Materials and Methods

Blackthorn fresh fruits were obtained from wild orchards of the Campobasso (Italy) area in October 2015. Taxonomic identification of plant material was confirmed by Prof. Elisabetta Brugiapaglia from Department of Agricultural, Environmental and Food Sciences, University of Molise, Campobasso, Italy.

2.1. Chemicals and Reagents

All solvents and chemicals were of analytical grade. Nutrient Broth (NB), Nutrient Agar (NA), Mueller Hinton Broth (MHB), Mueller Hinton Agar (MHA), McFarland standard were purchased from Oxoid (Basingstone, UK). 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid (Trolox), 2,2'-azobis (2-amidinopropane) dihydrochloride (AAPH), 2,7-dichlorodihydrofluorescein diacetate (DCFH-DA), dinitrophenylhydrazine (DNPH), trichloroacetic acid (TCA), perchloric acid (PCA), thiobarbituric acid (TBA), 1,1,3,3-tetramethoxypropane (TEP), guanidine hydrochloride, ortho-phthalaldehyde (OPA), reduced glutathione (GSH), phosphoric acid, potassium dihydrogen phosphate (KH₂PO₄), hydrochloric acid (HCl), streptozotocin (STZ), ethanol, ethyl acetate, and methanol from Sigma-Aldrich (St. Louis, MO, USA). All HPLC analytical standards, including protocatechuic acid, syringic acid, rutin, ellagic acid, cynaroside, daidzein, neochlorogenic acid, chlorogenic acid, vitexin, trans *p*-coumaric acid, trans-sinapic acid, trans ferulic acid, rosmarinic acid, resveratrol, apigenin, myricetin, quercetin, and kaempferol, were bought from Sigma-Aldrich (St. Louis, MO, USA). Phosphate buffer saline (PBS) was bought from VWR (Radnor, PA, USA).

2.2. Plant Material Preparation

After the pits were removed, the frozen fruits were lyophilized and crushed in a mortar, and the powder was stored at −20 °C. Briefly, 1 g of sample was extracted with 10 mL of water for 2 h on a horizontal shaker Unimax 2010 (Heidolph Instruments, GmbH, Schwabach, Germany) for *in vitro* antioxidant activity, polyphenols quantification, and antimicrobial activity. PSF extracts were

centrifuged (2300× g at 4 °C for 20 min) (Jouan CR3i centrifuge, Newport Pagnell, UK), and the supernatants were collected.

For the *in vivo* experiment, the lyophilized and powdered fruit was dissolved in water.

2.3. Total Phenolic Content (TPC) and Polyphenols Quantification by HPLC-DAD

For the determination of total phenolic content in water extract, we followed the Singleton and colleagues protocol [11]. The concentration of polyphenols was expressed as mg of gallic acid equivalents (GAE)/g of dry weight (d.w.).

Prior to HPLC analysis, the extracts were filtered through syringe filters Q-Max (0.22 µm, 25 mm, PVDF) (Frisenette ApS, Knebel, Denmark) into the HPLC vials. The HPLC apparatus consisted of an Agilent 1260 Infinity HPLC (Agilent Technologies GmbH, Waldbronn, Germany) quaternary solvent manager coupled with degasser (G1311B), sampler manager (G1329B), Diode Array Detector (G1315C), column manager (G1316A). The analytical column was a Waters Cortecs endcapped RP-C18 column (150 mm × 4.6 mm × 2.7 µm particle size; Waters Corp., Milford, MA, USA). The analyses were carried out at 30 °C by a gradient system with a mobile phase of 0.1% ortho-phosphoric acid in deionised water (C) and acetonitrile gradient grade (D) at a flow rate of 0.60 mL/min, and the injection volume was 5 µL. The gradient elution was as follows: 0–1 min (90% C and 10% D), 1–5 min (85% C and 15% D), 5–10 min. (80% C and 20% D), 10–12 min. (80% C and 20% D), 12–20 min (30% C and 70% D), and 20–25 min (30% C and 70% D). The post-run was set at 3 min. The samples were kept at 4 °C in the sampler manager. The detection wavelengths were set at 265 nm (gallic acid, 4-hydroxybenzoic acid, rutin, and genistein), 320 nm (chlorogenic acid, caffeic acid, trans-*p*-coumaric acid, trans-sinapic acid, trans-ferulic acid, rosmarinic acid, resveratrol), and 372 nm (myricetin, quercetin and kaempferol). Data were analyzed by Agilent Open Lab Chem Station software for LC 3D systems.

2.4. HPLC-DAD Method Validation

Reference phenolic compounds were dissolved in HPLC purity methanol and diluted to appropriate concentration ranges (5–50 µg/mL). The linearity of each calibration curve was assessed by linear regression analysis. The limit of detection (LOD) and quantification (LOQ) were estimated by measuring signal-to-noise ratio of the individual peak of each standard compound. The LOD and LOQ were calculated according to the International Conference on Harmonisation guidelines [12].

2.5. Antimicrobial Activity

2.5.1. Growth Conditions of Pathogenic Bacteria

The bacterial strains were supplied by the American Type Culture Collection (ATCC). The antimicrobial activity of PSF extract was studied on three Gram-negative bacteria, specifically *Escherichia coli* (ATCC 25922), *Salmonella enterica* ser. *typhimurium* (ATCC 14028), and *Enterobacter aerogenes* (ATCC 13048), and two Gram-positive bacteria, *Enterococcus faecalis* (ATCC 29212) and *Staphylococcus aureus* (ATCC 25923).

2.5.2. Antimicrobial Activity

The growth inhibition of selected bacteria exerted by PSF extract was determined according to Delgado Adamez and colleagues [13], with some modifications.

The tested bacteria were cultured in MHB at 37 °C for 16 h and diluted to match the turbidity of 0.5 McFarland standard. Fifty microliters of bacterial suspensions (about $1-5 \times 10^5$ CFU/mL) was added to 100 µL of MHB and to 100 µL of blackthorn extract (0, 0.25, 0.50, 0.75, and 1.00 mg/mL) in a 96-well plate. A negative control was included on each microplate. A positive control of bacterial growth inhibition consisting of two antibiotics, vancomycin (10 µg/mL) for Gram-positive and gentamicin (10 µg/mL) for Gram-negative bacteria was added to the microplate. The plates were incubated at 37 °C for 24 h. Afterwards, the optical density (OD) at 600 nm was determined by a

microplate reader (Eti-System fast reader Sorin Biomedica, Modena, Italy). The percentage of growth inhibition was calculated as follows:

$$\% \text{ growth inhibition} = 100 - (\text{OD}_s / \text{OD}_c) \times 100 \quad (1)$$

where OD_s is the optical density of the sample and OD_c is the optical density of the negative control (PSF 0 mg/mL).

2.6. In Vitro Antioxidant Activity in Red Blood Cells (CAA-RBC) and Hemolysis Test

According to the regulations of “Fondazione G. Monasterio CNR-Regione Toscana”, human blood samples were obtained from three healthy volunteers in ethylenediaminetetraacetic acid (EDTA)-treated tubes and centrifuged ($2300 \times g$ at 4°C for 10 min). Plasma and buffy coat were removed, and erythrocytes were washed twice with PBS pH 7.4.

The antioxidant activity of PSF extract (100 mg/mL) was evaluated in an in vitro system with a modified assay in red blood cells as described by Frassinetti and colleagues [14]. Each value was expressed as CAA units, as follows [12]:

$$\text{CAA unit} = 100 - (\text{JSA} / \text{JCA}) \times 100 \quad (2)$$

where JSA is the integrated area of the sample curve and JCA is the integrated area of the control curve.

Hemolysis of PSF extract (100 mg/mL) was analyzed according to the protocol described by Frassinetti and colleagues [15] using AAPH, a generator of peroxy radicals, to cause the red blood cell lysis. The values reported are the percentage of hemolysis compared with the control.

2.7. Animal Study

2.7.1. In Vivo experiment

Male Wistar rats (200–230 g b.w.) were maintained with ad libitum access to food and drinking water for a 12 h light/dark cycle in cages at room temperature with the 55% relative humidity. Rats were divided into two groups: the control (CTR) group ($n = 5$), fed a standard diet (64% carbohydrates, 19% proteins, 7% minerals and vitamins, 6% fibers, and 4% fats; the fats percentage corresponds to the 11% of the diet-derived energy) and the high-fat diet (HFD) group, fed a high fat/cholesterol diet (48.7% carbohydrates, 28% fats, including 2% cholesterol, 13.8% proteins, 4.4% fibers, 5.1% minerals and vitamins, the fats percentage corresponds to the 55% of diet-derived energy). After 5 weeks, the animals of HFD group were treated with a single i.p. injection of streptozotocin (40 mg/kg) [16]. Twenty four rats, resulted to be diabetic with a plasma glucose concentration $>250 \text{ mg/dL}$, continued to be fed a HFD diet for a further 4 weeks and were randomly divided into three groups: HFD ($n = 8$) group, PSF400 ($n = 8$) group, and PSF800 ($n = 8$) group (Figure 1). Rats from CTR and HFD groups were intragastrically administered the same volume of water; rats from PSF400 and PSF800 groups were intragastrically administered lyophilized PSF at different doses (400 mg/kg b.w. and 800 mg/kg b.w., respectively). The weight gain was calculated by initial and final weights. The rats were sacrificed and blood samples were collected by cardiac puncture under general anesthesia. Liver and brain tissues were stored at -80°C . Hepatic lipids were quantified and oxidative stress markers were analyzed in liver and brain. Blood was centrifuged ($2300 \times g$ for 15 min) to obtain serum samples for laboratory analysis. Local Ethical Committee approved all animal procedures in accordance with the European Communities Council Directive of 24 November, 1986 (86/609/EEC).




Figure 1. Study design of the *in vivo* experiment. CTR, control group; HFD, high-fat diet group; PSF400, PSF800,.

2.7.2. Biochemical Analysis

Serum analyses were performed by a semi-automatic analyzer for clinical chemistry (model ARCO, Biotecnica Instruments SPA, Rome, Italy) for aspartate aminotransferase (AST), alanine aminotransferase (ALT), total cholesterol, and triglycerides. Glucose levels were measured with a glucose meter (Accu-Chek® Roche, Mannheim, Germany), and insulin using a Rat Insulin ELISA kit (Mercodia, Uppsala, Sweden).

2.7.3. Hepatic Lipids Quantification

Fat content of the liver samples was determined by Folch and colleagues protocol [17], slightly modified. Liver samples from rats were homogenized with equal volumes of water and methanol. The resulting homogenate was subjected to three subsequent extractions in chloroform, followed by two washes with KCl 1 M and water. After complete evaporation and prolonged drying of the chloroform, fat content was weighed and expressed as mg/g tissue.

2.7.4. Oxidative Stress

Malondialdehyde (MDA) concentration of liver and brain samples was analyzed according to Seljeskog and colleagues [18], with some adaptations. An aliquot of 100 µL of homogenate sample was mixed with 0.1125 N PCA (300 µL) and 40 mM TBA (300 µL) for 10 sec and placed in a boiling water bath for 60 min. Methanol (600 µL) and 20% TCA (*w/v*) (200 µL) were added to the suspension and mixed for 10 sec, after cooling in a freezer at −20 °C for 20 min. The MDA content was quantified in the supernatant (7000× *g* for 6 min) by fluorimeter (Perkin Elmer LS-45, Perkin Elmer, Walham, MA, USA) ($\lambda_{\text{ex}} = 525$, $\lambda_{\text{em}} = 560$). A standard curve was prepared by dissolving hydrolyzed TEP in water at different concentrations (33.5, 16.8, 8.4, 4.20, 2.10, 1.05, and 0.52 µM). The results have been expressed as nmol MDA/g tissue.

The protein carbonylation was determined using the method adapted from Terevinto and colleagues [19]. Liver and brain samples were homogenized and incubated with 0.02 M DNPH in 2 M HCl. Proteins were then precipitated by adding 20% TCA and recovered by centrifugation (625× *g* for 10 min). Pellets were washed three times with ethanol:ethyl acetate (1:1, *v/v*), melted in 6 M guanidine HCl in 0.02 M KH₂PO₄ (pH 6.5), and centrifuged. The absorbance of the supernatant was measured at 390 nm. The results have been expressed as nmol/g tissue.

The GSH content in liver and brain samples was evaluated according to Browne and Armstrong [20], with slight modifications. Proteins were then precipitated by adding 10% TCA (*w/v*) at 4 °C for 30 min. An aliquot of 150 µL of the sample was incubated with an equal volume of *o*-phthaldehyde (1 mg/mL) in 10% methanol (*v/v*) by 15 min at 37 °C. After centrifugation (625× *g* for 3 min), fluorescence was measured (Perkin Elmer LS-45, Perkin Elmer, Walham, MA, USA) ($\lambda_{\text{ex}} = 350$, $\lambda_{\text{em}} = 420$). A calibration curve has been performed by dissolving GSH in water at different concentrations (50, 25, 12.5, 6.25, 3.13, 1.56, 0.78 µM), and GSH concentrations have been calculated as µmol GSH/g tissue.

2.8. Statistical Analysis

The statistical analyses have been performed by Statistical Package for Social Science (SPSS) 17 for Windows (SPSS, Inc., Chicago, IL, USA). The results are shown as the mean value \pm standard deviation (s.d.) and analyzed through a one-way ANOVA and Tukey's test for post-hoc with significance at $p \leq 0.05$.

3. Results and Discussion

3.1. Quantification of Total Polyphenols

The TPC of wild Italian blackthorn fruit extract was quantified by a spectrophotometric method, and the content was 5.50 ± 0.19 mg GAE/g d.w. To our knowledge, any other results have been found about total phenolic content of blackthorn fruit on dry weight, but some authors reported that TPC in blackthorn fruit on fresh weight ranged from 0.42–4.13 mg GAE/g [21–23].

The HPLC-DAD method validation was estimated by quantifying the limit of detection (LOD), limit of quantification (LOQ), and the recovery. All parameters indicate that the method exhibits a good sensitivity for identification as well as quantification of the polyphenols. All the parameters are listed in Table 1.

Table 1. Retention time (Rt), LOD (limit of detection), LOQ (limit of quantification), and recovery of phenolic compound quantification method by HPLC-DAD in the *Prunus spinosa* L. fruit (PSF) aqueous extract ($n = 3$).

Phenolic Compound	Rt (min)	LOD (ug/mL)	LOQ (ug/mL)	Recovery (%)
Gallic acid	2.860	0.012	0.033	98.2 \pm 0.81
Rutin	5.909	0.009	0.030	89.1 \pm 0.89
4-hydroxybenzoic acid	7.112	0.005	0.017	101.2 \pm 1.01
Caffeic acid	8.361	0.008	0.027	97.5 \pm 0.99
Trans p-coumaric acid	11.741	0.004	0.013	96.0 \pm 0.80
Trans -ferulic acid	12.981	0.003	0.010	97.8 \pm 1.08
Trans-sinapic acid	13.062	0.011	0.037	98.2 \pm 1.15
Myricetin	17.081	0.015	0.050	99.5 \pm 0.88
Rosmarinic acid	17.463	0.009	0.023	102.1 \pm 0.96
Quercetin	18.853	0.090	0.299	99.5 \pm 0.95
Genistein	19.811	0.009	0.031	91.5 \pm 0.77

The quality of phenolic profile of wild Italian blackthorn and the concentrations of single compounds are shown in Table 2. Rutin (183.94 mg/kg d.w.) was the principal phenolic component, followed by 4-hydroxybenzoic acid, gallic acid, trans-sinapic acid, quercetin, trans-ferulic acid, caffeic acid, rosmarinic acid, trans coumaric acid, genistin, and myricetin. Our findings are partially in accordance with those of some other authors that showed considerable quantities of phenolic acids (quercetin and caffeic acid) in blackthorn fruits from Southeast Serbia [3,24]; by contrast we did not find either neochlorogenic or kaempferol. HPLC-UV analysis of the methanolic extract of fresh blackthorn plums from Turkey recently allowed Baltas and colleagues to detect five phenolic acids, namely protocatechuic acid, p-OH benzoic acid, vanillic acid, syringic acid, and p-coumaric acid, as well as flavonoids, such as epicatechin and luteolin [25]. Another recent study about quantification

of phenolic compounds by HPLC-UV in methanolic extract of frozen blackthorn fruits from Romania showed high chlorogenic and neochlorogenic acid levels, followed by glycosides of quercetin [23]. Considering that the solubility of polyphenols in solvent of different polarity is determined by their structure, different types of extraction solvent and procedures may influence the efficiency of phenolic compounds extraction and their resultant content [3].

Table 2. Concentrations of phenolic compounds in the PSF aqueous extract.

Phenolic Compound	Concentration (mg/kg d.w.)
Gallic acid	41.10 ± 3.68
Rutin	183.94 ± 0.45
4-hydroxybenzoic acid	73.93 ± 0.06
Caffeic acid	3.36 ± 0.36
Trans <i>p</i> -coumaric acid	2.99 ± 0.02
Trans-ferulic acid	4.93 ± 0.07
Trans-sinapic acid	37.69 ± 0.05
Myricetin	1.47 ± 0.03
Rosmarinic acid	3.23 ± 0.03
Quercetin	9.94 ± 0.01
Genistin	1.74 ± 0.00

3.2. Antimicrobial Activity

The antimicrobial activity on selected Gram-negative (Figure 2A) and Gram-positive (Figure 2B) enteric bacteria was measured by evaluating the growth inhibition by increasing concentrations of PSF extract. The antimicrobial activities have been compared with the standard antibiotics, used as positive controls.

The lowest concentration of tested PSF extract (0.25 mg/mL) inhibited more than 50% of the Gram-negative bacteria *Escherichia coli* (70.19% ± 1.21%), *Salmonella typhimurium* (79.98% ± 0.54%), and *Enterobacter aerogenes* (83.02% ± 0.54%) growth (Figure 2A). The same concentration (0.25 mg/mL) was able to inhibit more than 50% of the Gram-positive bacteria *Enterococcus faecalis* (82.86% ± 1.94%) and *Staphylococcus aureus* (79.92% ± 1.23%) growth (Figure 2B). The antimicrobial activity of phenolic compounds occurring in plant foods has been widely studied against a wide range of microorganisms. The damage to the bacterial membrane and suppression of some virulence factors, including enzymes and toxins, are suggested to be the mechanisms of their antimicrobial action [26]. Some flavonoids (rutin, myricetin, and quercetin) and phenolic acids (gallic, caffeic, and ferulic acids) of PSF extract may be responsible for its antibacterial action [27,28].




Figure 2. Growth inhibition effect of PSF extract (0, 0.25, 0.50, 0.75, and 1.00 mg/mL) against Gram-negative bacteria (**A**) (*Escherichia coli* ATCC 25922, *Salmonella enterica* ser. *typhimurium* ATCC 14028, and *Enterobacter aerogenes* ATCC 13048) and Gram-positive bacteria (**B**) (*Enterococcus faecalis* ATCC 29212 and *Staphylococcus aureus* ATCC 25923). Significantly different from negative control (PSF 0 mg/mL): *** $p \leq 0.001$. Results are reported as means ($n = 3$) values \pm standard deviation.

3.3. In Vitro Antioxidant Activity

As shown in Figure 3A, pretreated erythrocytes with PSF aqueous extract (100 mg/mL) exhibited a significantly higher cellular antioxidant activity (CAA unit = 48.43 ± 1.68) compared with untreated cells (CAA = 0; $p \leq 0.001$), comparable to 100 μ M Trolox (CAA unit = 16.52 ± 3.60 ; $p \leq 0.001$) and 500 μ M Trolox (CAA unit = 36.67 ± 1.48 ; $p \leq 0.001$). Taking these results into consideration, the EC50 of PSF extract for antioxidant activity in red blood cells was 100 mg/mL.

The antihemolytic activity of PSF extract was screened in erythrocytes exposed to high doses of AAPH, causing a strong oxidative hemolysis. Figure 3B shows that PSF extract exerted a significant inhibition of AAPH-induced hemolysis compared with the control erythrocytes (AAPH-treated). PSF extract (100 mg/mL) pretreated cells demonstrated a marked antihemolytic effect (84% hemolysis inhibition) compared with AAPH-treated cells ($p \leq 0.001$), with a reduction of the hemolysis similar to that of the highest concentration of the reference standard (500 μ M Trolox). The antihemolytic EC50 of PSF extract was 10 mg/mL (data not shown). We found that PSF exerted a potent ROS-scavenger activity. Indeed, when intact human erythrocytes were pre-incubated with a PSF aqueous extract, a strong protective effect against AAPH-generated ROS production and hemolysis was observed. These antioxidant and antihemolytic effects of PSF are probably due to the activity of gallic acid, rutin, and quercetin in red blood cell [29,30].




Figure 3. (A) Effects of PSF extract (100 mg/mL) on cellular antioxidant activity (CAA) in human erythrocytes. Significantly different from untreated cells (CAA unit = 0): *** $p \leq 0.001$. (B) Effects of PSF extract (100 mg/mL) on dihydrochloride (AAPH)-induced oxidative hemolysis in human erythrocytes. Significantly different from CTR (AAPH-treated cells): *** $p \leq 0.001$. Trolox was used as reference standard. Results are reported as means ($n = 3$) values \pm standard deviation.

3.4. In Vivo Experiment

3.4.1. The Effect of Blackthorn on Body Weight and Liver Weight

In comparison with CTR group, rats of the HFD group exhibited a significant lower final body weight (396.8 ± 40.6 vs. 307.5 ± 23.3 g/rat, respectively) ($p \leq 0.001$). The administration of PSF did not induce significant changes in the final body weight, neither in PSF400 group (317.2 ± 27.6 g/rat), nor in PSF800 group (312.7 ± 42.5 g/rat), when compared with HFD group.

However, when compared with CTR rats, HFD rats exhibited a statistically significant increase in liver weight (8.9 ± 1.4 vs. 13.4 ± 1.2 g, respectively) ($p \leq 0.001$) and in relative liver weight (2.2 ± 0.2 vs. 4.1 ± 1.0 g liver/100 g b.w., respectively) ($p \leq 0.001$). No significant difference in liver weight was found between HFD-fed rats and PSF-treated rats of PSF400 group (14.8 ± 1.7 g) and PSF800 group (14.6 ± 1.8 g). The same trend was found in relative liver weight between HFD-fed rats and

PSF-treated rats of PSF400 group (4.5 ± 0.3 g liver/100 g b.w.) and PSF800 group (4.8 ± 0.7 g liver/100 g b.w.). HFD treatment caused hepatic lipid accumulation and increased liver weight and all the biochemical parameters in serum [16,31]. However, the PSF extract did not improve the liver weight and serum and liver biochemical parameters linked to steatosis.

3.4.2. The Effect of PSF on Serum and Liver Biochemical Parameters

Serum AST, ALT, glucose, total cholesterol, triglycerides, and total hepatic lipid content were significantly higher in the HFD group compared with the normal diet group (CTR group), while serum insulin was significantly lower. After four weeks of treatment with 800 mg of PSF/kg b.w., rats of the PSF800 group showed a significant decrease of total hepatic lipids content compared with HFD group (162.15 ± 35.52 vs. 209.90 ± 11.91 ; $p \leq 0.05$) (Table 3). Some studies have demonstrated that polyphenols decrease the hepatic lipid accumulation caused by high-fat diet [32]. Moreover, it was also reported that the single isolated polyphenol can improve the high liver lipids content due to a high-fat diet administration, as in the case of rutin [33], gallic acid [34], and quercetin [35]. The crude extracts can be more advantageous than the isolated components, since a single bioactive molecule can change its properties with the presence of other compounds in the extracts [36].

Table 3. Nutritional effect of PSF on biochemical parameters in serum and liver of rats ($n = 7$).

	CTR	HFD	PSF400	PSF800
AST (UI/dl)	93.98 ± 7.04	$194.33^{**} \pm 42.90$	$181.00^{**} \pm 22.23$	$168.50^{*} \pm 44.81$
ALT (UI/dl)	39.06 ± 10.09	$143.17^{**} \pm 35.15$	$146.52^{**} \pm 51.96$	$133.58^{**} \pm 36.90$
Insulin (μ g/l)	1.44 ± 0.80	$0.19^{**} \pm 0.15$	$0.22^{**} \pm 0.15$	$0.20^{**} \pm 0.03$
Glucose (mg/dl)	145.20 ± 20.80	$439.67^{***} \pm 70.21$	$432.57^{***} \pm 33.57$	$443.60^{***} \pm 43.32$
Total cholesterol (mg/dl)	109.30 ± 21.40	$236.20^{***} \pm 45.09$	$228.36^{***} \pm 29.13$	$219.18^{***} \pm 17.94$
Triglycerides (mg/dl)	75.73 ± 7.38	$179.80^{**} \pm 59.30$	$164.33^{**} \pm 30.52$	$170.25^{**} \pm 21.80$
Total hepatic lipids (mg/g)	65.36 ± 9.14	$209.90^{***} \pm 11.91$	$198.29^{***} \pm 34.52$	$162.15^{***} \pm 35.52$

Analyses were performed through one-way ANOVA and Tukey's test for post-hoc. *, $p \leq 0.05$ vs. CTR; **, $p \leq 0.01$ vs. CTR; ***, $p \leq 0.001$ vs. CTR; §, $p \leq 0.05$ vs. HFD. AST, aspartate aminotransferase; ALT, alanine aminotransferase.

3.4.3. The Effect of PSF on Liver and Brain Oxidative Stress of Rats

The high-fat diet was probably responsible for the decrease of GSH (Figure 4A) content and the increase of protein carbonylation (Figure 4C) and MDA (Figure 4E) levels in the liver samples of HFD rats, compared with the CTR group. Moreover, while hepatic GSH content was not affected by PSF treatment (Figure 4A), administration of PSF improved the oxidative stress status of rats according to the protein carbonylation, at the higher concentration of treatment (800 mg/kg b.w.) (Figure 4C), and to the MDA content, in a dose-dependent manner (Figure 4E). It has been demonstrated that plant polyphenols are related to the improvement of hepatic oxidative stress caused by a high-fat diet through the activation of Nrf2 transcription factor, which increases expression of antioxidant enzymes [37]. Moreover, it was reported that even the single polyphenol, if isolated, can improve the high-fat-diet-induced hepatic oxidative stress, as in the case of rutin [33] and gallic acid [38].

In comparison with the CTR group, HFD treatment promoted an increase of the brain oxidative stress parameters in rats, as shown by protein carbonylation (Figure 4D) and MDA assay (Figure 4F). The addition of 400 mg/kg b.w. and 800 mg/kg b.w. of PSF to the diet reversed the effect caused by the high-fat diet and, in particular, the MDA assay showed a dose-dependent response pattern. Nevertheless, both the HFD and the PSF treatments did not induce significant changes in rat brain GSH levels (Figure 4B). The intake of a high-fat diet is linked to an increased risk of neurodegenerative disease related to diabetes [39]. Considering that, the polyphenols-rich fruits could protect neurons against the oxidative stress induced by intake of saturated fatty acids [40]. Recently, Nabavi and colleagues demonstrated that gallic acid exerts a neuroprotective effect against sodium fluoride-induced oxidative stress in rat brain [41]. Moreover, it has been shown that other

polyphenols contained in PSF, such as rutin, ferulic acid, and trans-sinapic acid, should contribute to the prevention of brain oxidative stress in rats [42–44].

Our findings suggest an improved liver and brain antioxidant defense in rats treated with PSF.




Figure 4. Effect of PSF treatment at two different concentrations (PSF400 and PSF800) on GSH content of liver (A) and brain (B), protein carbonylation of liver (C) and brain (D), and malondialdehyde of liver (E) and brain (F). Results are reported as means ($n = 7$) values \pm standard deviation. Values within each row of different letters are significantly different ($p \leq 0.05$).

4. Conclusions

All in all, our findings indicated that wild Italian blackthorn fruit is rich in polyphenol compounds, shows an *in vitro* antioxidant activity, and exhibits a selective growth inhibition of some potentially pathogenic bacteria strains. Moreover, this study is the first to evaluate an *in vivo* antioxidant activity of PSF. In particular, our findings indicated that the oxidative stress arising in HFD group is decreased in liver and brain tissues by the intake of blackthorn fruit. The PSF supplementation demonstrated *in vivo* antioxidant capacities, reducing liver and brain oxidative

stress, probably due to the presence of polyphenols, such as rutin, 4-hydroxybenzoic acid, gallic acid, trans-sinapic acid, quercetin, trans-ferulic acid, caffeic acid, rosmarinic acid, trans coumaric acid, genistin, and myricetin, which were identified in the blackthorn fruit.

Thus, it is supposed that the regular consumption of wild Italian blackthorn fruit should increase the circulation of bioactive compounds, such as polyphenols, which could possibly improve the endogenous antioxidant system and protect tissues against oxidative stress damage induced by high-fat diet and hyperglycemia. Considering its beneficial properties, wild Italian blackthorn fruit can be potentially used to produce natural functional food, novel nutraceuticals, and it can also be employed in food processing.

Author Contributions: Investigation, L.P., R.R., F.V., A.V. and C.M.D.C.; Methodology, L.P., R.R., S.F., J.A., M.P. and C.M.D.C.; Supervision, D.C. and V.L.; Writing—original draft, L.P., R.R., F.V. and A.V.

Funding: This research received no external funding.

Acknowledgments: This study was supported by Institute of Biology and Agricultural Biotechnology of National Research Council (Italy), by Department of Agricultural, Environmental and Food Sciences, University of Molise (Italy) and by AgroBioTech Research Center (Slovakia), built in accordance with the project Building AgroBioTech Research Center ITMS 26220220180.

Conflicts of Interest: No conflict of interest was reported by the authors.

Abbreviations:

ALT	alanine aminotransferase
ANOVA	analysis of variance
AST	aspartate aminotransferase
AATCC	american type culture collection
AAPH	2,2'-azobis (2-amidinopropane) dihydrochloride
CAA-RBC	cellular antioxidant activity in red blood cells
CFU	colony-forming unit
CTR	control
DCFH-DA	2,7-dichlorodihydrofluorescein diacetate
DNPH	dinitrophenylhydrazine
EC50	half maximal effective concentration
EDTA	ethylenediaminetetraacetic acid
GAE	gallic acid equivalent
GSH	reduced glutathione
HCl	hydrochloric acid
HFD	high fat diet
KH ₂ PO ₄	potassium dihydrogen phosphate
LOD	limit of detection
LOQ	limit of quantification
MIC	minimum inhibitory concentrations
MDA	malondialdehyde
MHA	Mueller Hinton agar
MHB	Mueller Hinton broth
NA	nutrient agar
NB	nutrient broth
OD	optical density
PBS	phosphate buffer saline

References

1. Bhaskarachary, K.; Sudershan, R.V.; Subba Rao, M.G.; Apurva Kumar, R.J. Traditional foods, functional foods and nutraceuticals. *Proc. Indian Natn. Sci. Acad.* **2016**, *82*, 1565–1577.

2. Grochowicz, J.; Fabisiak, A.; Nowak, D. Market of functional food—legal regulations and development perspectives. *Zesz. Probl. Postepow Nauk Roln.* **2018**, *595*, 51–67.
3. Veličković, J.M.; Kostić, D.A.; Stojanović, G.S.; Mitić, S.S.; Mitić, M.N.; Randelović, S.S.; Đorđević, A.S. Phenolic composition, antioxidant and antimicrobial activity of the extracts from *Prunus spinosa* L. fruit. *Hem. Ind.* **2014**, *68*, 297–303.
4. Egea, I.; Sanchez-Bel, P.; Romorjaro, F.; Pretel, M.T. Six edible wild fruits as potential antioxidant additives or nutritional supplements. *Plant Foods Hum. Nutr.* **2010**, *65*, 121–129.
5. Pinacho, R.; Yolanda Caverro, R.; Astiasarán, I.; Ansorena, D.; Calvo, M.I. Phenolic compounds of blackthorn (*Prunus spinosa* L.) and influence of *in vitro* digestion on their antioxidant capacity. *J. Funct. Foods* **2015**, *19*, 49–62.
6. Pandey, K.B.; Rizvi, S.I. Plant polyphenols as dietary antioxidants in human health and disease. *Oxid. Med. Cell Longev.* **2009**, *2*, 270–278.
7. Vanzo, A.; Vrhovsek, U.; Tramer, F.; Mattivi, F.; Passamonti, S. Exceptionally fast uptake and metabolism of cyanidin 3-glucoside by rat kidneys and liver. *J. Nat. Prod.* **2011**, *74*, 1049–1054.
8. Marchelak, A.; Owczarek, A.; Matczak, M.; Pawlak, A.; Kolodziejczyk-Czepas, J.; Nowak, P.; Olszewska, M.A. Bioactivity potential of *Prunus spinosa* L. flower extracts: Phytochemical profiling, cellular safety, pro-inflammatory enzymes inhibition and protective effects against oxidative stress *in vitro*. *Front. Pharmacol.* **2017**, *8*, 680–695.
9. Palaniappan, K.; Holley, R.A. Use of natural antimicrobials to increase antibiotic susceptibility of drug resistant bacteria. *Int. J. Food Microbiol.* **2010**, *140*, 164–168.
10. Taleb, H.; Maddocks, S.E.; Morris, R.K.; Kanekanian, A.D. The antibacterial activity of date syrup polyphenols against *S. aureus* and *E. coli*. *Front. Microbiol.* **2016**, *7*, 1–9.
11. Singleton, V.; Ortoger, R.M.; Lamuela-Ravendo, R.M. Analysis of total phenols and others oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Methods Enzymol.* **1999**, *299*, 152–173.
12. International Conference on Harmonization of Technical Requirements for Registration of Pharmaceuticals for Human Use. Validation of Analytical Procedures: Text and Methodology, Q2 (R1), 2005. Available online: https://www.ema.europa.eu/en/documents/scientific-guideline/ich-q-2-r1-validation-analytical-procedure-s-text-methodology-step-5_en.pdf (accessed on June 1995).
13. Delgado-Adámez, J.; Fernández-León, M.F.; Velardo-Micharet, B.; González-Gómez, D. *In vitro* assays of the antibacterial and antioxidant activity of aqueous leaf extracts from different *Prunus salicina* Lindl. cultivars. *Food Chem. Toxicol.* **2012**, *50*, 2481–2486.
14. Frassinetti, S.; Gabriele, M.; Caltavuturo, L.; Longo, V.; Pucci, L. Antimutagenic and antioxidant activity of a selected lectin-free common bean (*Phaseolus vulgaris* L.) in two cell-based models. *Plant Foods Hum. Nutr.* **2015**, *70*, 35–41.
15. Wolfe, K.L.; Liu, R.H. Cellular antioxidant activity (CAA) assay for assessing antioxidants, foods, and dietary supplements. *J. Agric. Food Chem.* **2007**, *55*, 8896–8907.
16. Pozzo, L.; Vornoli, A.; Coppola, I.; Della Croce, C.M.; Giorgetti, L.; Gervasi, P.G.; Longo, V. Lipid, cholesterol and glucose metabolism gene expression in a high-fat-diet and low-dose-streptozotocin induced rat model. *Life Sci.* **2016**, *166*, 149–156.
17. Folch, J.; Lees, M.; Sloane Stanley, J.H. A simple method for the isolation and purification of total lipids from animal tissues. *J. Biol. Chem.* **1957**, *226*, 497–509.
18. Seljeskog, E.; Hervig, T.; Mansoor, M.A. A novel HPLC method for the measurement of thiobarbituric acid reactive substances (TBARS). A comparison with a commercially available kit. *Clin. Biochem.* **2006**, *39*, 947–954.
19. Terevinto, A.; Ramos, A.; Castroman, G.; Cabrera, M.C.; Saadoun, A. Oxidative status, *in vitro* iron-induced lipid oxidation and superoxide dismutase, catalase and glutathione peroxidase activities in rhea meat. *Meat Sci.* **2010**, *84*, 706–710.
20. Browne, R.W.; Armstrong, D. Reduced glutathione and glutathione disulfide. *Methods Mol. Biol.* **1998**, *108*, 347–352.
21. Kim, D.O.; Chun, O.K.; Kim, Y.J.; Moon, H.Y.; Lee, C.Y. Quantification of polyphenolics and their antioxidant capacity in fresh plums. *J. Agric. Food Chem.* **2003**, *51*, 6509–6515.
22. Erturk, Y.; Ercisli, Y.; Tosum, M. Physico-chemical characteristics of wild plum fruits (*Prunus spinosa* L.). *Int. J. Plant Prod.* **2009**, *3*, 89–92.

23. Varga, E.; Domokos, E.; Fogarasi, E.; Ștefănesu, R.; Fü-löp, I.; Croitoru, M.D.; Laczkó-Zöld, E. Polyphenolic compounds analysis and antioxidant activity in fruits of *Prunus spinosa* L. *Acta Pharm. Hung.* **2017**, *87*, 1–7.
24. Radovanović, B.C.; Anđelković, S.M.; Radovanović, A.B.; Anđelković, M.Z. Antioxidant and antimicrobial activity of polyphenol extracts from wild berry fruits grown in Southeast Serbia. *Trop. J. Pharm. Res.* **2013**, *12*, 813–819.
25. Baltas, N.; Parkyldiz, S.; Can, Z.; Dincer, B.; Kolayli, S. Biochemical properties of partially purified polyphenol oxidase and phenolic compounds of *Prunus spinosa* L. subsp. *Dasyphylla* as measured by HPLC-UV. *Int. J. Food Prop.* **2017**, *20*, 1377–1391.
26. Barbieri, R.; Coppo, E.; Marchese, A.; Daglia, M.; Sobarzo-Sánchez, E.; Nabavi, S.F.; Nabavi, S.M. Phytochemicals for human disease: An update on plant-derived compounds antibacterial activity. *Microbiol. Res.* **2017**, *196*, 44–68.
27. Hendra, R.; Ahmad, S.; Sukari, A.; Shukor, M.Y.; Oskoueian, E. Flavonoid analyses and antimicrobial activity of various parts of *Phaleria macrocarpa* (Scheff.) Boerl fruit. *Int. J. Mol. Sci.* **2011**, *12*, 3422–3431.
28. Saavedra, M.J.; Borges, A.; Dias, C.; Aires, A.; Bennett, R.N.; Rosa, E.S.; Simões, M. Antimicrobial activity of phenolics and glucosinolate hydrolysis products and their synergy with streptomycin against pathogenic bacteria. *Med. Chem.* **2010**, *6*, 174–183.
29. Asgary, S.; Naderi, G.H.; Askari, N. Protective effect of flavonoids against red blood cell hemolysis by free radicals. *Exp. Clin. Cardiol.* **2005**, *10*, 88–90.
30. Hatia, S.; Septembre-Malaterre, A.; Le Sage, F.; Badiou-Bénéteau, A.; Baret, P.; Payet, B.; Lefebvre d'hellencourt, C.; Gonthier, M.P. Evaluation of antioxidant properties of major dietary polyphenols and their protective effect on 3T3-L1 preadipocytes and red blood cells exposed to oxidative stress. *Free Radic. Res.* **2014**, *48*, 387–401.
31. Vornoli, A.; Pozzo, L.; Della Croce, C.M.; Gervasi, P.G.; Longo, V. Drug metabolism enzymes in a steatotic model of rat treated with a high fat diet and a low dose of streptozotocin. *Food Chem. Toxicol.* **2014**, *70*, 54–60.
32. Cheng, D.M.; Pogrebnyak, N.; Kuhn, P.; Poulev, A.; Watweman, C.; Rojas-Silva, P.; Johnson, W.D.; Raskin, I. Polyphenol-rich rutgers scarlet lettuce improves glucose metabolism and liver lipid accumulation in diet-induced obese C57BL/6 mice. *Nutrition* **2014**, *30*, 52–58.
33. Al-Rejaie, S.S.; Aleisa, A.M.; Sayed-Ahmed, M.M.; AL-Shabanah, O.A.; Abuhashish, H.M.; Ahmed, M.M.; Al-Hosaini, K.A.; Hafez, M.M. Protective effect of rutin on the antioxidant genes expression in hypercholesterolemic male Wistar rat. *BMC Complement. Altern. Med.* **2013**, *13*, 136–145.
34. Gandhi, G.R.; Jothi, G.; Antony, P.J.; Balakrishna, K.; Paulraj, M.G.; Ignacimuthu, S.; Stalin, A.; Al-Dhabi, N.A. Gallic acid attenuates high-fat diet fed-streptozotocin-induced insulin resistance via partial agonism of PPARgamma in experimental type 2 diabetic rats and enhances glucose uptake through translocation and activation of GLUT4 in PI3K/p-Akt signaling pathway. *Eur. J. Pharmacol.* **2014**, *745*, 201–216.
35. Ting, Y.; Chang, W.T.; Shiao, D.K.; Chou, P.H.; Wu, M.F.; Hsu, C.L. Antiobesity efficacy of quercetin-rich supplement on diet-induced obese rats: Effects on body composition, serum lipid profile, and gene expression. *J. Agric. Food Chem.* **2018**, *66*, 70–80.
36. Lee, O.H.; Lee, B.Y. Antioxidant and antimicrobial activities of individual and combined phenolics in *Olea europaea* leaf extract. *Bioresour. Technol.* **2010**, *101*, 3751–3754.
37. Tuzcu, Z.; Orhan, C.; Sahin, N.; Juturu, V.; Sahin, K. Cinnamon polyphenol extract inhibits hyperlipidemia and inflammation by modulation of transcription factors in high-fat diet-fed rats. *Oxid. Med. Cell Longev.* **2017**, *31*, 150–165.
38. Karimi-Khouzani, O.; Heidarian, E.; Amini, S.A. Anti-inflammatory and ameliorative effects of gallic acid on fluoxetine-induced oxidative stress and liver damage in rats. *Pharmacol. Rep.* **2017**, *69*, 830–835.
39. Winocur, G.; Greenwood, C.E. Studies of the effects of high fat diets on cognitive function in a rat model. *Neurobiol. Aging* **2005**, *26*, 46–49.
40. Charradi, K.; Mahmoudi, M.; Bedhiafi, T.; Kadri, S.; Elkahoui, S.; Limam, F.; Aouani, E. Dietary supplementation of grape seed and skin flour mitigates brain oxidative damage induced by a high-fat diet in rat: Gender dependency. *Biomed. Pharmacother.* **2017**, *87*, 519–526.
41. Nabavi, S.F.; Habtemariam, S.; Jafari, M.; Sureda, A.; Nabavi, S.M. Protective role of gallic acid on sodium fluoride induced oxidative stress in rat brain. *Bull. Environ. Contam. Toxicol.* **2012**, *89*, 73–77.
42. Nkpaa, K.W.; Onyeso, G.I. Rutin attenuates neurobehavioral deficits, oxidative stress,

- neuro-inflammation and apoptosis in fluoride treated rats. *Neurosci. Lett.* **2018**, *24*, 92–99.
43. Bais, S.; Kumari, R.; Prachar, Y. Ameliorative effect of trans-sinapic acid and its protective role in cerebral hypoxia in aluminium chloride induced dementia of Alzheimer's type. *CNS Neurol. Disord. Drug Targets* **2018**, *17*, 144–154.
44. Asano, T.; Matsuzaki, H.; Iwata, N.; Xuan, M.; Kamiuchi, S.; Hibino, Y.; Sakamoto, T.; Okazaki, M. Protective effects of ferulic acid against chronic cerebral hypoperfusion-induced swallowing dysfunction in rats. *Int. J. Mol. Sci.* **2017**, *18*, 550.



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).