

Supplementary Table 1. Ingredients and chemical composition of finisher diets administered to chicken belonging to the control group (CG) and chicken fed the dietary grape pomace supplementation of 2.5 % (EG1), 5 % (EG2) and 7 % (EG3).

| | Diet | | | |
|---|------|------|------|------|
| | CG | EG1 | EG2 | EG3 |
| Ingredients (%) | | | | |
| corn flour | 52.0 | 52.0 | 52.0 | 50.0 |
| soy flour | 32.0 | 32.0 | 32.0 | 32.0 |
| wheat flour | 8.50 | 6.00 | 3.50 | 3.50 |
| vegetable oil | 2.50 | 2.50 | 2.50 | 2.50 |
| gluten corn | 2.00 | 2.00 | 2.00 | 2.00 |
| GP | - | 2.50 | 5.00 | 7.00 |
| Minerals | 3.00 | 3.00 | 3.00 | 3.00 |
| Chemical Composition (%)¹ | | | | |
| Dry matter | 85.7 | 85.5 | 83.5 | 81.9 |
| crude protein | 24.3 | 24.1 | 24.5 | 25.1 |
| lipid | 6.53 | 6.52 | 6.53 | 6.55 |
| crude fiber | 5.25 | 5.22 | 6.04 | 6.76 |
| starch | 45.2 | 45.1 | 44.6 | 43.8 |
| ash | 6.81 | 6.77 | 7.20 | 7.60 |

¹ Data are reported on a dry matter (DM) basis. GP = grape pomace