

## Article

# Seasonal Variations in the Trace Elements and Mineral Profiles of the Bivalve Species, *Mytilus galloprovincialis*, *Chamelea gallina* and *Donax trunculus*, and Human Health Risk Assessment

Katya Peycheva <sup>1,\*</sup>, Veselina Panayotova <sup>1</sup>, Rositsa Stancheva <sup>1</sup>, Albena Merdzhanova <sup>1</sup>, Diana Dobрева <sup>1</sup>, Vincenzo Parrino <sup>2</sup>, Nicola Cicero <sup>3</sup>, Francesco Fazio <sup>4</sup> and Patrizia Licata <sup>4,\*</sup>

<sup>1</sup> Department of Chemistry, Medical University of Varna, 9002 Varna, Bulgaria

<sup>2</sup> Department of Chemical, Biological, Pharmaceutical and Environmental Sciences, University of Messina, 98168 Messina, Italy

<sup>3</sup> Department of Biomedical and Dental Sciences and Morphofunctional Imaging, University of Messina, 98168 Messina, Italy

<sup>4</sup> Department of Veterinary Sciences, Via Palatucci 13, Polo Universitario Annunziata, University of Messina, 98168 Messina, Italy

\* Correspondence: peytcheva@hotmail.com (K.P.); plicata@unime.it (P.L.)

**Abstract:** This study aimed to provide data on selected toxic (Cd, Pb and Ni), essential (Cr, Cu, Fe, Mn and Zn) and microelement (Na, K, Ca and Mg) concentrations in edible tissues of the Mediterranean mussel (*Mytilus galloprovincialis*), striped venus clam (*Chamelea gallina*) and the wedge clam (*Donax trunculus*). Samples were collected from the Black Sea (Bulgaria) four times over, a period of one year (2022). In comparison with the maximum permissible levels set by the EU and USFDA, all elemental concentration found in the bivalve species were lower than the prescribed limits. An estimation of the dietary metal intake through calculation of the target hazard quotients (THQ), hazard index (HI) and target risk (TR) was performed. The target hazard quotient (THQ) for individual metal and HI for combined metals were lower than 1, indicating no health risk for consumers due to the intake of either individual element or combined ones. The target risk value for toxic inorganic Pb and Cr was below  $10^{-6}$ , indicating no carcinogenic risk. According to these results, the consumption of these bivalve species is completely safe for human health.

**Keywords:** heavy metals; bivalves; target hazard quotient; hazard index; target risk

**Citation:** Peycheva, K.; Panayotova, V.; Stancheva, R.; Merdzhanova, A.; Dobрева, D.; Parrino, V.; Cicero, N.; Fazio, F.; Licata, P. Seasonal Variations in Trace Elements and Mineral Profiles of the Bivalve Species *Mytilus galloprovincialis*, *Chamelea gallina* and *Donax trunculus* and Human Health Risk Assessment. *Toxics* **2023**, *11*, 319. <https://doi.org/10.3390/toxics11040319>

Academic Editor: Joanna Burger

Received: 23 February 2023

Revised: 20 March 2023

Accepted: 27 March 2023

Published: 28 March 2023



**Copyright:** © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Over the past few decades, the consumption of shellfish has increased due to the potential beneficial properties [1–5]. Shellfish are excellent sources of proteins, fatty acids, vitamins (niacin, thiamine) and minerals (Ca, Mg, Fe, and Zn) [4]. Despite the numerous human health effects, various contaminants, such as heavy metals, could pose a risk to regular consumers [6]. All metals possess a negative impact on human health when their intake exceeds certain levels, but lead, cadmium and nickel are very toxic, even at low doses [7]. Other elements (such as Cu, Fe, Zn and Se) are essential for humans, but they may become toxic at high concentrations, as well [7,8]. This is the reason why it is of great importance to determine marine organisms' chemical composition, regarding the content of heavy metal, in order to evaluate the possible risks to human health [8].

Among mollusks, the Mediterranean mussel (*Mytilus galloprovincialis*), striped venus clam (*Chamelea gallina*) and the wedge clam (*Donax trunculus*) are the most harvested and cultivated species in the world [1,2]. *M. galloprovincialis* is widely distributed in the coastal

waters of the eastern Atlantic–Mediterranean region and is mainly farmed in the north-west coastal waters of Spain, Italy, Turkey and Bulgaria [9,10]. *C. gallina* is distributed throughout the entire Mediterranean, Black and Marmara Sea coasts, and is a filter-feeder, muddy sand species [1]. Black Sea *C. gallina* prefers greater depths of 5 to 18 m (rarely up to 25+ m) [11]. The benthic *D. trunculus* is an Atlantic–Mediterranean warm-temperate species distributed from Senegal to the northern Atlantic coast of France, the Mediterranean and the Black Sea, as well as along the Marmara Sea [11–13]. In the Bulgarian Black Sea, *D. trunculus* dominates the wave action upper infralittoral medium and fine sands, usually between 1 and 6.5 m (but is also observed from 0.9 to 9 m) [11].

The Black Sea is an enclosed sea isolated from the World Ocean and the large inflows of river water distinguish it from other sea basins [14]. One of the most polluted seas with higher degree of eutrophication in recent years in the world is the Black Sea [15]. The pollution of the Black Sea increases each year, which leads to a decline in fishery yield and has a serious impact on the tourist industry. The Mediterranean mussel (*Mytilus galloprovincialis*), striped venus clam (*Chamelea gallina*) and the wedge clam (*Donax trunculus*) are bivalve species of huge economic importance in the Black Sea region [15].

The aim of this study was to determine the seasonal dynamics of trace (Cd, Cr, Cu, Fe, Mn, Ni, Pb, Zn) and macro element (Ca, K, Mg, Na) content of three Black Sea bivalve species, *Donax trunculus*, *Mytilus galloprovincialis* and *Chamelea gallina*, over a period of one year, and to calculate the target hazard quotients (THQ), hazard index (HI) and target risk (TR) values for Pb and Cr as developed by the Environmental Protection Agency (EPA) in the US for the estimation of potential health risks associated with long-term exposure to chemical pollutants [16].

## 2. Materials and Methods

### 2.1. Sampling and Sample Treatment

The study was conducted four times during a one-year period (2022) on wild populations of the bivalves. The samples were collected from the area of Azalia resort zone at two points (43.24642° N 28.01972° E; 43.24781° N 28.02058° E), and from the area of Krapeç to the resort St. Konstantin and Elena at four points (43.587° N 28.609° E; 43.434° N 28.37° E; 43.246° N 28.02° E; 43.247° N 28.022° E). The *Donax trunculus* abundance was assessed, based on bottom trawling with a beam trawl in 1–3 m isobaths, while *Mytilus galloprovincialis* and *Chamelea gallina* within 1–8 m isobaths. Only clams with undamaged valves were studied. More than 2 kg of bivalves with comparable shell lengths were collected, placed into plastic bags and brought to the laboratory. The bivalves were cleaned, rinsed with Milli-Q water, brushed, shucked and soft tissues were removed with a Teflon knife and stored in polyethylene bags at −20 °C until analysis. Around two hundred specimens of each species were taken randomly for the determination of sample mean. The physical parameters of sampled shellfish were as follows (Table 1):

**Table 1.** Physical parameters of sampled shellfish (mean ± standard deviation).

	Spring	Summer	Autumn	Winter
Striped venus clam ( <i>C. gallina</i> )				
Total weight (g)	3.01 ± 0.12	3.15 ± 0.09	3.09 ± 0.02	3.32 ± 0.39
Soft tissue weight (g)	1.42 ± 0.68	1.46 ± 0.22	1.51 ± 0.26	1.54 ± 0.32
Total size (cm)	2.09 ± 0.34	2.11 ± 0.14	2.01 ± 0.84	2.31 ± 0.61
Mussel ( <i>M. galloprovincialis</i> )				
Total weight (g)	3.00 ± 0.59	3.06 ± 0.51	3.60 ± 0.29	3.90 ± 0.69
Soft tissue weight (g)	1.89 ± 0.12	1.76 ± 0.34	1.87 ± 0.05	1.95 ± 0.13
Total size (cm)	4.41 ± 1.52	4.61 ± 0.31	5.10 ± 0.9	4.20 ± 1.12
Wedge clam ( <i>D. trunculus</i> )				
Total weight (g)	4.02 ± 0.59	4.10 ± 0.49	4.32 ± 0.29	4.91 ± 0.05

Soft tissue weight (g)	1.56 ± 0.12	1.13 ± 0.03	0.89 ± 0.13	1.26 ± 0.13
Total size (cm)	2.9 ± 1.52	3.1 ± 0.19	3.3 ± 0.92	3.4 ± 1.11

## 2.2. Reagents and Standard Solutions

All solutions were prepared with analytical reagent grade chemicals and ultra-pure water (18 MΩ cm) generated by purified distilled water with a Millipore Milli-Q Gradient A-10 water purification system (Bedford, MA, USA). All the plastic and glassware were cleaned by soaking in 2 M HNO<sub>3</sub> for 48 h, and rinsed five times with distilled water, and then five times with deionized water prior to use. The calibration standard solutions were freshly prepared by dilution of Optima Family Multi-Element Standard, Matrix per Volume: 2% HNO<sub>3</sub>, and Multi-Element Calibration Standard 3, Matrix per Volume: 5% HNO<sub>3</sub> stock solution (Perkin Elmer®, Waltham, MA, USA).

## 2.3. Analytical Procedure

Wet digestions were performed in triplicate by weighing approximately 1.0 g of the tissues with a mixture of 8 mL HNO<sub>3</sub> (65% Merck, Suprapur, Darmstadt, Germany) and 2 mL H<sub>2</sub>O<sub>2</sub> (30–32% Optima™, for Ultra Trace Analysis, Fisher Chemical, Waltham, MA, USA) in a microwave closed-vessel digestion system MARS 6 (CEM Corporation, Matthews, NC, USA). The power system used provides continuous microwave emission at each level (Table 2). After digestion the samples were diluted to 25 mL in acid-washed polyethylene bottles with Milli-Q water, and stored prior to analysis. A blank digest was performed in the same way. The concentrations of Ca, K, Mg, Na, Cd, Cr, Cu, Fe, Mn, Ni, Pb, Zn in the samples were determined using ICP-OES Spectrometer (Optima 8000, Perkin Elmer, Waltham, MA, USA) with the following operating conditions: plasma gas flow—8 L/min; auxiliary gas flow—0.4 L/min; nebulizer gas flow—0.6 L/min; RF power—1500 watts; plasma view—axial/radial; nebulizer—concentric glass, MEINHARD® Type C.

**Table 2.** Microwave digestion system general and operational parameters.

Microwave Digestion System “MARS 6”, Acid Mixture		Step	Initial Power (W)	Time (min)	Final Power (W)	Fan
T °C (max)	200 °C	1	100	5	600	1
Pressure (max)	75 bar	2	600	5	600	1
Quartz vessels	EasyPrep™ Plus	3	600	5	800	1
Sample amount	1 g	4	800	15	800	1

## 2.4. Quality Assurance and Quality Control of the Method

Five sub-samples of each material were digested along with the blank sample, using the methods described above. All measurements were performed in triplicate for the samples and standard solutions. Metal contents were calculated and shown as mg/kg of wet weight. The accuracy of the procedure for the determination of trace metals in mollusks was tested using ERM-CE 278 k (mussel tissues from European Commission, joint research center, Belgium) certified reference material. The CRM was digested and analyzed in the same way as the analytical samples. The recovery values were between 86 and 103% for the individual elements.

## 2.5. Hazard Risk Estimation

### 2.5.1. Target Hazard Quotient (THQ)

The THQ, developed by USEPA [16], is used to assess the human health risk for non-carcinogenic elements in the local human population over a lifetime, in comparison with the reference oral dose (RfD) [17]. THQ was calculated as per USEPA Region III Risk Based Concentration Table [18], by using the following equation:

$$THQ = \frac{(M_c \cdot I_R \cdot 10^{-3} \cdot EF \cdot ED)}{RfD \cdot BW_a \cdot AT_n}$$

where  $M_c$  is the heavy metal concentration in bivalve species (mg/kg ww),  $I_R$  is the average daily consumption of shellfish (0.8 g/person/day) [19],  $EF$  is the exposure frequency (365 days/year),  $ED$  is the exposure duration (30 years or 10,950 days) for non-cancer risk as used by USEPA [18],  $RfD$  is the reference dose of individual metal (0.001 µg/g day for Cd, 0.04 µg/g day for Cu, 0.003 µg/g day for Cr, 0.009 µg/g day for Fe, 0.02 µg/g day for Ni, 0.004 mg/kg for Pb, 0.14 mg/kg for Mn and 0.3 µg/g day for Zn) [18],  $BW_a$  is an average adult body weight (70 kg average) and  $AT_n$  is the average exposure time for non-carcinogens [18]. Values of  $THQ$  below 1 show no harmful effect for human health.

### 2.5.2. Hazard Index (HI)

The hazard index from  $THQ$ s is expressed as the total of the hazard quotients [20]:

$$HI = THQ_{Cd} + THQ_{Cu} + THQ_{Cr} + THQ_{Fe} + THQ_{Ni} + THQ_{Pb} + THQ_{Mn} + THQ_{Zn}$$

### 2.5.3. Target Risk (TR)

Target cancer risk ( $TR$ ) indicates carcinogenic risks. The model for estimating  $TR$  is shown as follows:

$$TR = \frac{C \cdot C_R \cdot 10^{-3} \cdot CPS_o \cdot EF \cdot ED}{BW_a \cdot AT_c}$$

where  $C$  is the metal concentration in the mollusk species (mg/kg ww),  $C_R$  is the average daily consumption of shellfish (g/day) (0.8 g/person/day) [19],  $CPS_o$  is the carcinogenic potency slope, oral ( $Cr = 0.5$  mg/kg bw-day,  $Pb = 0.0085$  mg/kg bw-day),  $AT_c$  is the averaging time, carcinogens (day/years), and was calculated by multiplying exposure frequency in exposure duration over lifetime. The  $TR$  value for intake of  $Cr$ ,  $Ni$  and  $Pb$  was calculated to indicate the carcinogenic risk, since  $Cd$ ,  $Cu$ ,  $Fe$ ,  $Mn$  and  $Zn$  do not cause any carcinogenic hazards according to the USEPA. The acceptable values of  $TR$  are  $1 \times 10^{-6}$  according to the USEPA [18].

## 2.6. Statistical Analysis

All analyses were performed in 5 replicates and the results were expressed as the mean values  $\pm$  standard deviation (SD). The mean values of trace elements were compared by a one-way ANOVA, followed by a post hoc Tukey's test. Statistical significance was considered at  $p \leq 0.05$  (GraphPad Prism 5).

## 3. Results and Discussion

### 3.1. Trace Elements in Bivalve Species Tissues

Mean concentrations and standard deviations of toxic and essential elements of the three bivalve species from the region of the Black Sea Bulgaria are shown in Table 3. The summarized results of this study are expressed as means (mg/kg) of wet weight (ww).

In this study, significant differences in elemental content were recorded between different seasons (Table 3), especially between winter and summer samples. An increase in the concentration of some elements during cold months corresponds to the period of the high storage of energy reserves, before the reproduction period, as many authors reported [21,22].

Cd is a non-essential element for the human body and it is capable of producing chronic toxicity, present at minimal concentration of 1 mg/kg [22]. The concentration of Cd ranged between 0.22 mg/kg ww and 0.91 mg/kg ww for *Chamelea gallina*; between 0.28 mg/kg ww and 0.74 for *Mytilus galloprovincialis*; between 0.03 mg/kg ww and 0.09 for

*Donax trunculus*. The maximum Cd level permitted for bivalve mollusks intended for human consumption is 1.0 mg/kg ww according to the European Community and Bulgarian Food Regulation [23]. In the literature, cadmium concentration has been reported to range from 0.204 to 0.426 mg/kg ww for *C. gallina* [1], 0.024 to 0.159 mg/kg ww for *D. trunculus* [1], 0.292 to 0.970 mg/kg ww for *M. galloprovincialis* [2], all sampled from the Sea of Marmara, Turkey; between 0.17 and 0.64 mg/kg ww in wild and farmed *M. galloprovincialis* from Boka Kotorska Bay, south east Adriatic Sea [9]; between 0.34 and 1.71 mg/kg dw in whole soft tissue of mussels collected at 14 locations along the eastern Adriatic coast, Croatia [24]. Cadmium levels in the present study were in good agreement with the reported literature data and data from international organizations.

**Table 3.** Toxic and essential content (mg/kg ww) of the bivalve species (Black Sea, Bulgaria).

	Toxic Elements				Essential Elements			
	Cd	Pb	Ni	Zn	Cr	Cu	Fe	Mn
<i>Chamelea gallina</i>								
Spring	0.24 ± 0.01 <sup>c</sup>	0.03 ± 0.01 <sup>c</sup>	0.03 ± 0.01 <sup>b</sup>	8.68 ± 0.31 <sup>c</sup>	0.05 ± 0.03 <sup>b</sup>	1.34 ± 0.06 <sup>c</sup>	30.35 ± 2.83 <sup>b</sup>	0.69 ± 0.03 <sup>b</sup>
Summer	0.43 ± 0.08 <sup>b</sup>	0.11 ± 0.02 <sup>b</sup>	0.11 ± 0.09 <sup>b</sup>	13.96 ± 1.47 <sup>b</sup>	0.17 ± 0.05 <sup>b</sup>	1.89 ± 0.35 <sup>b</sup>	53.19 ± 18.27 <sup>a</sup>	1.42 ± 0.32 <sup>a</sup>
Autumn	0.22 ± 0.05 <sup>c</sup>	0.12 ± 0.02 <sup>b</sup>	0.13 ± 0.13 <sup>b</sup>	7.96 ± 3.41 <sup>c</sup>	0.07 ± 0.03 <sup>b</sup>	1.73 ± 0.51 <sup>b</sup>	16.62 ± 5.12 <sup>c</sup>	1.57 ± 0.19 <sup>a</sup>
Winter	0.91 ± 0.02 <sup>a</sup>	0.55 ± 0.03 <sup>a</sup>	0.51 ± 0.41 <sup>a</sup>	19.75 ± 0.38 <sup>a</sup>	1.53 ± 0.13 <sup>a</sup>	9.27 ± 0.34 <sup>a</sup>	48.76 ± 5.51 <sup>a</sup>	0.71 ± 0.02 <sup>b</sup>
<i>Mytilus galloprovincialis</i>								
Spring	0.28 ± 0.03 <sup>c</sup>	0.07 ± 0.02 <sup>b</sup>	0.04 ± 0.01 <sup>b</sup>	16.00 ± 3.75 <sup>b,c</sup>	0.01 ± 0.01 <sup>d</sup>	0.38 ± 0.05 <sup>d</sup>	13.98 ± 1.11 <sup>c</sup>	0.83 ± 0.04 <sup>b</sup>
Summer	0.44 ± 0.07 <sup>b</sup>	0.11 ± 0.02 <sup>b</sup>	0.15 ± 0.09 <sup>a,b</sup>	14.21 ± 1.38 <sup>c</sup>	0.17 ± 0.04 <sup>c</sup>	1.92 ± 0.32 <sup>b</sup>	53.91 ± 1.72 <sup>b</sup>	1.44 ± 0.30 <sup>b</sup>
Autumn	0.30 ± 0.03 <sup>b,c</sup>	0.11 ± 0.01 <sup>a,b</sup>	0.21 ± 0.10 <sup>a</sup>	24.97 ± 6.03 <sup>a</sup>	0.28 ± 0.06 <sup>b</sup>	0.90 ± 0.17 <sup>c</sup>	108.07 ± 7.33 <sup>a</sup>	3.27 ± 0.75 <sup>a</sup>
Winter	0.74 ± 0.06 <sup>a</sup>	0.41 ± 0.13 <sup>a</sup>	0.13 ± 0.03 <sup>a,b</sup>	21.41 ± 1.52 <sup>a,b</sup>	0.36 ± 0.05 <sup>a</sup>	2.72 ± 0.30 <sup>a</sup>	59.16 ± 3.54 <sup>b</sup>	0.97 ± 0.01 <sup>b</sup>
<i>Donax trunculus</i>								
Spring	0.04 ± 0.01 <sup>a,b</sup>	nd	nd	10.85 ± 1.07 <sup>b</sup>	nd	11.34 ± 2.71 <sup>c</sup>	13.92 ± 1.78 <sup>d</sup>	0.68 ± 0.06 <sup>a</sup>
Summer	0.04 ± 0.01 <sup>a,b</sup>	0.05 ± 0.01 <sup>b</sup>	nd	12.47 ± 0.79 <sup>b</sup>	0.11 ± 0.01 <sup>a</sup>	16.01 ± 1.29 <sup>b</sup>	39.78 ± 4.24 <sup>c</sup>	1.98 ± 0.73 <sup>a</sup>
Autumn	0.03 ± 0.02 <sup>b</sup>	0.05 ± 0.03 <sup>b</sup>	0.005 ± 0.008 <sup>a</sup>	7.91 ± 5.18 <sup>b</sup>	0.36 ± 0.43 <sup>a</sup>	14.13 ± 2.88 <sup>b,c</sup>	83.24 ± 1.63 <sup>b</sup>	2.47 ± 1.72 <sup>a</sup>
Winter	0.09 ± 0.05 <sup>a</sup>	0.17 ± 0.09 <sup>a</sup>	0.006 ± 0.131 <sup>a</sup>	27.46 ± 2.53 <sup>a</sup>	0.44 ± 0.10 <sup>a</sup>	26.53 ± 1.71 <sup>a</sup>	120.57 ± 13.52 <sup>a</sup>	0.37 ± 0.04 <sup>b</sup>

Legend: mean values ± standard deviation; values in a column not sharing a common superscript are significantly different between seasons ( $p < 0.05$ ); nd—not detected.

The lead concentration of *C. gallina* ranged from 0.02 mg/kg ww to 0.55 mg/kg ww; between 0.07 and 0.41 mg/kg ww for *M. galloprovincialis*; between 0.05 and 0.17 mg/kg ww for *Donax trunculus*. The maximum Pb concentration for all species was found in the winter season. According to the European Commission Regulation (1881/2006/EC) [23], the maximum acceptable concentration (MAC) of Pb in bivalve mollusks is 1.5 mg/kg ww. Özden et al. (2009) reported values of Pb up to 1.342 mg/kg ww for *C. gallina* [1], up to 0.916 mg/kg ww for *D. trunculus* [1], and between 0.292 and 0.970 mg/kg ww for *M. galloprovincialis* [2] from the Sea of Marmara. In their study of mussel tissues in the Eastern Black Sea, Turkey, Čevik et al. [25] found Pb levels between 5.9 and 45.9 µg/g. As a result of the Pearson correlation in the study, metal concentration in the tissues did not seem to depend on the mussel length and sampling site, according to the same authors [25]. Cohen et al. [26] reported values of 0.80 µg/g dw for the *M. galloprovincialis* species in three southern California coastal wetlands. Zhelyazkov et al. [6] declared the mean Pb levels of  $0.251 \pm 0.141$  for wild *M. galloprovincialis* from Varna Bay, Bulgaria. The results from the current study are within the MAC and the data in the literature.

Nickel normally occurs at very low levels in the environment and seafood is one of the major exposure paths. High amounts of Ni may lead to skin allergies, chronic bronchitis, reduced lung function, and cancer of the lung and nasal sinus. The minimum concentration was found in *D. trunculus* (0.005–0.006 mg/kg ww), while the other two species

are characterized with higher concentrations (0.003–0.51 mg/kg ww for *C. gallina* and 0.04–0.21 mg/kg ww for *M. galloprovincialis*). The maximum Ni level permitted for muscle meat of some marine fish is 0.5 mg/kg ww according to the Bulgarian Food Regulation [27]. Nickel concentration in the literature has been reported to range from 0.13 to 0.43 mg/kg ww in *M. galloprovincialis* from the Italian coast, West Adriatic Sea [28]; 67.8 to 1948 ng/g ww in blue mussels (*M. edulis* Complex) from the Baltic Sea [29]; 2194 to 6180 µg/kg dw in mussel samples from Durres and Vlora “hot spots”, Albania [30]; 0.673 to 1.618 mg/kg ww in *C. gallina* and 1.097 and 7.505 mg/kg ww in *D. trunculus* [1]. The concentrations of Ni in our study were below that reported in the literature and by various health organizations.

Cr, Cu, Fe, Mn and Zn are essential elements for humans and most of them occur naturally in many food sources. The content of Zn was different between species. The maximum Zn concentration was observed in the winter season for all species, following the order: *D. trunculus* (27.46 mg/kg ww) > *M. galloprovincialis* (21.41 mg/kg ww) > *C. gallina* (19.75 mg/kg ww). There was no difference among the species in the summer period. Özden et al. (2009) concluded that the concentrations of Zn were significantly higher in *D. trunculus* than in *C. gallina* ( $p < 0.05$ ) [1], similarly to other authors [4,31–33]. Concerning the Cu concentration of the species under analyses, it is with maximum value during the winter season (*D. trunculus* ( $26.53 \pm 1.71$  mg/kg); *C. gallina* ( $9.27 \pm 0.05$  mg/kg); *M. galloprovincialis* ( $2.72 \pm 0.30$  mg/kg)). These results are higher than the ones found by Özden et al. [1,2] and are in the range determined for *M. galloprovincialis* by other authors [24,25,32]. The Fe levels in bivalve species ranged from 13.78 mg/kg to 120.57 mg/kg for *D. trunculus*. There is no maximum permitted level for Fe in bivalve species according to European legislation [23], but the US Food and Drug Administration set 80 mg/kg ww as the limit [20]. The iron concentration found in the literature was in agreement with the data from this study [1,2,9,28]. Mn values in the literature vary between 41 and 59 µg/g ww in tissues of mussels from the Eastern Black Sea [25]; 0.865 and 11.306 mg/kg ww in wild *M. galloprovincialis* from Bosphorus of the Sea of Marmara [2]; 1.33 and 3.85 mg/kg w/w found in *M. galloprovincialis* at three locations in Boka Kotorska during four different seasons in 2015 [7]. There are no data regarding the maximum Mn level permitted according to the Bulgarian Food Codex [27], but the data from current study are clearly within those found in the literature.

### 3.2. Macro Elements

The macro elements studied were K, Na, Ca, and Mg (Table 4). Of all the elements, Na, K and Ca were the most abundant. The concentration range for these elements in the studied species was, in general, within the range reported by several authors [2,25,34]. Usually, seafood has lower contents of Na when compared to K, both in fish or bivalves from salt or fresh water [35].

Of all species, *C. gallina* displayed the highest levels of Mg ( $958.52 \pm 12.17$  mg/kg ww) during the autumn season, but still less than reported by Nekhoroshkov et al. [34] (1020–4320 µg/g ww for *M. galloprovincialis* along the South African coastline), and within the values reported by Özden et al. [2]. The minimum concentration of this element has been found for *Mytilus galloprovincialis* (341.51 mg/kgw).

Further, in winter, *Mytilus galloprovincialis* had the highest Ca, K, Mg and Na concentration. Such an increasing trend across the four seasons was not observed for the other species subject to this study.

**Table 4.** Concentration (mean  $\pm$  standard deviation) and nutritional contribution of 150 g serving portion for *Chamelea gallina*, *Mytilus galloprovincialis* and *Donax trunculus*, in terms of essential elements.

	DRI /mg/day/		<i>Chamelea gallina</i>			<i>Mytilus galloprovincialis</i>			<i>Donax trunculus</i>		
			mg/kg	EDI	%DRI	mg/kg	EDI	%DRI	mg/kg	EDI	%DRI
Ca	1000–1200 *	Sp	786.39 $\pm$ 91.04	118	9.8	1365.64 $\pm$ 30.16	205	17.1	1167.42 $\pm$ 37.18	175	14.6
		S	749.50 $\pm$ 96.75	112	9.4	588.27 $\pm$ 32.43	88	7.4	806.05 $\pm$ 37.53	121	10.1
		A	1759.14 $\pm$ 46.52	264	22.0	385.48 $\pm$ 37.08	58	4.8	1946.55 $\pm$ 37.08	292	24.3
		W	1173.00 $\pm$ 51.31	176	14.7	2282.43 $\pm$ 38.76	342	28.5	3905.96 $\pm$ 26.82	586	48.8
K	4700 *	Sp	1605.20 $\pm$ 66.03	241	5.1	2505.04 $\pm$ 19.54	376	8.0	2559.81 $\pm$ 58.16	384	8.2
		S	1527.33 $\pm$ 37.15	229	4.9	1581.99 $\pm$ 52.03	237	5.0	2820.50 $\pm$ 13.09	423	9.0
		A	2171.43 $\pm$ 29.29	326	6.9	1701.83 $\pm$ 66.07	255	5.4	1171.04 $\pm$ 17.67	176	3.7
		W	1891.00 $\pm$ 15.39	284	6.0	3346.48 $\pm$ 19.23	502	10.7	1702.34 $\pm$ 13.02	255	5.4
Mg	310–420 **	Sp	573.52 $\pm$ 31.95	86	20.5	341.51 $\pm$ 10.93	51	12.2	511.67 $\pm$ 12.30	77	18.3
		S	546.22 $\pm$ 36.59	82	19.5	388.29 $\pm$ 13.58	58	13.9	471.26 $\pm$ 13.77	71	16.8
		A	958.52 $\pm$ 12.17	144	34.2	426.98 $\pm$ 17.42	64	15.2	493.85 $\pm$ 12.95	74	17.6
		W	452.95 $\pm$ 14.73	68	16.2	773.45 $\pm$ 21.73	116	27.6	742.57 $\pm$ 15.13	111	26.5
Na	1200–1500 *	Sp	2089.37 $\pm$ 86.07	313	20.9	1835.89 $\pm$ 7.33	275	18.4	2223.96 $\pm$ 13.06	334	22.2
		S	1988.43 $\pm$ 66.81	298	19.9	1150.27 $\pm$ 64.09	173	11.5	2112.19 $\pm$ 14.63	317	21.1
		A	3804.41 $\pm$ 16.36	571	38.0	2019.48 $\pm$ 47.27	303	20.2	2668.44 $\pm$ 29.77	400	26.7
		W	2180.00 $\pm$ 0.96	327	21.8	3813.63 $\pm$ 18.36	572	38.1	2699.63 $\pm$ 14.76	405	27.0

Legend: Sp—spring; S—summer; A—autumn; W—winter; DRI (in bold)—recommended daily intake values [36], used in the calculation of % DRI; EDI—estimated daily intake; \* AI—adequate intake; \*\* RDA—recommended dietary allowances.

### 3.3. Nutritional Contribution of the Species in Terms of Essential Elements (Benefits)

The nutritional contribution (in DRI percent) of each essential element was calculated based on Dietary Reference Intakes (DRI) [36]. The estimated daily intake (EDI) for each element was calculated using the element average concentration (mg/kg) obtained for each mollusk, considering a meal of 150 g of bivalve [35].

Concerning Ca, *D. trunculus* is the bivalve that can provide the highest proportion (48.8% of the DRI) of the daily required intake of this element. *C. gallina* can contribute to the highest percent of Mg (34.2% of the DRI), while *M. galloprovincialis* contributes most to the DRI of Na (38.1% of the DRI) and K (10.7% of the DRI).

### 3.4. Potential Health Risk Assessment

The target hazard quotients for Cd, Cr, Cu, Fe, Mn, Ni, Pb and Zn hazard indexes and target risks for Cr and Pb, estimated through the consumption of the three bivalve species, are shown in Table 5.

THQ is an integrated risk index that compares the ingested amount of a contaminant with a standard reference dose [34]. The health risk assessment is done based on assumptions. The acceptable value for THQ equals to 1, according to the USEPA [18]. The elements Cd, Cr, Cu, Fe, Mn, Ni, Pb and Zn did not exceed 0.1 of THQ, and the THQ was less than 1 for all elements. Perošević et al. found that the  $THQ_{Fe}$  was between 0.014 and 0.044,  $THQ_{Zn}$  0.015–0.050,  $THQ_{Mn}$  0.003–0.011,  $THQ_{Cu}$  0.007–0.019,  $THQ_{Ni}$  0.004–0.024,  $THQ_{Pb}$  0.224–0.297,  $THQ_{Cr}$  0.017–0.071 and  $THQ_{Cd}$  0.051–0.187 for mussels *M. galloprovincialis* from Boka Kotorska Bay, Montenegro [7], and THQs in the range of 0.3–0.7 for the elements Cr, Co, Zn, and I for wild mussels along the South African coastline [34].

**Table 5.** Risk values (THQ, HI and TR) of each metal contaminant in bivalve species, Black Sea (Bulgaria).

	Target Hazard Quotient (THQ)								Hazard Index (HI)	Target Risk (TR)	
	Cd	Cr	Cu	Fe	Mn	Ni	Pb	Zn		Cr	Pb
Chamelea gallina											
Spring	0.0027	0.0002	0.0004	0.0005	0.0001	0.00002	0.0001	0.0003	0.004	$1.2 \times 10^{-7}$	$1.2 \times 10^{-9}$
Summer	0.0049	0.0007	0.0005	0.0009	0.0001	0.00009	0.00003	0.0005	0.008	$4.2 \times 10^{-7}$	$4.4 \times 10^{-9}$
Autumn	0.0025	0.0003	0.0005	0.0003	0.0001	0.00007	0.0001	0.0003	0.004	$1.8 \times 10^{-7}$	$7.7 \times 10^{-10}$
Winter	0.0129	0.0058	0.0026	0.0008	0.0001	0.00029	0.00018	0.0008	0.025	$3.7 \times 10^{-6}$	$2.3 \times 10^{-8}$
Mytilus galloprovincialis											
Spring	0.0032	0.0001	0.0001	0.0002	0.0001	0.00002	0.0002	0.0006	0.004	$3.5 \times 10^{-8}$	$2.9 \times 10^{-9}$
Summer	0.0050	0.0007	0.0005	0.0009	0.0001	0.00009	0.00004	0.0005	0.008	$4.3 \times 10^{-7}$	$4.7 \times 10^{-9}$
Autumn	0.0035	0.0011	0.0003	0.0018	0.0003	0.00012	0.0001	0.0010	0.008	$6.8 \times 10^{-7}$	$7.8 \times 10^{-10}$
Winter	0.0084	0.0014	0.0008	0.0010	0.0001	0.00007	0.0013	0.0008	0.014	$8.9 \times 10^{-7}$	$1.7 \times 10^{-8}$
Donax trunculus											
Spring	0.0004	-	0.0032	0.0002	0.00006	-	-	0.0004	0.004	-	-
Summer	0.0005	0.0004	0.0017	0.0006	0.00016	-	0.0002	0.0005	0.004	$2.7 \times 10^{-7}$	$2.2 \times 10^{-9}$
Autumn	0.0003	0.0014	0.0012	0.0014	0.00020	0.000003	0.0002	0.0003	0.005	$8.7 \times 10^{-7}$	$2.0 \times 10^{-9}$
Winter	0.0010	0.0017	0.0076	0.0020	0.00003	0.000003	0.0006	0.0010	0.014	$1.1 \times 10^{-6}$	$7.4 \times 10^{-9}$

The HI values, which evaluate an additive risk from different elements, were lower than 1 for each bivalve sample, which indicates that there is no concern for potential health effects.

The Environmental Protection Agency (EPA) defines TR as “the incremental probability of an individual to develop cancer over a lifetime, as a result of exposure to a potential carcinogen” [17]. The cancer slope factor (CSF) value ( $\mu\text{g kg}^{-1} \text{day}^{-1}$ ), according to USEPA, is only available for As, Pb and Cr [17]. A risk level of  $1 \times 10^{-6}$  has been considered the point of excess cancer risk, indicating 1 per 1,000,000 chance of getting cancer via consumption of shellfish toxic metals, estimated for 70 years [37]. The safe point for carcinogenic risks must be lower than this level [37]. The range of risks set by the EPA is  $1 \times 10^{-4}$  to  $1 \times 10^{-6}$ , and it is deemed unacceptable if the risks are surpassing  $1 \times 10^{-4}$ . A carcinogenic risk of  $1 \times 10^{-4}$  poses health hazards; therefore, if sufficiently large, health hazards need some sort of intervention and remediation [8]. The calculated values in this study are within the range of  $10^{-6}$ , suggesting that the intake of Cr and Pb by consumption of these bivalve species would not result in appreciable hazard risk of the human body. As a conclusion, the levels of Cd, Cr, Cu, Fe, Mn, Ni, Pb and Zn in mussels could be considered relatively safe for consumption according to the hazardous quotients and indexes.

#### 4. Conclusions

The toxic, essential and macro element concentrations in Mediterranean mussel (*Mytilus galloprovincialis*), striped venus clam (*Chamelea gallina*) and the wedge clam (*Donax trunculus*) from Bulgarian coast of the Black Sea, at different seasons, did not exceed the European Union and US Food and Drug Administration requirements, and are within the maximum acceptable limits set by those health organizations. In terms of the THQ and HI values of the elements, the continuous consumption of these shellfish species in this area may not create health risk over a prolonged period. The target risk value for Cr and Pb was below  $10^{-6}$ , indicating no carcinogenic risk. Based on the results from this study, the consumption of these bivalve species is safe for human health.

**Author Contributions:** Conceptualization, K.P., P.L. and F.F.; Investigation, K.P., V.P. (Vincenzo Parrino), R.S., D.D. and A.M.; Writing—original draft preparation, K.P. and V.P. (Veselina Panayotova); Methodology, K.P., V.P. (Veselina Panayotova) and N.C.; Data curation, K.P., V.P. (Vincenzo Parrino), R.S., A.M. and F.F.; Supervision, F.F.; Writing—review and editing, A.M., F.F. and N.C.;



Project administration, P.L. and A.M. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Conflicts of Interest:** The authors declare no conflicts of interest.

## References

1. Özden, Ö.; Erkan, N.; Deval, M.C. Trace mineral profiles of the bivalve species *Chamelia gallina* and *Donax trunculus*. *Food Chem.* **2009**, *113*, 222–226.
2. Özden, Ö.; Ulusoy, S.; Erkan, N. Study on the behavior of the trace metal and macro minerals in *Mytillus galloprovincialis* as a bioindicator species; the case of Marmara Sea, Turkey. *J. Verbr. Lebensm.* **2010**, *5*, 407–412.
3. Stanković, C.; Jović, M.; Milanov, R.; Joksimović, D. Trace elements concentrations (Zn, Cu, Pb, Cd, As and Hg) in the Mediterranean mussel (*Mytilus galloprovincialis*) and evaluation of mussel quality and possible human health risk from cultivated and wild sites of the southeastern Adriatic Sea, Montenegro. *J. Serb. Chem. Soc.* **2011**, *76*, 1725–1737.
4. Peycheva, K.; Panayotova, V.; Stancheva, R.; Makedonski, L.; Merdzhanova, A.; Cicero, N.; Camilleri, G.; Fazio, F. Trace elements and omega-3 fatty acids of black sea (Bulgaria) bivalve species *Mytilus galloprovincialis*, *Chamelea gallina* and *Donax trunculus*. Human health risk. *Nat. Prod. Res.* **2022**, *36*, 2735–2742.
5. Bat, L.; Şahin, F.; Öztekin, A.; Arici, E. Toxic Metals in Seven Commercial Fish from the Southern Black Sea: Toxic Risk Assessment of Eleven-Year Data Between 2009 and 2019. *Biol. Trace Elem. Res.* **2022**, *200*, 832–843.
6. Zhelyazkov, G.; Yankovska-Stefanova, T.; Mineva, E.; Stratev, D.; Vashin, I.; Dospatliev, L.; Valkova, E.; Popova, T. Risk assessment of some heavy metals in mussels (*Mytilus galloprovincialis*) and veined rapa whelks (*Rapana venosa*) for human health. *Mar. Pollut. Bull.* **2018**, *128*, 197–201.
7. Perošević, A.; Joksimović, D.; Đurović, D.; Milašević, I.; Radomirović, M.; Stanković, S. Human exposure to trace elements via consumption of mussels *Mytilus galloprovincialis* from Boka Kotorska Bay, Montenegro. *J. Trace Elem. Med. Biol.* **2018**, *50*, 554–559.
8. Mejdoub, Z.; Zaid, Y.; Hmimid, F.; Kabine, M. Assessment of metals bioaccumulation and bioavailability in mussels *Mytilus galloprovincialis* exposed to outfalls pollution in coastal areas of Casablanca. *J. Trace Elem. Med. Biol.* **2018**, *48*, 30–37.
9. Jović, M.; Stanković, S. Human exposure to trace metals and possible public health risks via consumption of mussels *Mytilus galloprovincialis* from the Adriatic coastal area. *Food Chem. Toxicol.* **2014**, *70*, 241–251.
10. Eurostat. Production from Aquaculture Excluding Hatcheries and Nurseries (from 2008 Onwards). 2020. Available online: <https://appsso.eurostat.ec.europa.eu/nui/submitViewTableAction.do> (accessed on 13 March 2023).
11. Gumus, M.R.; Todorova, V.R.; Panayotova, M.D. Recent Observations on the Size Structure of *Donax trunculus* Linnaeus, 1758 and *Chamelea gallina* (Linnaeus, 1758) in the Bulgarian Black Sea as Status Indicators of Commercially Exploited Shellfish under the Marine Strategy Framework Directive (MSFD). *Ecol. Balk* **2020**, *3*, 63–71.
12. Bayed, A.; Guillou, J. Contribution a l'e-tude des populations du genre Donax: La population de *Donax trunculus* L. (Mollusca, Bivalvia) de Mehdiia (Maroc). *Ann. Inst. Oceanogr.* **1985**, *61*, 139–147.
13. Mazé, R.A.; Laborda, A.J. Aspectos de la dinámica de población de *Donax trunculus* L. (Bivalvia: Donacidae) en la ría de El Barquero (Lugo, NW España). *Inv. Pesq.* **1988**, *52*, 299–312.
14. Panayotova, V.; Merdzhanova, A.; Dobrev, D.A.; Bratoeva, K.; Makedonski, L. Nutritional composition, bioactive compounds and health-beneficial properties of Black Sea shellfish. *J. IMAB* **2020**, *26*, 3293–3297.
15. Makedonski, L.; Peycheva, K.; Stancheva, M. Determination of some heavy metal of selected black sea fish species. *Food Control* **2017**, *72*, 313–318.
16. USEPA (United States Environmental Protection Agency). *Risk Assessment Guidance for Superfund I: Human Health Evaluation Manual*; EPA/540/1-89/002; Office of Emergency and Remedial Response: Washington, DC, USA. 1989. Available online: [https://www.epa.gov/sites/production/files/2015-09/documents/rags\\_a.pdf](https://www.epa.gov/sites/production/files/2015-09/documents/rags_a.pdf) (accessed on 22 October 2022).
17. EPA. *Integrated Risk Information System*; U.S. Environmental Protection Agency: Washington, DC, USA. 2016. Available online: <https://www.epa.gov/iris/> (accessed on 14 October 2016).
18. USEPA (United States Environmental Protection Agency). Risk-Based Concentration Table. 2010. Available online: <https://archive.epa.gov/region9/superfund/web/html/index-23.html> (accessed on 13 June 2022).
19. EAFA (Executive Agency of Fisheries and Aquaculture). *Annual Report for Fish and Aquaculture Consumption in Bulgarian Territory*; Ministry of Agriculture and Forestry: Sofia, Bulgaria, 2012. (In Bulgarian).
20. US FDA/CFR; ISSC. National Shellfish Sanitation Program. Guide for the Control of Molluscan Shellfish 2007. Guidance Documents Chapter II. Growing Areas: 04. Action Levels, Tolerances and Guidance Levels for Poisonous or Deleterious Substances in Seafood. 2007. Available online: [http://www.issc.org/client\\_resources/2007%20nssp%20guide/section%20iv%20chap%20ii%20.04.pdf](http://www.issc.org/client_resources/2007%20nssp%20guide/section%20iv%20chap%20ii%20.04.pdf) (accessed 31 on January 2023).

21. Rouane-Hacene, O.; Boutiba, Z.; Belhaouari, B.; Guibbolini-Sabatier, M.E.; Francour, P.; Risso-de Faverney, C. Seasonal assessment of biological indices bioaccumulation and bioavailability of heavy metals in mussels *Mytilus galloprovincialis* from Algerian west coast, applied to environmental monitoring. *Oceanologia* **2015**, *57*, 362–374.
22. Sarkar, T.; Alam, M.M.; Parvin, N.; Fardous, Z.; Chowdhury, A.Z.; Hossain, S.; Haque, M.E.; Biswas, N. Assessment of heavy metals contamination and human health risk in shrimp collected from different farms and rivers at Khulna-Satkhiraregion, Bangladesh. *Toxicol. Rep.* **2016**, *3*, 346–350.
23. European Commission. Commission Regulation (EC) No 1881/2006 of 19 December 2006 Setting Maximum Levels for Certain Contaminants in Foodstuff. 2006. Available online: <https://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2006:364:0005:0024:EN:PDF> (accessed on 15 January 2022).
24. Kljaković-Gaspić, Z.; Herceg-Romanić, S.; Kozul, D.; Veza, J. Biomonitoring of organochlorine compounds and trace metals along the Eastern Adriatic coast (Croatia) using *Mytilus galloprovincialis*. *Mar. Pollut. Bull.* **2010**, *60*, 1879–1889.
25. Çevik, U.; Damla, N.; Kobya, A.I.; Bulut, V.N.; Duran, C.; Dalgıç, G.; Bozacı, R. Assessment of metal element concentrations in mussel (*M. galloprovincialis*) in Eastern Black Sea, Turkey. *J. Hazard. Mater.* **2008**, *160*, 396–401.
26. Cohen, T.; Hee, S.S.Q.; Ambrose, R.F. Trace Metals in Fish and Invertebrates of Three California Coastal Wetlands. *Mar. Pollut. Bull.* **2001**, *42*, 224–232.
27. Anonymous. Regulation of setting maximum levels of certain contaminants in foodstuff. *Darjaven Vestn.* **2004**, *31*, 2–3. (In Bulgarian).
28. Desideri, D.; Meli, M.A.; Roselli, C. A biomonitoring study  $^{210}\text{Po}$  and heavy metals in marine organisms from the Adriatic Sea (Italy). *J. Radioanal. Nucl. Chem.* **2010**, *285*, 373–382.
29. Knopf, B.; Flieger, A.; Radermacher, G.; Rüdel, H.; Paulus, M.; Pirntke, U.; Koschorreck, J. Seasonal variability in metal and metalloid burdens of mussels: Using data from the German Environmental Specimen Bank to evaluate implications for long-term mussel monitoring programs. *Environ. Sci. Eur.* **2020**, *32*, 7.
30. Cullaj, A.; Lazo, P.; Duka, S. Heavy metals and metallone levels in mussel samples from the Albanian sea coast, MAP/Med Pol. In *Biological Effects Monitoring Programme—Achievements and Future Orientations, Proceedings of the Workshop, Alessandria, Italy, 20–21 December 2006*; MAP Technical Reports Serial No. 166; UNEP/MAP: Athens, Greece, 2007; pp. 141–151.
31. Usero, J.; Morillo, J.; Gracia, I. Heavy metal concentrations in molluscs from the Atlantic coast of southern Spain. *Chemosphere* **2005**, *59*, 1175–1181.
32. Cammilleri, G.; Galluzzo, P.; Pulvirenti, A.; Giangrosso, I.E.; Dico, G.M.L.; Montana, G.; Lampiasi, N.; Mobilia, M.A.; Lastra, A.; Vazzana, M.; et al. Toxic mineral elements in *Mytilus galloprovincialis* from Sicilian coasts (Southern Italy). *Nat. Prod. Res.* **2020**, *34*, 177–182.
33. Locatelli, C.; Fabbri, D.; Torsi, G. Heavy metal determinations in algae and clams and their possible employment for assessing the sea water quality criteria. *Ann. Chim.* **2001**, *91*, 425–434.
34. Nekhoroshkov, P.S.; Bezuidenhout, J.; Frontasyeva, M.V.; Zinikovskaia, I.I.; Yushin, N.S.; Vergel, K.N.; Petrik, L. Trace elements risk assessment for consumption of wild mussels along South Africa coastline. *J. Food Compos. Anal.* **2021**, *98*, 103825.
35. Afonso, C.; Lourenço, H.M.; Cardoso, C.; Bandarra, N.M.; Carvalho, M.L.; Castro, M.; Nunes, M.L. From fish chemical characterisation to the benefit-risk assessment—Part A. *Food Chem.* **2013**, *137*, 99–107.
36. IOM. *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids*; National Academic Press: Washington, DC, USA, 2005.
37. Alidadi, H.; Tavakoly Sany, S.B.; Zarif Garaati Oftadeh, B.; Mohamad, T.; Shamszade, H.; Fakhari, M. Health risk assessments of arsenic and toxic heavy metal exposure in drinking water in northeast Iran. *Environ. Health Prev. Med.* **2019**, *24*, 59.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.