Supplementary Materials

Enhancing Arid Mining Area Restoration with a Lignite-Based Superabsorbent Gel

Zhaojun Yang[1], Naeem Akram[2], Lei Zhou[3], Saman Khawaja[2], Yi Zhang[1], Jia Guo[4,*]

[1] College of Chemistry and Chemical Engineering, Central South University, Changsha 410083, P. R. China; 232311053@csu.edu.cn and yzhangcsu@csu.edu.cn

[2] School of Chemical Engneering, Minhaj University Lahore, (54000) Lahore, Pakistan. drnaeem.ce@mul.edu.pk and saman.ce@mul.edu.pk

[3] Urumqi Midong District Chemical Industry Park Administrative Committee, Urumqi 830019, China 18199134666@163.com

[4] Xinjiang Hami Santanghu Energy Development and Construction Co., Ltd; Hami 839207, China gj459824057@126.com

* Correspondence: gj459824057@126.com




Fig. S1. Images of dry and swollen superabsorbent gel




Fig. S2. Vegetation Restoration Status in Experimental Fields
(a)1 day; (b)1 month; (c)3 months;(d)5 months




Fig. S3. Plant Growth in the Experimental Field. (a) Ammopiptanthus mongolicus; (b) Corethrodendron scoparium; (c) Calligonum; (d) Suaeda salsa