

Lactate metabolism and microbiome composition are affected by nitrogen gas supply in continuous lactate-based chain elongation

Carlos A. Contreras-Dávila, Arielle Ali, Cees J.N. Buisman and David P.B.T.B. Strik*

Environmental Technology, Wageningen University & Research, Bornse Weilanden 9, 6708
WG, Wageningen, the Netherlands.

*Correspondence: david.strik@wur.nl

4 figures

- 1 Feed tank
- 2 Feeding pump
- 3 Liquid sampling ports
- 4 Reactor column
- 5 Inverted funnel
- 6 Gas sampling port
- 7 Gas flow meter
- 8 pH probe
- 9 pH controller
- 10 Acid pump
- 11 Recirculation pump
- 12 Acid bottle
- 13 Effluent tank
- 14 Nitrogen gas line




Figure S1. Schematic of upflow anaerobic reactor set-up.




Figure S2. Biomass appearance (A) before continuous gas supply (day 57.7); (B) during continuous gas supply at $12 \text{ mL}\cdot\text{min}^{-1}$ (day 84.7); (C) during continuous gas supply at $120 \text{ mL}\cdot\text{min}^{-1}$; (D) intermittent gas supply at HRT 2 days (day 140.8); (E) intermittent gas supply at HRT 1 day (day 171.7); and (F) without gas supply (day 219).




Figure S3. Carboxylates concentrations over operation time. (A) Metabolites concentrations, (B) residual lactate concentrations and L-lactate enantiomeric excess (ee). Shaded areas I-V indicate stabilized operation time periods.




Figure S4. Inoculum composition at genus level. Taxonomy was assigned based on SILVA 138 SSU Ref NR 99 database.