

Supplementary Materials

Layer Inoculation as a New Technology to Resist Volatile Fatty Acid Inhibition during Solid-State Anaerobic Digestion: Methane Yield Performance and Microbial Responses

Xingliang Gao ^{1,†}, Zhuowu Li ^{2,†}, Keqiang Zhang ^{2,3}, Dewang Kong ⁴, Wenxuan Gao ², Junfeng Liang ^{2,3}, Fuyuan Liu ^{1,*} and Lianzhu Du ^{2,3,*}

1 State Key Laboratory of Sheep Genetic Improvement and Healthy Production, Xinjiang Academy of Agricultural Reclamation Sciences, 221 Wuyi Road, Shihezi 832000, China; gaoxingliang2020@outlook.com

2 Agro-Environmental Protection Institute, Ministry of Agriculture and Rural Affairs, 31 Fukang Road, Tianjin 300191, China; li2227634950@163.com (Z.L.); kqzhang68@126.com (K.Z.); wenxuanguao@hotmail.com (W.G.); liangjunfeng2008@126.com (J.L.)

3 Key Laboratory of Low-Carbon Green Agriculture in North China, Ministry of Agriculture and Rural Affairs, Beijing 100193, China

4 Hangzhou Energy and Environmental Engineering Co., Ltd., 207 East Fengqi Road, Hangzhou 310020, China; hnkongdewang@126.com

* Correspondence: xjliufuyuan@sohu.com (F.L.); dulianzhu99@163.com (L.D.);

Tel.: +86-22-23616673 (L.D.)

Figure S1. Variations in SCOD and pH values during SS-AD

Figure S2. TAN concentration during SS-AD in reactors P, M, and L




Figure S1. Variations in SCOD and pH values during SS-AD




Figure S2. TAN concentration during SS-AD in reactors P, M, and L