Article

Composition of *Zingiber officinale* Roscoe (Ginger), Soil Properties and Soil Enzyme Activities Grown in Different Concentration of Mineral Fertilizers

Dilfuza Jabborova ^{1,*} , Ravish Choudhary ² , Abdulahat Azimov ¹, Zafarjon Jabbarov ³, Samy Selim ⁴ , Mohammed Abu-Elghait ⁵ , Said E. Desouky ⁵, Islam H. El Azab ⁶ , Amnah Mohammed Alsuhailani ⁷, Adel Khattab ⁸ and AlBaraa ElSaied ⁵

- ¹ Laboratory of Medicinal Plants Genetics and Biotechnology, Institute of Genetics and Plant Experimental Biology, Uzbekistan Academy of Sciences, Kibray 111208, Uzbekistan; ahat2021@yahoo.com
 - ² Division of Seed Science and Technology, ICAR-Indian Agricultural Research Institute, New Delhi 110012, India; ravianu1110@gmail.com
 - ³ Department of Soil Science, National University of Uzbekistan, Tashkent 100125, Uzbekistan; zafarjonjabbarov@gmail.com
 - ⁴ Department of Clinical Laboratory Sciences, College of Applied Medical Sciences, Jouf University, P.O. Box 2014, Sakaka 72388, Saudi Arabia; sabdulsalam@ju.edu.sa
 - ⁵ Department of Botany and Microbiology, Faculty of Science, Al-Azhar University, Cairo 11884, Egypt; dr_abuelghait@azhar.edu.eg (M.A.-E.); said.desouky@azhar.edu.eg (S.E.D.); albraa.mahmoud@azhar.edu.eg (A.E.)
 - ⁶ Food Science & Nutrition Department, College of Science, Taif University, P.O. Box 11099, Taif 21944, Saudi Arabia; i.helmy@tu.edu.sa
 - ⁷ Department of Physical Sport Science, College of Education, Princess Nourah bint Abdulrahman University, P.O. Box 84428, Riyadh 11671, Saudi Arabia; amalsuhailani@pnu.edu.sa
 - ⁸ Department of Botany and Microbiology, Faculty of Science, Cairo University, Giza 12613, Egypt; adelkhattab01@gmail.com
- * Correspondence: dilfuzajabborova@yahoo.com

Citation: Jabborova, D.; Choudhary, R.; Azimov, A.; Jabbarov, Z.; Selim, S.; Abu-Elghait, M.; Desouky, S.E.; Azab, I.H.E.; Alsuhailani, A.M.; Khattab, A.; et al. Composition of *Zingiber officinale* Roscoe (Ginger), Soil Properties and Soil Enzyme Activities Grown in Different Concentration of Mineral Fertilizers. *Horticulturae* **2022**, *8*, 43. <https://doi.org/10.3390/horticulturae8010043>

Academic Editors:
Sławomir Michałek and
Marzena Błazewicz-Woźniak

Received: 18 November 2021

Accepted: 23 December 2021

Published: 2 January 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Ginger is rich in different chemical compounds such as phenolic compounds, terpenes, polysaccharides, lipids, organic acids, minerals, and vitamins. The present study investigated the effect of mineral fertilizers on the content of mineral elements in the rhizomes of *Zingiber officinale* Roscoe, soil enzymes activity, and soil properties in Surkhandarya Region, Uzbekistan. To the best of our knowledge, the present study is the first in Uzbekistan to investigate the mineral elements of ginger rhizome inhabiting Termez district, Surkhandarya region. A Field experiment was conducted at the Surkhandarya experimental station research Institute. Four treatments have been studied (Control with no fertilizers (T-1), N₇₅P₅₀K₅₀ kg/ha (T-2), N₁₂₅P₁₀₀K₁₀₀ kg/ha (T-3) and N₁₀₀P₇₅K₇₅ + B₃Zn₆Fe₆ kg/ha (T-4)). Results showed that T-4 treatment significantly increased ginger rhizome K, Ca, P, Mg, Fe, Na, Mn, Zn, Si, Li, and V content as compared to all other treatments and control. T-3 treatment significantly increased Mo, Ga, and Ag content in comparison to other treatments. Soil enzymes showed a significant increase for all treatments against control, while T-4 treatment has recorded the highest enzyme activity in comparison to all other treatments in urease, invertase, and catalase content. Soil chemical properties have significantly changed for all treatments against the non-cultivated soil and the zero fertilizers plantation with variation among different treatments. Results showed that ginger root is rich in minerals and can be used as a great potential for nutritional supplements and soil enrichment. This study suggest that combination of macro-microelements have the potential to increase the content of mineral elements in the rhizomes of ginger in field conditions.

Keywords: ginger; rhizome nutrients; soil nutrients; catalase; urease; invertase

1. Introduction

During the last five decades, the face of agriculture has been changed because of new technologies in both developed and developing countries [1]. Spices are the main commodities that are useful to increase the agricultural economy of any country due to their high impact value and export potentialities [2]. India is the top country for spices supply in the global market, followed by several other countries, including Indonesia, Malaysia, Pakistan, Australia, Spain, Egypt, and Tanzania. Herbal plants are a major source of traditional medicine and modern medicine and play a major role in the world [3–7]. Ginger is one of the most popular, high valued spices and is well known in human communities of the world [8]. The wet and dry root of ginger is widely used in the medicine and food industry [9]. Among all the spices, ginger is an important cash crop helpful in improving the economic level and livelihood of many farmers and stakeholders, including ginger growers from several countries such as India, Indonesia, Nigeria, and Bangladesh [10].

Ginger (*Zingiber officinale* Roscoe) belongs to the *Zingiberaceae* family, which is a medicinal, herbaceous perennial spice indigenous cash crop native to warm tropical sub-tropical and humid climates, particularly Southeastern Asia, for that placid environment is suitable to flourish it [11]. Presently, India is the top ginger-growing country, followed by China, Japan, Jamaica, Indonesia, Fiji [12]. Ginger is a very much interested and economically important crop to all farmers, consumers, stakeholders, dieticians have attracted growing interest among both dietitians and consumers due to its large amount of uses, such as medicinal and high nutritional value ayurvedic medicine [13]. It is propagated through rhizome and can be grown up at an altitude of 1500 m with well-distributed rainfall [14]. Ginger rhizome crop has been commonly known as Ada, Adrak, Ingwer, and Zingiber in Asia since ancient times, for nearly 3000 years [15–20]. Ginger is an essential and valuable 1.2 m tall perennial herb such as a reed with erect, slender leaves annually grown along with one or more aerial leafy stems having up to 2 cm thick robust, fibrous, and fleshy rhizome with a striated texture. The ginger flesh color varies between red, white, and yellow, while the outer skin is brown in color [21]. Rain-fed and irrigated conditions along with well-drained soils, sandy loam, clay loam, red loam, or lateritic loam are most suitable for ginger cultivation [22]. It is consumed as a fresh and dried form which is very useful in both conventional and herbal medicine for the treatment of digestive tract, vomiting, motion sickness, headache, and nausea [23–25]. Due to antioxidant properties, ginger possesses several health benefits as it has edible oils and fats which protect them against oxidative rancidity [26]. It serves as an antitumorigenic, immunomodulatory effect, antimicrobial, antiviral agent, potent analgesic, a powerful stimulant, heating agent and also controls various diseases such as high blood cholesterol level, blood pressure, nausea and insomnia [20,27–29]. Ginger rhizome is a natural remedy for the treatment of migraine, headaches, and lower LDL cholesterol levels, and increases peristalsis due to its health-promoting properties [30]. As ginger has its medicinal, nutritional and therapeutic value, the demand for supply of ginger is increasing day by day, which is putting a heavy load on the market [31].

The ginger rhizome consists of a high amount of minerals such as phosphorus, potassium, and calcium, which is useful to control many physiological processes and also for health benefits in the human body [32,33]. Finding the best combination of fertilizers that enrich rhizome nutrients and improve soil quality is of high importance to fulfill the market need. Several studies have revealed that NPK fertilizers are important fertilizers for crop yield, production, and enhancement of whole agriculture production and regulate metabolic and regulatory activities of plants [34]. Few studies have examined the impact of mineral fertilizers on the mineral content of ginger rhizomes and the activity of enzymes in the soil.

The present study aims at investigating the effect of mineral fertilizers on macro, micro, and ultra-micro elements in ginger rhizomes. Evaluating the impact of the mineral fertilizers' application on soil chemical properties and enzyme activity is another major goal of the present study. We hypothesized that the combined macro-microelements

would facilitate beneficial effects on plant growth, plant nutrients, soil properties and soil enzyme activities.

2. Materials and Methods

2.1. Experimental Design

A field experiment was conducted to study the effect of mineral fertilizers on the mineral nutrients of ginger and soil properties. The experiment was carried out in randomized block design with five replications at the Surkhandarya scientific experimental station of the vegetable, melon crops and potato research Institute (N 37°13'44", E 67°16'34"), Uzbekistan (Figure 1). Experimental treatments included: Control with no fertilizers (T-1), N₇₅P₅₀K₅₀ kg/ha (T-2), N₁₂₅P₁₀₀K₁₀₀ kg/ha (T-3) and N₁₀₀P₇₅K₇₅ + B₃Zn₆Fe₆ kg/ha (T-4).

Figure 1. Location of the Surkhandarya scientific experimental station of vegetable, melon crops and potato research Institute, Termiz district, Uzbekistan (google.maps.com, (14 May 2020)).

2.2. Measurement of Plant Nutrients

Ginger rhizomes were harvested after 240 days of cultivation (Figure 2). For six hours, plant samples were disintegrated using hydrogen peroxide and nitric acid in a special microwave oven until they were reduced to their atomic elements for examination. Accurate volumetric measurements were made for the samples before being subjected to the nitric acid solution. The analysis was carried out using an optical emission spectrometer with an inductively coupled argon plasma (2100DV Santa Clara, CA, USA).

Figure 2. Field experiment plantation and rhizomes of ginger after harvest. (A) Ginger plant in the field after 20 days. (B) Fresh harvested ginger plants. (C) Ginger rhizomes after dryness.

2.3. Analysis of Soil Nutrients

Soil samples were collected from a non-cultivated soil (T-0) and from the other four treatments (T-1, T-2, T-3, and T-4). Carbon and organic matter contents of soil were determined according to the method of Tyurin modified by CINA0 (CINA0, 2003). Mobile compounds of phosphorus and potassium were determined by Machigin method modified by CINA0 [35]. Compounds were extracted by ammonium carbonate in the ratio of 1:20. Total nitrogen, phosphorus, and potassium contents were determined using wet combustion with sulfuric acid, and the measurement was conducted by photo-colorimeter and a flame photometer, respectively [36,37]. The salinity level of soil was determined by water extraction methods [38], with 1:20 ratio, and salinity levels were calculated as the dry mass of salts, including the cations and anions expressed as water dissolved salts.

2.4. Analysis of Soil Enzymes

Urease activity of soil were assayed using the method by [39]. 2.5 g soil sample were added with 0.5 mL of toluene for 15min. Then mixed and added to 2.5 mL of 10% urea and 5mL citrate buffer in incubator at 38 °C for 24 h. Then with 38 °C distilled water diluted, filtered, then 4 mL of sodium phenate and 3 mL of sodium hypochlorite were added to 1 mL filtrate and diluted to 50 mL for 20 min. Urease activity of soil was measured at wavelengths of 578 nm using a spectrophotometer [40]. Urease activity was expressed as NH_4/g of soil/h.

Catalase activity was measured using H_2O_2 as a substrate, shaken for 20 min and the filtrate was titrated with 0.1 mol/L KMnO_4 ; catalase activity was expressed as ml 0.1 mol/L KMnO_4/g /h. Catalase activity of soil was measured at wavelengths of 480 nm using spectrophotometer [40].

Invertase activity was determined using sucrose as a substrate and incubation at 37 °C for 24 h, measuring the produced glucose with the colorimetric method, and invertase activity was expressed as $\text{mg glucose kg}^{-1}\text{h}^{-1}$. Invertase activity of soil was measured at wavelengths of 508 nm using spectrophotometer [40]. Invertase activity was taken as mg of glucose liberated/g soil.

2.5. Statistical Analyses

Data on plant macroelements, microelements and ultra-microelements, soil chemical properties, and enzyme activity have been analyzed using Excel 365 and Minitab 19 (Coventry, UK). Data were cleaned before running any statistical analyses. Data gaps and typos have been addressed. For each level, five replicates have been used to compute descriptive statistics, such as the mean standard deviation. All Box-Cox assumptions have been tested using the optimal λ method for non-normal dependent variables was applied whenever needed. Different comparisons were carried out using one-way analysis of variance (ANOVA) under fit General linear model in Minitab ver.19.

3. Results

The present study aimed at assessing the effect of cultivating ginger under different treatments on the mineral composition of the growing plants as well as on the soil chemical properties and enzymatic activity.

3.1. Measurement of Plant Nutrients

Different macroelements including K, Ca, P, Mg, and Na have been detected and results showed that ginger rhizome macronutrients have increased under all treatments against control treatment (no fertilizers) from one to two and half folds for some elements. Results showed that T-4 treatment ($\text{N}_{125}\text{P}_{100}\text{K}_{100} + \text{B}_3\text{Zn}_6\text{Fe}_6$ kg/ha) has increased all elements the most, followed by T-3 ($\text{N}_{100}\text{P}_{75}\text{K}_{75}$ kg/ha) and T-2 ($\text{N}_{75}\text{P}_{50}\text{K}_{50}$ kg/ha), respectively, except for calcium increase of T-2 over T-3 treatment. Macroelements results are presented in Table 1.

Different microelements including Fe, Mn, Zn, Cu, Cr, Mo, and Si have been detected in ginger rhizomes and results showed that all elements have increased under all treatment conditions except for chromium where only T-2 and T-4 treatments have produced almost the same chromium concentrations as of the control treatment (T-1) while T-3 showed significant lower concentration than control. All other elements showed significant increase with the highest ratio for the T-4 treatment with over 100% increase in comparison to the control treatment. Results of the microelements under different conditions is presented in Table 2.

Table 1. Mean $\pm$ Standard deviation and One-Way ANOVA results of the ginger rhizome macroelements content for the four treatments T-1 (No fertilizers), T-2 (N75P50K50), T-3 (N100P75K75), and T-4 (N₁₂₅P₁₀₀K₁₀₀ + B₃Zn₆Fe₆). Treatments that do not share superscript letters are significantly different.

Parameter	T-1	T-2	T-3	T-4
K	11,232 $\pm$ 4.31 ^A	13,322 $\pm$ 3.38 ^B	15,406 $\pm$ 1.85 ^C	27,474 $\pm$ 9.65 ^D
Ca	14,721 $\pm$ 2.39 ^A	24,570 $\pm$ 10 ^B	20,719 $\pm$ 4.15 ^C	35,885 $\pm$ 7.21 ^D
P	5209.6 $\pm$ 6 ^A	5544.2 $\pm$ 2.1 ^B	6592.4 $\pm$ 10.3 ^C	7542.7 $\pm$ 9.02 ^D
Mg	5534 $\pm$ 2.97 ^A	5774.3 $\pm$ 8.71 ^B	6374.3 $\pm$ 0.058 ^C	9055 $\pm$ 4.58 ^D
Na	1555.7 $\pm$ 0.972 ^A	1890 $\pm$ 0.998 ^B	1923.4 $\pm$ 2.78 ^C	3771.4 $\pm$ 3.46 ^D

Table 2. Mean $\pm$ Standard deviation and One-Way ANOVA results of the ginger rhizome microelements content for the four treatments T-1 (No fertilizers), T-2 (N75P50K50), T-3 (N100P75K75), and T-4 (N₁₂₅P₁₀₀K₁₀₀ + B₃Zn₆Fe₆). Treatments that do not share superscript letters are significantly different.

Parameter	T-1	T-2	T-3	T-4
Fe	375.61 $\pm$ 0.995 ^A	574.7 $\pm$ 7.9 ^B	602.35 $\pm$ 0.568 ^C	645.42 $\pm$ 3.06 ^D
Mn	69.146 $\pm$ 1.001 ^A	113.93 $\pm$ 2.04 ^B	134.18 $\pm$ 0.989 ^C	171.41 $\pm$ 1.58 ^D
Zn	3.371 $\pm$ 0.04 ^A	4.273 $\pm$ 0.115 ^B	5.262 $\pm$ 0.106 ^C	5.493 $\pm$ 0.005 ^D
Cu	2.779 $\pm$ 0.009 ^A	6.413 $\pm$ 0.09 ^B	5.242 $\pm$ 0.018 ^C	5.905 $\pm$ 0.005 ^D
Cr	1.663 $\pm$ 0.095 ^A	1.667 $\pm$ 0.015 ^A	1.452 $\pm$ 0.011 ^B	1.624 $\pm$ 0.005 ^A
Mo	0.16 $\pm$ 0.01 ^A	0.161 $\pm$ 0.009 ^A	0.261 $\pm$ 0.001 ^B	0.21 $\pm$ 0.01 ^C
Si	0.155 $\pm$ 0.011 ^A	0.213 $\pm$ 0.015 ^B	0.212 $\pm$ 0.012 ^B	0.309 $\pm$ 0.001 ^C

Different Ultra microelements including Li, Be, V, Co, Ga, Ge, Nb, Ag, Cd, In, Sn, Sb, Cs, Ta, W and Re have been investigated in ginger rhizomes. Results showed that indium, titanium, and rhenium were not detected under any treatments while tungsten, cesium, antimony, tin, beryllium, germanium, cadmium, neon remained non significantly different under all treatments. On the other side, lithium, gallium, vanadium, cobalt, and silver were significantly different under different conditions. T-4 has the highest increase in lithium, vanadium, cobalt while T-3 recorded the highest gallium and silver content. Results of the Ultra microelements under different conditions is presented in Table 3.

3.2. Analysis of Soil Nutrients

Soil chemical properties including total and active P₂O₅, total and active K₂O, NO₃, N, Humus, C, C:N ratio, CO₂, total HCO₃, chlorides, sulphates, calcium, magnesium have been determined for all treatments and for a non-cultivated soil. Results showed significant variation under different conditions. Carbon%, sulphates%, Humus%, N%, total and active P₂O₅ showed a constant increase for all treatments against T-0 and T-1 for the favor of T-4 followed by T-3 and T-2, respectively. C:N ratio, CO₂, and chlorides have significantly decreased under all treatments against both T-0 and T-1. Total and active K₂O, NO₃, HCO₃, calcium, and magnesium on the other side showed significant decrease under all treatment conditions against T-0 treatment. Total and active K₂O, NO₃, HCO₃, significantly increased under T2, 3, 4 treatments against T-1 treatment. Calcium and magnesium content under T-2 and T-3 conditions showed significant decrease against T-1 while under T-4 conditions

both calcium and magnesium showed significant increase against T-1. Results of the soil chemical properties under different conditions is presented in Table 4.

Table 3. Mean $\pm$ Standard deviation and One-Way ANOVA results of the ginger rhizome ultra-microelements content for the four treatments T-1 (No fertilizers), T-2 (N75P50K50), T-3 (N100P75K75), and T-4 (N₁₂₅P₁₀₀K₁₀₀ + B₃Zn₆Fe₆). Treatments that do not share superscript letters are significantly different.

Parameter	T-1	T-2	T-3	T-4
Li	0.297 $\pm$ 0.012 ^A	0.29 $\pm$ 0.01 ^A	0.367 $\pm$ 0.002 ^B	0.44 $\pm$ 0.001 ^C
Be	0.02 $\pm$ 0.01 ^A	0.012 $\pm$ 0.002 ^A	0.012 $\pm$ 0.001 ^A	0.014 $\pm$ 0.004 ^A
V	0.332 $\pm$ 0.001 ^A	0.489 $\pm$ 0.01 ^B	0.568 $\pm$ 0.002 ^C	0.625 $\pm$ 0.005 ^D
Co	0.119 $\pm$ 0.003 ^A	0.144 $\pm$ 0.001 ^B	0.135 $\pm$ 0.003 ^C	0.155 $\pm$ 0.002 ^D
Ga	0.313 $\pm$ 0.005 ^A	0.355 $\pm$ 0.005 ^B	0.367 $\pm$ 0.003 ^B	0.314 $\pm$ 0.017 ^A
Ge	0.002 $\pm$ 0.001 ^A	0.002 $\pm$ 0.001 ^A	0.002 $\pm$ 0.001 ^A	0.002 $\pm$ 0.001 ^A
Nb	0.001 $\pm$ 0.001 ^A	0.001 $\pm$ 0.001 ^A	0.001 $\pm$ 0.001 ^A	0.001 $\pm$ 0.001 ^A
Ag	0.015 $\pm$ 0.001 ^A	0.025 $\pm$ 0.002 ^{AB}	0.03 $\pm$ 0.01 ^B	0.019 $\pm$ 0.001 ^{AB}
Cd	0.008 $\pm$ 0.001 ^A	0.008 $\pm$ 0.001 ^A	0.008 $\pm$ 0.001 ^A	0.008 $\pm$ 0.001 ^A
In	0 $\pm$ 0 ^A	0 $\pm$ 0 ^A	0 $\pm$ 0 ^A	0 $\pm$ 0 ^A
Sn	0.033 $\pm$ 0 ^A	0.033 $\pm$ 0 ^A	0.033 $\pm$ 0 ^A	0.033 $\pm$ 0 ^A
Sb	0.013 $\pm$ 0.006 ^A	0.013 $\pm$ 0.006 ^A	0.013 $\pm$ 0.006 ^A	0.013 $\pm$ 0.006 ^A
Cs	0.003 $\pm$ 0.001 ^A	0.003 $\pm$ 0.001 ^A	0.003 $\pm$ 0.001 ^A	0.003 $\pm$ 0.001 ^A
Ta	0 $\pm$ 0 ^A	0 $\pm$ 0 ^A	0 $\pm$ 0 ^A	0 $\pm$ 0 ^A
W	0.003 $\pm$ 0.001 ^A	0.003 $\pm$ 0.001 ^A	0.003 $\pm$ 0.001 ^A	0.003 $\pm$ 0.001 ^A
Re	0 $\pm$ 0 ^A	0 $\pm$ 0 ^A	0 $\pm$ 0 ^A	0 $\pm$ 0 ^A

Table 4. Mean $\pm$ Standard deviation and One-Way ANOVA results of the soil chemical properties under the four treatments T-1 (No fertilizers), T-2 (N75P50K50), T-3 (N100P75K75), and T-4 (N₁₂₅P₁₀₀K₁₀₀ + B₃Zn₆Fe₆) and non-cultivated soil (T-0). Treatments that do not share superscript letters are significantly different.

Parameter	T-0	T-1	T-2	T-3	T-4
Active P ₂ O ₅ mg/kg	31.33 $\pm$ 0.58 ^A	35.3 $\pm$ 1 ^B	37.6 $\pm$ 0.1 ^{BC}	39.5 $\pm$ 1 ^C	42.6 $\pm$ 1.1 ^D
Active K ₂ O mg/kg	351.67 $\pm$ 0.76 ^D	120.57 $\pm$ 1.03 ^C	135.66 $\pm$ 1.88 ^B	140.69 $\pm$ 0.09 ^B	152.45 $\pm$ 1.05 ^A
N-NO ₃ mg/kg	89.13 $\pm$ 1.03 ^A	10.25 $\pm$ 0.05 ^E	28.483 $\pm$ 0.9 ^D	33.44 $\pm$ 0.06 ^C	35.12 $\pm$ 1.02 ^B
Total P ₂ O ₅ %	0.19 $\pm$ 0.02 ^A	0.19 $\pm$ 0.01 ^D	0.241 $\pm$ 0.001 ^C	0.21 $\pm$ 0.01 ^B	0.32 $\pm$ 0.02 ^B
Total K ₂ O%	1.87 $\pm$ 0.02 ^{CD}	0.837 $\pm$ 0.012 ^C	0.92 $\pm$ 0.02 ^B	0.96 $\pm$ 0.02 ^{BC}	0.98 $\pm$ 0.02 ^A
N%	0.09 $\pm$ 0 ^A	0.124 $\pm$ 0.002 ^D	0.128 $\pm$ 0.001 ^C	0.137 $\pm$ 0.012 ^{BC}	0.195 $\pm$ 0.002 ^B
Humus, %	1.7 $\pm$ 0 ^{CD}	1.996 $\pm$ 0.005 ^B	2.02 $\pm$ 0.01 ^B	2.013 $\pm$ 0.015 ^B	2.41 $\pm$ 0.01 ^A
C%	0.98 $\pm$ 0 ^{CD}	1.157 $\pm$ 0.004 ^B	1.172 $\pm$ 0.002 ^B	1.166 $\pm$ 0.006 ^B	1.396 $\pm$ 0.005 ^A
C/N	10.4 $\pm$ 0.1 ^D	9.367 $\pm$ 0.116 ^C	9.1 $\pm$ 0.3 ^B	8.9 $\pm$ 0.1 ^{BC}	7.733 $\pm$ 0.153 ^A
CO ₂ %	8.92 $\pm$ 0.07 ^A	6.45 $\pm$ 0.05 ^B	6.4 $\pm$ 0.01 ^{BC}	6.01 $\pm$ 0.01 ^C	6.013 $\pm$ 0.006 ^D
Total HCO ₃ %	0.02 $\pm$ 0 ^A	0.014 $\pm$ 0.004 ^B	0.013 $\pm$ 0.001 ^B	0.011 $\pm$ 0.002 ^C	0.021 $\pm$ 0.001 ^C
Cl%	0.09 $\pm$ 0 ^A	0.008 $\pm$ 0.001 ^B	0.006 $\pm$ 0.001 ^D	0.004 $\pm$ 0.001 ^D	0.005 $\pm$ 0.001 ^C
SO ₄ %	0.28 $\pm$ 0.01 ^C	1.003 $\pm$ 0.006 ^A	0.98 $\pm$ 0.01 ^B	0.743 $\pm$ 0.006 ^C	1.122 $\pm$ 0.001 ^C
Ca%	0.24 $\pm$ 0 ^A	0.14 $\pm$ 0.01 ^A	0.12 $\pm$ 0.01 ^B	0.133 $\pm$ 0.058 ^B	0.213 $\pm$ 0.006 ^C
Mg%	0.06 $\pm$ 0 ^A	0.032 $\pm$ 0.002 ^B	0.03 $\pm$ 0.01 ^C	0.013 $\pm$ 0.006 ^D	0.054 $\pm$ 0.001 ^C

3.3. Analysis of Soil Enzymes

Soil enzymes including catalase, invertase and urease have been detected for all treatments. Results showed significant variation ranged between 1.5 and almost five folds increase under different conditions against T-1 for the favor of T-4 followed by T-3 and T-2, respectively. Results of the soil enzyme activity under different conditions is presented in Figure 3.

Figure 3. Mean $\pm$ Standard deviation of the soil enzyme activity under the four treatments T-1 (No fertilizers), T-2 (N75P50K50), T-3 (N100P75K75), and T-4 (N₁₂₅P₁₀₀K₁₀₀ + B₃Zn₆Fe₆).

4. Discussion

Many reporters have studied macro-micro elements in plants [41–46]. For the first time, the study of the content of macro-elements and microelements of ginger in the soil climatic conditions of the Surkhandarya Region, Uzbekistan, revealed that N, P, K, Ca, Mg, Na, Mn, Fe, Zn, and Cu are high amount in the rhizome (Tables 1 and 2). Numerous studies have been conducted on analyzing the essential and non-essential metal content of ginger in Nigeria [47,48], India [49], Saudi Arabia [50,51], and Ethiopia [52]. Macro and micronutrient fertilizer NPK + BZnFe applications rate (100:75:75:3:6:6 kg/ha) showed a significant increase in ginger rhizome K, Ca, P, Mg, Na contents over the control (Table 1). Similar findings corroborate increased ginger nutrients such as N content, P content, and K content by the NPK applications rate (100:60:60 kg/h) [53]. Similar results were also observed by Yanthan et al. [54]. Thakur and Sharma [55] reported that mineral fertilizers significantly increased the nutrients in ginger. A large amount of minerals, especially K are beneficial for the successful cultivation of ginger in China, and an imbalance of N, P and K applications revealed low rhizome yield and inferior ginger quality [56,57]. Similar results were obtained in turmeric rhizome after using mineral fertilizer revealed increased nutrients, namely, Na, K, Mg, Ca, Fe in rhizome, which showed the significant effect in plants [58]. Likewise, Kavitha [59] revealed in their study in Kacholam crop and found that maximum rhizome yield obtained with extra nutrients. Earlier, Chhibba et al. [60] also used Fe fertilizer to increase the yield in fenugreek. Halder et al. [61] were observed better growth and yield in turmeric using Zn, Fe, and B, similarly in ginger by Roy et al. [62]. A positive effect of inorganic fertilizers on the uptake of nutrients in ginger was observed by Singh [25].

The purpose of the study was to examine the influence of mineral fertilizers on soil enzyme activity, agrochemical and chemical characteristics in the Termez area of Uzbekistan. According to a number of studies, mineral fertilizers can boost the nutritional content of plants and soils [18,43,45]. The soil's agrochemical qualities were improved by applying mineral fertilizer at various amounts. Soil with mineral fertilizer treatments had considerably lower levels of total P, total K, humus and active phosphorus and potassium than soil with the NPK + BZnFe application rate (100:75:75:3:6:6 kg/ha). Inorganic fertilizers have been shown to have a negative impact on soil nutrients, according to a number of publications [56–58]. Numerous researchers have found that inorganic fertilizers have an effect on soil agrochemical and chemical characteristics [63–65]. According to Dinesh et al. [66], chemical fertilizer management had a favorable effect on the soil's nitrate content in rainfed conditions (*Zingiber officinale* Rosc.). The NPK application rate (100:60:60 kg/h)

was found to enhance soil N, P, and K concentrations, according to Yanthan et al. [54]. High mineral fertilizer reduced the N, P, K, Ca, Mg, and Fe concentration in the soil according to Srinivasan et al. [67].

Some enzyme activities such as urease, alkaline phosphatase, and catalase are sensitive to changes in environmental conditions. The hydrolysis of organic matter having nitrogen promote urease activity and show a positive effect with available nitrogen in soil as urease activity is much sensitive for alkaline and saline environment so that it can suggest the soil quality [68,69]. The urease, invertase, and catalase activity in soil was greatly boosted by applying NPK + BZnFe at a rate of 100:75:75:3:6:6 kg/ha (Figure 3). According to Srinivasan et al., the NPK treatment rate of 75:50:50 kg/h was observed to increase soil urease activity by 27.0% [67]. Urease enzyme activity in soil was shown to be stimulated by the application of mineral fertilizers, according to Singh [70] and Allison et al. [71]. The NPK application rate (75:50:50 kg/ha) decreased urease activity compared to T-3 and T-4. A previous study by Dinesh et al. [66] showed a reduction in urease activity when the NPK treatment rate was 75:50:50 kg/h. Researchers found that mineral fertilizer boosted soil urease and catalase activity considerably [72,73]. Similarly, Liu et al. [74] revealed that organic fertilizer enhanced urease and catalase activity as compared to chemical fertilizer alone. Chen and Huang (2020) [75] reported that mineral fertilizers increase the invertase enzyme activity in soil. In addition, they observed a positive correlation between soil organic matter or available nitrogen in soil with urease, alkaline phosphatase, and catalase activity, while it was non-significant with soil salinity and pH. However, negative effect was observed with available phosphorus from catalase activity

5. Conclusions

Results of the present study have showed that the NPK + BZnFe application rate of (100:75:75:3:6:6 kg/ha) has significantly enhanced the content macro and micronutrients of the ginger rhizome. The combined application of the NPK + BZnFe (100:75:75:3:6:6 kg/ha) significantly improved soil nutrients and soil enzyme activity. These results suggest that the NPK + BZnFe applications rate of (100:75:75:3:6:6 kg/ha) can be used for producing larger yield with high-quality ginger plants as well as improving soil properties under soil-climate conditions. Results showed that ginger root is rich in minerals and can be used as a great potential for nutritional supplements and soil enrichment. This study suggest that combination of macro-microelements have the potential to increase the content of mineral elements in the rhizomes of ginger in field conditions.

Author Contributions: D.J. have contributed in developing the research ideas, analyzing the data, designed the experiments, conducting the research and writing and editing the manuscript; A.A. and Z.J. performed the methodology; R.C., S.S., A.K., A.M.A., I.H.E.A. and A.A. have contributed to manuscript writing; A.E. has conducted the statistical analyses and participated in manuscript writing and editing and visualization. S.S., A.K., M.A.-E. and S.E.D. interpreted the data and edited the paper; S.S., I.H.E.A. and A.M.A. have conceived the funding. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by the Ministry of Innovational Development of the Republic of Uzbekistan [project number A-ΦA-2019-33, 2019–2022].

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Acknowledgments: The authors extend their appreciation to the Deanship of Scientific Research at Taif University for funding this work through Researchers Supporting Project number (TURSP-2020/27), Taif University, Taif, Saudi Arabia. We thank colleagues at the Institute of Genetics and Plant Experimental Biology, Kibray, Tashkent region, Uzbekistan, for providing necessary support and laboratory and field facilities.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. FAO. *The Future of Food and Agriculture—Trends and Challenges*; FAO: Rome, Italy, 2017; p. 296.
2. Fouad, A.; Hegazy, A.E.; Azab, E.; Khojah, E.; Kapiel, T. Boosting of Antioxidants and Alkaloids in *Catharanthus roseus* Suspension Cultures Using Silver Nanoparticles with Expression of CrMPK3 and STR Genes. *Plants* **2021**, *10*, 2202. [\[CrossRef\]](#) [\[PubMed\]](#)
3. Egamberdieva, D.; Jabborova, D. Medicinal plants of Uzbekistan and their traditional uses. In *Vegetation of Central Asia and Environs*; Egamberdieva, D., Öztürk, M., Eds.; Springer Nature: Cham, Switzerland, 2018; pp. 211–237. [\[CrossRef\]](#)
4. Jabborova, D.; Wirth, S.; Halwani, M.; Ibrahim, M.F.M.; El Azab, I.H.; El-Mogy, M.M.; Elkelish, A. Growth Response of Ginger (*Zingiber officinale*), Its Physiological Properties and Soil Enzyme Activities after Biochar Application under Greenhouse Conditions. *Horticulturae* **2021**, *7*, 250. [\[CrossRef\]](#)
5. Jabborova, D.; Ma, H.; Bellingrath-Kimura, S.D.; Wirth, S. Impacts of biochar on basil (*Ocimum basilicum*) growth, root morphological traits, plant biochemical and physiological properties and soil enzymatic activities. *Sci. Hortic.* **2021**, *290*, 110518. [\[CrossRef\]](#)
6. Jabborova, D.; Sayyed, R.Z.; Azimova, A.; Jabbarov, Z.; Matchanov, A.; Baazeeme, A.; Sabagh, A.E.; Danish, S.; Datta, R. Impact of mineral fertilizers on mineral nutrients in the ginger rhizome and on soil enzymes activities and soil properties. *Saudi J. Biol. Sci.* **2021**, *28*, 5268–5274. [\[CrossRef\]](#) [\[PubMed\]](#)
7. Jabborova, D.; Enakiev, Y.; Sulaymanov, K.; Kadirova, D.; Ali, A.; Annapurna, K. Plant growth promoting bacteria *Bacillus subtilis* promote growth and physiological parameters of *Zingiber officinale* Roscoe. *Plant Sci. Today* **2021**, *8*, 66–71. [\[CrossRef\]](#)
8. Ahmed, H.I.S.; Badr, A.; El-Shazly, H.H.; Watson, L.; Fouad, A.S.; Ellmouni, F.Y. Molecular Phylogeny of *Trifolium* L. Section *Trifolium* with Reference to Chromosome Number and Subsections Delimitation. *Plants* **2021**, *10*, 1985. [\[CrossRef\]](#)
9. Jabborova, D.; Davranov, K.; Egamberdieva, D. Antibacterial, antifungal and antiviral properties of medicinal plants. In *Medically Important Plant Biomes: Source of Secondary Metabolites*; Egamberdieva, D., Tiezzi, A., Eds.; Springer Nature: Singapore, 2019; pp. 51–65.
10. Choekwan, S. Mysterious ginger: Enclaves of a boom crop in Thailand. *For. Soc.* **2017**, *1*, 144–153. [\[CrossRef\]](#)
11. Kumar, K.M.P.; Asish, G.R.; Sabu, M.; Balachandran, I. Significance of gingers (*Zingiberaceae*) in Indian system of medicine—Ayurveda: An overview. *Anc. Sci. Life* **2013**, *32*, 253–261. [\[CrossRef\]](#) [\[PubMed\]](#)
12. Asafa, R.F.; Akanbi, W.B. Growth and Rhizome Yield of Ginger (*Zingiber officinale* L.) as Influenced by Propagule Size and Nitrogen Levels in Ogbomoso, Southwestern Nigeria. *Int. Lett. Nat. Sci.* **2018**, *67*, 35–45. [\[CrossRef\]](#)
13. Schwertner, H.A.; Rio, D.C. High performance liquid chromatographic analysis of 6-gingerol, 8-gingerol, 10-gingerol and 6-shogaol in ginger containing dietary supplements, spices teas and beverages. *J. Chromatogr. B* **2007**, *856*, 41–47. [\[CrossRef\]](#)
14. Rafiq, A.; Mohammad, A.; Naeem, A. Productivity of ginger (*Zingiber officinale*) by amendment of vermicompost and biogas slurry in Saline soils. *Pak. J. Bot.* **2009**, *41*, 3107–3116.
15. Chamani, G.; Zarei, M.R.; Mehrabani, M.; Taghiabadi, Y. Evaluation of Effects of *Zingiber officinale* on salivation in rats. *Acta Med. Iran.* **2011**, *49*, 336–340. [\[PubMed\]](#)
16. Moustafa-Farag, M.; Mahmoud, A.; Arnao, M.B.; Sheteiwy, M.S.; Dafea, M.; Soltan, M.; Elkelish, A.; Hasanuzzaman, M.; Ai, S. Melatonin-Induced Water Stress Tolerance in Plants: Recent Advances. *Antioxidants* **2020**, *9*, 809. [\[CrossRef\]](#) [\[PubMed\]](#)
17. Gaber, A.; Alsanie, W.F.; Kumar, D.N.; Refat, M.S.; Saied, E.M. Novel Papaverine Metal Complexes with Potential Anticancer Activities. *Molecules* **2020**, *25*, 5447. [\[CrossRef\]](#)
18. Sharar, M.; Saied, E.M.; Rodriguez, M.C.; Arenz, C.; Montes-Bayón, M.; Linscheid, M.W. Elemental Labelling and Mass Spectrometry for the Specific Detection of Sulfenic Acid Groups in Model Peptides: A Proof of Concept. *Anal. Bioanal. Chem.* **2017**, *409*, 2015–2027. [\[CrossRef\]](#) [\[PubMed\]](#)
19. Gaber, A.; Refat, M.S.; Belal, A.A.M.; El-Deen, I.M.; Hassan, N.; Zakaria, R.; Alhomrani, M.; Alamri, A.S.; Alsanie, W.F.; Saied, M.E. New Mononuclear and Binuclear Cu(II), Co(II), Ni(II), and Zn(II) Thiosemicarbazone Complexes with Potential Biological Activity: Antimicrobial and Molecular Docking Study. *Molecules* **2021**, *26*, 2288. [\[CrossRef\]](#) [\[PubMed\]](#)
20. Newerli-Guz, J.; Pych, A. Antioxidant properties of ginger (*Zingiber officinale* Roscoe). *Zesz. Nauk. AMG* **2012**, *73*, 28–33. (In Polish)
21. Abdou, R.H.; Abdel-Daim, M.M. Alpha-Lipoic Acid Improves Acute Deltamethrin-Induced Toxicity in Rats. *Can. J. Physiol. Pharmacol.* **2014**, *92*, 773–779. [\[CrossRef\]](#)
22. NAIR, K.P.P. *The Agronomy and Economy of Turmeric and Ginger: The Invaluable Medicinal Spice Crops*; Elsevier: Amsterdam, The Netherlands, 2013.
23. Abdel-Daim, M.M.; El-Tawil, O.S.; Bungau, S.G.; Atanasov, A.G. Applications of Antioxidants in Metabolic Disorders and Degenerative Diseases: Mechanistic Approach. *Oxid. Med. Cell. Longev.* **2019**, *2019*, 4179676. [\[CrossRef\]](#) [\[PubMed\]](#)
24. Borrelli, F.; Capasso, R.; Pinto, A.; A Izzo, A. Inhibitory effect of ginger (*Zingiber officinale*) on rat ileal motility in vitro. *Life Sci.* **2004**, *74*, 2889–2896. [\[CrossRef\]](#)
25. Singh, M.; Masroor, A.M.; Naeem, K.M. Effect of nitrogen on growth, nutrient assimilation, essential oil content, yield and quality attributes in *Zingiber officinale* Rosc. *J. Saudi Soc. Agric. Sci.* **2016**, *15*, 171–178. [\[CrossRef\]](#)
26. Ippoushi, K.; Azuma, K.; Ito, H.; Horie, H.; Higashio, H. [6]-Gingerol inhibits nitric oxide synthesis in activated J774.1 mouse macrophages and prevents peroxynitrite-induced oxidation and nitration reactions. *Life Sci.* **2003**, *73*, 3427–3437. [\[CrossRef\]](#) [\[PubMed\]](#)
27. Reddy, A.C.; Lokesh, B.R. Studies on spice principles as antioxidants in the inhibition of lipid peroxidation of rat liver microsomes. *Mol. Cell. Biochem.* **1992**, *111*, 117–124. [\[PubMed\]](#)

28. Vimala, S.; Norhanom, A.W.; Yadav, M. Anti-tumor promoter activity in Malaysian ginger rhizome used in traditional medicine. *Br. J. Cancer*. **1999**, *80*, 110–116. [\[CrossRef\]](#)
29. Akoachere, J.-F.T.K.; Ndip, R.N.; Chenwi, E.B.; Ndip, L.M.; Njock, T.E.; Anong, D.N. Antibacterial effect of *Zingiber officinale* and *Garcinia kola* on respiratory tract pathogens. *East Afr. Med. J.* **2002**, *79*, 588–592. [\[CrossRef\]](#) [\[PubMed\]](#)
30. Ghosh, A.K.; Banerjee, S.; Mullick, H.I.; Banerjee, J. *Zingiber officinale*: A natural gold. *Int. J. Pharma Bio Sci.* **2011**, *2*, 283–294.
31. Baliga, M.S.; Haniadka, R.; Pereira, M.M.; D'Souza, J.J.; Pallaty, P.L.; Bhat, H.P.; Popuri, S. Update on the chemopreventive effects of ginger and its phytochemicals. *Crit. Rev. Food Sci. Nutr.* **2011**, *51*, 499–523. [\[CrossRef\]](#)
32. Akhiani, S.P.; Vishwakarma, S.L.; Goyal, R.K. Anti-diabetic activity of *Zingiber officinale* in streptozotocin-induced type I diabetic rats. *J. Pharm. Pharmacol.* **2011**, *56*, 101–105. [\[CrossRef\]](#)
33. Gupta, S.K.; Sharma, A. Medicinal properties of *Zingiber officinale* Roscoe—A Review. *IOSR J. Pharm. Biol. Sci.* **2014**, *9*, 124–129. [\[CrossRef\]](#)
34. Singh, M. Influence of Triacantanol, Nitrogen and Phosphorus on the Growth, Yield and Quality of Ginger (*Zingiber officinale* Rosc.) and Turmeric (*Curcuma longa* L.). Ph.D. Thesis, Aligarh Muslim University, Aligarh, India, 2009.
35. Tursunov, L. Soil Physics. In *Determination of the Mechanical Composition of Soils by the Kachinsky Method*; National University of Uzbekistan Press: Tashkent, Uzbekistan, 2010; pp. 1–358.
36. GOST 26261-84; Soils. *Methods for Determining Total Phosphorus and Total Potassium*; Publishing House of Standards: Minsk, Belarus, 2005.
37. GOST 26107-84; Soils. *Methods for Determination of Total Nitrogen*; Publishing House of Standards: Minsk, Belarus, 2005.
38. Pancu, M.; Gautheyrou, J. *Handbook of Soil Analysis Mineralogical, Organic and Inorganic Methods*; Springer: Berlin/Heidelberg, Germany, 2006; p. 800.
39. Guo, H.; Yao, J.; Cai, M.; Qian, Y.; Guo, Y.; Richnow, H.; Blake, R.E.; Doni, S.; Ceccanti, B. Effects of petroleum contamination on soil microbial numbers, metabolic activity and urease activity. *Chemosphere* **2012**, *87*, 1273–1280. [\[CrossRef\]](#) [\[PubMed\]](#)
40. Xaziev, F.X. *Methods of Soil Enzymology*; Publishing Nauka: Moscow, Russia, 2005; p. 252. ISBN 5020339407.
41. Egamberdieva, D.; Jabborova, D.; Berg, G. Synergistic interactions between *Bradyrhizobium japonicum* and the endophyte *Stenotrophomonas rhizophila* and their effects on growth and nodulation of soybean under salt stress. *Plant. Soil.* **2016**, *405*, 35–45. [\[CrossRef\]](#)
42. Egamberdieva, D.; Wirth, S.; Jabborova, D.; Räsänen, L.A.; Liao, H. Coordination between *Bradyrhizobium* and *Pseudomonas* alleviates salt stress in soybean through altering root system architecture. *J. Plant. Interact* **2017**, *12*, 100–107. [\[CrossRef\]](#)
43. Egamberdieva, D.; Jabborova, D.; Wirth, S.; Pravej, A.; Alyemeni, M.N.; Parvaiz, A. Interaction of magnesium with nitrogen and phosphorus modulates symbiotic performance of soybean with *Bradyrhizobium japonicum*, and its root architecture. *Front. Microbiol.* **2018**, *9*, 1000. [\[CrossRef\]](#)
44. Sarabekov, A.; Matchanov, A.; G'ofurov, M.B.; Xamidova, G.; Maulyanov, S.; Babaev, B.; Jabborova, D. Element analysis of *Helichrysum maracandicum* collected in different regions of Uzbekistan. *Plant. Cell Biotechnol. Mol. Biol.* **2021**, *22*, 53–59.
45. Jabborova, D.; Enakiev, Y.I.; Kakhramon, D.; Begmatov, S. Effect of coinoculation with *Bradyrhizobium japonicum* and *Pseudomonas putida* on root morph-architecture traits, nodulation and growth of soybean in response to phosphorus supply under hydroponic conditions. *Bulg. J. Agric. Sci.* **2018**, *24*, 1004–1011.
46. Jabborova, D.; Annapurna, K.; Al-Sadi, A.M.; Alharbi, S.A.; Datta, R.; Zuan, A.T.K. Biochar and Arbuscular mycorrhizal fungi mediated enhanced drought tolerance in Okra (*Abelmoschus esculentus*) plant growth, root morphological traits and physiological properties. *Saudi J. Biol. Sci.* **2021**, *28*, 5490–5499. [\[CrossRef\]](#) [\[PubMed\]](#)
47. Obiajunwa, E.I.; Adebajo, A.C.; Omobuwajo, O.R. Essential and trace element contents of some Nigerian medicinal plants. *J. Radioanal. Nucl. Chem.* **2002**, *252*, 473–476. [\[CrossRef\]](#)
48. Ogunwanda, I.A.; Olawore, N.O. Heavy trace metals and macronutrients status in herbal plants of Nigeria. *Food Chem.* **2004**, *85*, 67–71.
49. Aiwonegbe, A.E.; Ikhuoria, E.U. Levels of selected heavy metals in some Nigerian vegetables. *Trends App. Sci. Res.* **2007**, *2*, 76–79.
50. Devi, K.N.; Sarma, H.N.; Kumar, S. Estimation of essential and trace elements in some medicinal plants by PIXE and PIGE techniques. *Nucl. Instr. Methods Phys. Res. B* **2008**, *266*, 1605–1610. [\[CrossRef\]](#)
51. Al-Eed, M.A.; Assubaie, F.N.; El-Garawany, M.M.; El-Hamshary, H.; Eltayeb, Z.M. Determination of heavy metal levels in common spices. *J. Appl. Sci.* **2002**, *17*, 87–98.
52. Alwakeel, S.S. Microbial and heavy metals accumulation of herbal medicines. *Res. J. Microbiol.* **2008**, *3*, 683–691. [\[CrossRef\]](#)
53. Wagesho, Y.; Chandravanshi, B.S. Levels of essential and non-essential metals in ginger (*Zingiber officinale*) cultivated in Ethiopia. *SpringerPlus* **2015**, *4*, 107. [\[CrossRef\]](#) [\[PubMed\]](#)
54. Yanthan, L.; Singh, A.K.; Singh, V.B. Effect of INM on yield, quality, and uptake of N, P and K by ginger. *Agropedology* **2010**, *20*, 74–79.
55. Thakur, S.K.; Sharma, S.K. Response of ginger to N and P in sub-tropical zone of H.P. *Ind. J. Agric. Res.* **1997**, 3195–3198.
56. Majumdar, B.; Venkatesh, M.S.; Kumar, K. Effect of nitrogen and farmyard manure on yield and nutrient uptake of turmeric (*Curcuma longa*) and different forms of inorganic N build up in an acidic alfisol of Meghalaya. *Ind. J. Agric. Sci.* **2002**, *72*, 528–531.
57. Lujiu, L.; Fang, C.; Jiajia, W.; Dianli, Y.; Pingping, W. Ginger Yield and Quality Influenced by Potassium Fertilization. *Better Crops Plant Food* **2015**, *99*, 14–15.
58. Hassan, S.A.; Mijin, S.; Yusoff, U.K.; Ding, P.; Wahab, P.E. Nitrate, ascorbic acid, mineral and antioxidant activities of *cosmos caudatus* in response to organic and mineral-based fertilizer rates. *Molecules* **2012**, *17*, 7843–7853. [\[CrossRef\]](#)
59. Kavitha, P.R. Nutrient Management for Yield and Quality Improvement in Kacholam (*Kaempferia galanga* L.). Master's Thesis, Kerala Agricultural University, Thrissur, India, 2012.

60. Chhibba, I.M.; Nayyar, V.K.; Kanwar, J.S. Influence of mode and source of applied iron on fenugreek (*Trigonella corniculata* L.) in a Typic Ustochrept in Punjab, India. *Int. J. Agric. Biol.* **2007**, *9*, 254–256.
61. Halder, N.K.; Shill, N.C.; Siddiky, M.A.; Sarkar, J.; Gomes, R. Response of turmeric to zinc and boron fertilization. *J. Biol. Sci.* **2007**, *7*, 182–187. [[CrossRef](#)]
62. Roy, A.; Cahatterjee, R.; Hassan, A.; Maitra, S.K. Effect of Zn, Fe and B on growth, yield and nutrient content in leaf of ginger. *Ind. Cocoa Arecanut Spices J.* **1992**, *15*, 99–101.
63. McDowell, W.H.; Magill, A.H.; Aitkenhead-Peterson, J.A.; Aber, J.D.; Merriam, J.L.; Kaushal, S.S. Effects of chronic nitrogen amendment on dissolved organic matter and inorganic nitrogen in soil solution. *Forest Ecol. Manag.* **2004**, *196*, 29–41. [[CrossRef](#)]
64. Fang, H.J.; Yu, G.R.; Cheng, S.L.; Mo, J.M.; Yan, J.H.; Li, S. 13C abundance, water-soluble, and microbial biomass carbon as potential indicators of soil organic carbon dynamics in subtropical forests at different successional stages and subject to different nitrogen loads. *Plant. Soil.* **2009**, *320*, 243–254. [[CrossRef](#)]
65. Wang, Q.K.; Wang, S.L.; Liu, Y.X. Responses to N and P fertilization in a young Eucalyptus dunnii plantation: Microbial properties, enzyme activities and dissolved organic matter. *Appl. Soil Ecol.* **2008**, *40*, 484–490. [[CrossRef](#)]
66. Dinesh, R.; Srinivasan, V.; Hamza, S.; Manjusha, A.; Kumar, P.S. Short-term effects of nutrient management regimes on biochemical and microbial properties in soils under rainfed ginger (*Zingiber officinale* Rosc.). *Geoderma* **2012**, *173*, 192–198. [[CrossRef](#)]
67. Srinivasan, V.; Thankamani, C.K.; Dinesh, R.; Kandiannan, K.; Hamza, S.; Leela, N.K.; Zachariah, T.J. Variations in soil properties, rhizome yield, and quality as influenced by different nutrient management schedules in rainfed ginger. *Agric. Res.* **2019**, *8*, 218–230. [[CrossRef](#)]
68. Liang, Y.; Yang, Y.; Yang, C.; Shen, Q.; Zhou, J.; Yang, L. Soil enzymatic activity and growth of rice and barley as influenced by organic manure in an anthropogenic soil. *Geoderma* **2003**, *115*, 149–160. [[CrossRef](#)]
69. Zhang, T.; Wan, S.; Kang, Y.; Feng, H. Urease activity and its relationships to soil physiochemical properties in a highly saline-sodic soil. *J. Soil Sci. Plant. Nutr.* **2014**, *14*, 302–313. [[CrossRef](#)]
70. Singh, S.P. Nutrient supplementation through organic manures for growth and yield of ginger (*Zingiber officinale* Rose). *J. Eco-friendly Agric.* **2015**, *10*, 28–31.
71. Allison, V.J.; Condron, L.M.; Peltzer, D.A.; Richardson, S.J.; Turner, B.L. Changes in enzyme activities and soil microbial community composition along carbon and nutrient gradients at the Franz Josef chronosequence, New Zealand. *Soil Biol. Biochem.* **2007**, *39*, 1770–1781. [[CrossRef](#)]
72. Jabborova, D.; Choudhary, R.; Karunakaran, R.; Ercisli, S.; Ahlawat, J.; Sulaymanov, K.; Jabbarov, Z. The Chemical Element Composition of Turmeric Grown in Soil–Climate Conditions of Tashkent Region, Uzbekistan. *Plants* **2021**, *10*, 1426. [[CrossRef](#)] [[PubMed](#)]
73. Jabborova, D.; Sulaymanov, K.; Sayyed, R.Z.; Alotaibi, S.H.; Enakiev, Y.; Azimov, A.; Datta, R. Effect of Different Mineral Fertilizers on Quality of Turmeric and Soil Properties. *Sustainability* **2021**, *13*, 9437. [[CrossRef](#)]
74. Liu, G.; Zhang, X.; Wang, X.; Shao, H.; Yang, J.; Wang, X. Soil enzymes as indicators of saline soil fertility under various soil amendments. *Agric. Ecosyst. Environ.* **2017**, *237*, 274–279.
75. Chen, H.W.; Huang, L. Correlation between long-term fertilization and soil enzyme activity in the rhizosphere of halophytes. *Appl. Ecol. Environ. Res.* **2020**, *18*, 2669–2685. [[CrossRef](#)]