



Review

Biotechnologies and Strategies for Grapevine Improvement

Anca Butiuc-Keul ^{1,2} and Ana Coste ^{3,*}

¹ Department of Molecular Biology and Biotechnology, Faculty of Biology and Geology, Babes-Bolyai University, 400084 Cluj-Napoca, Romania

² Centre for Systems Biology, Biodiversity and Bioresource, Babes-Bolyai University, 400006 Cluj-Napoca, Romania

³ National Institute of Research and Development for Biological Sciences, Institute of Biological Research Cluj-Napoca, 400015 Cluj-Napoca, Romania

* Correspondence: ana.coste@icbcluj.ro

Abstract: Grapevine (*Vitis vinifera* subsp. *vinifera*) is one of the most widespread and economically important perennial fruit crops in the world. Viticulture has changed over the years in response to changing environmental conditions and market demands, triggering the development of new and improved varieties to ensure the crop's sustainability. The aim of this review is to provide a perspective on the recent developments in biotechnology and molecular biology and to establish the potential of these technologies for the genetic improvement of grapevine. The following aspects are discussed: (i) the importance of molecular marker-based methods for proper cultivar identification and how NGS-based high-throughput technologies have greatly benefited the development of genotyping techniques, trait mapping, and genomic selection; (ii) the recent advances in grapevine regeneration, genetic transformation, and genome editing, such as new breeding technology approaches for enhanced grapevine yield, quality improvement, and the selection of valuable varieties and cultivars. The specific problems and challenges linked to grapevine biotechnology, along with the importance of integrating classical and new technologies, are highlighted.

Keywords: biotechnology; crop improvement; genome editing; germplasm management; molecular markers; organogenesis; somatic embryogenesis



Citation: Butiuc-Keul, A.; Coste, A. Biotechnologies and Strategies for Grapevine Improvement. *Horticulturae* **2023**, *9*, 62. <https://doi.org/10.3390/horticulturae9010062>

Academic Editor: Ilaria Filippetti

Received: 26 November 2022

Revised: 22 December 2022

Accepted: 26 December 2022

Published: 4 January 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Domesticated grapevine (*Vitis vinifera* subsp. *vinifera*), one of the most economically important perennial fruit crops in the world [1,2], had a worldwide production of approximately 78 million tons in 2020 [3]. In 2020, the International Organization of Vine and Wine (OIV) estimated the world area under vines at 7.3 million ha [4]. The European Union (EU) had 3.2 million hectares of vines in 2020, equivalent to approximately 44% of the world's total wine-growing areas [5]. Over half of these areas were cultivated with red wine varieties. Most of the harvested grapes are used for wine production but also for consumption as fresh fruit, raisins, juices, vinegar, seed oils, and spirit drinks [6,7]. Grape extracts are also used as food additives, in cosmetics, and in the pharmaceutical industry, while some species are grown for ornamental purposes [8,9].

Grapevine cultivars are susceptible to biotic (bacteria, fungi, viruses, and insects) and abiotic stressors (drought, extreme temperatures, and salinity) that reduce both the yield and lifespan of vineyards, causing substantial economic losses to grape production and the wine industry [10]. Vegetative propagation and growth on rootstocks ensure the grapevine tolerance to phylloxera, an insect pest that almost destroyed European viticulture in the late nineteenth century [11]. Other pathogens, such as gray mold, downy, and powdery mildew, as well as bacterial diseases (e.g., black rot and Pierce's disease), cause serious yield losses [11,12]. Due to continuous climate change, viticulture faces outbreaks of these diseases, while thermal stress and severe dryness registered during

the last decade harmfully impacted the whole winemaking sector, especially in southern European winemaking regions [13–15]. Thus, a reduction in table quality vines and wine grape production was anticipated [13]. All these threats triggered the need to improve viticulture's sustainability by imposing new varieties versus traditional varieties [15–17].

Conventional breeding, based on the hybridization of valuable grapevine genotypes, is difficult due to the long life cycle and heterozygosity [18]. Moreover, numerous diseases require specific solutions found in the natural variations of the *Vitis* genus [19] but also in genetic improvements via molecular manipulation. Thus, in recent years, biotechnology research proved to be a powerful complement to conventional breeding methods playing an increasing role in the improvement of the existing grapevine varieties and rootstocks. The novel approaches include the development of molecular markers for fingerprinting, genetic mapping, genetic diversity assessments in populations, gene tagging for breeding purposes (marker-assisted selection), and gene cloning; these technologies aim to improve on the current plant transformation strategies and genetic editing to enhance disease tolerance and improve berry quality [9,17].

In the precision breeding of grapevine, only genetic elements encoding desirable traits are used; thus, the results are more predictable than conventional breeding. In the last decades, whole genome sequencing by next-generation sequencing (NGS) platforms and bioinformatics have allowed the rapid selection of plants for propagation and manipulation for different purposes.

Although modern techniques of precision breeding have taken a large scale and numerous valuable varieties have been obtained, genetically modified plants, however, are not easily accepted on the market. Edited plants were considered a solution to genetically transformed plants, mostly because the edited plants did not contain a transgene; thus, these plants were not subject to legal restrictions. Depending on the specific legislations in different countries, such as the USA, Argentina, Australia, and Brazil, even genetically modified organisms (GMOs) are accepted on the market because it is considered that edited plants do not contain foreign genes; thus, risk assessments are unnecessary. On the other hand, the European Union also considers that organisms obtained via mutagenesis, as well as those obtained via genome editing, are GMOs, as stated in the GMO European Directive 2001/18/EC [20] and should pass through the regulatory process of classical GMOs [21,22]. It is important to mention that point mutations induced by gene editing are practically impossible to distinguish from the natural ones or those induced by mutagenesis. This uncertainty could have negative consequences not only for agriculture but also for the economy; most researchers claim that organisms with only small genetic modifications without any foreign genetic material should not be considered GMOs [21,22]. Nevertheless, the latest technologies, i.e., genetic transformation and genome editing, are currently developing, the main limitation in grapevine being related to the complexity of some important agronomic traits [23].

In this review, we discuss the most recent achievements, specific problems, and challenges linked to grapevine biotechnology, along with the importance of integrating classical and new technologies for the genetic improvement of grapevine.

2. Grapevine Genetic Diversity and Molecular Markers Used in the Identification of Cultivars

Archaeological and archaeobotanical data showed that grapevine domestication began 6000–8000 years ago in the Transcaucasian region [24]. The dispersal of cultivated varieties from this primary center throughout the Near East and Europe relied upon (i) cultivars, the later clonal selection, and (ii) vegetative propagation [6]. In addition, secondary domestication events in other areas were also reported [25–28].

The genus *Vitis* [$2n = 38$] exhibits significant genetic diversity among cultivars, wild subspecies, and hybrids [29,30]. This is mainly due to their asexual reproduction, wide range of suitable planting, and frequent communication among grape accessions [31]. There are approximately 60–70 inter-fertile wild *Vitis* species widespread throughout Eurasia

and Northern America [32]. *Vitis vinifera* subsp. *sylvestris*, the only wild *Vitis* taxon native to Europe and the Near East, is considered the ancestor of almost 10,000 domesticated grapevine cultivars [31]. Moreover, approximately 1200 commercial grapevine cultivars are interspecific hybrids of the domesticated grapevines and other wild *Vitis* species [33]. Consequently, there is a huge number of named cultivars (15,000), including several synonyms (identical genotypes but different names) and homonyms (same names but different genotypes) [34]. The working group on *Vitis*, referring to the European cooperative programme for plant genetic resources (ECPGR), reports 27,000 accessions of grapes held only in European collections [35]. In spite of the passport data available for approximately 35,000 accessions from European countries, the problem of cultivar synonyms and the presence of duplications need to be solved [35]. Therefore, a proper identification system, cultivar registration and protection, seed certification, and plant variety rights are essential in grapevine germplasm management for breeding programs but also for economic interests, trade, and scientific knowledge [30,36].

2.1. Morphological Markers

Grapevine cultivars have traditionally been identified based on their morphological characteristics (ampelography) jointly provided by the International Organization of Vine and Wine (OIV), the International Union for the Protection of New Varieties of Plants (UPOV), and the International Plant Genetic Resources Institute (IPGRI) [37]. Although efficient for the assessment of qualitative traits, the application of morphological markers is limited in the evaluation of quantitative traits. Moreover, this method is time-consuming, requiring extensive field trials and the identification of closely related cultivars is difficult [38].

To complement the morphological identification of grapevine varieties and overcome classification errors and double designations, cytogenetic, biochemical, as well as DNA and RNA-based technologies were developed for the analysis of the existing grape germplasm diversity [30].

2.2. Cytological Markers

In earlier studies, following morphological markers, cytological markers (karyotypes, banding patterns, deletions, repeats, translocations, and inversions) were developed. The chromosome number and morphology, and the DNA amount and composition, are characterized with Giemsa staining, fluorochrome banding, silver staining, and fluorescence in situ hybridisation (FISH) [39]. However, only a few cytogenetic studies are available in grapevine, mainly due to the large number of small chromosomes and the difficulty of obtaining good chromosome preparations from the roots or anthers [30]. Sequential silver nitrate staining and FISH were used to study ribosomal DNA (rDNA) loci [39–41] to localize the retrotransposon Gret1 [40], BAC clones [42], and telomeric sequences [43]. However, FISH is considered a niche technique because it allows the analysis of only a few samples at a time, and its accuracy is highly dependent on excellent quality confocal microscopy and image analysis procedures [30]. However, with the advent of sequencing technologies (next-generation sequencing (NGS)) and the availability of high-quality de novo reference genomes for grapes, new horizons were opened for modern cytogenomics [30]. Thus, the physical mapping of DNA sequences on chromosomes facilitated comparative plant genomics, improving the genome and chromosome assemblies.

2.3. Molecular Markers

Molecular markers include protein-based markers (products of gene expression) and DNA-based markers derived from the direct analysis of polymorphisms in DNA sequences [44]. Depending on the detection method used, DNA markers are categorized as hybridization-based markers and polymerase chain reaction (PCR) and DNA sequence-dependent molecular markers.

Since their discovery, different molecular markers, such as (RFLPs) [45], random amplified polymorphic DNA (RAPD) [46], sequence characterized amplified regions (SCARs) [47,48], simple sequence repeats (SSRs) [31,49], inter-simple sequence repeats (ISSRs) [50,51], amplified fragment length polymorphisms (AFLPs) [52], single nucleotide polymorphisms (SNPs) [53], expressed sequence tags (ESTs) [54,55], and random amplified microsatellite polymorphisms (RAMPs) [56] have been widely used for the genetic diversity characterization of grapevine cultivars, molecular mapping, parentage analysis, clone's identification, and the detection of synonyms.

Among these, SSRs and SNPs have become the preferred markers for the characterization of grapevine genetic resources and varietal identification in germplasm collections [30]. Several studies reported nuclear and chloroplastidial SSR loci (nSSR and cpSSR) as useful to demonstrate the multiple origins of *V. vinifera* spp. *sativa* (cultivated grapevine), to reveal synonyms, homonyms, as well as inter and intra-specific genetic variations and phylogenetic relationships among wild and cultivated grapevines [27,57–62]. In recent years, SNP markers have also gained high popularity for evaluating allelic variations throughout grapevine genomes and dissecting complex traits via QTL (quantitative trait loci) for breeding programs [63,64]. SNPs are highly abundant across plant genomes and offer higher reproducibility than microsatellite data, facilitating the integration and interpretation of genotyping data throughout grape gene banks and databases [65,66]. The rise of NGS and resequencing techniques have facilitated the release of an extensive number of SNPs [31,67] and the development of reliable platforms, such as VitisGDB (the multi-functional database for grapevine breeding and genetics), for comparing and mining *Vitis* genomic information [68]. There are currently many reference *Vitis* databases, including simple genetic information or only descriptive information (i. e., the species name, country of origin, cultivar names, usage, etc.) Among these, we mention the International Variety Catalogue (VIVC) [69], Instituto de Ciencias de la Vid y del Vino [70], the European *Vitis* database [71], the Greek *Vitis* database [72], the Italian *Vitis* database [49], the molecular pathway database, the transcriptome database, grape sRNA atlas [68], etc.

In the last years, DNA marker systems used in grapevine characterization have evolved from interrogating small numbers of loci and individuals to tens of thousands of loci in studies of large populations [73]. Moreover, *Vitis* spp. genome sequencing has led to significant progress in the development of large-scale high-throughput DNA markers and the identification of QTL, allowing the confirmation of candidate genes and the development of breeding programs based on marker-assisted selection (MAS). The earliest method, restriction site-associated DNA [RAD] sequencing, was successfully applied to identify significant traits in elite grape cultivars [74,75]. Other sequencing-based methods, such as whole genome resequencing approaches, have been applied for the characterization of somaclonal variations within cultivars [76,77]. Moreover, RNA-seq has been widely applied in *Vitis vinifera* to study different aspects such as bud development [78], berry development and ripening [79], and the response to disease or pathogens [80]. Recently, a set of 2000 DNA low-cost marker panels transferrable across the entire *Vitis* genus was designed and implemented using rhAmpSeq (RNase H2 enzyme-dependent amplicon sequencing), which could be easily adapted for other taxa for ecological and evolutionary studies, QTL mapping, a genome-wide association study (GWAS), and molecular breeding [81].

In general, each technology provides results with a different resolution and accuracy, and the degree of detected genetic divergence depends on the marker system applied and the scope and type of plant samples used. The different applications of molecular markers in grapevine are presented in Table 1.

Table 1. Application of the most widely used molecular markers in grapevine.

Application	Molecular Marker *	References
Genetic diversity population structure	RAPD, ISSR, SSR, retrotransposon-based markers, SRAP, SNP, RAMP, REMAP, and IRAP	[49,50,82–88]
Cultivars, rootstocks, and clone identification	RAPD, cpSSR; SCAR; RFLP, SSR, SNP, AFLP, SAMPL, M-AFLP, MSAP, CAPS, IRAP, REMAP, SSAP, EST, and retrotransposon-based markers	[31,38,45,47,89–98]
Synonymies and homonymies clarification	RAPD and SSR	[99–101]
Origins of cultivated grapevine/phylogeographic patterns	cpSSR, SSR, and SNP	[27,53,57,60,62,102–104]
Genetic linkage maps	RFLP, SRAP, and SNP	[105,106]
Disease diagnostics	RFLP, SCAR, SSR, SSCP, ITS, and RNA sequencing	[48,80,107–110]
Transcriptome analysis and new gene discovery	EST	[54,55,111]
Genetic stability and somaclonal variation	RAPD, AFLP, SSR, CDDP, ISSR, and MSAP	[112–117]
QTL mapping	RAPD, CAPS, AFLP, SCAR, SSR, SNP, RAD sequencing, rhAmpSeq markers, SLAF-seq, and sequencing	[70,74,75,81,118–127]

* RFLP, restriction fragment length polymorphism; RAPD, randomly amplified polymorphic DNA; SCAR, sequence characterized amplified region; AFLP, amplified fragment length polymorphism; M-AFLP, microsatellite-amplified fragment length polymorphism; SSR, simple sequence repeat; cpSSR, chloroplast simple sequence repeat microsatellite; ISSR, inter simple sequence repeat; SAMPL, selectively amplified microsatellite polymorphic loci; SRAP, sequence-related amplified polymorphism; SSCP, single strand conformational polymorphism; SSAP, sequence-specific amplification polymorphism; CAPS, cleaved amplified polymorphic sequence; SNP, single nucleotide polymorphism; EST, expressed sequence tag; RAMP, randomly amplified microsatellite polymorphism; REMAP, retrotransposon-microsatellite amplified polymorphism; IRAP, interretrotransposon amplified polymorphism; SSAP, sequence-specific amplified polymorphism; ITS region sequences-ribosomal DNA internal transcribed spacer; CDDP, conserved DNA derived polymorphism; RAD sequencing, restriction site-associated DNA sequencing; rhAmpSeq, RNase H2 enzyme-dependent amplicon sequencing; SLAF-seq, specific length amplified fragment sequencing.

Although high-throughput sequencing technologies provide enormous potential to improve our way of understanding and accessing grapevine biodiversity, downstream bioinformatic analysis requires reliability to be ensured. There is not just one perfect array for all different research questions; therefore, the choice of the genotyping tool should be based on the purpose, sample size, resolution, accuracy, and budget available.

3. Grapevine Plant Regeneration Methods

Biotechnology research offers the potential to improve the yield and quality of grapes. The development of in vitro plant regeneration methods is essential to overcome the difficulties in conventional breeding studies, preserve and propagate valuable genotypes, as well as to increase genetic variability through genetic engineering (transgenic, cisgenic, and gene-edited plants) [128,129]. In addition, in vitro mass multiplication represents an alternative to the current greenhouses or outdoor repositories and allows the exposure of genotypes to in vitro-induced stresses (i.e., biotic and abiotic risks) [82].

To date, grapevine regeneration has been obtained with two fundamental pathways of propagation and regeneration through organogenesis and somatic embryogenesis [7,128]. Organogenesis is based on the ability of competent tissues to form whole plants directly from the meristematic regions of explants (direct organogenesis) or intervened with callus formation (indirect organogenesis). Indirect organogenesis induces somaclonal variation

involving both genetic and epigenetic changes in in vitro regenerated plants. In applied studies involving commercial scale multiplication or transgenic plants, the plants must be “true to type”, meaning high genetic uniformity of the regenerated plants [130,131]. Therefore, to demonstrate the genetic fidelity as well as the somaclonal variation of the in vitro plants, molecular marker approaches were applied. Somaclonal variation is an alternative source of genetic variability in horticultural crops with a narrow genetic base or difficult breeding [132].

Grapevine, similar to other woody species, has revealed a genotype and explant type dependent recalcitrance to in vitro regeneration techniques [17,129]. Many studies have reported that rootstock varieties reveal higher organogenesis and somatic embryogenesis potential than hybrids and varieties belonging to the *V. vinifera* species [133]. However, several other factors were reported to influence the efficiency of grapevine in vitro plant regeneration, such as culture medium composition, especially the type and concentrations of plant growth regulators (PGRs) [134,135]; explants’ developmental stage [136] and phyllotactic position [137]; light regime [138]; pH value [139], etc. Therefore, over the years, numerous studies were focused mainly on optimizing protocols for efficient regeneration across different grapevine genotypes. A thorough review of successful reports via organogenesis and somatic embryogenesis in several grapevine cultivars and rootstock species using different explants was published by Zhang et al. [17]. Further on, we briefly discuss some aspects related to the applications of in vitro regeneration systems in grapevine improvements.

The first attempts in grapevine biotechnologies started in the 1960s with in vitro propagation for mass production and healthy plant regeneration [23]. Over the years, different studies have focused on inflorescence culture [140,141] to study the mechanisms of floral induction [44], hairy root cultures [142–144] to study plant-pathogen interactions, the efficiency against nematodes [145,146], or phylloxera [147,148], and shoot tip culture combined with thermotherapy, chemotherapy, or cryotherapy to eliminate viruses [149–151].

Nowadays, the rapid technological advancements in molecular and cell biology include a wide range of new plant breeding technologies (NPBTs), which in association with the new genomic data available, offer the opportunity: (i) to develop new grapevine varieties with enhanced yields, quality, stress tolerance, and disease resistance through genetic manipulation [17] and (ii) to have a clearer picture of the molecular regulation of plant cells, tissue culture, and regeneration processes [128].

Due to their morphogenetic competence, embryogenic tissues are mostly preferred in genetic transformation studies for the application of new genomic techniques, such as cisgenesis and intragenesis, genome editing, and RNAi [7], and these could be a possible tool for virus and viroid elimination [152]. Moreover, somatic embryogenesis could prevent the development of chimerism, allowing the regeneration of genetically transformed embryos under selective culture conditions [153]. However, some genotypes have proven to be very recalcitrant to somatic embryogenesis; thus, genetic engineering techniques are difficult to apply [154].

The acquisition of embryogenic competence is related to different patterns of gene expression involving internal cell reprogramming leading to a reversion of the differentiation state [128,155,156]. Somatic embryos may be obtained through the development of an embryogenic callus (indirect embryogenesis) followed by the emergence of pro-embryogenic masses (PEMs), from which new somatic embryos are formed [157], or may occur directly from the explants without the callus developmental phase. In most cases, direct somatic embryogenesis is used for clonal propagation rather than indirect somatic embryogenesis, which is characterized by a high incidence of somaclonal variation [158]. The explants mostly employed for somatic embryogenesis induction are anthers, ovaries, leaves, petioles, tendrils, and nodal sections [159]. It was demonstrated that, amongst other factors, the type and concentrations of plant growth regulators (PGRs) play a crucial role in the induction of an embryogenic callus [160]. In particular, 2,4-dichlorophenoxyacetic acid (2,4-D) was reported as the most effective compound for the induction of somatic embryos [161,162].

Considering the importance of somatic embryogenesis in the genetic improvement of grapevine and the critical factors affecting its success, the development of efficient and reproducible genotype-specific protocols for all major grapevine, table grape cultivars, and rootstocks is required.

4. Somaclonal Variation

Plants cultivated in vitro could develop different modified characteristics and genetic variability due to somaclonal variation [163]. The genetic bases of somaclonal variation are gene mutations and also the rearrangements of chromosomes, karyotype changes [164–166], or epigenetic modifications driven by hyper or hypo-methylation [167]. Most of these are induced by oxidative stress [168], as shown in Figure 1.




Figure 1. Mechanism of somaclonal variation induced by oxidative stress in plants cultivated in vitro.

The genetic variation of micro-propagated plants is generally considered obstructive, and the loss of genetic fidelity was often observed. However, increased genetic variability has applications in the improvement of horticultural crops. The most important advantage of somaclonal variations is the reduced time and space for the screening of valuable genitors and traits than the crossing of perennial crops. Moreover, different somaclones could be used in breeding applications and genetic improvements with in vitro selection [132].

Several in vitro procedures imposing oxidative stress, such as protoplast culture, callus induction, or somatic embryogenesis, are more frequently followed by somaclonal variation due to the epigenetic changes in plant tissues [169]. Thus, polyploids were obtained with somatic embryogenesis in six Spanish *V. vinifera* cultivars [170]. High methylation was detected in two cultivars during somatic embryogenesis, and the AFLP markers showed higher variability in these plants, but the SSR patterns were similar in plants derived from somatic embryos and control plants [117]. Somaclonal variation in grapevine could also appear by several major genetic changes by the spatial arrangement of periclinal, mericlinal, and sectorial cell layers that are genetically different [171]. Periclinal chimeras originated from mutations in one of the cell layers, are stable plants producing axillary buds and have the same apical organization as the terminal meristem from which they were generated [172]. Somatic embryos in grapevine derive from a single cell; thus, the clonal characteristics are not transferred to the progeny, as shown with microsatellite markers in Pinot Meunier. Nevertheless, different accessions of Pinot Meunier have three alleles in the VVS2 locus and the accessions of Pinot Noir, Pinot Gris, and Pinot Blanc have two alleles in this locus, explained by a mutation in one of the two alleles in one of the cell layers in Pinot Meunier and its maintenance through the vegetative propagation of a periclinal chimera [173]. A similar variation was also observed in the VVS19 locus in the embryogenic callus induced from the cell layers of anther filaments in Primmitivo ist [173,174].

Other studies showed that the intracultivar variability within Pinot Noir and chardonnay are also due to mutations in the by a mutation in one of the two alleles in one of the cell layers in Pinot Meunier and its maintenance through vegetative propagation of a periclinal chimera [174,175]. Thus, somatic embryogenesis is a valuable tool to understand the origins, genetic structures, and relationships between ancient cultivars and should be considered before using them for Micropropagation, genetic conservation, or transformation [175].

5. Genome Sequencing and Applications

Grapevine is not only an important fruit crop but also a plant model for genetic studies due to its small genome size of 475–500 Mb and 38 chromosomes ($n = 19$). Most of the *Vitis* species are diploids, but there are also fertile interspecies hybrids [9]. The international grape genome program (IGGP) generated the first genome sequence for the Pinot Noir clone ENTAV 115 with Sanger and shotgun sequencing, which was important to understand the genome organization and the 19 linkage groups of *Vitis vinifera*. The genomic sequences of Pinot Noir clone ENTAV 115 (477.1 Mb) were assembled in 2093 metacontigs (approximately 28,352 genes and pseudogenes), of which 96.1% were assigned to linkage groups and candidate genes encoding relevant traits were predicted. In the NCBI taxonomy web portal for *V. vinifera*, there are 29,971 listed unique coding genes, and the information about these genes and the metabolic pathways in which they are involved are available at the TIGR site [176]. A consensus map was developed based on the genetic maps [177–181] and physical maps [182] previously developed. Other facilities are also available for different genomic and transcriptomic analyses, such as the grape BAC library from the French national resources center for plant genomics [183]; 14,000 transcripts from *V. vinifera* and 1700 transcripts from other *Vitis* species were released by The GeneChip® *Vitis vinifera* genome array (Affymetrix). An array-ready oligo set contains 14,562 probes of 70-mers representing grape gene transcripts released by Qiagen (<http://www1.qiagen.com>, accessed on 12 September 2022). The design of the grape oligo set was based on the sequence information from TIGR's grape gene index (<http://www.tigr.org/tdb/tgi>, accessed on 12 September 2022) [9].

Advancements in NGS technology allowed the development of genome-wide approaches for the genetic characterization of complex traits or for marker-assisted selection, such as genome-wide association studies (GWAS) or genomic selection (GS).

GWAS was used to understand the genetic bases of the important traits and to identify the polymorphic molecular markers associated with these traits [184] that could be further used in marker-assisted selection programs.

In contrast to the GWAS method, which identifies polymorphisms linked to the variations for selected traits, GS allows the prediction of a breeding value for the genotypes tested [185] based on large sets of markers. Thus, GS could significantly reduce costs for the marker-assisted selection of valuable variants by limiting the size and number of field experiments. Genotype-based prediction also allows selection in breeding schemes when the phenotyping of breeding candidates is impossible or difficult [186,187].

Unfortunately, GWAS and GS methods, which use genome-wide marker data for phenotype prediction, are difficult to use in highly heterozygous species such as grapevine [188]. Moreover, the efficiency of GWAS also depends on the genetic architecture of the trait; thus, the detection of molecular markers associated with polygenic traits depends on the size of the sample and the density of the molecular marker used [189]. In grapevine, there are no available valuable lines from complex breeding schemes; the breeders use highly diverse and heterozygous parental genotypes ($H = 0.76$) [190]. This is the result of strong inbreeding depression and vegetative propagation [190,191]. In addition, this panel of parental genotypes is also characterized by a low level of linkage disequilibrium (LD) between marker loci ($r^2 \sim 0.2$ at 5–10 Kb) [67,191]; most cultivars are interconnected by a series of first-degree relationships (i.e., Pinot Noir—Chardonnay—Gouais Blanc, Cabernet Franc—Merlot [192,193]), but the number of connected generations is relatively low [194,195]. However, GWAS and GS have become more relevant in grapevine since the number of molecular tools is con-

stantly increasing due to the high demand for new grapevine cultivars adapted to climate change [196,197].

6. Genetic Transformation

As conventional culture and the selection of new valuable varieties are time and resource-consuming, genetic transformation provided an alternative for developing new varieties with increased productivity, higher quality, and tolerance to different stress factors. Conventional breeding cannot provide resistance to diseases or pests to elite cultivars of *Vitis* [198]; thus, these cultivars are currently maintained through vegetative propagation [199] and require the frequent use of pesticides to control diseases [200]. To overcome these concerns, modern biotechnology proposed so-called precision breeding [201] and the genetic improvement of elite cultivars, which was previously known as cisgenic or intragenic improvements [202].

Unfortunately, the grapevine is considered a recalcitrant species in terms of genetic transformation due to several aspects, including (i) genes involved in grapevine transformation, (ii) vectors used for gene delivery and protocols for grapevine transformation, and (iii) protocols for transgenic plant regeneration [7].

The insertion of specific genes into plants with different methods and vectors was developed for over thirty years in perennial crops. In grapevine, physical and chemical delivery methods were tested over the years, and transgene delivery was mediated by *Agrobacterium* and viruses [23,203–206]. Several grapevine varieties were transformed with biolistic bombardment [203] and *Agrobacterium*-mediated transformation [204]. There are also several other methods of transformation, such as electroporation or protoplast transfection [23,205]. Viral vectors were used for the heterologous gene expression or the silencing of host genes (i.e., virus-induced gene silencing -VIGS) [206]. Cloning strategies and tools for the genetic engineering of grapevine were detailed and reviewed by [17,23].

Several grapevine infectious viruses, such as Vitivirus Grapevine Virus A (GVA) [207], the Closterovirus Grapevine leafroll-associated virus-2 (GLRaV-2) [208], and the Foveavirus Grapevine rupestris stem pitting-associated virus (GRSPaV) [209] were used for the silencing of PDS (phytoene desaturase) or ChII (subunit I of magnesium protoporphyrin IX chelatase), which was observed with the development of the albino phenotype [207,208]. A diagram showing the most important steps in genetic manipulation with transformation or editing of grapevine is shown in Figure 2.




Figure 2. Diagram of grapevine genetic manipulation with transformation or editing.

The success of a transformation is facilitated by the marker genes as selectable marker genes and reporter genes that ensure the rapid and accurate selection of modified cells from non-modified cells [210]. Some of the marker genes encode resistance to antibiotics

or herbicides, and the reporter genes encode proteins such as green fluorescence protein (GFP) or glucuronidase (GUS) [211]. Previously, most transgenic plants were obtained with vectors containing the *NPTII* gene as a selectable marker gene and GFP as a reporter gene, ensuring the visual selection of transformed plants [210]. For the elimination of the marker gene from transgenic plants, a complex system is used, which consists of two strains of *Agrobacterium*, one of them carrying a binary vector which contains the target gene, and another containing a binary vector with the selection marker genes as *codA/nptII*. The cells containing the target gene and the *codA/nptII* genes grow on the media with kanamycin, and then the cells containing the marker gene are eliminated with negative selection based on the *codA* function [212].

The analysis of the gene expression was performed using the *VvMybA1* gene involved in the anthocyanin biosynthesis pathway [213]. This gene allows the visual identification of transformed cells without kanamycin selection [210]. Unfortunately, the transformed plants of grapevines (Thompson Seedless) containing this visual reporter gene were not vigorous due to the intense pigmentation and curly and highly brittle leaves [213], and the viability was reduced. These inconveniences were mitigated by placing the *VvMybA1* gene under the control of tissue-specific and developmentally regulated promoters, for example, the promoter of the *Dc3* gene, expressed in late embryogenesis in carrots (*Daucus carota*), followed by the production of anthocyanin exclusively in embryos [214]. Moreover, being a visible marker, *VvMybA1* could be used for monitoring transgene expression in the whole plant. Transgene expression could also be monitored by reporter genes such as GUS and GFP [215]. Anthocyanin has a pink-to-red color which is easily discerned, and it is suitable for the analysis of gene expression in experiments involving the selection of hundreds of transgenic plants. [216]. The marker gene *VvMybA1*, placed under the control of the ubiquitin gene promoter, was used for the high-throughput analysis of genes and promoters [217].

The employment of appropriate promoters highly influences the development of valuable traits. Unfortunately, the progress of functional analyses in grapevine is reduced in comparison with *Arabidopsis* and rice, mainly due to the limited ability to obtain enough explants and difficulties in the analysis of the gene expression. The functional annotation of at least 18,725 genes in the grapevine allowed the evaluation of their promoters, as well as the gene expression under developmental and environmental factors. For proper uses, a detailed characterization of these promoters is required regarding their sequences and activation. First, constitutive viral promoters were used for most of the transgenes in grapevines [218] and then after, these promoters were replaced by ubiquitin gene promoters. The gene expression was monitored with an anthocyanin-based assay mentioned previously [214].

Precision breeding requires appropriate promoters for gene expression in particular tissues or certain stages of plant development. Thus, the discovery and characterization of potential promoters are extremely important for precision breeding in grapevine [218].

Successful genetic transformation also depends on the efficient regeneration of transgenic plants. Factors such as genotype, explant source, acceptor material, culture medium, bacterial strains, selectable markers, and selection methods affect the efficiency of plant transformation and regeneration. The genetic transformation of several grapevine cultivars, such as Thompson Seedless, Silcora, and Chardonnay, was obtained with shoot organogenesis from meristematic tissue [219–221]. The in vitro organogenesis of some grapevine cultivars and rootstocks was obtained from different types of explants, such as petioles, leaf internodes, and shoot apices [137,138,220–222]. One limitation of direct organogenesis is the regeneration of chimeras explained by the induction of adventitious buds from multiple cells [223]. Plant regeneration from somatic embryos induced from a single cell could be used to avoid such chimeras. Somatic embryogenesis was also used for grapevine micropropagation and genetic transformation. Unfortunately, the induction of somatic embryogenesis is generally low and dependent on the type of explants [224]. Moreover, the maintenance of embryogenic masses on calluses and somatic embryos is

also very important [225]. Therefore, transgenesis in grapevine is mostly based on the *Agrobacterium* system, and the regeneration of transformants is generally achieved with somatic embryogenesis. Improved *Agrobacterium*-mediated transformation protocols were published to enhance the fruit quality and tolerance to different stress factors [226,227].

Several factors influence the regeneration of transformants via somatic embryogenesis, such as the grapevine genotype [228], explant source [137,138,229], and culture medium [222,224,230]. All of these factors were reviewed by Zhang et al. [17].

Important achievements in grapevine using direct transformation methods and *Agrobacterium*-mediated transformations are shown in Table 2. Several experiments were also carried out in order to develop an efficient transformation method; these were reviewed by Zhang et al. [17].

Table 2. Grapevine improvements with genetic transformations.

Method	Cultivar	Target Gene	Trait	References
Transient expression assays using direct transformation methods, modified from Jelly et al. [231]				
Biolistics	Cabernet Sauvignon	VvAdh1, VvAdh2 VvAdh2	Abiotic stress	[232,233]
Biolistics	Chardonnay	VvMYBA1, -F1, -PA1, -PA2 and VvCHS1, VvCHS2, VvCHS3 promoters	Flavonoid synthesis	[234]
Biolistics	Chardonnay	VvMYB5a,-5b and VvANR, VvANS, VvCHI, VvF30 50 H, VvLAR1 promoters	Flavonoid synthesis	[235]
Biolistics	Chardonnay	VvMYBA1, -A2 and VvUGT promoter Flavonoids GFP, dual Luc Walker et al. [225] VvMYBF1 and VvANR, VvCHI, VvFLS1, VvLDOX promoters	Flavonoid synthesis	[236]
Biolistics	Chardonnay	VvMYBF1 and VvANR, VvCHI, VvFLS1, VvLDOX promoters	Flavonoid synthesis	[237]
Biolistics	Chardonnay	VvMYBPA1 and VvANR, VvCHI, VvF30 50 H, VvLAR1, VvLDOX promoters	Flavonoid synthesis	[238]
Biolistics	Chardonnay	VvMYC1 and VvMYB5a, -5b, -A1, -A2, -PA1 and VvANR, VvCHI, VvMYC1, VvUGT promoters	Flavonoid synthesis	[239]
Biolistics	Chardonnay Pinot Noir	VvMYB14, -15 and VvSTS29, -41 promoters	Stilbene synthesis	[240]
Biolistics	Chardonnay	VvPGIP1 promoter	Resistance to <i>B. cinerea</i>	[241]
Biolistics	Thompson Seedless	VvPGIP1 promoter	Resistance to <i>B. cinerea</i>	[241]
PEG treatment	Cabernet Sauvignon	VvMSA	Abiotic stress	[242]
PEG treatment	Cabernet Sauvignon	VvWRKY1 and VvJAZ1.1, VvLOX promoters	JA defence pathway	[243]
PEG treatment	Cabernet Sauvignon	VvMYC1	Flavonoid synthesis	[239]
PEG treatment	Cabernet Sauvignon	VvMSA	Abiotic stress	[244]

Table 2. Cont.

Method	Cultivar	Target Gene	Trait	References
Transient expression assays using <i>Agrobacterium</i>-mediated transformation, modified from Jelly et al. [220]; Zhang et al. [17]				
<i>Agrobacterium</i>	Superior Seedless	hpRNA against VvPDS	Gene silencing	[245]
<i>Agrobacterium</i>	Cabernet Franc, Syrah, Zinfandel	GLRaV-2 cDNA	-	[208]
<i>Agrobacterium</i>	Thompson Seedless	D4E1 (synthetic AMP)	Resistance to <i>A. vitis</i> , <i>X. ampelinus</i>	[246]
<i>Agrobacterium</i>	Cabernet Franc	hpRNA against VvPGIP1	Resistance to <i>B. cinerea</i>	[247]
<i>Agrobacterium</i>	Cabernet Sauvignon, Cinsault, Muscat Ottonel, Syrah	VvVST1	Resistance to <i>P. viticola</i>	[248]
<i>Agrobacterium</i>	Carignane	VpGLOX	Resistance to <i>E. necator</i>	[249]
<i>Agrobacterium</i>		VpPR10.2	Resistance to <i>P. viticola</i>	[250]
<i>Agrobacterium</i>		VpSTS	Resistance to <i>E. necator</i>	[251]
<i>Agrobacterium</i>		VpPR10.1	Resistance to <i>P. viticola</i>	[252]
<i>Agrobacterium</i>	Syrah	VvNPR1	Resistance to <i>P. viticola</i>	[253]
<i>Agrobacterium</i>	Grenache	GLRaV-2 cDNA	-	[254]
<i>Agrobacterium</i>	Prime	GVA cDNA	-	[207]
<i>Agrobacterium</i>	Prime, Thompson Seedless	GRSPaV cDNA	-	[209]
<i>Agrobacterium</i>	Gamay Red	VvDFR	Flavonoid synthesis	[255]
<i>Agrobacterium</i>	Chardonnay	amiRNAs against Grapevine fanleaf virus and GUS sensor	Resistance to GFLV	[256]
<i>Agrobacterium</i>	Thompson Seedless	CaMV35S, CsVMV, <i>Arabidopsis</i> ACT2 promoters	-	[257]
<i>Agrobacterium</i>	Thompson Seedless	BDDPs with CaMV35S, CsVMV promoters and enhancers	-	[213]
<i>Agrobacterium</i>	Thompson Seedless	31 grapevine promoters (PR1, PAL, Ubiquitin etc.)	-	[217]
<i>Agrobacterium</i>	Russalka and Rupestris du lot	GFLV CP (grape fanleaf virus coat protein) and four encoding antifreeze proteins (Atf11, Atf62, Atf78, B5) for Russalka, GUS (β -glucuronidase gene) for Rupestris du Lot	Fan Leaf Virus resistance and cold resistance	[258]
<i>Agrobacterium</i>	Chancellor	tfdA gene (a 2,4-D α -ketoglutarate dioxygenase)	Tolerance to 2,4-D	[259]

Table 2. Cont.

Method	Cultivar	Target Gene	Trait	References
<i>Agrobacterium</i>	Pusa Seedless	A rice chitinase gene	Resistance to powdery mildew	[260]
<i>Agrobacterium</i>	Thompson Seedless	vvtl-1 (a thaumatin-like protein)	Broad spectrum fungal disease resistance	[261]
<i>Agrobacterium</i>	Thompson Seedless	VvMybA1 (regulatory gene for the last metabolic step of anthocyanin biosynthesis)	Development of an anthocyanin-based quantitative reporter system	[213]
<i>Agrobacterium</i>	Crimson Seedless	Chitinase and β -1,3-glucanase genes	Tolerance to downy mildew	[262]
<i>Agrobacterium</i>	Thompson Seedless	VpPUB23 (a ubiquitin ligase gene)	Overexpression of VpPUB23 decreased powdery mildew resistance	[225]
<i>Agrobacterium</i>	Chardonnay	VpSTSgDNA2	Developed a protocol and increased powdery mildew resistance	[263]
<i>Agrobacterium</i>	Thompson Seedless and Freedom	LIMA-A (a synthetic gene encoding lytic peptide)	Durable Pierce's disease resistance	[264]
<i>Agrobacterium</i>	Thompson Seedless; Red Globe	VqSTS6 in Thompson Seedless; VpPR4-1 (pathogenesis-related proteins) in Red globe	Resistance to powdery mildew	[265,266]
<i>Agrobacterium</i>	Brachetto	Knockdown through RNA interference of VvMLO6, 7, 11 and 13 (mildew locus O)	Reduced susceptibility to powdery mildew	[267]
<i>Agrobacterium</i>	Thompson Seedless	Overexpression of VaTLP (thaumatin-like protein)	Resistance to downy mildew	[268]
<i>Agrobacterium</i>	Thompson Seedless	Overexpression of VaPUB (a stress-responsive U-box protein gene)	Disease resistance	[269]
<i>Agrobacterium</i>	Thompson Seedless	Overexpression of VpPR10.1	Resistance to downy mildew	[270,271]
<i>Agrobacterium</i>	Thompson Seedless	Overexpression of AgNHX1 (Na^+/H^+ antiporter gene in <i>Atriplex gmelini</i>)	Increased salt tolerance	[272]
<i>Agrobacterium</i>	Thompson Seedless	Overexpression of VvWRKY8	Repressed VvSTS15/21 expression and resveratrol biosynthesis	[273]

Table 2. Cont.

Method	Cultivar	Target Gene	Trait	References
<i>Agrobacterium</i>	Thompson Seedless	Overexpression of VpSTS29/STS2	Resistance to powdery mildew	[274]
<i>Agrobacterium</i>	Thompson Seedless	Overexpression of VlbZIP30 (a basic region/leucine zipper transcription factor)	Improved drought resistance	[275]

7. Genome Editing

Genome editing consists of precise genetic modifications with different purposes, such as gene inactivation, that allow the possibility to explore the function of a particular gene and the insertion or replacement of genes at specific sites for genetic improvements. Strategies involving genome editing are known as new breeding technologies [23]. Artificially engineered nucleases, such as zinc-finger nucleases (ZFNs) [276], transcription activator-like effector nucleases (TALENs) [277,278] and clustered regularly interspersed short palindromic repeats (CRISPR) in association with the Cas9 nuclease [279] are capable of inducing specific double-stranded breaks (DSBs) of DNA molecules. The DSBs are repaired with natural mechanisms present in all cells, such as non-homologous end joining (NHEJ), which is followed by point mutations due to the insertion or deletion of some nucleotides (INDELS) in the target gene, or homologous recombination (HDR) if a DNA sequence is available for recombination. CRISPR-Cas9 technology is considered the most efficient, among the genome editing tools, due to the high specificity and minimal nontarget effects [280]. Gene editing with the CRISPR-Cas9 system requires a guide RNA (gRNA) containing a spacer sequence complementary with the desired DNA sequence. The complex formed by guide RNA and Cas9 scans the genome, searching for complementary double-stranded DNA [281]. The nuclease recognizes the protospacer-adjacent motif (PAM) and generates a DSB in the specific gene sequence. Thus, genome editing through CRISPR-Cas9 technology requires the PAM sequence downstream of the target gene and proper guide RNAs, designed based on the gene sequences encoding important traits [282]. The description of the genome editing technologies based on ZFNs, TALENs and the CRISPR-Cas9 system was provided by Bortesi and Fischer [283] and Butiuc-Keul et al. [284]. Despite the many advantages of CRISPR-Cas9 technology over ZFNs and TALENs, the occurrence of off-target mutations is one of the shortcomings [285,286], being influenced by different parameters, such as the recognition of the target, the design of guide RNAs, the frequency of repair events with homologous recombination, and anti-CRISPR proteins that inactivate Cas9 [280].

Another limitation of using CRISPR-Cas9 technology is related to the delivery of the system in plant tissues. Generally, gRNAs and Cas9 are delivered into plant cells by *Agrobacterium*, viral vectors, PEG-mediated transformation, biolistic methods, and nanoparticles [280]. The CRISPR-Cas9 system could be released in tissues in its DNA, mRNA, or ribonucleoprotein forms and incorporated in different biomaterials for proper delivery [287], but these studies were carried out mostly in animal and human cells for cancer therapies. The structure of the plant cell wall limits the delivery of the system; thus, the most used system for delivery into plant tissues is *Agrobacterium* [288]. Nevertheless, the direct delivery of the purified Cas9 protein and gRNAs was also applied for the editing of the plant genome [289,290].

Most of the studies regarding the genome editing of grapevine were conducted in order to increase the resistance to powdery mildew [136,289,291,292] and *Botrytis cinerea* [227], the production of tartaric acid [293], the manipulation of the carotenoid biosynthesis pathway, and the induction of the albino phenotype [294,295] [Table 3].

Table 3. Grapevine improvements with CRISPR-Cas9 technology (modified from Zhang et al. [17]; Butiuc-Keul et al. [284]).

Technology	Cultivar	Target Gene	Trait	References
CRISPR/Cas9	Chardonnay	MLO-7	Resistance to powdery mildew	[289]
CRISPR/Cas9	Neo Muscat	VvPDS	Albino phenotype	[294]
CRISPR/Cas9	Chardonnay, 41B	VvPDS	Albino phenotype and dwarf morphology	[295]
CRISPR/Cas9	Chardonnay	L-idonate dehydrogenase gene (IdnDH)	Biosynthesis of tartaric acid	[293]
CRISPR/Cas9	Neo Muscat	VvPDS (phytoene desaturase gene)	CRISPR-Cas9-mediated protocol development	[294]
CRISPR/Cas9	Thompson Seedless	VvWRKY52	Resistance to <i>B. cinerea</i>	[227]
CRISPR/Cas9	Thompson Seedless	VvPR4b	VvPR4b knockout decreased downy mildew resistance	[291]
	Thompson Seedless	VvMLO3 and VvMLO4	Resistance to powdery mildew	[293]

8. Conclusions and Perspectives

The selection of grapevine rootstocks and scion varieties with improved fruit qualities, resistance to herbicides, and tolerance to biotic and abiotic stressors requires different biotechnologies, such as in vitro plant regeneration and multiplication, mutagenesis, the induction of somaclonal variability and selection of new valuable genotypes, and genetic transformation and genome editing. The strategies based on genome editing have the advantage of speeding up crop improvement and reducing the cost of the process, but the implementation of these technologies needs government support. Usually, the edited plants are not considered transgenic even though the delivery of the CRISPR-Cas9 system is mediated by *Agrobacterium*. Thus, edited plants could be easily accepted on the market. Improving grapevine tolerance to diseases and pests is the most promising contribution of new breeding technologies because little is known about genes encoding disease resistance and their functions, and the QTLs are not identified. The development of a multi-resistant genotype is complicated, and *V. vinifera* is susceptible to different fungi. Thus, new breeding technology could be considered a significant alternative to the classical selection and breeding of grapevine varieties resistant to biotic and abiotic stress.

Author Contributions: Conceptualization, A.B.-K. and A.C.; writing—original draft preparation, A.B.-K. and A.C.; writing—review and editing, A.B.-K. and A.C.; supervision, A.B.-K.; funding acquisition, A.B.-K. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the grant AGC nr.332066/23.06.2022 and Ministry of Research, Innovation and Digitization through the Core Project BioClimpact nr.7/30.12.2022, code 23020401.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: We would like to thank Matei Dobrescu, for his support in formatting the figures.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Basso, M.F.; Fajardo, T.V.; Saldarelli, P. Grapevine virus diseases: Economic impact and current advances in viral prospection and management. *Rev. Bras. Frutic.* **2017**, *39*, e-411. [[CrossRef](#)]
2. Bi, W.L.; Pan, C.; Hao, X.Y.; Cui, Z.H.; Kher, M.M.; Marković, Z.; Wang, Q.C.; Teixeira da Silva, J.A. Cryopreservation of grapevine [*Vitis* spp.]—A review. *Vitr. Cell. Dev. Biol. Plant.* **2017**, *53*, 449–460. [[CrossRef](#)]
3. FAOSTAT. 2020. Available online: <https://www.fao.org/faostat/en/#data> (accessed on 5 January 2021).
4. OIV 2020. State of the World Vitivinicultural Sector in 2020. Available online: <https://www.oiv.int/> (accessed on 13 April 2021).
5. EUROSTAT 2020. Available online: https://ec.europa.eu/eurostat/statistics-explained/index.php?title=Vineyards_in_the_EU_-_statistics&oldid=566726. (accessed on 5 March 2021).
6. Reynolds, A.G. The Grapevine, Viticulture, and Winemaking: A Brief Introduction. In *Grapevine Viruses: Molecular Biology, Diagnostics and Management*; Meng, B., Martelli, G., Golino, D., Fuchs, M., Eds.; Springer: Berlin/Heidelberg, Germany, 2017; pp. 3–29. [[CrossRef](#)]
7. Campos, G.; Chialva, C.; Miras, S.; Lijavetzky, D. New technologies and strategies for grapevine breeding through genetic transformation. *Front. Plant. Sci.* **2021**, *25*, 767522. [[CrossRef](#)] [[PubMed](#)]
8. Abiri, K.; Rezaei, M.; Tahanian, H.; Heidari, P.; Khadivi, A. Morphological and pomological variability of a grape (*Vitis vinifera* L.) germplasm collection. *Sci. Hortic.* **2020**, *266*, 1–12. [[CrossRef](#)]
9. Kambiranda, D.; Obuya, J.; Snowden, J. Grapevine Improvement through Biotechnology. In *Genetic Transformation in Crops*; To, K.-Y., Ed.; IntechOpen: London, UK, 2020. [[CrossRef](#)]
10. Chacón-Vozmediano, J.L.; Gramaje, D.; León, M.; Armengol, J.; Moral, J.; Izquierdo-Cañas, P.M.; Martínez-Gascueña, J. Cultivar Susceptibility to Natural Infections Caused by Fungal Grapevine Trunk Pathogens in La Mancha Designation of Origin (Spain). *Plants* **2021**, *10*, 1171. [[CrossRef](#)]
11. Galet, P. Apoplexie. In: Les maladies et les parasites de la vigne. *Imp. Paysan du Midi Montp.* **1977**, *1*, 409–430.
12. Santos, R.B.; Figueiredo, A. Two sides of the same story in grapevine-pathogen interactions. *J. Exp. Bot.* **2021**, *72*, 3367–3380. [[CrossRef](#)]
13. Cardell, M.; Amengual, A.; Romero, R. Future effects of climate change on the suitability of wine grape production across Europe. *Reg. Environ. Change* **2019**, *19*, 1–12. [[CrossRef](#)]
14. Santos, J.A.; Fraga, H.; Malheiro, A.C.; Moutinho-Pereira, J.; Dinis, L.-T.; Correia, C.; Moriondo, M.; Leolini, L.; Dibari, C.; Costafreda-Aumedes, S.; et al. A Review of the Potential Climate Change Impacts and Adaptation Options for European Viticulture. *Appl. Sci.* **2020**, *10*, 3092. [[CrossRef](#)]
15. Töpfer, R.; Trapp, O. A cool climate perspective on grapevine breeding: Climate change and sustainability are driving forces for changing varieties in a traditional market. *Appl. Genet.* **2022**, *7*, 1–4. [[CrossRef](#)]
16. Fraga, H. Climate Change: A New Challenge for the Winemaking Sector. *Agronomy* **2020**, *10*, 1465. [[CrossRef](#)]
17. Zhang, X.-M.; Wu, Y.-F.; Li, Z.; Song, C.-B.; Wang, X.-P. Advancements in plant regeneration and genetic transformation of grapevine (*Vitis* spp.). *J. Integrat. Agric.* **2021**, *20*, 1407–1434. [[CrossRef](#)]
18. Gray, D.J.; Meredith, C.P. *Biotechnology of Perennial Fruit Crops*; Hammerschlag, F.A., Litz, R.E., Eds.; CAB International: Wallingford, UK, 1992; pp. 229–262.
19. Vezzulli, S.; Doligez, A.; Bellin, D. Molecular Mapping of Grapevine Genes. In *The Grape Genome. Compendium of Plant Genomes*; Cantu, D., Walker, M., Eds.; Springer: Cham, Switzerland, 2019. [[CrossRef](#)]
20. Directive 2001/18/ec of the European Parliament and of the Council of 12 March 2001 on the Deliberate Release into the Environment of Genetically Modified Organisms and Repealing Council Directive 90/220/EEC (OJ L 106, 17.4.2001, pp. 1–60). Available online: <https://www.legislation.gov.uk/eudr/2001/18/contents> (accessed on 21 September 2022).
21. Regulation (EC) No 1829/2003 of the European Parliament and of the Council of 22 September 2003 on Genetically Modified Food and Feed (Text with EEA Relevance). Official Journal L 268, 18/10/2003 P. 0001–0023. Available online: <https://eur-lex.europa.eu/legal-content/EN/ALL/?uri=CELEX%3A32003R1829> (accessed on 21 September 2022).
22. Regulation (EC) No 1830/2003 of the European Parliament and of the Council of 22 September 2003 Concerning the Traceability and Labelling of Genetically Modified Organisms and the Traceability of Food and Feed Products Produced from Genetically Modified Organisms and Amending Directive 2001/18/EC. Official Journal L 268, 18/10/2003 P. 0024–0028. Available online: <https://eur-lex.europa.eu/legal-content/EN/ALL/?uri=CELEX%3A32003R1829> (accessed on 21 September 2022).
23. Dalla Costa, L.; Malnoy, M.; Lecourieux, D.; Deluc, L.; Ouaked-Lecourieux, F.; Deluc, L.; Ouaked-Lecourieux, F.; Thomas, M.; Torregrosa, L.J.-M. The state-of-the-art of grapevine biotechnology and new breeding technologies (NBTS). *OENO One* **2019**, *53*, 189–212. [[CrossRef](#)]
24. Grassi, F.; De Lorenzis, G. Back to the origins: Background and perspectives of grapevine domestication. *Int. J. Mol. Sci.* **2021**, *22*, 4518. [[CrossRef](#)] [[PubMed](#)]
25. Rivera Nuñez, D.; Walker, M.J. A review of palaeobotanical findings of early *Vitis* in the Mediterranean and the origins of cultivated grape-vines, with special reference to new pointers to prehistoric exploitation in the Western Mediterranean. *Rev. Palaeobot. Palynol.* **1989**, *61*, 205–237. [[CrossRef](#)]
26. Grassi, F.; Labra, M.; Imazio, S.; Spada, A.; Sgorbati, S.; Scienza, A.; Sala, F. Evidence of secondary grapevine domestication centre detected by SSR analysis. *Theor. App. Genet.* **2003**, *107*, 1315–1320. [[CrossRef](#)]

27. Arroyo-García, R.; Ruiz-García, L.; Bolling, L.; Ocete, R.; López, M.A.; Arnold, C.A.; Ergul, A.; Söylemezoglu, G.; Uzun, H.I.; Cabello, F.; et al. Multiple origins of cultivated grapevine (*Vitis vinifera* L. ssp. *sativa*) based on chloroplast DNA polymorphisms. *Mol. Ecol.* **2006**, *15*, 3707–3714. [CrossRef]
28. Terral, J.F.; Tabard, E.; Bouby, L.; Ivorra, S.; Pastor, T.; Figueiral, I.; Picq, S.; Chevance, J.B.; Jung, C.; Fabre, L.; et al. Evolution and history of grapevine (*Vitis vinifera*) under domestication: New morphometric perspectives to understand seed domestication syndrome and reveal origins of ancient European cultivars. *Ann. Bot.* **2010**, *105*, 443–455. [CrossRef]
29. Rusjan, D. Genetic and phenotypic diversity and relations between grapevine varieties: Slovenian germplasm. In *The Mediterranean Genetic Code-Grapevine and Olive*; Poljuha, D., Sladonja, B., Eds.; IntechOpen: London, UK, 2013. [CrossRef]
30. Villano, C.; Aiese Cigliano, R.; Esposito, S.; D’Amelia, V.; Iovene, M.; Carputo, D.; Aversano, R. DNA-Based Technologies for grapevine biodiversity exploitation: State of the art and future perspectives. *Agronomy* **2022**, *12*, 491. [CrossRef]
31. Emanuelli, F.; Lorenzi, S.; Grzeskowiak, L.; Catalano, V.; Stefanini, M.; Troggio, M.; Myles, S.; Martinez-Zapater, J.M.; Zyprian, E.; Moreira, F.M.; et al. Genetic diversity and population structure assessed by SSR and SNP markers in a large germplasm collection of grape. *BMC Plant. Biol.* **2013**, *13*, 39. [CrossRef]
32. Zecca, G.; Grassi, F.; Tabidze, V.; Pipia, I.; Kotorashvili, A.; Kotaria, N.; Beridze, T. Dates and rates in grape’s plastomes: Evolution in slow motion. *Curr. Genet.* **2020**, *66*, 123–140. [CrossRef]
33. Lacombe, T.; Audeguin, L.; Boselli, M.; Buccetti, B.; Cabello, F.; Chatelet, P.; Crespan, M.; D’Onofrio, C.; Eiras Dias, J.; Ercisli, S.; et al. Grapevine European catalogue: Towards a comprehensive list. *Vitis* **2011**, *50*, 65–68. [CrossRef]
34. Jackson, R.S. Grape Species and Varieties. In *Food Science and Technology, Wine Science*, 4th ed.; Ronald, S.J., Ed.; Academic Press: Cambridge, MA, USA, 2014; pp. 21–67. ISBN 9780123814685. [CrossRef]
35. Panara, F.; Bergamini, C.; Palliotti, A.; Calderini, O. Use of Molecular Markers (Ssrs) and Public Databases in *Vitis vinifera* L. as the Main Case of Efficient Crop Cultivar Identification. *JOJ Hortic. Arboric.* **2018**, *2*, 555576. [CrossRef]
36. Sefc, K.; Pejić, I.; Maletić, E.; Thomas, M.; Lefort, F. Microsatellite Markers for Grapevine: Tools for Cultivar Identification & Pedigree Reconstruction. In *Grapevine Molecular Physiology & Biotechnology*; Roubelakis-Angelakis, K.A., Ed.; Springer: Dordrecht, The Netherland, 2009. [CrossRef]
37. Nasiri, A.; Taheri-Garavand, A.; Fanourakis, D.; Zhang, Y.-D.; Nikoloudakis, N. Automated Grapevine Cultivar Identification via Leaf Imaging and Deep Convolutional Neural Networks: A Proof-of-Concept Study Employing Primary Iranian Varieties. *Plants* **2021**, *10*, 1628. [CrossRef] [PubMed]
38. Cho, K.H.; Noh, J.H.; Park, S.J.; Kim, S.H.; Kim, D.; Chun, J.A. Development of sequence characterized amplified region markers for the identification of grapevine cultivars. *Hortic. Sci.* **2015**, *50*, 1744–1750. [CrossRef]
39. Castro, C.; Carvalho, A.; Pavia, I.; Leal, F.; Moutinho-Pereira, J.; Lima-Brito, J. Nucleolar Activity and Physical Location of Ribosomal DNA Loci in *Vitis vinifera* L. by Silver Staining and Sequential FISH. *Sci. Hortic.* **2018**, *232*, 57–62. [CrossRef]
40. Pereira, H.S.; Barão, A.; Delgado, M.; Morais-Cecílio, L.; Viegas, W. Genomic Analysis of Grapevine Retrotransposon 1 (Gret1) in *Vitis vinifera*. *Theor. Appl. Genet.* **2005**, *111*, 871–878. [CrossRef]
41. Falistocco, E.; Passeri, V.; Marconi, G. Investigations of 5S rRNA of *Vitis vinifera* L.: Sequence Analysis and physical mapping. *Genome* **2007**, *50*, 927–938. [CrossRef]
42. Giannuzzi, G.; D’Addabbo, P.; Gasparro, M.; Martinelli, M.; Carelli, F.N.; Antonacci, D.; Ventura, M. Analysis of High-Identity Segmental Duplications in the Grapevine Genome. *BMC Genom.* **2011**, *12*, 436. [CrossRef]
43. Pereira, H.S.; Delgado, M.; Avó, A.P.; Barão, A.; Serrano, I.; Viegas, W. Pollen grain development is highly sensitive to temperature stress in *Vitis vinifera*. *Aust. J. Grape Wine Res.* **2014**, *20*, 474–484. [CrossRef]
44. Goswami, M.; Attri, K.; Goswami, I. Applications of Molecular Markers in Fruit Crops: A Review. *IJEP* **2022**, *9*, 121–126. Available online: <http://www.pphouse.org/ijep-article-details.php?art=323> (accessed on 12 September 2022). [CrossRef]
45. Bourquin, J.; Otten, L.; Walter, B. PCR-RFLP analysis of *Vitis*, *Ampelopsis* and *Parthenocissus* and its application to the identification of rootstocks. *Vitis* **1995**, *34*, 103–108.
46. Karataş, H.; AĞAOĞLU, Y.S. RAPD analysis of selected local Turkish grape cultivars (*Vitis vinifera*). *GMR* **2010**, *9*, 1980–1986. [CrossRef]
47. Vidal, J.; Delavault, P.; Coarer, M.; Defontaine, A. Design of grapevine (*Vitis vinifera* L.) cultivar-specific SCAR primers for PCR fingerprinting. *Appl. Genet.* **2000**, *101*, 1194–1201. [CrossRef]
48. Pollastro, S.; Dongiovanni, C.; Abbatecola, A.; de Guido, M.A.; de Miccolis Angelini, R.M.; Faretra, F. Specific SCAR Primers for Fungi Associated with Wood Decay of Grapevine. *Phytopathol. Mediterr.* **2001**, *40*, S362–S368.
49. Villano, C.; Carpoto, D.; Frusciante, L.; Santoro, X.; Aversano, R. Use of SSR and Retrotransposon-Based Markers to Interpret the Population Structure of Native Grapevines from Southern Italy. *Mol. Biotechnol.* **2014**, *56*, 1011–1020. [CrossRef] [PubMed]
50. Karimi, M.R.; Dehvari, V.; Hajiyani, M. Genetics Diversity of Some Grape Genotypes by ISSR and RAPD Markers. *Eur. J. Hortic. Sci.* **2011**, *76*, 201–207.
51. Nookaraju, A.; Agrawal, D.C. Genetic homogeneity of in vitro raised plants of grapevine cv. Crimson Seedless revealed by ISSR and microsatellite markers. *S. Afr. J. Bot.* **2012**, *78*, 302–306. [CrossRef]
52. Stajner, N.; Jakse, J.; Javornik, B.; Masuelli, R.W.; Martínez, L.E. Highly variable AFLP and S-SAP markers for the identification of ‘Malbec’ and ‘Syrah’ clones. *Vitis* **2009**, *48*, 145–150.

53. Péros, J.P.; Cousins, P.; Launay, A.; Cubry, P.; Walker, A.; Prado, E.; Peressotti, E.; Wiedemann-Merdinoglu, S.; Laucou, V.; Merdinoglu, D.; et al. Genetic diversity and population structure in *Vitis* species illustrate phylogeographic patterns in eastern North America. *Mol. Ecol.* **2021**, *30*, 2333–2348. [[CrossRef](#)]
54. Peng, F.Y.; Reid, K.E.; Liao, N.; Schlosser, J.; Lijavetzky, D.; Holt, R.; Martínez Zapater, J.M.; Jones, S.; Marra, M.; Bohlmann, J.; et al. Generation of ESTs in *Vitis vinifera* wine grape (Cabernet Sauvignon) and table grape (Muscat Hamburg) and discovery of new candidate genes with potential roles in berry development. *Gene* **2007**, *402*, 40–50. [[CrossRef](#)]
55. Ji, X.N.; Li, F.; Yang, C.J.; Li, B.; Wang, J.; Zhang, W. Expressed sequence tags (ESTs) analysis of the ripening *Vitis amurensis* cv. Shuang Hong berry skins. *J. For. Res.* **2013**, *24*, 495–502. [[CrossRef](#)]
56. Lü, X.-L.; Zhang, G.-L.; Liao, M.-A.; Gong, R.-G.; Zeng, X.-L. Random Amplified Microsatellite Polymorphism (RAMP) Analysis of Grape Breeds Genetic Relationships. *JSAU* **2004**, *22*, 133–137. [[CrossRef](#)]
57. Arroyo-García, R.; Lefort, F.; de Andrés, M.T.; Ibáñez, J.; Borrego, J.; Jouve, N.; Cabello, F.; Martínez-Zapater, J.M. Chloroplast microsatellite polymorphisms in *Vitis* species. *Genome* **2002**, *45*, 1142–1149. [[CrossRef](#)]
58. Veloso, M.M.; Almadanim, M.C.; Baleiras-Couto, M.; Pereira, H.S.; Carneiro, L.C.; Fevereiro, P.; Eiras-Dias, J.E. Microsatellite database of grapevine (*Vitis vinifera* L.) cultivars used for wine production in Portugal. *Cien. Tecn. Vitivinic.* **2010**, *25*, 53–61.
59. Castro, I.; Martín, J.P.; Ortiz, J.M.; Pinto-Carnide, O. Varietal discrimination and genetic relationships of *Vitis vinifera* L. cultivars from two major controlled appellation (DOC) regions in Portugal. *Sci. Hortic.* **2011**, *127*, 507–514. [[CrossRef](#)]
60. Castro, I.; Pinto-Carnide, O.; Ortiz, J.M.; Martin, J.P. Chloroplast genome diversity in Portuguese grapevine (*Vitis vinifera* L.) cultivars. *Mol. Biotechnol.* **2013**, *54*, 528–540. [[CrossRef](#)]
61. Ferreira, V.; Pinto-Carnide, O.; Mota, T.; Martín, J.P.; Ortiz, J.M.; Castro, I. Identification of minority grapevine cultivars from Vinhos Verdes Portuguese DOC Region. *Vitis* **2015**, *54*, 53–58. [[CrossRef](#)]
62. De Michele, R.; La Bella, F.; Gristina, A.S.; Fontana, I.; Pacifico, D.; Garfi, G.; Motisi, A.; Crucitti, D.; Abbate, L.; Carimi, F. Phylogenetic Relationship Among Wild and Cultivated Grapevine in Sicily: A Hotspot in the Middle of the Mediterranean Basin. *Front. Plant. Sci.* **2019**, *10*, 1506. [[CrossRef](#)] [[PubMed](#)]
63. Bianchi, D.; Brancadoro, L.; De Lorenzis, G. Genetic Diversity and Population Structure in a *Vitis* spp. Core Collection Investigated by SNP Markers. *Diversity* **2020**, *12*, 103. [[CrossRef](#)]
64. Flutre, T.; Le Cunff, L.; Fodor, A.; Launay, A.; Romieu, C.; Berger, G.; Bertrand, Y.; Terrier, N.; Beccavin, I.; Bouckenooghe, V.; et al. A genome-wide association and prediction study in grapevine deciphers the genetic architecture of multiple traits and identifies genes under many new QTLs. *G3 Genes | Genomes | Genet.* **2022**, *12*, jkac103. [[CrossRef](#)] [[PubMed](#)]
65. Ghaffari, S.; Hasnaoui, N.; Zinelabidine, L.H.; Ferchichi, A.; Martínez-Zapater, J.M.; Ibáñez, J. Genetic diversity and parentage of Tunisian wild and cultivated grapevines (*Vitis vinifera* L.) as revealed by single nucleotide polymorphism (SNP) markers. *Tree Genet. Genomes* **2014**, *10*, 1103–1112. [[CrossRef](#)]
66. Augusto, D.; Ibáñez, J.; Pinto-Sintra, A.L.; Falco, V.; Leal, F.; Martínez-Zapater, J.M.; Oliveira, A.A.; Castro, I. Grapevine Diversity and Genetic Relationships in Northeast Portugal Old Vineyards. *Plants* **2021**, *10*, 2755. [[CrossRef](#)] [[PubMed](#)]
67. Lijavetzky, D.; Cabezas, J.A.; Ibáñez, A.; Rodríguez, V.; Martínez-Zapater, J.M. High throughput SNP discovery and genotyping in grapevine (*Vitis vinifera* L.) by combining a re-sequencing approach and SNplex technology. *BMC Genom.* **2007**, *8*, 424. [[CrossRef](#)]
68. Dong, X.; Chen, W.; Liang, Z.; Li, X.; Nick, P.; Chen, S.; Dong, Y.; Li, S.; Sheng, J. VitisGDB: The Multifunctional Database for Grapevine Breeding and Genetics. *Mol. Plant.* **2020**, *13*, 1098–1100. [[CrossRef](#)]
69. Maul, E.; Töpfer, R. Vitis International Variety Catalogue (V IVC): A cultivar database referenced by genetic profiles and morphology. *BIO Web Conf.* **2015**, *5*, 01009. [[CrossRef](#)]
70. Cabezas, J.A.; Cabezas, M.T.; Cervera, M.T.; Cervera, L.; Ruiz-García, L.; Ruiz-García, J.; Carreño, J.; Martínez-Zapater, C.J.M. A genetic analysis of seed and berry weight in grapevine. *Genome* **2006**, *49*, 1572–1585. [[CrossRef](#)]
71. Maul, E.; Sudharma, K.N.; Kecke, S.; Marx, G.; Müller, C.; Audeguin, L.; Boselli, M.; Boursiquot, J.M.; Buccetti, B.; Cabello, F.; et al. The European Vitis Database (www.Eu-Vitis.De): A Technical Innovation through An Online Uploading and Interactive Modification System. *Vitis* **2012**, *51*, 79–85.
72. Lefort, F.; Roubelakis-Angelakis, K.A. The Greek Vitis Database: A Multimedia Web-Backed Genetic Database for Germplasm Management of *Vitis* Resources in Greece. *J. Wine Res.* **2000**, *11*, 233–242. [[CrossRef](#)]
73. Davey, J.W.; Hohenlohe, P.A.; Etter, P.D.; Boone, J.Q.; Catchen, J.M.; Blaxter, M.L. Genome-wide genetic marker discovery and genotyping using next-generation sequencing. *Nat. Rev. Genet.* **2011**, *12*, 499–510. [[CrossRef](#)]
74. Marrano, A.; Birolo, G.; Prazzoli, M.L.; Lorenzi, S.; Valle, G.; Grando, M.S. SNP-Discovery by RAD-Sequencing in a Germplasm Collection of Wild and Cultivated Grapevines (*V. vinifera* L.). *PLoS ONE* **2017**, *12*, e0170655. [[CrossRef](#)] [[PubMed](#)]
75. Zhu, J.; Guo, Y.; Su, K.; Liu, Z.; Ren, Z.; Li, K.; Guo, X. Construction of a highly saturated genetic map for *Vitis* by Next-generation restriction site-associated DNA sequencing. *BMC Plant. Biol.* **2018**, *18*, 347. [[CrossRef](#)] [[PubMed](#)]
76. Gambino, G.; Dal Molin, A.; Boccacci, P.; Minio, A.; Chitarra, W.; Avanzato, C.G.; Tononi, P.; Perrone, I.; Raimondi, S.; Schneider, A. Whole-Genome Sequencing and SNV Genotyping of ‘Nebbiolo’ (*Vitis vinifera* L.) Clones. *Sci. Rep.* **2017**, *7*, 1–15. [[CrossRef](#)] [[PubMed](#)]
77. Carbonell-Bejerano, P.; Royo, C.; Torres-Pérez, R.; Grimplet, J.; Fernandez, L.; Franco-Zorrilla, J.M.; Lijavetzky, D.; Baroja, E.; Martínez, J.; García-Escudero, E. Catastrophic unbalanced genome rearrangements cause somatic loss of berry color in grapevine. *Plant. Physiol.* **2017**, *175*, 786–801. [[CrossRef](#)]

78. Pucker, B.; Schwandner, A.; Becker, S.; Hausmann, L.; Viehöver, P.; Töpfer, R.; Weisshaar, B.; Holtgräwe, D. RNA-Seq time series of *Vitis vinifera* bud development reveals correlation of expression patterns with the local temperature profile. *Plants* **2020**, *9*, 1548. [[CrossRef](#)]
79. Ma, Q.; Yang, J. Transcriptome profiling and identification of the functional genes involved in berry development and ripening in *Vitis vinifera*. *Gene* **2019**, *680*, 84–96. [[CrossRef](#)] [[PubMed](#)]
80. Toffolatti, S.L.; De Lorenzis, G.; Brilli, M.; Moser, M.; Shariati, V.; Tavakol, E.; Maddalena, G.; Passera, A.; Casati, P.; Pindo, M. Novel aspects on the interaction between grapevine and *Plasmopara Viticola*: Dual-RNA-Seq analysis highlights gene expression dynamics in the pathogen and the plant during the battle for infection. *Genes* **2020**, *11*, 261. [[CrossRef](#)] [[PubMed](#)]
81. Zou, C.; Karn, A.; Reisch, B.; Nguyen, A.; Sun, Y.; Bao, Y.; Campbell, M.S.; Church, D.; Williams, S.; Xu, X. Haplotyping the *Vitis* collinear core genome with rhAmpSeq improves marker transferability in a diverse genus. *Nat. Commun.* **2020**, *11*, 413. [[CrossRef](#)] [[PubMed](#)]
82. Alizadeh, M.; Singh, S.K.; Jhang, T.; Sharma, T.R. Inter Simple Sequence Repeat Analysis to Confirm Genetic Stability of Micropropagated Plantlets in Three Grape (*Vitis* spp.) Rootstock Genotypes. *J. Plant. Biochem. Biotechnol.* **2008**, *17*, 77–80. [[CrossRef](#)]
83. Hameed, U.K.A.; Abdelaziz, K.; El Sherif, N. Genetic diversity of grapevine (*Vitis vinifera* L.) cultivars in Al-Madinah Al-Munawara based on molecular markers and morphological traits. *Bangladesh J. Plant. Taxon.* **2020**, *27*, 113–127. [[CrossRef](#)]
84. Guo, Y.; Lin, H.; Liu, Z.; Zhao, Y.; Guo, X.; Li, K. SSR and SRAP marker-based linkage map of *Vitis vinifera* L. *Biotechnol. Biotechnol. Equip.* **2014**, *28*, 221–229. [[CrossRef](#)] [[PubMed](#)]
85. Zhang, X.; Gong, P.; Shi, Y.; Wang, Y.; Zhang, C. Genetic inter-relationships among Chinese wild grapes based on SRAP marker analyses. *Vitis* **2018**, *57*, 151–157. [[CrossRef](#)]
86. Pelsy, F. Untranslated leader region polymorphism of *Tvv1*, a retrotransposon family, is a novel marker useful for analyzing genetic diversity and relatedness in the genus *Vitis*. *Appl. Genet.* **2007**, *116*, 15–27. [[CrossRef](#)]
87. Pelsy, F.; Bevilacqua, L.; Blanc, S.; Merdinoglu, D.A. Molecular marker set combining a retrotransposon insertion and SSR polymorphisms is useful for assessing diversity in *Vitis*. *OENO One* **2021**, *55*, 403–414. [[CrossRef](#)]
88. Roach, M.J.; Johnson, D.L.; Bohlmann, J.; van Vuuren, H.J.J.; Jones, S.J.M.; Pretorius, I.S.; Schmidt, S.A.; Borneman, A.R. Population sequencing reveals clonal diversity and ancestral inbreeding in the grapevine cultivar Chardonnay. *PLoS Genet.* **2018**, *14*, e1007807. [[CrossRef](#)] [[PubMed](#)]
89. Butiuc-Keul, A.; Crăciunaş, C.; Coste, A.; Farago, M. Discrimination and genetic polymorphism in several cultivar of grapevine by RAPD markers. *Rom. Biotechnol. Lett.* **2010**, *15*, 110–115.
90. Cretazzo, E.; Meneghetti, S.; De Andrés, M.T.; Gaforio, L.; Frare, E.; Cifre, J. Clone Differentiation and Varietal Identification by means of SSR, AFLP, SAMPL and M-AFLP in order to assess the clonal selection of grapevine: The case study of Manto Negro, Callet and Moll, autochthonous cultivars of Majorca. *Ann. Appl. Biol.* **2010**, *157*, 213–227. [[CrossRef](#)]
91. Coste, A.; Postolache, D.; Popescu, F.; Butiuc-Keul, A. Authentication of valuable grapevine varieties from Romania, through molecular markers. *Rom. Biotechnol. Lett.* **2010**, *15*, 3–10.
92. Baránková, K.; Sotolář, R.; Baránek, M. Identification of Rare Traditional Grapevine Cultivars Using SSR Markers and Their Geographical Location within the Czech Republic. *Czech. J. Genet. Plant. Breed.* **2020**, *56*, 71–78. [[CrossRef](#)]
93. Jiménez-Cantizano, A.; Muñoz-Martín, A.; Amores-Arrocha, A.; Sancho-Galán, P.; Palacios, V. Identification of red grapevine cultivars (*Vitis vinifera* L.) preserved in ancient vineyards in Axarquia (Andalusia, Spain). *Plants* **2020**, *9*, 1572. [[CrossRef](#)]
94. Ocaña, J.; Walter, B.; Schellenbaum, P. Stable MSAP markers for the distinction of *Vitis vinifera* cv Pinot Noir clones. *Mol. Biotechnol.* **2013**, *55*, 236–248. [[CrossRef](#)]
95. Emanuelli, F.; Sordo, M.; Lorenzi, S.; Battilana, J.U.R.I.; Grando, M. Development of user-friendly functional molecular markers for VvDXS gene conferring muscat flavor in grapevine. *Mol. Breed.* **2014**, *33*, 235–241. [[CrossRef](#)] [[PubMed](#)]
96. Castro, I.; D’Onofrio, C.; Martín, J.P.; Ortiz, J.M.; De Lorenzis, G.; Ferreira, V.; Pinto-Carnide, O. Effectiveness of AFLPs and retrotransposon-based markers for the identification of Portuguese grapevine cultivars and clones. *Mol. Biotechnol.* **2012**, *52*, 26–39. [[CrossRef](#)] [[PubMed](#)]
97. Fan, X.C.; Chu, J.Q.; Liu, C.H.; Sun, X.; Fang, J.G. Identification of grapevine rootstock cultivars using expressed sequence tag-simple sequence repeats. *GMR* **2014**, *13*, 7649–7657. [[CrossRef](#)] [[PubMed](#)]
98. D’Onofrio, C.; De Lorenzis, G.; Giordani, T.; Natali, L.; Cavallini, A.; Scalabrelli, G. Retrotransposon-based molecular markers for grapevine species and cultivars identification. *Tree Genet. Genomes* **2010**, *6*, 451–466. [[CrossRef](#)]
99. Ulanovsky, S.; Gogorcena, Y.; Martinez de Toda, F.; Ortiz, J. Use of molecular markers in detection of synonymies and homonymies in grapevines (*Vitis vinifera* L.). *Sci. Hortic.* **2002**, *92*, 241–254. [[CrossRef](#)]
100. Moreno-Sanz, P.; Suárez, B.; Loureiro, M.D. Identification of synonyms and homonyms in grapevine cultivars (*Vitis vinifera* L.) from Asturias (Spain). *J. Hortic. Sci. Biotechnol.* **2008**, *83*, 683–688. [[CrossRef](#)]
101. Castro, I.; Martín, J.P.; Ortiz, J.M.; Mota, M.T.; Pinto-Carnide, O.; Martin, J.; Ortiz, J.M.; Mota, M.T.; Pinto-Carnide, O. The Portuguese grapevine cultivar ‘Amaral’: Synonymies, homonymies and misnames. *Vitis* **2012**, *51*, 61–63. [[CrossRef](#)]
102. Crespan, M.; Migliaro, D.; Larger, S.; Pindo, M.; Palmisano, M.; Manni, A.; Manni, E.; Polidori, E.; Sbaffi, F.; Silvestri, Q.; et al. Grapevine (*Vitis vinifera* L.) varietal assortment and evolution in the Marche region (central Italy). *OENO One* **2021**, *55*, 4628. [[CrossRef](#)]

103. D’Onofrio, C.; Tumino, G.; Gardiman, M.; Crespan, M.; Bignami, C.; de Palma, L.; Barbagallo, M.G.; Muganu, M.; Morcia, C.; Novello, V.; et al. Parentage atlas of Italian grapevine varieties as inferred from SNP genotyping. *Front. Plant. Sci.* **2021**, *11*, 2265. [[CrossRef](#)]
104. Nebish, A.; Tello, J.; Ferradás, Y.; Aroutiounian, R.; Martínez-Zapater, J.M.; Ibáñez, J. SSR and SNP Genetic Profiling of Armenian Grape Cultivars Gives Insights into Their Identity and Pedigree Relationships. *OENO One* **2021**, *55*, 101–114. [[CrossRef](#)]
105. Botstein, D.; White, R.; Skolnick, M.; Davis, R. Construction of a genetic linkage map in man using restriction fragment length polymorphisms. *Am. J. Hum. Genet.* **1980**, *32*, 314–331. [[PubMed](#)]
106. Vervalle, J.A.; Costantini, L.; Lorenzi, S.; Pindo, M.; Mora, R.; Bolognesi, G.; Marini, M.; Lashbrooke, J.G.; Tobutt, K.R.; Vivier, M.A.; et al. A high-density integrated map for grapevine based on three mapping populations genotyped by the Vitis18K SNP chip. *TAG. Theor. Appl. Genet.* **2022**, *21*, 1–20. [[CrossRef](#)]
107. Duduk, B.; Botti, S.; Ivanović, M.; Krstić, B.; Dukić, N.; Bertaccini, A. Identification of *Phytoplasmas* associated with grapevine yellows in Serbia. *J. Phytopathol.* **2004**, *152*, 575–579. [[CrossRef](#)]
108. Wang, Z.; Wang, Y.; Cao, X.; Wu, D.; Hui, M.; Han, X.; Yao, F.; Li, Y.; Li, H.; Wang, H. Screening and validation of ssr molecular markers for identification of downy mildew resistance in intraspecific hybrid F1 progeny (*V. vinifera*). *Horticulturae* **2022**, *8*, 706. [[CrossRef](#)]
109. Goszczynski, D.E.; Jooste, A.E.C. The application of single-strand conformation polymorphism (SSCP) technique for the analysis of molecular heterogeneity of grapevine virus A. *Vitis* **2002**, *41*, 77–82.
110. Aroca, A.; Raposo, R. PCR-based strategy to detect and identify species of *Phaeoacremonium* causing grapevine diseases. *Appl. Environ. Microbiol.* **2007**, *73*, 2911–2918. [[CrossRef](#)]
111. Wang, X.C.; Guo, L.; Shangguan, L.F.; Wang, C.; Yang, G.; Qu, S.C.; Fang, J.G. Analysis of expressed sequence tags from grapevine flower and fruit and development of simple sequence repeat markers. *Mol. Biol. Rep.* **2012**, *39*, 6825–6834. [[CrossRef](#)]
112. Crăciunaş, C.; Coste, A.; Farago, M.; Iliescu, M.; Iuorăş, R.; Butiuc-Keul, A. Genetic stability of several cultivars of grapevine cultivated in vitro. *Acta Hortic.* **2009**, *812*, 515–520. [[CrossRef](#)]
113. Schneider, S.; Reustle, G.; Zyprian, E. Detection of somaclonal variation in grapevine regenerants from protoplasts by RAPD-PCR. *Vitis* **1996**, *35*, 99–100.
114. Baránek, M.; Raddová, J.; Krizan, B.; Pidra, M. Genetic changes in grapevine genomes after stress induced by in vitro cultivation, thermotherapy and virus infection, as revealed by AFLP. *Genet. Mol. Biol.* **2009**, *32*, 834–839. [[CrossRef](#)] [[PubMed](#)]
115. Gribaudo, I.; Torello Marinoni, D.; Gambino, G.; Mannini, F.; Akkak, A.; Botta, R. Assessment of genetic fidelity in regenerants from two *Vitis vinifera* cultivars. *Acta Hortic.* **2009**, *827*, 131–136. [[CrossRef](#)]
116. Aljuaid, B.S.; Ismail, I.A.; Attia, A.O.; El Dessoky, S. Genetic Stability of in vitro Propagated Grapevine (*Vitis vinifera* L.) cv. Al-Bayadi. *J. Agric. Crops* **2022**, *8*, 12–19. [[CrossRef](#)]
117. Schellenbaum, P.; Mohler, V.; Wenzel, G.; Walter, B. Variation in DNA methylation patterns of grapevine somaclones (*Vitis vinifera* L.). *BMC Plant. Biol.* **2008**, *8*, 78. [[CrossRef](#)] [[PubMed](#)]
118. Dalbò, M.A. Genetic mapping, QTL Analysis, and Marker-Assisted Selection for Disease Resistance Loci in Grapes. Ph.D. Thesis, Cornell University, Ithaca, NY, USA, August 1998.
119. Doligez, A.; Bouquet, A.; Danglot, Y.; Lahogue, F.; Riaz, S.; Meredith, P.; Edwards, J.; This, P. Genetic mapping of grapevine (*Vitis vinifera* L.) applied to the detection of QTLs for seedlessness and berry weight. TAG. Theoretical and applied genetics. *Theor. Und Angew. Genet.* **2002**, *105*, 780–795. [[CrossRef](#)]
120. Fanizza, G.; Lamaj, F.; Costantini, L.; Chaabane, R.; Grando, M.S. QTL analysis for fruit yield components in table grapes (*Vitis vinifera*). *Appl. Genet.* **2005**, *111*, 658–664. [[CrossRef](#)] [[PubMed](#)]
121. Costantini, L.; Moreira, F.; Zyprians, E.; Martínez-Zapater, J.; Grando, M. *Molecular Maps, QTL Mapping & Association Mapping In Grapevine Molecular Physiology & Biotechnology*; Roubelakis-Angelakis, K.A., Ed.; Springer: Dordrecht, The Netherland, 2009. [[CrossRef](#)]
122. Demmings, E.M.; Williams, B.R.; Lee, C.-R.; Barba, P.; Yang, S.; Hwang, C.-F.; Reisch, B.I.; Chitwood, D.H.; Londo, J.P. Quantitative trait locus analysis of leaf morphology indicates conserved shape loci in grapevine. *Front. Plant. Sci.* **2019**, *10*, 1373. [[CrossRef](#)]
123. Su, K.; Guo, Y.; Zhong, W.; Lin, H.; Liu, Z.; Li, K.; Li, Y.; Guo, X. High-Density Genetic Linkage Map Construction and White Rot Resistance Quantitative Trait Loci Mapping for Genus *Vitis* Based on Restriction Site-Associated DNA Sequencing. *Phytopathology* **2021**, *111*, 659–670. [[CrossRef](#)]
124. Su, K.; Xing, H.; Guo, Y.; Zhao, F.; Liu, Z.; Li, K.; Li, Y.; Guo, X. High-density genetic linkage map construction and cane cold hardiness QTL mapping for *Vitis* based on restriction site-associated DNA sequencing. *BMC Genom.* **2020**, *21*, 419. [[CrossRef](#)]
125. Reshef, N.; Karn, A.; Manns, D.C.; Mansfield, A.K.; Cadle-Davidson, L.; Reisch, B.; Sacks, G.L. Stable QTL for malate levels in ripe fruit and their transferability across *Vitis* species. *Hortic. Res.* **2022**, *9*, uhac009. [[CrossRef](#)]
126. Wang, H.; Yan, A.; Sun, L.; Zhang, G.; Wang, X.; Ren, J.; Xu, H. Novel stable QTLs identification for berry quality traits based on high-density genetic linkage map construction in table grape. *BMC Plant. Biol.* **2020**, *20*, 411. [[CrossRef](#)] [[PubMed](#)]
127. Fu, P.; Wu, W.; Lai, G.; Li, R.; Peng, Y.; Yang, B.; Wang, B.; Yin, L.; Qu, J.; Song, S.; et al. Identifying *Plasmopara viticola* resistance loci in grapevine (*Vitis amurensis*) via genotyping-by-sequencing-based QTL mapping. *PPB* **2020**, *154*, 75–84. [[CrossRef](#)]
128. Fehér, A. Callus, dedifferentiation, totipotency, somatic embryogenesis: What these terms mean in the era of molecular plant biology? *Front. Plant. Sci.* **2019**, *10*, 536. [[CrossRef](#)] [[PubMed](#)]

129. Nuzzo, F.; Gambino, G.; Perrone, I. Unlocking grapevine in vitro regeneration: Issues and perspectives for genetic improvement and functional genomic studies. *Plant. Physiol. Biochem.* **2022**, *193*, 99–109. [[CrossRef](#)]
130. Butiuc-Keul, A.; Coste, A.; Oltean, B.; Crăciunaş, C.; Halmagyi, A.; Deliu, C.; Farago, M.; Iliescu, M.; Iuoraş, R. In vitro clonal propagation of several grapevine cultivars. *Acta Hortic.* **2009**, *843*, 151–156. [[CrossRef](#)]
131. Butiuc-Keul, A.; Coste, A.; Crăciunaş, C. Molecular characterization and in vitro preservation of some grapevine cultivars. *Rom. Biotechnol. Lett.* **2011**, *16*, 6226–6233.
132. Krishna, H.; Alizadeh, M.; Singh, D.; Singh, U.; Chauhan, N.; Eftekhari, M.; Sadh, R.K. Somaclonal variations and their applications in horticultural crops improvement. *3 Biotech* **2016**, *6*, 54. [[CrossRef](#)] [[PubMed](#)]
133. Atak, A. New Perspectives in Grapevine (*Vitis* spp.) Breeding. In *Plant Breeding—New Perspectives*; Wang, H., Ed.; IntechOpen: London, UK, 2022; Available online: <https://www.intechopen.com/online-first/82151> (accessed on 15 November 2022). [[CrossRef](#)]
134. Mukherjee, P.; Husain, N.; Misra, S.C.; Rao, V.S. In vitro propagation of a grape rootstock, deGrasset *Vitis champinii* Planch. Effects of medium compositions and plant growth regulators. *Sci. Hortic.* **2010**, *126*, 13–19. [[CrossRef](#)]
135. Yerbolova, L.S.; Ryabushkina, N.A.; Oleichenko, S.N.; Kampitova, G.A.; Galiakparov, N. The effect of growth regulators on in vitro culture of some *Vitis vinifera* L. cultivars. *World Appl. Sci. J.* **2013**, *23*, 76–80. [[CrossRef](#)]
136. Capriotti, L.; Baraldi, E.; Mezzetti, B.; Limera, C.; Sabbadini, S. Biotechnological approaches: Gene overexpression, gene silencing, and genome editing to control fungal and oomycete diseases in grapevine. *Int. J. Molec. Sci.* **2020**, *21*, 5701. [[CrossRef](#)]
137. Nicholson, K.L.; Tarlyn, N.; Armour, T.; Swanson, M.E.; Dhingra, A. Effect of phyllotactic position and cultural treatments toward successful direct shoot organogenesis in dwarf ‘Pixie’ grapevine (*Vitis vinifera* L.). *Plant. Cell Tiss. Organ. Cult.* **2012**, *111*, 123–129. [[CrossRef](#)]
138. Zhang, P.; Yu, Z.Y.; Cheng, Z.M.; Zhang, Z.; Tao, J.M. In vitro explants regeneration of the grape ‘Wink’ (*Vitis vinifera* L. ‘Wink’). *J. Plant Breed. Crop Sci.* **2011**, *3*, 276–282.
139. Park, H.J.; Lee, H.R.; Pyee, J.; Cha, H.C. Regeneration of grape (*Vitis labruscana* cv. Kyoho) by shoot-tip culture. *J. Plant. Biol.* **2001**, *44*, 185–192. [[CrossRef](#)]
140. Pool, R.M. Effect of cytokinin on in vitro development of ‘concord’ flowers. *Am. J. Enol. Vitic.* **1975**, *26*, 43–46.
141. Lilov, D.; Isvorska, N. Flower bud initiation from isolated meristem tissues of grapevine tendrils. *Fisiol. Rast.* **1978**, *4*, 73–78.
142. Barreto, M.S. In Vitro Plant Regeneration and Transformation Studies in Grape (*Vitis vinifera* L.). Ph.D. Thesis, University of Pune, Pune, India.
143. Torregrosa, L. Culture in vitro et Transformation Genetique de la Vigne. These, Ecole Nationale Supérieure Agronomique de Montpellier, France, 1994. Available online: <https://www.theses.fr/1994ENSA0024> (accessed on 12 September 2022).
144. Torregrosa, L.; Bouquet, A. *Agrobacterium tumefaciens* and *A. rhizogenes-rhizogenes* cotransformation to obtain grapevine hair roots producing coat protein of grapevine chrome mosaic nepo virus. *Plant. Cell Tiss. Org. Cult.* **1997**, *49*, 59–63. [[CrossRef](#)]
145. Loubser, J.T.; Meyer, A.J. Dual cultures of *Meloidogyne javanica* and grapevine rootstocks on artificial media. *S. Afr. J. Enol. Vitic.* **1990**, *11*, 42–45. [[CrossRef](#)]
146. Bavaresco, L.; Walker, M.A. Techniques for successfully establishing Xiphinema index in dual culture with grapes. *Am. J. Enol. Vitic.* **1994**, *45*, 273–277.
147. Forneck, A.; Merkt, N.; Blaich, R. A tripartite asceptic culture system for grapes (*Vitis* spp.), phylloxera (*Daktulosphaira vitifoliae*) and mites (*Tarsanemus* sp.). *Vitis* **1998**, *37*, 95–96.
148. Kellow, A.V. A Study of The Interaction between Susceptible and Resistant Grapevine and Phylloxera. Ph.D. Thesis, Natural and Agricultural Resource Sciences, Department of Horticulture, Viticulture and Oenology, The University of Adelaide, Adelaide, Australia, June 2000.
149. Saygaç, S.; Önder, S. Effects of shoot tip size on in vitro regeneration and virus elimination of grapevine cv. Superior Seedless. *Plant. Prot. Bull.* **2021**, *61*, 5–9. [[CrossRef](#)]
150. Mourad, A. Elimination of grapevine bois noir phytoplasma by tissue culture coupled or not with heat therapy or hot water treatment. *Adv. Crop. Sci. Technol.* **2013**, *1*, 1–4. [[CrossRef](#)]
151. Miljanić, V.; Rusjan, D.; Škvarč, A.; Chatelet, P.; Štajner, N. Elimination of eight viruses and two viroids from preclonal candidates of six grapevine varieties (*Vitis vinifera* L.) through in vivo thermotherapy and in vitro meristem tip micrografting. *Plants* **2022**, *11*, 1064. [[CrossRef](#)] [[PubMed](#)]
152. Malenica, N.; Jagić, M.; Pavletić, B.; Bauer, N.; Vončina, D.; Zdunić, G.; Levanić, D.L. Somatic embryogenesis as a tool for virus elimination in Croatian indigenous grapevine cultivars. *Acta Bot. Croat.* **2020**, *79*, 26–34. [[CrossRef](#)]
153. Pérez-Núñez, M.T.; Chan, J.L.; Sáenz, L.; González, T.; Verdeil, J.L.; Oropeza, C. Improved somatic embryogenesis from *Cocos nucifera* (L.) plumule explants. *Vitr. Cell. Dev. Biol. Plant.* **2006**, *42*, 37–43. [[CrossRef](#)]
154. Forleo, L.R.; D’Amico, M.; Basile, T.; Marsico, A.D.; Cardone, M.F.; Maggiolini, F.A.M.; Velasco, R.; Bergamini, C. Somatic embryogenesis in *Vitis* for genome editing: Optimization of protocols for recalcitrant genotypes. *Horticulturae* **2021**, *7*, 511. [[CrossRef](#)]
155. Dal Santo, S.; De Paoli, E.; Pagliarani, C.; Amato, A.; Celii, M.; Boccacci, P.; Zenoni, S.; Gambino, G.; Perrone, I. Stress responses and epigenomic instability mark the loss of somatic embryogenesis competence in grapevine. *Plant. Physiol.* **2022**, *188*, 490–508. [[CrossRef](#)]

156. Capriotti, L.; Limera, C.; Mezzetti, B.; Ricci, A.; Sabbadini, S. From induction to embryo proliferation: Improved somatic embryogenesis protocol in grapevine for Italian cultivars and hybrid *Vitis* rootstocks. *Plant. Cell Tiss. Organ. Cult.* **2022**, *151*, 221–233. [[CrossRef](#)]
157. Horstman, A.; Bemer, M.; Boutilier, K. A transcriptional view on somatic embryogenesis. *Regeneration* **2017**, *4*, 201–216. [[CrossRef](#)]
158. Ghadirzadeh-Khorzoghi, E.; Jahanbakhshian-Davaran, Z.; Seyedi, S.M. Direct somatic embryogenesis of drought resistance pistachio (*Pistacia vera* L.) and expression analysis of somatic embryogenesis-related genes. *S. Afr. J. Bot.* **2019**, *121*, 558–567. [[CrossRef](#)]
159. Catalano, C.; Abbate, L.; Motisi, A.; Crucitti, D.; Cangelosi, V.; Pisciotta, A.; Di Lorenzo, R.; Carimi, F.; Carra, A. Autotetraploid emergence via somatic embryogenesis in *Vitis vinifera* induces marked morphological changes in shoots, mature leaves, and stomata. *Cells* **2021**, *10*, 1336. [[CrossRef](#)] [[PubMed](#)]
160. Corredoira, E.; Merkle, S.A.; Martínez, M.T.; Toribio, M.; Canhoto, J.M.; Correia, S.I.; Ballester, A.; Vieitez, A. Non-zygotic embryogenesis in hardwood species. *Crit Rev. Plant. Sci.* **2019**, *38*, 29–97. [[CrossRef](#)]
161. Maillet, P.; Lebel, S.; Schellenbaum, P.; Jacques, A.; Walter, B. Differential regulation of SERK, LEC1-like and pathogenesis-related genes during indirect secondary somatic embryogenesis in grapevine. *Plant. Physiol. Biochem.* **2009**, *47*, 743–752. [[CrossRef](#)]
162. Ricci, A.; Sabbadini, S.; Prieto, H.; Padilla, I.M.; Dardick, C.; Li, Z.; Scorza, R.; Limera, C.; Mezzetti, B.; Perez-Jimenez, M.; et al. Genetic transformation in peach (*Prunus persica* L.): Challenges and ways forward. *Plants* **2020**, *9*, 971. [[CrossRef](#)]
163. Phillips, R.L.; Kaepller, S.M.; Olhoff, P. Genetic instability of plant tissue cultures: Breakdown of normal controls. *Proc. Natl. Acad. Sci. USA* **1994**, *91*, 5222–5226. [[CrossRef](#)] [[PubMed](#)]
164. Muler, E.; Brown, P.T.H.; Hartke, S.; Lorz, H. DNA variation in tissue-culture-derived rice plants. *Theor. Appl. Genet.* **1990**, *80*, 673–679. [[CrossRef](#)]
165. Kuksova, V.B.; Piven, N.M.; Gleba, Y.Y. Somaclonal variation and in vitro induced mutagenesis in grapevine. *Plant. Cell Tiss. Organ. Cult.* **1997**, *49*, 17–27. [[CrossRef](#)]
166. Palombi, M.A.; Damiano, C. Comparison between RAPD and SSR molecular markers in detecting genetic variation in kiwifruit (*Actinidia deliciosa* A. Chev.). *Plant. Cell Rep.* **2002**, *20*, 1061–1066. [[CrossRef](#)]
167. Kaepller, S.M.; Phillips, R.L. DNA methylation and tissue culture-induced variation in plants. *Vitr. Cell Dev. Biol. Plant.* **1993**, *29*, 125–130. [[CrossRef](#)]
168. Nivas, S.K.D.; Souza, L. Genetic fidelity in micropropagated plantlets of *Anacardium occidentale* L. (Cashew) an important fruit tree. *Int. J. Sci. Res.* **2014**, *3*, 2142–2146.
169. Smulders, M.; de Klerk, G. Epigenetics in plant tissue culture. *Plant. Growth Regul.* **2011**, *63*, 137–146. [[CrossRef](#)]
170. Prado, M.J.; Rodriguez, E.; Rey, L.; González, M.V.; Santos, C.; Rey, M. Detection of somaclonal variants in somatic embryogenesis-regenerated plants of *Vitis vinifera* by flow cytometry and microsatellite markers. *Plant. Cell Tiss. Organ. Cult.* **2010**, *103*, 49–59. [[CrossRef](#)]
171. Marcotrigiano, M.; Bernatzky, R. Arrangement of cell layers in the shoot apical meristems of periclinal chimeras influences cell fate. *Plant. J.* **1995**, *7*, 193–202. [[CrossRef](#)]
172. Burge, G.K.; Morgan, E.R.; Seelye, J.F. Opportunities for synthetic plant chimeral breeding: Past and future. *Plant. Cell Tiss. Organ. Cult.* **2002**, *70*, 13–21. [[CrossRef](#)]
173. Franks, T.; Botta, R.; Thomas, M.R.; Franks, J. Chimerism in grapevines: Implications for cultivar identity, ancestry and genetic improvement. *Theor. Appl. Genet.* **2002**, *104*, 192–199. [[CrossRef](#)] [[PubMed](#)]
174. Bertsch, C.; Kieffer, F.; Maillot, P.; Farine, S.; Butterlin, G.; Merdinoglu, D.; Walter, B. Genetic chimerism of *Vitis vinifera* cv. Chardonnay 96 is maintained through organogenesis but not somatic embryogenesis. *BMC Plant. Biol.* **2005**, *5*, 20. [[CrossRef](#)]
175. Riaz, S.; Garrison, K.E.; Dangl, G.S.; Boursiquot, J.M.; Meredith, C.P. Genetic divergence and chimerism within ancient asexually propagated winegrape cultivars. *J. Amer. Soc. Hort. Sci.* **2002**, *127*, 508–514. [[CrossRef](#)]
176. Quackenbush, J.; Liang, F.; Holt, I.; Pertea, G.; Upton, J. The TIGR gene indices: Reconstruction and representation of expressed gene sequences. *Nucleic Acids Res.* **2000**, *28*, 141–145. [[CrossRef](#)]
177. Siemann, E.; Creasy, L. Concentration of the phytoalexin resveratrol in wine. *Am. J. Enol. Viticult.* **1992**, *43*, 49–52.
178. Sotheeswaran, S.; Pasupathy, V. Distribution of resveratrol oligomers in plants. *Photochemistry* **1993**, *32*, 1083–1092. [[CrossRef](#)]
179. Kikkert, J.K.; Ali, G.S.; Wallace, P.G.; Reisch, B.; Reustle, G.M. Expression of fungal chitinase in *Vitis vinifera* L. ‘Merlot’ and ‘Chardonnay’ plants produced by biolistic transformation. *Acta Hort.* **2000**, *528*, 297–303. [[CrossRef](#)]
180. Tamaoki, M.; Imai, H.; Takahashi, H.; Toda, Y.; Niwa, Y.; Nakajima, N.; Aono, M.; Kubo, A.; Saji, H. Development of visible markers for transgenic plants and their availability for environmental risk assessment. *Z. Für Nat.* **2006**, *61*, 377–386. [[CrossRef](#)] [[PubMed](#)]
181. Braidot, E.; Zancani, M.; Petruzza, E.; Peresson, C.; Bertolini, A.; Patui, S.; Macri, F.; Vianello, A. Transport and accumulation of flavonoids in grapevine (*Vitis vinifera* L.). *Plant. Signal. Behav.* **2008**, *3*, 626–632. [[CrossRef](#)] [[PubMed](#)]
182. Basha, S.M.; Musingo, M.; Colova, V.S. Compositional differences in the phenolics compounds of muscadine and bunch grape wines. *Afr. J. Biotechnol.* **2004**, *10*, 523–528. [[CrossRef](#)]
183. INRA-CNRCV The French Plant Genomic Resource Center. Available online: <http://cnrcv.toulouse.inra.fr> (accessed on 10 October 2022).
184. Huang, X.; Wei, X.; Sang, T.; Zhao, Q.; Feng, Q.; Zhao, Y.; Li, C.; Zhy, C.; Lu, T.; Zhang, Z.; et al. Genome-wide association studies of 14 agronomic traits in rice landraces. *Nat. Genet.* **2010**, *42*, 961–967. [[CrossRef](#)] [[PubMed](#)]

185. Meuwissen, T.H.E.; Hayes, B.J.; Goddard, M.E. Prediction of Total Genetic Value Using Genome-Wide Dense Marker Maps. *Genetics* **2001**, *157*, 1819–1829. [CrossRef]
186. Jannink, J.-L.; Lorenz, A.J.; Iwata, H. Genomic selection in plant breeding: From theory to practice. *Brief. Funct. Genom.* **2010**, *9*, 166–177. [CrossRef]
187. Nakaya, A.; Isobe, S.N. Will genomic selection be a practical method for plant breeding? *Ann. Bot.* **2012**, *110*, 1303–1316. [CrossRef]
188. Fodor, A.; Segura, V.; Denis, M.; Neuenschwander, S.; Fournier-Level, A.; Chatelet, P.; Homa, F.A.; Lacombe, T.; This, P.; Le Cunff, L. Genome-wide prediction methods in highly diverse and heterozygous species: Proof-of-concept through simulation in grapevine. *PLoS ONE* **2014**, *9*, e110436. [CrossRef]
189. Wang, M.; Jiang, N.; Jia, T.; Leach, L.; Cockram, J.; Waugh, R.; Ramsay, L.; Thomas, B.; Luo, Z. Genome-wide association mapping of agronomic and morphologic traits in highly structured populations of barley cultivars. *Theor. Appl. Genet.* **2012**, *124*, 233–246. [CrossRef]
190. Laucou, V.; Lacombe, T.; Dechesne, F.; Siret, R.; Bruno, J.-P.; Dessup, M.; Dessup, T.; Ortigosa, P.; Parra, P.; Roux, C. High throughput analysis of grape genetic diversity as a tool for germplasm collection management. *Theor. Appl. Genet.* **2011**, *122*, 1233–1245. [CrossRef] [PubMed]
191. Myles, S.; Chia, J.-M.; Hurwitz, B.; Simon, C.; Zhong, G.Y.; Buckler, E.; Ware, D. Rapid Genomic Characterization of the Genus *Vitis*. *PLoS ONE* **2010**, *5*, e8219. [CrossRef] [PubMed]
192. Bowers, J.; Boursiquot, J.M.; This, P.; Chu, K.; Johansson, H.; Meredith, C. Historical Genetics: The Parentage of Chardonnay, Gamay, and Other Wine Grapes of Northeastern France. *Science* **1999**, *285*, 1562–1565. [CrossRef]
193. Boursiquot, J.-M.; Lacombe, T.; Laucou, V.; Julliard, S.; Perrin, F.-X.; Lanier, N.; Legrand, D.; Meredith, C.; This, P. Parentage of Merlot and related winegrape cultivars of southwestern France: Discovery of the missing link. *Aust. J. Grape Wine Res.* **2009**, *15*, 144–155. [CrossRef]
194. Myles, S.; Boyko, A.R.; Owens, C.L.; Brown, P.J.; Grassi, F.; Aradhya, M.K.; Prins, B.; Reynolds, A.; Chia, I.-M.; Ware, D.; et al. Genetic Structure and Domestication History of the Grape. *Proc. Natl. Acad. Sci. USA* **2011**, *108*, 3530–3535. [CrossRef] [PubMed]
195. Lacombe, T.; Boursiquot, J.-M.; Laucou, V.; Di Vecchi-Staraz, M.; Péros, J.-P.; This, P. Large-scale parentage analysis in an extended set of grapevine cultivars (*Vitis vinifera* L.). *Theor. Appl. Genet.* **2012**, *126*, 1–14. [CrossRef]
196. Hannah, L.; Roehrdanz, P.R.; Ikegami, M.; Shepard, A.V.; Shaw, M.R.; Tabor, G.; Zhi, L.; Marquet, P.A.; Hijmans, R.J. Climate change, wine, and conservation. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, 6907–6912. [CrossRef] [PubMed]
197. Moriondo, M.; Jones, G.V.; Bois, B.; Dibari, C.; Ferrise, R.; Trombi, G.; Bindi, M. Projected shifts of wine regions in response to climate change. *Clim. Change* **2013**, *119*, 825–839. [CrossRef]
198. Gray, D.J.; Jayasankar, S.; Li, Z.T. Vitaceae (Grape *Vitis* spp.). In *Biotechnology of Fruit and Nut Crops. Biotechnology in Agriculture Series*; No. 29; Litz, R.E., Ed.; CAB International: Wallingford, UK, 2005; Volume Chapter 22, pp. 672–706.
199. Maraš, V.; Tello, J.; Gazivoda, A.; Mugoša, M.; Perišić, M.; Raičević, J.; Štajner, N.; Ocete, R.; Božović, V.; Popović, T.; et al. Population genetic analysis in old Montenegrin vineyards reveals ancient ways currently active to generate diversity in *Vitis vinifera*. *Sci. Rep.* **2020**, *10*, 15000. [CrossRef] [PubMed]
200. Čuš, F.; Cesnik, H.B.; Bolta, S.V.; Gregorcic, A. Pesticide residues and microbio-logical quality of bottled wines. *Food Control* **2010**, *21*, 150–154. [CrossRef]
201. Pretorius, I.S.; Hej, P.B. Grape and wine biotechnology: Challenges, opportunities and potential benefits. *Aust. J. Grape Wine Res.* **2006**, *11*, 83–108. [CrossRef]
202. Gray, D.J.; Li, Z.T.; Dhekney, S.A. Precision breeding of grapevine (*Vitis vinifera* L.) for improved traits. *Plant. Sci.* **2014**, *228*, 3–10. [CrossRef] [PubMed]
203. Li, Z.T.; Dhekney, S.A.; Dutt, M.; Van Aman, M.; Tattersall, J.; Kelley, K.T.; Gray, D.J. Optimizing *Agrobacterium*-mediated transformation of grapevine. *Vitr. Cell. Dev. Biol. Plant.* **2006**, *42*, 220–227. [CrossRef]
204. Li, Z.T.; Dhekney, S.A.; Dutt, M.; Gray, D.J. An improved protocol for *Agrobacterium*-mediated transformation of grapevine (*Vitis vinifera* L.). *Plant. Cell Tiss. Organ. Cult.* **2008**, *93*, 311–321. [CrossRef]
205. Torregrosa, L.; Rienth, M.; Luchaire, N.; Novelli, F.; Bigard, A.; Chatbanyong, R.; Lopez, G.; Farnos, M.; Roux, C.; Adivèze, A.; et al. The Microvine, A Biological Model, Very Versatile and Efficient to Boost Grapevine Research in Physiology and Genetics. In Proceedings of the 39th OIV Meeting, Bento Goncalvez, Brazil, 24–28 October 2016.
206. Dalla Costa, L.; Piazza, S.; Campa, M.; Flachowsky, H.; Hanke, M.V.; Malnoy, M. Efficient heat-shock removal of the selectable marker gene in genetically modified grapevine. *Plant. Cell Tiss. Organ. Cult.* **2016**, *124*, 471–481. [CrossRef]
207. Muruganantham, M.; Moskovitz, Y.; Haviv, S.; Horesh, T.; Fenigstein, A.; Preez, J.D.; Stephan, D.; Burger, J.T.; Mawassi, M. Grapevine virusA-mediated gene silencing in Nicotiana benthamiana and *Vitis vinifera*. *J. Virol. Methods* **2009**, *155*, 167–174. [CrossRef] [PubMed]
208. Kurth, E.G.; Peremyslov, V.V.; Prokhnevsky, A.I.; Kasschau, K.D.; Miller, M.; Carrington, J.C.; Dolja, V.V. Virus-derived gene expression and RNA interference vector for grapevine. *J. Virol.* **2012**, *86*, 6002–6009. [CrossRef] [PubMed]
209. Meng, B.; Venkataraman, S.; Li, C.; Wang, W.; Dayan-Glick, C.; Mawassi, M. Construction and biological activities of the first infectious cDNA clones of the genus Foveavirus. *Virology* **2013**, *435*, 453–462. [CrossRef]
210. Wood, K. Marker proteins for gene expression. *Curr. Opin. Biotechnol.* **1995**, *6*, 50–58. [CrossRef]

211. Rosellini, D. Selectable markers and reporter genes. A well furnished tool-box for plant science and genetic engineering. *Crit. Rev. Plant. Sci.* **2012**, *31*, 401–453. [[CrossRef](#)]
212. Dutt, M.; Li, Z.T.; Dhekney, S.A.; Gray, D.J. A co-transformation system to produce transgenic grapevines free of marker genes. *Plant. Sci.* **2008**, *175*, 423–430. [[CrossRef](#)]
213. Li, Z.T.; Dhekney, S.A.; Gray, D.J. Use of the *VvMybA1* gene for non-destructive quantification of promoter activity via color histogram analysis in grapevine (*Vitis vinifera*) and tobacco. *Transgenic Res.* **2011**, *20*, 1087–1097. [[CrossRef](#)]
214. Dutt, M.; Li, Z.T.; Gray, D.J.; Gmitter, F.; Grosser, J.W. Purple citrus? Utilization of a Myb-related transcription factor gene for Anthocyanin production. *AMER Soc. Hortic. Sci.* **2013**, *48*, 10–11.
215. Li, Z.T.; Jayasankar, S.; Gray, D.J. Expression of a bifunctional green fluorescent protein (GFP) fusion marker under the control of three constitutive promoters and enhanced derivatives in transgenic grape (*Vitis vinifera*). *Plant. Sci.* **2001**, *160*, 877–887. [[CrossRef](#)] [[PubMed](#)]
216. Conn, S.; Zhang, W.; Franco, C. Anthocyanic vacuolar inclusions (AVIs) selectively bind acylated anthocyanins in *Vitis vinifera* L. (grapevine) suspension culture. *Biotechnol. Lett.* **2003**, *25*, 835–839. [[CrossRef](#)] [[PubMed](#)]
217. Li, Z.T.; Kim, K.-H.; Jasinski, J.R.; Creech, M.R.; Gray, D.J. Large-scale characterization of promoters from grapevine (*Vitis* spp.) using quantitative anthocyanin and GUS assay systems. *Plant. Sci.* **2012**, *196*, 132–142. [[CrossRef](#)]
218. Pilati, S.; Perazzolli, M.; Malossini, A.; Cestaro, A.; Damattè, L.; Fontana, P.; Dal Ri, A.; Viola, R.; Velasco, R.; Moser, C. Genome-wide transcriptional analysis of grapevine berry ripening reveals a set of genes similarly modulated during three seasons and the occurrence of an oxidative burst at veraison. *BMC Genom.* **2007**, *8*, 428. [[CrossRef](#)]
219. Mezzetti, B.; Pandolfini, T.; Navacchi, O.; Landi, L. Genetic transformation of *Vitis vinifera* via organogenesis. *BMC Biotechnol.* **2002**, *2*, 18. [[CrossRef](#)]
220. Xie, X.; Agüero, C.B.; Wang, Y.; Andrew Walker, M. Genetic transformation of grape varieties and rootstocks via organogenesis. *Plant. Cell Tiss. Organ. Cult.* **2016**, *126*, 541–552. [[CrossRef](#)]
221. Sabbadini, S.; Capriotti, L.; Molesini, B.; Pandolfini, T.; Navacchi, O.; Limera, C.; Ricci, A.; Mezzetti, B. Comparison of regeneration capacity and *Agrobacterium*-mediated cell transformation efficiency of different cultivars and rootstocks of *Vitis* spp. via organogenesis. *Sci. Rep.* **2019**, *9*, 582. [[CrossRef](#)]
222. de Carvalho, D.C.; da Silva, A.L.L.; Schuck, M.R.; Purcino, M.; Tanno, G.N.; Biasi, L.A. Fox grape cv. Bordô (*Vitis labrusca* L.) and grapevine cv. Chardonnay (*Vitis vinifera* L.) cultivated in vitro under different carbohydrates, amino acids and 6-benzylaminopurine levels. *Brazil. Arch. Biol. Technol.* **2013**, *56*, 191–201. [[CrossRef](#)]
223. Colby, S.M.; Juncosa, A.M.; Meredith, C.P. Cellular differences in *Agrobacterium* susceptibility and regenerative capacity restrict the development of transgenic grapevines. *J. Amer. Soc. Hort. Sci.* **1991**, *116*, 356–361. [[CrossRef](#)]
224. Maillot, P.; Deglène-Benbrahim, L.; Walter, B. Efficient somatic embryogenesis from meristematic explants in grapevine (*Vitis vinifera* L.) cv. Chardonnay: An improved protocol. *Trees* **2016**, *30*, 1377–1387. [[CrossRef](#)]
225. Zhou, Q.; Dai, L.; Cheng, S.; He, J.; Wang, D.; Zhang, J.; Wang, Y. A circulatory system useful both for long-term somatic embryogenesis and genetic transformation in *Vitis vinifera* L. cv. Thompson Seedless. *Plant. Cell Tiss. Organ. Cult.* **2014**, *118*, 157–168. [[CrossRef](#)]
226. Saporta, R.; San Pedro, T.; Gisbert, C. Attempts at grapevine (*Vitis vinifera* L.) breeding through genetic transformation: The main limiting factors. *Vitis* **2016**, *55*, 173–186. [[CrossRef](#)]
227. Wang, X.; Tu, M.; Wang, D.; Liu, J.; Li, Y.; Li, Z.; Wang, Y.; Wang, X. CRISPR/Cas9-mediated efficient targeted mutagenesis in grape in the first generation. *Plant. Biotechnol. J.* **2018**, *16*, 844–855. [[CrossRef](#)]
228. Carra, A.; Sajeva, M.; Abbate, L.; Siragusa, M.; Pathirana, R.; Carimi, F. Factors affecting somatic embryogenesis in eight Italian grapevine cultivars and the genetic stability of embryo-derived regenerants as assessed by molecular markers. *Sci. Hort.* **2016**, *204*, 123–127. [[CrossRef](#)]
229. de Carvalho, D.C.; da Silva, A.L.L.; Tanno, G.N.; Purcino, M.; Biasi, L.A. Organogenesis from leaf segments and internodes of grapevine cv. Merlot. *Ciência Agropecuária* **2011**, *35*, 108–114. [[CrossRef](#)]
230. Perrin, M.; Martin, D.; Joly, D.; Demangeat, G.; This, P.; Masson, J.E. Medium-dependent response of grapevine somatic embryogenic cells. *Plant. Sci.* **2001**, *161*, 107–116. [[CrossRef](#)]
231. Jelly, N.S.; Valat, L.; Walter, B.; Maillot, P. Transient expression assays in grapevine: A step towards genetic improvement. *Plant. Biotechnol. J.* **2014**, *12*, 1231–1245. [[CrossRef](#)]
232. Torregrosa, L.; Verries, C.; Tesniere, C. Grapevine (*Vitis vinifera* L.) promoter analysis by biolistic-mediated transient transformation of cell suspensions. *Vitis* **2002**, *41*, 27–32. [[CrossRef](#)]
233. Verries, C.; Pradal, M.; Chatelet, P.; Torregrosa, L.; Tesniere, C. Isolation and analysis of the promoter of *VvAdh2*, a grapevine *Vitis vinifera* L. ripening-related gene. *Plant. Sci.* **2004**, *167*, 1067–1074. [[CrossRef](#)]
234. Harris, N.N.; Luczo, J.M.; Robinson, S.P.; Walker, A.R. Transcriptional regulation of the three grapevine chalcone synthase genes and their role in flavonoid synthesis in Shiraz. *Aus. J. Grape Wine Res.* **2013**, *19*, 221–229. [[CrossRef](#)]
235. Deluc, L.; Bogs, J.; Walker, A.R.; Ferrier, T.; Decendit, A.; Merillon, J.M.; Robinson, S.P.; Barrieu, F. The transcription factor VvMYB5b contributes to the regulation of anthocyanin and proanthocyanidin biosynthesis in developing grape berries. *Plant. Physiol.* **2008**, *147*, 2041–2053. [[CrossRef](#)] [[PubMed](#)]
236. Walker, A.R.; Lee, E.; Bogs, J.; McDavid, D.A.; Thomas, M.R.; Robinson, S.P. White grapes arose through the mutation of two similar and adjacent regulatory genes. *Plant. J.* **2007**, *49*, 772–785. [[CrossRef](#)]

237. Czermel, S.; Stracke, R.; Weisshaar, B.; Cordon, N.; Harris, N.N.; Walker, A.R.; Robinson, S.P.; Bogs, J. The grapevine R2R3-MYB transcription factor VvMYBF1 regulates flavonol synthesis in developing grape berries. *Plant. Physiol.* **2009**, *151*, 1513–1530. [[CrossRef](#)]
238. Bogs, J.; Jaffe, F.W.; Takos, A.M.; Walker, A.R.; Robinson, S.P. The grapevine transcription factor VvMYBPA1 regulates proanthocyanidin synthesis during fruit development. *Plant. Physiol.* **2007**, *143*, 1347–1361. [[CrossRef](#)]
239. Hichri, I.; Heppel, S.C.; Pillet, J.; Leon, C.; Czermel, S.; Delrot, S.; Lauvergeat, V.; Bogs, J. The basic helix-loop-helix transcription factor MYC1 is involved in the regulation of the flavonoid biosynthesis pathway in grapevine. *Mol. Plant.* **2010**, *3*, 509–523. [[CrossRef](#)]
240. Höll, J.; Vannozzi, A.; Czermel, S.; D’Onofrio, C.; Walker, A.R.; Rausch, T.; Lucchin, M.; Boss, P.K.; Dry, I.B.; Bogs, J. The R2R3-MYB transcription factors MYB14 and MYB15 regulate stilbene biosynthesis in *Vitis vinifera*. *Plant. Cell* **2013**, *25*, 4135–4144. [[CrossRef](#)]
241. Joubert, D.A.; de Lorenzo, G.; Vivier, M.A. Regulation of the grapevine polygalacturonase-inhibiting protein encoding gene: Expression pattern, induction profile and promoter analysis. *J. Plant. Res.* **2013**, *126*, 267–281. [[CrossRef](#)]
242. Saumonneau, A.; Laloi, M.; Lallemand, M.; Rabot, A.; Atanassova, R. Dissection of the transcriptional regulation of grape ASR and response to glucose and abscisic acid. *J. Exp. Bot.* **2012**, *63*, 1495–1510. [[CrossRef](#)] [[PubMed](#)]
243. Marchive, C.; Leon, C.; Kappel, C.; Coutos-Thevenot, P.; Corio-Costet, M.F.; Delrot, S.; Lauvergeat, V. Over-expression of VvWRKY1 in grapevines induces expression of jasmonic acid pathway-related genes and confers higher tolerance to the downy mildew. *PLoS ONE* **2013**, *8*, e54185. [[CrossRef](#)]
244. Saumonneau, A.; Agasse, A.; Bidoyen, M.T.; Lallemand, M.; Cantereau, A.; Medici, A.; Laloi, M.; Atanassova, R. Interaction of grape ASR proteins with a DREB transcription factor in the nucleus. *FEBS Lett.* **2008**, *582*, 3281–3287. [[CrossRef](#)] [[PubMed](#)]
245. Urso, S.; Zottini, M.; Ruberti, C.; Schiavo, F.L.; Stanca, A.M.; Cattivelli, L.; Vale, G. An *Agrobacterium tumefaciens*-mediated gene silencing system for functional analysis in grapevine. *Plant. Cell Tiss. Organ. Cult.* **2013**, *114*, 49–60. [[CrossRef](#)]
246. Visser, M.; Stephan, D.; Jaynes, J.M.; Burger, J.T. A transient expression assay for the *in planta* efficacy screening of an antimicrobial peptide against grapevine bacterial pathogens. *Lett. Appl. Microbiol.* **2012**, *54*, 543–551. [[CrossRef](#)] [[PubMed](#)]
247. Bertazzon, N.; Raiola, A.; Castiglioni, C.; Gardiman, M.; Angelini, E.; Borgo, M.; Ferrari, S. Transient silencing of the grapevine gene VvPGIP1 by agroinfiltration with a construct for RNA interference. *Plant. Cell Rep.* **2012**, *31*, 133–143. [[CrossRef](#)]
248. Santos-Rosa, M.; Poutaraud, A.; Merdinoglu, D.; Mestre, P. Development of a transient expression system in grapevine via agro-infiltration. *Plant. Cell Rep.* **2008**, *27*, 1053–1063. [[CrossRef](#)]
249. Guan, X.; Zhao, H.; Xu, Y.; Wang, Y. Transient expression of glyoxal oxidase from the Chinese wild grape *Vitis pseudoreticulata* can suppress powdery mildew in a susceptible genotype. *Protoplasma* **2011**, *248*, 415–423. [[CrossRef](#)]
250. He, M.; Xu, Y.; Cao, J.; Zhu, Z.; Jiao, Y.; Wang, Y.; Guan, X.; Yang, Y.; Xu, W.; Fu, Z. Subcellular localization and functional analyses of a PR10 protein gene from *Vitis pseudoreticulata* in response to *Plasmopara viticola* infection. *Protoplasma* **2013**, *250*, 129–140. [[CrossRef](#)]
251. Xu, W.; Yu, Y.; Ding, J.; Hua, Z.; Wang, Y. Characterization of a novel stilbene synthase promoter involved in pathogen- and stress-inducible expression from Chinese wild *Vitis pseudoreticulata*. *Planta* **2010**, *231*, 475–487. [[CrossRef](#)]
252. Xu, T.F.; Zhao, X.C.; Jiao, Y.T.; Wei, J.Y.; Wang, L.; Xu, Y. A pathogenesis related protein, VpPR-10.1, from *Vitis pseudoreticulata*: An insight of its mode of antifungal activity. *PLoS ONE* **2014**, *9*, e95102. [[CrossRef](#)] [[PubMed](#)]
253. Le Henaff, G.; Heitz, T.; Mestre, P.; Mutterer, J.; Walter, B.; Chong, J. Characterization of *Vitis vinifera* NPR1 homologs involved in the regulation of pathogenesis-related gene expression. *BMC Plant. Biol.* **2009**, *9*, 54. [[CrossRef](#)] [[PubMed](#)]
254. Liu, Y.P.; Peremyslov, V.V.; Medina, V.; Dolja, V.V. Tandem leader proteases of grapevine leafroll-associated virus-2: Host-specific functions in the infection cycle. *Virology* **2009**, *383*, 291–299. [[CrossRef](#)] [[PubMed](#)]
255. Gollop, R.; Even, S.; Colova-Tsolova, V.; Perl, A. Expression of the grape dihydroflavonol reductase gene and analysis of its promoter region. *J. Exp. Bot.* **2002**, *53*, 1397–1409. [[CrossRef](#)]
256. Jelly, N.S.; Schellenbaum, P.; Walter, B.; Maillot, P. Transient expression of artificial microRNAs targeting Grapevine fanleaf virus and evidence for RNA silencing in grapevine somatic embryos. *Transgenic Res.* **2012**, *21*, 1319–1327. [[CrossRef](#)]
257. Li, Z.T.; Jayasankar, S.; Gray, D.J. Bi-directional duplex promoters with duplicated enhancers significantly increase transgene expression in grape and Tobacco. *Transgenic Res.* **2004**, *13*, 143–154. [[CrossRef](#)]
258. Gutoranov, G.P.; Tsvetkov, I.J.; Colova-Tsolova, V.M.; Atanassov, A.I. Genetically engineered grapevines carrying GFLV coat protein and antifreeze genes. *Agricul. Conspect. Sci.* **2001**, *66*, 71–76.
259. Mulwa, R.M.S.; Norton, M.A.; Farrand, S.K.; Skirvin, R.M. Agrobacterium-mediated transformation and regeneration of transgenic ‘Chancellor’ wine grape plants expressing the tfdA gene. *Vitis* **2007**, *46*, 110–115.
260. Nirala, N.K.; Das, D.K.; Srivastava, P.S.; Sopory, S.K.; Upadhyaya, K.C. Expression of a rice chitinase gene enhances antifungal potential in transgenic grapevine (*Vitis vinifera* L.). *Vitis* **2010**, *49*, 181–187. [[CrossRef](#)]
261. Dhekney, S.A.; Li, Z.T.; Gray, D.J. Grapevines engineered to express cisgenic *Vitis vinifera* thaumatin-like protein exhibit fungal disease resistance. *Vitr. Cell. Dev. Biol. Plant.* **2011**, *47*, 458–466. [[CrossRef](#)]
262. Nookaraju, A.; Agrawal, D.C. Enhanced tolerance of transgenic grapevines expressing chitinase and β -1,3-glucanase genes to downy mildew. *Plant. Cell Tiss. Organ. Cult.* **2012**, *111*, 15–28. [[CrossRef](#)]

263. Dai, L.; Zhou, Q.; Li, R.; Du, Y.; He, J.; Wang, D.; Cheng, S.; Zhang, J.; Wang, Y. Establishment of a picloram-induced somatic embryogenesis system in *Vitis vinifera* cv. Chardonnay and genetic transformation of a stilbene synthase gene from wild-growing *Vitis* species. *Plant. Cell Tiss. Organ. Cult.* **2015**, *121*, 397–412. [[CrossRef](#)]
264. Li, Z.T.; Hopkins, D.L.; Gray, D.J. Overexpression of antimicrobial lytic peptides protects grapevine from Pierce's disease under greenhouse but not field conditions. *Transgenic Res.* **2015**, *24*, 821–830. [[CrossRef](#)] [[PubMed](#)]
265. Cheng, S.; Xie, X.; Xu, Y.; Zhang, C.; Wang, X.; Zhang, J.; Wang, Y. Genetic transformation of a fruit-specific, highly expressed stilbene synthase gene from Chinese wild *Vitis quinquangularis*. *Planta* **2016**, *243*, 1041–1053. [[CrossRef](#)] [[PubMed](#)]
266. Dai, L.; Wang, D.; Xie, X.; Zhang, C.; Wang, X.; Xu, Y.; Wang, Y.; Zhang, J. The novel gene VpPR4-1 from *Vitis pseudoreticulata* increases powdery mildew resistance in transgenic *Vitis vinifera* L. *Front. Plant. Sci.* **2016**, *7*, 695. [[CrossRef](#)] [[PubMed](#)]
267. Pessina, S.; Lenzi, L.; Perazzoli, M.; Campa, M.; Dalla Costa, L.; Urso, S.; Valè, G.; Salamini, F.; Velasco, R.; Malnoy, M. Knockdown of MLO genes reduces susceptibility to powdery mildew in grapevine. *Hortic. Res.* **2016**, *3*, 16016. [[CrossRef](#)]
268. He, R.; Wu, J.; Zhang, Y.; Agüero, C.B.; Li, X.; Liu, S.; Wang, C.; Andrew Walker, M.; Lu, J. Overexpression of a thaumatin-like protein gene from *Vitis amurensis* improves downy mildew resistance in *Vitis vinifera* grapevine. *Protoplasma* **2017**, *254*, 1579–1589. [[CrossRef](#)]
269. Jiao, L.; Zhang, Y.; Lu, J. Overexpression of a stressresponsive U-box protein gene VaPUB affects the accumulation of resistance related proteins in *Vitis vinifera* 'Thompson Seedless'. *Plant. Physiol. Biochem.* **2017**, *112*, 53–63. [[CrossRef](#)]
270. Ma, H.; Xiang, G.; Li, Z.; Wang, Y.; Dou, M.; Su, L.; Yin, X.; Liu, R.; Wang, Y.; Xu, Y. Grapevine VpPR10.1 functions in resistance to *Plasmopara viticola* through triggering a cell death-like defense response by interacting with VpVDAC3. *Plant. Biotechnol. J.* **2018**, *16*, 1488–1501. [[CrossRef](#)]
271. Su, H.; Jiao, Y.T.; Wang, F.F.; Liu, Y.E.; Niu, W.L.; Liu, G.T.; Xu, Y. Overexpression of VpPR10.1 by an efficient transformation method enhances downy mildew resistance in *V. vinifera*. *Plant. Cell Rep.* **2018**, *37*, 819–832. [[CrossRef](#)]
272. Soliman, H.I.A. Production of genetically modified grape (*Vitis vinifera* L.) plants. *Int. J. Hortic. Agric. Food Sci.* **2018**, *2*, 111–120. [[CrossRef](#)]
273. Jiang, J.; Xi, H.; Dai, Z.; Lecourieux, F.; Yuan, L.; Liu, X.; Patra, B.; Wei, Y.; Li, S.; Wang, L. VvWRKY8 represses stilbene synthase genes through direct interaction with VvMYB14 to control resveratrol biosynthesis in grapevine. *J. Exp. Bot.* **2019**, *70*, 715–729. [[CrossRef](#)] [[PubMed](#)]
274. Xu, W.; Ma, F.; Li, R.; Zhou, Q.; Yao, W.; Jiao, Y.; Zhang, C.; Zhang, J.; Wang, X.; Xu, Y.; et al. VpSTS29/STS2 enhances fungal tolerance in grapevine through a positive feedback loop. *Plant. Cell Environ.* **2019**, *42*, 2979–2998. [[CrossRef](#)] [[PubMed](#)]
275. Tu, M.; Wang, X.; Yin, W.; Wang, Y.; Li, Y.; Zhang, G.; Li, Z.; Song, J.; Wang, X. Grapevine VlbZIP30 improves drought resistance by directly activating VvNAC17 and promoting lignin biosynthesis through the regulation of three peroxidase genes. *Hortic. Res.* **2020**, *7*, 150. [[CrossRef](#)]
276. Carroll, D. Genome engineering with zinc-finger nucleases. *Genetics* **2011**, *188*, 773–782. [[CrossRef](#)]
277. Mahfouz, M.M.; Li, L.; Shamimuzzaman, M.; Wibowo, A.; Fang, X.; Zhu, J.-K. De novo-engineered transcription activator-like effector (TALE) hybrid nuclease with novel DNA binding specificity creates double-strand breaks. *Proc. Nat. Acad. Sci. USA* **2011**, *108*, 2623–2628. [[CrossRef](#)]
278. Li, L.; Piatek, M.J.; Atef, A.; Piatek, A.; Wibowo, A.; Fang, X.; Sabir, J.S.M.; Zhu, J.-K.; Mahfouz, M.M. Rapid and highly efficient construction of TALE-based transcriptional regulators and nucleases for genome modification. *Plant. Molec. Biol.* **2012**, *78*, 407–416. [[CrossRef](#)]
279. Jinek, M.; Chylinski, K.; Fonfara, I.; Hauer, M.; Doudna, J.A.; Charpentier, E. Programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science* **2012**, *337*, 816–821. [[CrossRef](#)]
280. Kaul, T.; Sony, S.K.; Verma, R.; Motelb, K.F.A.; Prakash, A.T.; Eswaran, M.; Bharti, J.; Nehra, J.; Kaul, R. Revisiting CRISPR/Cas-mediated crop improvement: Special focus on nutrition. *J. Biosci.* **2020**, *45*, 1–37. [[CrossRef](#)]
281. van der Oost, J.; Westra, E.R.; Jackson, R.N.; Wiedenheft, B. Unravelling the structural and mechanistic basis of CRISPR-Cas systems. *Nat. Rev. Microbiol.* **2014**, *12*, 479–492. [[CrossRef](#)]
282. Xie, K.; Zhang, J.; Yang, Y. Genome-wide prediction of highly specific guide RNA spacers for CRISPR-Cas9-mediated genome editing in model plants and major crops. *Mol. Plant.* **2014**, *7*, 923–926. [[CrossRef](#)] [[PubMed](#)]
283. Bortesi, L.; Fischer, R. The CRISPR/Cas9 system for plant genome editing and beyond. *Biotechnol. Adv.* **2015**, *33*, 41–52. [[CrossRef](#)] [[PubMed](#)]
284. Butiuc-Keul, A.; Farkas, A.; Carpa, R.; Dobrota, C.T.; Iordache, D. Development of smart fruit crops by genome editing. *Turk. J. Agric. For.* **2022**, *46*, 129–140. [[CrossRef](#)]
285. Cong, L.; Ran, F.A.; Cox, D.; Lin, S.; Barretto, R.; Habib, N.; Hsu, P.D.; Wu, X.; Jiang, W.; Marraffini, L.A.; et al. Multiplex genome engineering using CRISPR/Cas systems. *Science* **2013**, *339*, 819–823. [[CrossRef](#)]
286. Hsu, P.D.; Scott, D.A.; Weinstein, J.A.; Ran, F.A.; Konermann, S.; Agarwala, V.; Li, Y.; Fine, E.J.; Wu, X.; Shalem, O.; et al. DNA targeting specificity of RNA-guided Cas9 nucleases. *Nat. Biotechnol.* **2013**, *31*, 827–832. [[CrossRef](#)]
287. Carpa, R.; Remizovschi, A.; Culda, C.A.; Butiuc-Keul, A.L. Inherent and composite hydrogels as promising materials to limit antimicrobial resistance. *Gels* **2022**, *8*, 70. [[CrossRef](#)]
288. Ali, Z.; Abulfaraj, A.; Idris, A.; Ali, S.; Tashkandi, M.; Mahfouz, M.M. CRISPR/ Cas9-mediated viral interference in plants. *Genome Biol.* **2015**, *16*, 238. [[CrossRef](#)]

289. Malnoy, M.; Viola, R.; Jung, M.-H.; Koo, O.-J.; Kim, J.-S.; Velasco, R.; Kanchiswamy, C.N. DNAfree genetically edited grapevine and apple protoplast using CRISPR/Cas9 ribonucleoproteins. *Front. Plant. Sci.* **2016**, *7*, 1904. [[CrossRef](#)]
290. Osakabe, Y.; Liang, Z.; Ren, C.; Nishitani, C.; Osakabe, K.; Wada, M.; Komori, S.; Malnoy, M.; Velasco, R.; Poli, M.; et al. CRISPR-Cas9-mediated genome editing in apple and grapevine. *Nat. Prot.* **2018**, *13*, 2844–2863. [[CrossRef](#)]
291. Li, M.Y.; Jiao, Y.T.; Wang, Y.T.; Zhang, N.; Wang, B.B.; Liu, R.Q.; Yin, X.; Xu, Y.; Liu, G.T. CRISPR/Cas9-mediated VvPR4b editing decreases downy mildew resistance in grapevine (*Vitis vinifera* L.). *Hortic. Res.* **2020**, *7*, 149. [[CrossRef](#)]
292. Wan, D.Y.; Guo, Y.; Cheng, Y.; Hu, Y.; Xiao, S.Y.; Wang, Y.J.; Wen, Y.Q. CRISPR/Cas9-mediated mutagenesis of VvMLO3 results in enhanced resistance to powdery mildew in grapevine (*Vitis vinifera*). *Hortic. Res.* **2020**, *7*, 116. [[CrossRef](#)] [[PubMed](#)]
293. Ren, C.; Liu, X.; Zhang, Z.; Wang, Y.; Duan, W.; Li, S.; Liang, Z. CRISPR/Cas9-mediated efficient targeted mutagenesis in Chardonnay (*Vitis vinifera* L.). *Sci. Rep.* **2016**, *6*, 32289. [[CrossRef](#)] [[PubMed](#)]
294. Nakajima, I.; Ban, Y.; Azuma, A.; Onoue, N.; Moriguchi, T.; Yamamoto, T.; Toki, S.; Endo, M. CRISPR/Cas9-mediated targeted mutagenesis in grape. *PLoS ONE* **2017**, *12*, e0177966. [[CrossRef](#)]
295. Ren, F.; Ren, C.; Zhang, Z.; Duan, W.; Lecourieux, D.; Li, S.; Liang, Z. Efficiency Optimization of CRISPR/Cas9-Mediated Targeted Mutagenesis in Grape. *Front. Plant. Sci.* **2019**, *10*, 612. [[CrossRef](#)] [[PubMed](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.