

Table S6
Percentage of Total LDH Released with Lower PEG Concentrations

Abbreviations are defined on page 2

		Hours after addition of rickettsiae			
		1	2	3	4
RpM IFN	Peg 4000-30 mM d/a	0.04	18.57	29.4	32.04
		0	18.89	29.89	34.25
		1	15.7	34.2	40.6
		0.4	14.7	32.7	43.1
		3.1	20.7	30.3	33.4
		1.5	20.1	28.9	32.2
Mean		1.0	18.1	30.9	35.9
Std Dev		1.2	2.4	2.1	4.7
RpM IFN	Peg 4000-15 mM d/a	1.4	32.9	54.8	62.8
		2.1	32.7	50.2	57.8
Mean		1.8	32.8	52.5	60.3
Std Dev		0.5	0.1	3.3	3.5
RpM IFN	Peg 4000-7.5 mM d/a	2.4	45	63.6	75.2
		3.6	36.7	54.1	62.6
Mean		3.0	40.9	58.9	68.9
Std Dev		0.8	5.9	6.7	8.9
RpM IFN	No PEG	3.76	41.75	52.12	55.62
		2.81	43.3	54.58	58.46
		5	43.3	57.1	65.4
		3.9	47.5	63	74.3
		8.9	38.6	47.5	49.1
		9.5	36.9	43.8	46.1
		2.26		60.3	66.1
		3.02		58	61.4
		1.3		61.5	67.5
		1.2		58.6	62.1
		1.3		57.2	50
				59.9	49.4
				50.8	50
				53.1	
				53.2	
				52.4	
Mean		3.91	41.89	55.19	58.11
Std Dev		2.88	3.77	5.23	8.82

Hours after addition of rickettsiae

		1	2	3	4
RpM IFN	Peg 8000-15 mM d/a	2.8	4.1	3.3	4.8
		2.2	3.6	3.2	3.9
		2	2.5	3	3.5
		2	2.5	3.3	4.3
				3.9	4.7
				5.1	5.7
				5.5	5.1
				3.9	5.2
				4.6	8.5
				4.7	7.3
				5.2	5.7
				5	7.3
				5.1	3.4
				5.8	2.9
				6.5	3
				4.9	

Mean	2.3	3.2	4.6	5.0
Std Dev	0.4	0.8	1.0	1.7

RpM IFN	Peg 8000-7.5 mM d/a	1.4	28.4	44.2	58.8
		1.6	21.7	37.6	49

Mean	1.5	25.1	40.9	53.9
Std Dev	0.1	4.7	4.7	6.9

DEFINITION OF ABBREVIATIONS IN THIS TABLE

"d/a" indicates that PEGs were added along with the rickettsiae.

RpM IFN=IFN-treated, RpMedium infected

See Figure 3 and the results section for additional information about RpL, RpM, and RpH. These indicate the use of different concentrations of rickettsiae.