Quality Assessment of Avocado Pulp Oils during Storage [†]

Rita Brzezińska *, Agata Górka, Kinga Gotowicka, Joanna Bryś, Ewa Ostrowska-Ligeza and Magdalena Wirkowska-Wojdyla

Department of Chemistry, Institute of Food Sciences, Warsaw University of Life Sciences—SGGW, 02-787 Warszawa, Poland; agata_gorska@sggw.edu.pl (A.G.); kinga.gotowicka@gmail.com (K.G.); joanna_brys@sggw.edu.pl (J.B.); ewa_ostrowska_ligeza@sggw.edu.pl (E.O.-L.); magdalena_wirkowska@sggw.edu.pl (M.W.-W.)

* Correspondence: rita_brzezinska@sggw.edu.pl

[†] Presented at the 1st International Electronic Conference on Food Science and Functional Foods, 10–25 November 2020; Available online: https://foods_2020.sciforum.net/.

Abstract: Recently, consumers' awareness is becoming a crucial aspect driving the food industry to develop new products with high nutritional value. The oil industry explores the use of less well-known plant materials such as avocado fruit which is a rich source of bioactive compounds. The objective of this study was to assess the quality and oxidative stability of avocado pulp oils during a 2-month storage period. Two avocado varieties, Hass and Reed, were selected and oil extraction was performed with the use of hexane. The extent of oxidative deterioration and oil stability were tested by measuring the acid value and peroxide value. The PDSC method was applied to evaluate the oxidative induction time. The composition of fatty acids and their distribution in internal (sn-2) and external (sn-1 and sn-3) positions in triacylglycerols were also analyzed. The acid value and the peroxide value of fresh extracted avocado oils reached approximately 0.6 mg KOH g⁻¹ fat and 5 meq O₂ kg⁻¹ fat, respectively. Generally, during avocado oil storage both the acid values and the peroxide values were in accordance with Codex Alimentarius requirements (the limit for acid value is 4 mg KOH g⁻¹, while for peroxide value it is 15 meq O₂ kg⁻¹). The Hass avocado pulp oil was characterized by a higher value of the oxidative induction time, about 111 min, compared to the oil extracted from Reed avocado pulp (61 min). The GC analysis revealed that avocado pulp oil could be considered a source of monounsaturated fatty acids. The dominant fatty acid found in this group was oleic acid with a percentage share of above 60%. In accordance with the results of fatty acids distribution in triacylglycerol molecules, the main fatty acids in the sn-2 position were linoleic acid and oleic acid and their percentage share in this item was up to 59% and 34%, respectively. It was also noticed that after a 2-month storage period, the acid value and the peroxide value increased about 7-fold and 2-fold, the oxidative induction time decreased about 2-fold and the percentage shares of fatty acid groups changed. In conclusion, the results obtained in this research indicate that storage period has a significant impact on avocado pulp oil quality.

Citation: Brzezińska, R.; Górka, A.; Gotowicka, K.; Bryś, J.; Ostrowska-Ligeza, E.; Wirkowska-Wojdyla, M. Quality Assessment of Avocado Pulp Oils during Storage. *Proceedings* **2021**, *70*, 14. https://doi.org/10.3390/foods_2020-07694

Published: 9 November 2020

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Copyright: © 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

Keywords: avocado oil; acid value; peroxide value; fatty acids composition; oxidative stability

1. Introduction

In recent years, the demand for unprocessed, “healthful” foods and beverages has increased rapidly [1]. Functional foods are edible products with additional health benefits beyond the essential nutrients, which are enhanced by the addition of bioactive compounds, or by changes in the production processes. A plethora of functional foods throughout the market diminish the disparity between pharmaceuticals and nutrition [2].

Vegetable oils are an original source of fat-soluble compounds such as free fatty acids, vitamins or phenolics. Due to the abundance of bioactive compounds, vegetable oils

have numerous functional properties: mainly, antioxidant, antibacterial, anti-inflammatory and anticancer capacities [3]. Nowadays manufacturers seek out new sources of vegetable oils.

Avocado oil is a rich source of unsaturated fatty acids which are invaluable in cancer prevention and in cardiovascular disease prophylaxis. Additionally, the unsaponifiable fraction has a high concentration of compounds such as phytosterols and policosanols which can decrease low-density lipoprotein cholesterol in the blood. Oils containing high monounsaturated fatty acids are less susceptible to oxidation, which is a main cause of deteriorative changes in sensory, chemical and nutritional properties during storage [4].

Taking the above into the account, the aim of this research was to monitor the quality of avocado pulp oils during a two-month storage period.

2. Materials and Methods

2.1. Preparation of Avocado Oil Samples

To obtain avocado oil samples, Hass and Reed cultivars pulps were selected. The oil samples were extracted with hexane in a 6:10 ratio (*w/v*; homogenized pulp sample:hexane). Subsequently, the mixtures were put in an Elpin Plus type 357 (Lubawa, Poland) laboratory shaker for one hour. After the shaking process, the samples were filtrated, dried with anhydrous magnesium sulfate, evaporated under vacuum in a BÜCHI laboratory evaporator and finally dried under nitrogen atmosphere.

2.2. Analytical Methods

2.2.1. Determination of Acid Value

Acid value was determined with regard to the procedure described in PN-EN ISO 660:2010 [5]. It was calculated as the mg of potassium hydroxide needed to neutralize the organic acids present in 1 g of oil samples, with the use of phenolphthalein as an indicator.

2.2.2. Determination of Peroxide Value

Peroxide value was computed as meq O₂·kg⁻¹ (miliequivalents of oxygen per kg of oil samples). It was determined in pursuance of PN-EN ISO 3960:2012 [6], after dissolving the oil samples in an acetic acid–chloroform mixture (3:2; *v/v*) in the presence of a potassium iodide solution. The mixture was then titrated with sodium thiosulfate using starch as an indicator.

2.2.3. Determination of Oxidative Stability by Means of PDSC Method

A differential scanning calorimeter DSC Q20, TA Instruments, coupled with a high-pressure cell was used. Oil samples of 3–4 mg were put into the aluminum pan and placed in the sample chamber. The analyses were performed with an initial pressure of 1400 kPa, under oxygen atmosphere, with the temperature set to 120 °C (isothermal process). Oxidative induction time was determined based on the PDSC diagrams.

2.2.4. Determination of Fatty Acid Profiles

The fatty acids' composition was determined in accordance with the method described in PN-EN ISO 5508:1996 [7]. To prepare the samples for GC analysis, esterification with potassium hydroxide in methanol (1 M) according to PN-EN ISO 5509:2001 [8] was performed. After the esterification process, samples were analyzed in YL6100 GC chromatograph coupled with a flame ionization detector. A BPX-70 capillary column (0.2 mm i.d × 60 m length × 0.25 µm film thickness) was used. Samples were injected into the GC with an inlet temperature of 225 °C and with a split ratio of 1:100. Initial oven temperature of 60 °C was maintained for 5 min, rising to 180 °C with an increment of 10 °C min⁻¹, from 180 °C to 230 °C at a rate of 3 °C min⁻¹ and finally kept at 230 °C for 15 min. The temperature of the detector was set to 250 °C. Nitrogen was used as a carrier gas at a flow rate of

1 mL min⁻¹. The area under the curve for each peak was manually integrated and the results were expressed as a relative percentage of each fatty acid. Identification of fatty acids was performed by comparing the relative retention times of peaks within the sample with the chemical standard.

2.2.5. Distribution of Fatty Acids in the sn-2 and sn-1,3 Positions of Triacylglycerols

TAGs positions of oil samples were analyzed by use of the Brockerhoff method which involves partial hydrolysis of triacylglycerol carried out in the presence of the pancreatic lipase enzyme [9,10]. The content of sn-1,3 fatty acids was calculated based on the information on the initial composition of TAGs and the composition of the sn-2 MAG:

$$\text{sn-1,3} = [3 \times (\text{FA in TAGs}) - (\text{FA in sn-2 MAG})] \times 2^{-1}, \quad (1)$$

where:

sn-1,3—content of a given FA in the sn-1 and sn-3 positions [%];

FA in TAGs—content of a given FA in starting TAGs [%];

FA in sn-2 MAG—content of a given FA in sn-2 MAGs [%].

The percentage of selected fatty acids in sn-2 positions in relation to the total content of given fatty acid was calculated based on the formula:

$$\text{sn-2} = (\text{FA in sn-2 MAG}) \times [3 \times (\text{FA in TAGs})]^{-1} \quad (2)$$

2.3. Statistical Analysis

The results were analyzed using STATISTICA 13 software. Multivariate ANOVAs with Tukey's post hoc tests were conducted for significant differences between samples at $p < 0.05$.

3. Results and Discussion

The acid values (AV) and peroxide values (PV) are important indicators to assess edible oil quality. In Table 1, the changes of AV and PV of avocado oil during storage are shown. The AV and PV of freshly extracted avocado oils did not exceed 0.6 mg KOH g⁻¹ oil and 5 meq O₂ kg⁻¹ oil, respectively. The examined oils were of good quality being in agreement with Codex Alimentarius [11] recommendations which state that the AV and PV for cold pressed oils should not exceed 4 mg KOH g⁻¹ oil and 15 meq O₂ kg⁻¹ oil.

Table 1. Changes of acid value (AV), peroxide value (PV) and oxidation induction time (OIT) of oil samples during storage. Different letters indicate that the samples are significantly different during storage at $p < 0.05$.

	Oil Sample	Storage Time		
		0 Month	1 Month	2 Months
AV	Hass	0.62 ± 0.08 ^A	0.82 ± 0.04 ^A	3.91 ± 0.05 ^C
[mg KOH*g oil ⁻¹]	Reed	0.57 ± 0.01 ^A	1.31 ± 0.16 ^B	4.66 ± 0.13 ^D
PV	Hass	3.99 ± 0.08 ^A	7.01 ± 0.06 ^C	9.13 ± 0.33 ^E
[meq O ₂ *kg oil ⁻¹]	Reed	4.90 ± 0.08 ^B	7.48 ± 0.08 ^C	8.29 ± 0.05 ^D
OIT	Hass	111.42 ± 2.57 ^B	59.55 ± 1.08 ^A	49.44 ± 4.00 ^A
[min]	Reed	60.63 ± 8.24 ^A	50.62 ± 1.94 ^A	48.06 ± 0.21 ^A

After 2 months of storage time, AV and PV increased significantly: over six times for Hass oil and over eight times for Reed oil. After that time, only oil extracted from the Reed cultivar exceeded the recommended limit for AV. This substantial increase in AV and PV was probably induced by the ambient temperature of the oils' storage [12,13].

According to the literature, PV values of avocado oils might differ greatly, affected mostly by the extraction process and source material used. Bora et al. [14] reported a PV of 1.4 meq O₂ kg⁻¹ oil for the Fuerte cultivar, while Salgado et al. [15] found PV at a level

of 20.58 meq O₂ kg⁻¹ oil for the Margarida variety. Our results are in accordance with Ortega et al. [16] who reported a PV of 3.79 meq O₂ kg⁻¹ oil for the oil extracted from Hass cultivars.

Freshly extracted avocado oils were characterized by a relatively long oxidation induction time (OIT) (111 min for the Hass cultivar and 60 min for the Reed cultivar). This disparity might be the result of source fruits being in different ripening stages or higher SFA content in Hass oil. After a 2-month period of storage, the OIT of Hass oil decreased significantly by about 56%.

The beneficial influence on human health of avocado oil could be attributed to its fatty acid profiles. The fatty acids composition of Hass and Reed avocado pulp oil were shown in Table 2. It can be seen that tested avocado pulp oils were characterized by a high content of monounsaturated fatty acids (63–70%). Among MUFA, oleic acid is the dominant fatty acid and its percentage share accounted for approximately 60%. The results agree with the findings of Krumreich et al. [17].

Table 2. Changes of fatty acid profiles of tested avocado oils during storage. Different letters indicate that the samples are significantly different during storage at $p < 0.05$.

		Hass			Reed		
		0 Month	1 Month	2 Months	0 Month	1 Month	2 Months
SFA	C 16:0	17.12 ± 0.01 ^B	18.01 ± 0.01 ^C	17.86 ± 0.46 ^{BC}	14.39 ± 0.04 ^A	17.25 ± 0.03 ^{BC}	17.29 ± 0.11 ^{BC}
	C 18:0	0.63 ± 0.01 ^A	0.65 ± 0.01 ^A	0.64 ± 0.06 ^A	0.92 ± 0.03 ^B	0.93 ± 0.08 ^B	0.91 ± 0.03 ^B
	C 20:0	0.13 ± 0.02 ^A	0.12 ± 0.02 ^A	0.12 ± 0.04 ^A	0.15 ± 0.08 ^B	0.18 ± 0.01 ^D	0.17 ± 0.01 ^C
	ΣSFA	17.88	18.78	18.62	15.46	18.36	18.37
MUFA	C 16:1	7.34 ± 0.05 ^C	7.40 ± 0.01 ^C	7.41 ± 0.22 ^C	3.88 ± 0.05 ^A	5.48 ± 0.02 ^B	5.58 ± 0.01 ^B
	C 17:1	0.14 ± 0.01 ^A	0.13 ± 0.02 ^A	0.12 ± 0.01 ^A	0.14 ± 0.01 ^A	0.14 ± 0.01 ^A	0.11 ± 0.01 ^A
	C 18:1	62.14 ± 0.02 ^B	61.88 ± 0.01 ^B	61.92 ± 0.66 ^B	64.39 ± 0.11 ^C	56.97 ± 0.15 ^A	56.81 ± 0.20 ^A
	C 20:1	0.26 ± 0.02 ^A	0.25 ± 0.01 ^A	0.26 ± 0.02 ^A	0.28 ± 0.01 ^A	0.28 ± 0.01 ^A	0.29 ± 0.05 ^A
	ΣMUFA	69.87	69.66	69.70	68.69	62.87	62.78
PUFA	C 18:2	11.11 ± 0.01 ^C	10.59 ± 0.02 ^A	10.73 ± 0.01 ^B	14.82 ± 0.05 ^D	17.30 ± 0.02 ^E	17.40 ± 0.03 ^F
	C 18:3	0.84 ± 0.02 ^B	0.72 ± 0.02 ^A	0.73 ± 0.07 ^A	0.77 ± 0.01 ^A	1.17 ± 0.03 ^C	1.14 ± 0.03 ^C
	C 20:5	0.12 ± 0.05 ^A	0.06 ± 0.03 ^A	0.06 ± 0.01 ^A	0.08 ± 0.01 ^A	0.12 ± 0.01 ^A	0.12 ± 0.02 ^A
	ΣPUFA	12.06	11.37	11.51	15.67	18.58	18.66
ΣUFA		81.93	81.03	81.21	84.36	81.45	81.43
Unidentified		0.19	0.19	0.18	0.18	0.20	0.21

The quality of oils depends on the distribution of fatty acids in the triacylglycerols. Saturated fatty acids occupying the sn-2 position positively affect the absorption of fats [18]. According to the results presented in Table 3, it can be seen that the sn-2 position is occupied mainly by oleic acid. The content of oleic acid in the sn-2 position was about 63% for Hass and 55.3% for the Reed cultivar, and its share in this position reached 34% and 29% for Hass and Reed, respectively.

Table 3. Positional distribution of selected fatty acids in avocado oil samples.

Fatty Acids	Oil Sample	Fatty Acid Content in TAG [%]	Fatty Acid Content in Position [%]		Percentage Distribution of Fatty Acid in sn-2 [%]
			sn-2	sn-1,3	
C 16:0	Hass	17.12	7.77	21.8	15%
	Reed	14.39	11.8	15.69	27%
C 16:1	Hass	7.34	5.95	8.03	27%
	Reed	3.89	3.85	3.91	33%
C 18:1	Hass	62.14	62.95	61.73	34%
	Reed	64.39	55.29	68.94	29%
C 18:2	Hass	11.11	19.49	6.91	59%
	Reed	14.82	24.64	9.91	55%

4. Conclusions

Our results indicate that avocado oil could be considered a rich source of unsaturated fatty acids, mainly monounsaturated fatty acids. A two-month period of storage at ambient temperature had a significant effect on the quality and composition of researched samples. Acid and peroxide values increased after storage, but these parameters in most cases did not exceed the limits recommended by the Codex Alimentarius. The oxidative induction time had decreased significantly in the case of Hass avocado oil, which suggests oil oxidation processes.

Author Contributions: Conceptualization, A.G.; methodology, A.G.; validation, R.B. and K.G.; formal analysis, R.B. and K.G.; investigation, M.W.-W.; resources, K.G. and J.B.; writing—original draft preparation, R.B.; writing—review and editing, E.O.-L. and J.B.; visualization, R.B.; supervision, A.G.; project administration, A.G. All authors have read and agreed to the published version of the manuscript.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data presented in this study are available in the article.

References

1. Nazir, M.; Arif, S.; Khan, R.S.; Nazir, W.; Khalid, N.; Maqsood, S. Opportunities and challenges for functional and medicinal beverages: Current and future trends. *Trends Food Sci. Technol.* **2019**, *88*, 513–526.
2. Eussen, S.R.B.M.; Verhagen, H.; Klungel, O.H.; Garssen, J.; van Loveren, H.; van Kranen, H.J.; Rempelberg, C.J.M. Functional foods and dietary supplements: Products at the interface between pharma and nutrition. *Eur. J. Pharmacol.* **2011**, *668* (Suppl. 1), S2–S9.
3. Ruiz Ruiz, J.C.; Ortiz Vazquez, E.L.; Segura Campos, M.R. Encapsulation of vegetable oils as source of omega-3 fatty acids for enriched functional foods. *Crit. Rev. Food Sci.* **2017**, *57*, 1423–1434.
4. Flores, M.A.; Perez-Camino, M.D.C.; Troca, J. Preliminary studies on composition, quality and oxidative stability of commercial avocado oil produced in Chile. *J. Food Sci. Eng.* **2014**, *4*, 21–26.
5. Polish norm PN-EN ISO 660:2010. Oleje i Tłuszcze Roślinne Oraz Zwierzęce. Oznaczanie Liczby Kwasowej i Kwasowości.
6. Polish norm PN-EN ISO 3960:2012. Oleje i Tłuszcze Roślinne Oraz Zwierzęce. Oznaczanie Liczby Nadtlencowej. Jodometryczne (Wizualne) Oznaczanie Punktu Końcowego.
7. Polish norm PN-EN ISO 5508:1996. Oleje i Tłuszcze Roślinne Oraz Zwierzęce. Analiza Estrów Metylowych Kwasów Tłuszczowych Metodą Chromatografii Gazowej.
8. Polish norm PN-EN ISO 5509:2001. Oleje i Tłuszcze Roślinne Oraz Zwierzęce. Przygotowanie Estrów Metylowych Kwasów Tłuszczowych.
9. Brockerhoff, H. A stereospecific analysis of triglycerides. *J. Lipid Res.* **1965**, *6*, 10–15.
10. Hazuka, Z.; Ledóchowska, E.; Jurkowska, A. Porównanie metod oznaczania struktury triacylogliceroli z wykorzystaniem lipazy trzustkowej—metody Brockerhoffa i metody ISO. *Tłuszcze Jadalne* **2003**, *38*, 31–41.
11. FAO-WHO Codex Stan 210. **1999**. Standard for named vegetable oils. Available at <http://www.fao.org/fao-who-codexalimentarius/codex-texts/list-standards/en/>
12. Kondratowicz-Pietruszka, E. Zmiany jakościowe wybranych olejów roślinnych przechowywanych w warunkach chłodniczych. *Zeszyty Naukowe Towaroznawstwo. Uniw. Ekon. Krak.* **2013**, *912*, 49–72.
13. Masłowski, A.; Andrejko, D.; Ślaska-Grzywna, B.; Sagan, A.; Szmigielski, M.; Mazur, J.; Rydzak, L.; Sobczak, P. Wpływ temperatury i czasu przechowywania na wybrane cechy jakościowe oleju rzepakowego, lnianego i lniankowego. *Inż. Rol.* **2013**, *141*, 115–124.
14. Bora, S.P.; Narain, N.; Rocha, R.V.M.; Paulo, M.Q. Characterization of the oils from the pulp and seeds of avocado (cultivar: Fuerte) fruits. *Grasas Aceites* **2001**, *52*, 171–174.
15. Salgado, J.M.; Danieli, F.; Regitano-D’arce, M.A.B.; Frias, A.; Mansi, D.N. O oleo de abacate (*Persea americana* Mill.) como matéria-prima para a indústria alimentícia. *Cienc. Technol. Alime* **2008**, *28*, 20–26.
16. Ortega, J.A.A.; Lopez, M.R.; Torre, R.R.R. Effect of electric field treatment on avocado oil. *Int. J. Res. Agric. Food Sci.* **2013**, *1*, 13–22.

17. Krumreich, F.D.; Borges, C.D.; Mendonca, C.R.B.; Jansen-Alves, R.; Zambiasi, C. Bioactive compounds and quality parameters of avocado oil obtained by different processes. *Food Chem.* **2018**, *257*, 376–381.
18. Wirkowska, M.; Bryś, J.; Górka, A.; Ostrowska-Ligeza, E.; Żubrzycka, K. Wzbogacanie tłuszczu mlecznego w kwasy n-3 na drodze przeestryfikowania enzymatycznego. *Bromat. Chem. Toksykol.* **2012**, *45*, 699–704.