clinical and translational neuroscience

MDPI

Article

Associations Between P300 Latency and Reaction Time on Event-Related Potentials in Children with Varying Levels of Fluid Intelligence

Nikolaos C. Zygouris [1,*], Irini Dermitzaki [2], Panayiotis Patrikelis [3], Lambros Messinis [3] and Eugenia I. Toki [4]

[1] Laboratory of Digital Neuropsychological Assessment, Department of Informatics and Telecommunications, University of Thessaly, 35100 Lamia, Greece
[2] Department of Early Childhood Education, University of Thessaly, 38221 Volos, Greece; idermitzaki@uth.gr
[3] Laboratory of Neuropsychology and Behavioural Neuroscience, School of Psychology, Aristotle University of Thessaloniki, 57001 Thessaloniki, Greece; patrikelis@psy.auth.gr (P.P.); lmessinis@psy.auth.gr (L.M.)
[4] Department of Speech and Language Therapy, School of Health Sciences, University of Ioannina, 45110 Ioannina, Greece; toki@uoi.gr
* Correspondence: nzygouris@uth.gr

Abstract: Exploring cognitive abilities is necessary in educational contexts, where such insights shape decisions about student placement and teaching methods. Traditionally, educational assessments have been leaned on academic performance to guide decisions related to grading and student placement. This study examines the relationships among specific neuropsychological measures, namely the Event Related Potentials (ERPs), P300 waveform, reaction time, and fluid intelligence in children. Raven's Standard Progressive Matrices (RSPM) was utilized to assess intelligence levels. Based on their RSPM scores, participants were grouped into two categories: those with "high mental abilities" and those with "average mental abilities". It was hypothesized that children with higher RSPM scores might display reduced P300 latencies and quicker reaction times, potentially reflecting greater neural efficiency. Electrophysiological data collected using ERPs, focusing on the P300 component. The results suggest a possible association between higher intelligence scores and shorter P300 latencies and quicker reaction times, which could support the concept of neural efficiency and the significance of cognitive speed in understanding intelligence. This investigation into the neuropsychological foundations of cognitive ability in children is in the same line with studies supporting how brain activity, connectivity, and processing efficiency vary. These differences could help develop educational strategies that are more tailored to individual cognitive processing styles.

Keywords: raven standard progressive matrices; intelligence; highly intelligent children; P300 latency; reaction time; neural efficiency hypothesis

check for updates

Academic Editor: Claudio Bassetti

Received: 10 March 2025
Revised: 5 April 2025
Accepted: 23 April 2025
Published: 13 May 2025

Citation: Zygouris, N.C.; Dermitzaki, I.; Patrikelis, P.; Messinis, L.; Toki, E.I. Associations Between P300 Latency and Reaction Time on Event-Related Potentials in Children with Varying Levels of Fluid Intelligence. *Clin. Transl. Neurosci.* **2025**, *9*, 24. https://doi.org/10.3390/ctn9020024

Copyright: © 2025 by the authors. Published by MDPI on behalf of the Swiss Federation of Clinical Neuro-Societies. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https://creativecommons.org/licenses/by/4.0/).

1. Introduction

Fluid intelligence, which enables thinking abstractly, reasoning rapidly, and solving new problems independently of learned knowledge, generally peaks in early adulthood and might decline as we age. This decline is linked to changes in brain areas critical for attention and short-term memory, such as the dorsolateral prefrontal cortex and the anterior cingulate cortex [1]. In contrast, crystallized intelligence, encompassing accumulated knowledge and skills, tends to stay stable or may increase as we grow older. The interaction between fluid and crystallized intelligence is essential for effective problem-solving and continuous learning throughout life [2].

Fluid intelligence is commonly used to predict a child's potential and likelihood of academic success [3]. Assessing this type of intelligence involves complex mental tasks designed to solve problems that go beyond rote memorization. Additionally, the relationship between fluid intelligence and learning abilities suggests that it could influence educational practices [4]. Children who are intellectually gifted typically show more efficient neural functioning, allowing them to perform better than their peers on basic cognitive tasks and on tasks requiring cognitive control, such as those involving executive functions and inhibitory control [5]. Thus, intelligence assessments can predict children's performance in cognitive control tasks [6].

The Raven's Standard Progressive Matrices (RSPM) is a non-verbal test that measures fluid intelligence by evaluating the ability to break down complex problems into smaller, manageable parts and to think abstractly [7,8]. The RSPM test challenges participants to identify the correct piece to complete a visual pattern, focusing on attributes like sameness, symmetry, and analogy, as well as form, color, and linear arrangement. Its language-free format helps reduce cultural and linguistic biases, making it a valuable tool for assessing intelligence across different cultural backgrounds [8].

This test is reliant on the visual processing abilities mediated by the occipital cortex, particularly Brodmann areas 18 and 19. These areas are crucial for processing complex visual stimuli through two distinct pathways: the ventral stream, which runs from the occipital cortex to the temporal lobe and is mainly involved in object recognition, and the dorsal stream, which extends to the parietal lobe and handles the processing of spatial relationships and motion [9]. The performance on the RSPM likely depends on the functionality and efficiency of these visual processing areas, as the test requires both abstract pattern recognition and spatial reasoning skills [10,11].

Neuroimaging research has revealed two primary models that describe the neuroanatomical and functional foundations of human fluid intelligence. The Parie-to-Frontal Integration Theory (P-FIT), as described by Jung and Haier [12], suggests that general intelligence arises from coordinated activity across distributed cortical networks. This theory highlights the importance of integrative processing across several brain regions, beginning with the initial sensory inputs processed in the occipital and temporal cortices, which handle basic perceptual and semantic tasks. This sensory data were then forwarded to the parietal cortex, which plays a role in abstracting and integrating this information into higher-order cognitive representations. The dorsolateral prefrontal cortex is thought to manage complex problem-solving, working memory, and executive functions that are critical for fluid reasoning. Additionally, the anterior cingulate cortex is believed to aid cognitive control by regulating response selection and conflict monitoring [13].

An alternate model, the neural efficiency hypothesis, is primarily supported by findings from Positron Emission Tomography (PET) studies. This hypothesis, as evidenced by the work of Haier et al. [14], indicates that repeated exposure to a task leads to a decrease in regional cerebral glucose metabolism, suggesting a more efficient neural function. This reduction in metabolic activity implies that individuals with higher intelligence may utilize more focused and task-specific neural activations, thereby optimizing the allocation of cognitive resources by only activating the essential brain regions for task performance. Conversely, those with lower intelligence may engage broader and less efficient networks, possibly as a compensatory or less-specialized processing approach. This hypothesis is further supported by neurophysiological methods like electroencephalography (EEG), which assesses patterns of cortical activation. Studies analyzing event-related synchronization (ERS) and desynchronization (ERD) in EEG data have shown that individuals with higher intelligence levels display distinct patterns of neural oscillatory activity, aligning with efficient cognitive processing and resource utilization [15].

Event-related potentials (ERPs) are a non-invasive electrophysiological tool for examining brain activity related to cognitive information processing. ERPs measure the average electrophysiological response to a stimulus, with a significant P300 component emerging as the third positive deflection when stimuli are attentively processed and a specific target is consciously identified [16]. This method is considered highly effective for exploring the correlations between brain electrical activity and the dynamic processes involved in handling cognitive stimuli [17]. The P300 component is particularly notable for its association with higher-level cognitive processing and the attentional mechanisms that are crucial for evaluating the context when focusing on specific stimuli [18].

The latency of the P300 component is thought to mirror complex cognitive functions such as categorization and sensory evaluation, acting as a temporal marker of brain activity that facilitates more efficient allocation of attention and refreshes working memory processes [4]. Longer P300 latencies often suggest slower cognitive processing and are commonly associated with cognitive impairments [4,19]. Additionally, the P300 component has shown relevance to learning and memory, as studies involving memory recall and new learning tasks have found notable correlations between P300 responses and performance [20].

In educational settings, the relationship between P300 latencies and intelligence is particularly significant because these latencies can reflect the efficiency of cognitive processing, which correlates closely with intelligence and, by extension, academic performance [21]. By linking P300 latencies with intelligence metrics, educators and psychologists can better understand a student's cognitive strengths or challenges, paving the way for tailored educational support and interventions [22,23]. Research by Walhovd and Fjell [24] suggests that individuals with higher cognitive capabilities tend to have shorter P300 latencies, reinforcing the notion that rapid neural processing is associated with greater intelligence and learning capacity. Polich [25] also supports this, highlighting that the P300 component is indicative of cognitive capacity, with shorter latencies associated with higher cognitive functions, thus presenting a potential objective measure of intellectual capacity that aligns with educational achievements.

Furthermore, understanding cognitive processes such as attention and decision-making, which can be evaluated through reaction time measures, is crucial. Reaction time is closely linked to intelligence. It is suggested that reaction time can be a predictor of academic success that may vary in impact at different educational levels [26]. P300 latency and reaction time are vital neurophysiological markers for defining students' cognitive profiles, offering essential insights for crafting interventions to enhance learning and memory in environments that demand rapid cognitive processing and strong attentional control. Although systematic reviews indicate that the correlation between intelligence and academic achievement tends to decrease as students' progress in their education [27], integrating core cognitive metrics with intelligence assessments provides a robust framework for educational psychologists and neuroeducators to examine cognitive dynamics within learning environments [27,28]. This approach aids in refining teaching strategies to better match individual neural and cognitive profiles, enhancing neural efficiency and optimizing educational outcomes. Leveraging intelligence research and cognitive assessments can help educators devise precise interventions tailored to diverse cognitive processing needs, supporting more effective neuroeducational practices [28].

Several studies have employed P300 latency within the oddball paradigm to explore the dynamics of intelligence, e.g., [28–33]. It is suggested that individuals with higher cognitive capabilities tend to have shorter P300 latency compared to those with standard cognitive function, aligning with the mental speed theory that posits quicker intellectual processing and better attentional control in those with greater intellectual abilities [25,28,34–36].

However, this association between intelligence and P300 latency does not consistently extend to other cognitive tasks. For instance, conflicting results have emerged from studies employing the Sternberg memory scanning task and response-conflict paradigms [37,38].

In contrast, the linkage between intelligence scores and reaction time when responding to oddball target stimuli appears more stable. Consistently, research indicates that individuals with higher intelligence levels process information more swiftly, as evidenced by shorter reaction times during straightforward cognitive tasks—a conclusion supported by meta-analytic findings [39]. Additionally, these individuals often show faster reaction times than control groups, reflecting their superior attentional mechanisms, working memory capacity, and engagement with tasks [37].

While previous research has often focused on P300 latency or reaction time separately in relation to intelligence (e.g., [23,33] for P300 latency and [40,41] for reaction time), no studies to date have concurrently examined these metrics within the context of intelligence categorization. This combined approach could provide a more comprehensive view of cognitive processing speed and attentional dynamics. Furthermore, while existing studies on P300 latency and reaction time have primarily targeted adult or mixed-age populations (e.g., [33,35]), the current study is different in its focus on children. This emphasis on a developmental group allows for an analysis of cognitive profiles during pivotal educational phases. Additionally, this research aims to establish a direct correlation between scores on Raven's Standard Progressive Matrices (RSPM) and both P300 latency and reaction times. Previous studies generally analyzed these variables separately or linked them to broader intelligence measures (e.g., [11,29]). The present study seeks to bridge this gap by associating specific fluid intelligence scores with both electrophysiological and behavioral indicators.

The present study aims to examine variations in electrophysiological brain activity, measured by ERPs, among children categorized by cognitive abilities into groups labeled as possessing "high mental abilities" or "average mental abilities" based on RSPM performance. Building on earlier neuropsychological investigations into electrophysiological markers across varying intelligence levels [24,40,41], the first hypothesis of this study expects significant differences in P300 latency between the groups, reflecting variations in cognitive processing speed and efficiency. The second hypothesis assumes these groups show different reaction times, indicative of differences in attentional and executive functions. Lastly, the third hypothesis suggests that both P300 latency and reaction time will correlate with RSPM scores across groups, proposing these measures as potential indicators of fluid intelligence in the developing brain.

2. Materials and Methods

2.1. Participants

Twenty-four male students aged between 10 and 12 years (mean age = 11.33, SD = 1.03) participated in this study. The cohort was divided into two groups based on their scores on the Raven's Standard Progressive Matrices (RSPM). The first group consisted of twelve right-handed children who scored highly on the RSPM (M = 137.33, SD = 11.65), and the second group also included twelve right-handed children but with average RSPM scores (M = 100.33, SD = 3.50). Selection for each group was strictly based on these RSPM performance metrics. It is important to note that none of the participants had undergone prior assessments of mental ability or electrophysiological evaluations. Additionally, all participants were screened to ensure they had no psycho-pathological disorders, learning difficulties, developmental disorders, or significant visual or hearing impairments, verified through detailed interviews with both the children and their parents.

Recruitment for the study occurred in primary schools, starting with interviews with teachers to confirm good academic performance among potential participants. The RSPM tests were then administered by a licensed psychologist [39], while another psychologist conducted interviews with the children's parents to gather further developmental and health-related information. Based on the outcomes of the RSPM, the children were assigned to their respective groups. Subsequently, EEG recordings were administered to explore electrophysiological correlations of cognitive functioning.

Participation in the study required informed consent from all participants' parents or guardians, in line with ethical standards. All human data included in this manuscript were collected in compliance with the Helsinki Declaration and the guidelines established by the Internal Research Ethics Committee (EHDE) under protocol code 227092023. Lastly, this study employed the use of ChatGPT 3.5 [42] to enhance the clarity and quality of the English writing in the manuscript. Table 1 outlines the stepwise procedure employed in the study.

Table 1. Procedural flow of study implementation.

Step	Procedure	Description
Step 1: Informed Consent	1.1 Ethical Briefing	Parents/guardians receive a detailed explanation of the study's aims, procedures, and potential risks.
	1.2 Consent Form Signing	Written informed consent is obtained from parents/guardians in accordance with ethical guidelines.
	1.3 Clinical interview	Children, parents/guardians, and educators.
Step 2: Cognitive Assessment	2.1 Instruction Phase	Children are given instructions and sample items to familiarize them with RSPM format.
	2.2 RSPM Test Completion	Children complete the RSPM to assess fluid intelligence and abstract reasoning.
	2.3 Break (if needed)	A short break is provided to ensure sustained attention and optimal performance.
Step 3: EEG Data Acquisition	3.1 EEG Preparation	Electrode placement, impedance checks, and EEG system calibration are conducted.
	3.2 Auditory Oddball Paradigm (ERP Task)	Children perform an auditory oddball task to elicit the P300 component while EEG data are recorded.
	3.3 Reaction Time Recording	Behavioral responses (button presses) are recorded concurrently with EEG to measure reaction time.
	3.4 Data Quality Check	EEG data undergo visual inspection to ensure artifact-free, high-quality recordings.

2.2. Implementation

All children participating in the study were assessed using Raven's Standard Progressive Matrices (RSPM) to evaluate their non-verbal IQ. The RSPM comprises 36 items divided into three sets—A, Ab, and B—with each set containing 12 matrices that increase progressively in complexity. In each matrix, participants encounter an in-complete visual pattern and must choose the correct piece from six options to logically complete the pattern.

Set A introduces participants to basic principles of pattern recognition and visual-spatial reasoning, featuring simple and easily discernible designs. Moving to Set Ab, the complexity escalates, bridging the gap between Sets A and B by demanding greater attention to detail and enhanced relational reasoning. Set B challenges participants with intricate visual relationships and abstract patterns, requiring advanced cognitive processing and sophisticated problem-solving strategies.

Participants were instructed to select the option that logically completes each pattern. Scoring is based on the total number of correct responses, which are then compared to age-specific normative data to gauge the child's cognitive performance relative to their peers. High scores on this scale are indicative of robust abstract reasoning abilities and cognitive

flexibility, whereas lower scores may suggest difficulties in visual-spatial processing or problem-solving capabilities.

The assessment was carried out at the Digital Neuropsychological Assessment Laboratory of the University of Thessaly, within a specially designated research space. A psychologist was present throughout the session to provide instructions and ensure adherence to the timing specifications outlined in the test manual [7,8,43], ensuring a standardized testing environment conducive to accurate measurement of cognitive abilities.

2.3. Electrophysiological Assessment

2.3.1. Electrode Placement and Data Recording

ERPs were measured using a Medtronic device (710 Medtronic Parkway, Minneapolis, MN, USA). Ag-AgCl electrodes were placed at 15 sites (FP1, FPz, FP2, F3, Fz, F4, T3, T4, C3, Cz, C4, P3, Pz, P4, and Oz) in accordance with the 10–20 International System [44]. The ground electrode was placed at the nasion, and all channels were referenced to linked mastoids. Also, EOG electrodes were placed below the eye (for vertical movements) and the other at the outer canthi (for horizontal movements). Electrode impedance was maintained at <5 kΩ. Recordings were made at a sampling rate of 256 Hz, with a bandpass filter set between 0.16 Hz and 70 Hz. EEG data were segmented into epochs spanning 200 ms pre-stimulus to 800 ms post-stimulus.

2.3.2. P300 Component Detection

The P300 component latency was measured for both target and non-target stimuli across all channels. The P300 waveform, identified by its long positive peak occurring 250–500 ms after the N200 waveform, was visually inspected for accurate detection. It is noteworthy that the researchers did not employ algorithmic approaches for the identification of P300 latencies. While automated methods offer certain advantages, such as increased efficiency, they also present limitations that can be influenced by factors including data quality, signal filtering parameters, the nature of the cognitive task, and the specific research objectives. To address potential shortcomings associated with automated detection, the researchers conducted visual inspections of the electrophysiological data and implemented manual adjustments to ensure the accuracy and reliability of P300 latency measurements. If a reliable P300 peak was not observed, the data for that ERP were marked invalid [45]. Participants were seated comfortably with their eyes closed to minimize visual distractions and reduce ocular artifacts during EEG recording.

2.3.3. Auditory Stimuli

A passive auditory oddball paradigm was employed to generate the P300 component of ERPs. Auditory stimuli were delivered binaurally through headphones, featuring 200 auditory tones at 85 dB SPL. Each tone lasted for 40 ms with 10 ms rise and fall times, aimed at reducing sudden auditory shifts. The stimuli were distributed with a fixed interstimulus interval (ISI) of 1500 ms, enhancing the time frame for cognitive processing and minimizing the overlap of neural responses. The standard (non-target) stimuli were tones at 1000 Hz, occurring with an 80% probability, while the oddball stimuli were 2000 Hz tones, occurring less frequently (20%) to elicit the P300 response. Participants were instructed to sit comfortably and respond to the target tones by pressing a button with their right hand quickly and accurately upon recognition. The response times were recorded with high precision using a response box, which was synchronized with the stimulus presentation software, ensuring millisecond precision in capturing both stimulus onset and response timing within the EEG data stream. Following EEG data collection, reaction times were obtained from the event markers and assessed together with P300 latencies. Software

designed to identify correct and erroneous responses (provided by Medtronic was used to selectively remove epochs associated with incorrect responses from the analysis dataset.

2.3.4. Data Preprocessing and Artifact Removal

To ensure high-quality data, trials were excluded if the voltage exceeded 70 μV in any of the 15 channels (excluding EOG) or if participants provided incorrect responses. A notch filter was applied at 50 Hz to remove power line artifacts, as recommended in EEG processing guidelines [44]. Baseline correction was performed using the 200 ms pre-stimulus period, and epochs were averaged separately for target and non-target stimuli [46,47]. Only children with at least 30 artifact-free trials for both conditions were included in the analysis [48]. Recordings took place in a soundproof room to prevent external interruptions. This methodology ensured reliable electrophysiological measurements and established correlations between cognitive ability and P300 latency [49].

2.4. Statistical Analysis

Initially, descriptive statistics were conducted to outline the demographic and clinical characteristics of the participants. The Kolmogorov–Smirnov test was then utilized to check the distribution of the data, confirming their conformity to a normal distribution. Following this, a *t*-test was employed to explore the differences in P300 latency between groups of children categorized as having high versus average mental abilities. The Benjamini–Hochberg (BH) procedure was calculated to manage the False Discovery Rate (FDR), which measures the expected rate of incorrect rejections of the null hypothesis among all the hypotheses tested. This approach was used in the present study as it involved multiple comparisons. Subsequently, a power analysis was conducted to determine the required sample sizes for achieving statistical power using alpha = 0.05. This power analysis aimed to assess the capability of the study design to detect true effects, thereby minimizing the risk of Type II errors. The analysis was performed assuming equal sample sizes across groups, which is a common scenario in neuropsychological research. Additionally, reaction times were compared between the two groups using another *t*-test. To quantify the size of the observed differences, effect sizes were calculated using Cohen's d [49]. This measure provides a standardized method of assessing the practical significance of the findings, offering a clearer understanding of the impact of cognitive abilities on these neurophysiological responses. Finally, correlation analyses were conducted to examine the relationships between P300 latency and the children's scores on Raven's Standard Progressive Matrices (RSPM). Given that RSPM scores might exhibit non-normal distribution due to developmental variability and the relatively small sample sizes typical in such studies, these analyses were crucial for understanding how these scores related to neurophysiological measurements. This approach aimed to elucidate further the interplay between standardized intelligence measures and underlying brain activities.

3. Results

In assessing cognitive processing through electrophysiological metrics, a *t*-test was employed to elucidate differences between cohorts of children exhibiting high mental abilities and those demonstrating average mental abilities according to RSPM. The analytic focus included mean scores and variability, expressed as standard deviations, of P300 latency across 15 electroencephalographic sites. Additionally, the magnitude of observed differences was quantified using Cohen's d to present effect sizes. The statistical analysis also included calculations to determine the observed power of the P300 latency for participants in both groups. Table 2 summarizes these results, offering details on mean

scores, standard deviations, levels of statistical significance, effect sizes, and the observed power for P300 latency at each electroencephalographic site.

Table 2. Summary of descriptive statistics, *t*-test outcomes, effect sizes, and power analysis for P300 latencies among children with high and average mental abilities.

Electro/ Encephalographic Sites	P300 Latency of Children with High Mental Abilities	SD	P300 Latency of Children with Average Mental Abilities	SD	t	*p*	Cohen's d	Observed Power
Fp1	304.48	6.21	316.58	3.46	−5.89	<0.001	2.41	1.00
FPz	305.44	7.64	318.41	4.63	−5.89	<0.001	2.05	0.99
Fp2	307.55	6.33	320.53	5.92	−5.03	<0.001	2.12	0.99
F3	307.00	7.14	326.65	7.56	−5.03	<0.001	2.67	1.00
Fz	307.87	6.05	325.53	2.21	−5.18	<0.001	3.88	1.00
F4	313.15	10.31	336.38	1.99	−5.18	<0.001	3.13	1.00
T3	306.28	10.03	329.00	4.82	−6.54	<0.001	2.89	1.00
T4	308.32	11.85	325.85	2.45	−6.54	<0.001	2.05	0.99
C3	306.78	13.10	330.45	6.37	−9.49	<0.001	2.30	1.00
Cz	311.69	15.84	337.53	3.22	−9.49	<0.001	2.26	1.00
C4	317.77	9.41	338.19	2.89	−7.66	<0.001	2.93	1.00
P3	309.51	10.04	329.96	5.21	−7.66	<0.001	2.56	1.00
Pz	311.98	14.98	336.37	5.45	−7.07	<0.001	2.16	0.99
P4	313.39	17.56	339.68	3.67	−7.07	<0.001	2.07	0.99
Oz	316.91	10.38	337.87	5.71	−5.01	<0.001	2.50	1.00

P300 latency exhibited statistically significant variations correlating with scores on RSPM. Specifically, children characterized by higher mental abilities demonstrated shorter P300 latencies compared to their peers with average mental abilities. The negative values observed, ranging from −9.49 to −5.89, consistently indicated longer mean P300 latencies for children with average mental abilities across all electroencephalographic recording sites. Moreover, Cohen's d values, ranging from 3.88 to 2.05, present the differences in cognitive response times between the two cohorts. The results of the power analysis reveal that the statistical power for detecting differences in P300 latencies between children with high and average mental abilities at all 15 electroencephalographic sites was high. Specifically, the observed power ranged from 0.99 to 1.00 across the different sites. These values indicate that the present study was highly effective in identifying true differences when they exist, minimizing the risk of Type II errors.

For a comprehensive visual representation, median P300 latencies were calculated for each child across all 15 electrodes, providing a measure of cognitive response time. The resulting boxplots in Figure 1 effectively illustrate the median, interquartile range, and variability within each group, enhancing the interpretability of how cognitive processing speeds differ between children with high and average mental abilities.

The findings facilitate an examination of neural processing speeds, revealing that children with high mental abilities typically demonstrate faster and more equal neural responses. This is reflected in shorter P300 latencies along with reduced variability in these measurements, suggesting enhanced efficiency in cognitive processing.

To ensure the findings of these observations, the Benjamini–Hochberg procedure was employed to control the False Discovery Rate (FDR) among multiple comparisons across various EEG recording sites. This statistical approach aids in observing the true differences in P300 latencies between groups, categorized by differing mental abilities, minimizing the risk of Type I errors in the analysis. Table 3 encapsulates the outcomes of applying this procedure, offering a detailed account of the controlled FDR values across the multiple EEG sites involved in the study.




Figure 1. Comparison of P300 latency of children with high and average mental abilities.

Table 3. Summary of 15 EEG site analysis using the Benjamini–Hochberg test.

Electro/ Encephalographic Sites	*p*-Value	BH Critical Value
Fp1	<0.05	0.02
FPz	<0.05	0.01
Fp2	<0.05	0.03
F3	<0.05	0.01
Fz	<0.05	0.01
F4	<0.05	0.01
T3	<0.05	0.01
T4	<0.05	0.05
C3	<0.05	0.03
Cz	<0.05	0.04
C4	<0.05	0.01
P3	<0.05	0.02
Pz	<0.05	0.04
P4	<0.05	0.04
Oz	<0.05	0.02

Each of the 15 EEG sites demonstrated statistically significant differences in P300 latency, indicating consistent differences in neural processing speeds across the complete spectrum of measured sites between the two groups. The low *p*-values observed offer substantive evidence against the null hypothesis, which posits no difference in latencies between the groups. Further examination through comparison of these *p*-values against the Benjamini–Hochberg (BH) values confirms that the observed differences are indeed statistically significant, even after adjusting for multiple comparisons. This methodology confirms that the conclusions are not merely the result of random chance, thereby supporting the reliability of the findings.

Next, the analysis focused on examining the reaction times of children with high mental abilities compared to those with average mental abilities. It is important to remember that all participants were asked to press a button as quickly and accurately as possible when they detected the target stimulus (oddball). Table 4 displays the mean scores, standard deviations, statistical significance, and effect sizes for the reaction times from both groups of participants.

Table 4. Mean scores, standard deviations, *t*-test results, and significance and Cohen's d of reaction time for children that participated in each group of mental abilities.

Reaction Time	High Mental Abilities		Average Mental Abilities				
	M	**SD**	**M**	**SD**	**t**	***p***	**Cohen's d**
	319.70	6.54	352.30	11.76	−8.39	<0.001	3.43

As shown in Table 4, children with high mental abilities demonstrated shorter reaction times than their peers with average mental abilities, with statistical significance differences. Additionally, the negative t-value presents that the average reaction times for children with average mental abilities were generally longer than those for children with higher mental abilities. Furthermore, a high Cohen's D value suggests a significant effect size in these differences.

To further explore the relationship between cognitive metrics, a Spearman's ρ correlation analysis was conducted. This analysis assessed the associations between the latency of the P300 waveform and the RSPM scores of children, covering all 15 topographic brain areas. The findings from this correlation analysis are detailed in Table 5.

Table 5. Correlation analysis between P300 latency and RSPM scores.

Electro/ Encephalographic Sites	Correlation ρ	Sign
FP1	−0.844	0.001
FPZ	−0.804	0.001
FP2	−0.742	0.001
F3	−0.862	0.001
FZ	−0.889	0.001
F4	−0.813	0.001
T3	−0.818	0.001
T4	−0.893	0.001
C3	−0.844	0.001
CZ	−0.865	0.001
C4	−0.770	0.001
P3	−0.853	0.001
PZ	−0.803	0.001
P4	−0.790	0.001
OZ	−0.781	0.001

The results showed a range of findings, with correlations varying from ρ = −0.742 at the FP2 electrode site to ρ = −0.893 at the T4 electrode site. The analysis highlighted a strong negative correlation, suggesting that higher RSPM scores are associated with shorter P300 latencies.

Following the same statistical approach, the correlation between reaction time and RSPM scores was also examined. A significant negative correlation (ρ = −0.877) was observed, indicating that higher RSPM scores tend to correlate with quicker reaction times.

4. Discussion

In this study, we aimed to explore differences in electrophysiological brain activity between children classified by Raven's Standard Progressive Matrices as having "high mental abilities" and those with "average mental abilities". Our first hypothesis, in the same line of earlier research, e.g., [23,24,30–33], examined whether P300 latency differs between these two groups. Our findings confirmed this, showing that children with high mental abilities consistently had shorter P300 latencies across various electroencephalographic sites

compared to their peers with average abilities. The observed power results contribute to a more reliable understanding of the neural mechanisms underlying cognitive differences in children, as measured by P300 latencies. The implications of such power reinforce the validity of our conclusions and support the potential application of these findings in both neuropsychological and educational settings.

This pattern of shorter P300 latencies in children with higher cognitive abilities, consistent with results from other studies, e.g., [30–33], might explain their superior performance on the RSPM. A shorter P300 latency suggests better memory and attentional functions [50], reflecting quicker stimulus assessment and more efficient cognitive processing. This efficiency is significant as studies have found that individuals with better memory capabilities often exhibit shorter auditory P300 latencies [51,52], while a longer P300 latency might indicate attentional disorders [53].

Overall, longer P300 latencies are generally associated with lower cognitive performance [47]. These results not only support our hypothesis but also align with the neural efficiency model, which suggests that cognitive efficiency varies with task complexity. This model is further supported by our use of the auditory oddball paradigm, a task that requires selective attention and swift decision-making. This task seems particularly effective at highlighting cognitive differences, supporting the suggestions that brain efficiency is task-dependent and varies between individuals with different levels of cognitive abilities [5].

The Benjamini–Hochberg (BH) correction was used to minimize the likelihood of Type I statistical errors that could arise from the multiple comparisons across 15 EEG sites. This adjustment of the significance threshold helps ensure the reliability and validity of the study's results, given the extensive number of statistical tests performed. Significant differences in P300 latency between children with high and average mental abilities were noted at all EEG sites, with p-values falling well below the adjusted threshold. These results highlight notable differences in cognitive processing speed and efficiency between the two groups, providing further evidence in support of the neural efficiency theory. Additionally, these findings emphasize the value of P300 latency as a neurophysiological marker for discerning and understanding variations in cognitive abilities, reinforcing its practical importance in educational and developmental settings.

The second hypothesis of this study was encouraged by prior research suggesting that individuals with higher mental abilities often exhibit shorter reaction times [18,52–57]. Consistent with these findings, our study observed that children with higher mental abilities demonstrated statistically significant shorter reaction times compared to their peers with average abilities.

This reduction in reaction time may reflect superior memory and attentional functions, as it has been proposed that children and adolescents with higher mental abilities are better at inhibiting premature responses, allowing them to answer with greater confidence [58]. Additionally, shorter reaction times are thought to be indicative of enhanced self-regulation capabilities, suggesting that highly intelligent children may possess more effective self-regulation strategies [6].

By integrating both electrophysiological and behavioral data, as it is suggested, children with higher intelligence levels display more mature and efficient neural functioning in cognitive tasks [15,56]. The ability for rapid stimulus detection and response proposes improved sensory processing and motor execution, facilitated by reduced neural "noise" and more optimized cortical activation patterns in those with elevated cognitive abilities [59,60]. These observations align with neuroscientific research indicating that as intelligence increases, reaction time variability decreases, pointing to more stable and efficient cognitive processing across various tasks [38].

The considerable Cohen's d values reported for both P300 latency and reaction time not only reinforce the statistical significance of these findings but also their practical relevance. These large effect sizes illuminate the profound impact of cognitive differences on neural processing speeds and serve as crucial metrics in this study, highlighting the practical implications of our findings, particularly in the contexts of cognitive and educational development.

The third hypothesis anticipated a correlation between P300 latency and reaction time with RSPM scores, based on other studies that noted a positive correlation between P300 amplitude and RSPM scores, and a negative correlation between P300 latency and RSPM scores [31,61]. Our analysis suggested negative correlations between both P300 latency and reaction time with RSPM scores across the two groups, linking higher intelligence scores to shorter P300 latencies and reaction times.

4.1. Psychoeducational Implications of the Study Using ERPs and RSPM Results in Identifying Children's Mental Abilities

The combination of P300 and RSPM in cognitive classification has meaningful applications in psychoeducational contexts. By considering both P300 latency and RSPM scores, it becomes possible to identify gifted individuals at an early stage, which can support timely educational interventions [62]. Distinguishing between children with high and average mental abilities allows educators to develop personalized learning strategies that cater to their cognitive profiles [63]. For instance, children with shorter P300 latencies may benefit from accelerated learning programs, while those with average latencies might need more structured, scaffolded instruction to optimize their learning experience [63,64].

Additionally, longitudinal research that incorporates both P300 waveforms and RSPM scores provides a valuable framework for monitoring cognitive development over time. These studies contribute to understanding neurodevelopmental trajectories by examining how cognitive processing and executive functioning mature across different age groups [24]. The integration of electrophysiological markers such as P300 latency with behavioral assessments from the RSPM facilitates an evaluation of how educational interventions impact attentional control, working memory, and fluid intelligence at various developmental stages [65]. This approach provides insights into the ways in which cognitive functions influence learning, ultimately guiding evidence-based teaching practices. By classifying children based on cognitive abilities, such methodologies help acknowledge the diversity of intelligence and inform educational policies that support the development of all learners [66–70]. In this way, neuroscience contributes to practical solutions in education, bridging the gap between research and real-world applications [68].

4.2. Limitations

There are several methodological considerations to consider when interpreting the findings of this study. The first notable limitation is the inclusion of only male participants, which significantly restricts the generalizability of the results. The absence of female participants means it remains unclear whether the observed relationships between P300 latency, reaction time, and fluid intelligence hold true across genders. This lack of gender diversity limits the applicability of the findings to the broader population of children, potentially overlooking important gender-specific cognitive differences. Additionally, the study's relatively small sample size, a result of the stringent criteria required to recruit children with high scores on the Raven's Standard Progressive Matrices (RSPM), further limits the generalizability. The restrictive age range of participants due to these criteria could also impact the developmental generalizability of the results, as it may not accurately reflect the cognitive processes across different developmental stages. The reliance on the RSPM as the sole measure of fluid intelligence is another limitation. Since the RSPM assesses primarily

non-verbal and abstract problem-solving abilities, it does not evaluate verbal intelligence, which is a crucial component of cognitive functioning. Future research should include a broader array of cognitive assessments to capture a more comprehensive view of intelligence. Moreover, the focus on the P300 waveform as the primary event-related potential (ERP) component analyzed limits the scope of the study. While the P300 is significant for its relevance to cognitive abilities and higher-order cognitive functions, other ERP components, such as the N200, might also provide valuable insights into cognitive processes. A more extensive examination of these components could deepen the understanding of the neural mechanisms involved. To address these limitations, future research should aim to include both male and female participants to enhance the representativeness of the findings. Employing a larger and more diverse sample, incorporating a wider range of cognitive assessments, and exploring additional ERP components would not only strengthen the statistical power but also broaden the applicability and depth of the research findings.

5. Conclusions

In summary, although this study has some limitations, its main findings are in line with previous research. These results suggest that children with higher levels of intelligence often show shorter P300 latencies and quicker reaction times. Rather than simply being faster because they are better at the tasks, this pattern might point to more efficient brain activity, as it is suggested by the neural efficiency hypothesis [5]. This idea is supported by the fact that quicker brain responses, both in processing incoming information (P300 latency) and in responding behaviorally (reaction time), are often linked to better performance in areas like attention, memory, and self-control. These abilities are important for learning and doing well in school and are usually stronger in children who score higher on cognitive tests [34,71]. Because of this, the study offers a possible connection between brain activity and academic success. The results also reflect what earlier studies have found. Earlier studies have shown that children with higher IQs or higher scores on the Raven's Standard Progressive Matrices (RSPM) tend to have shorter P300 latencies, e.g., [21]. This could mean that these children are processing the information they are given more quickly and perhaps with more organized brain activity [72]. By grouping children based on their fluid intelligence, this study looked closely at how cognitive skills develop during middle childhood. This is a key stage where attention and other executive functions grow rapidly [73], and understanding how efficiently the brain works during this period can be especially helpful as children face increasing academic challenges.

Measuring both reaction time and P300 latency allowed the study to explore how efficiently different children process information. These two measurements work well together because they help show not just how quickly a child can respond, but also how consistently and accurately their brain is working. Research has shown that slower or more variable responses are often linked with more difficult tasks, while faster and steadier responses tend to show up in children with higher intelligence [57,59]. The fact that both of these brain measures were linked with RSPM scores gives further support to using them as tools for understanding cognitive ability. The size of the differences found, measured using effect sizes, suggests that these results could be meaningful in real-world settings, not just statistically. If used carefully, these types of brain measurements might help with early identification of children who could benefit from extra cognitive challenges or support. Looking to the future, this study may enable long-term research on how brain efficiency develops over time. It would be useful to know how different factors, like a child's home environment, school experience, or access to learning resources, shape their cognitive development [74]. Tracking these patterns over the years could help understanding whether the brain advantages seen in higher-ability children continue, change, or even fade. It could

also show whether specific types of cognitive training might help other children to improve how their brains process information.

In conclusion, this study supports previous findings presenting how intelligence relates to brain function in children. Using tools like P300 latency and reaction time gives researchers and educators a broader way to explore how children think and learn. It supports the idea that smarter brain processing may look different on a neural level and provides new ways to think about helping all children reach their potential.

Author Contributions: Data curation, I.D., P.P., L.M. and E.I.T.; Formal analysis, N.C.Z. and I.D.; Investigation, N.C.Z.; Methodology, N.C.Z. and E.I.T.; Visualization, N.C.Z.; Writing—original draft, N.C.Z. and I.D.; Writing—review and editing, P.P., L.M. and E.I.T. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: All participants' parents/guardians were required to sign the consent form allowing their child to participate in the research. Finally, all human data included in this manuscript were obtained in compliance with the Helsinki Declaration and the guidelines of the University of Thessaly Internal Research Ethics Committee (EHDE) protocol (code 227092023).

Informed Consent Statement: Informed consent was obtained from the parents or guardians of all children involved in the study.

Data Availability Statement: The original contributions presented in the study are included in the article; further inquiries can be directed to the corresponding author.

Acknowledgments: The authors acknowledge the use of ChatGPT [48], an AI-powered language model, for leveraging advanced natural language processing (NLP) techniques to enhance English writing. ChatGPT assisted in refining grammar, sentence structure, and overall coherence, benefiting both native and non-native speakers. Its application contributed to improving the clarity of the manuscript.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Mitchell, D.J.; Mousley, A.L.; Shafto, M.A.; Duncan, J. Neural contributions to reduced fluid intelligence across the adult lifespan. *J. Neurosci.* **2023**, *43*, 293–307. [CrossRef] [PubMed]
2. Scherrer, V.; Breit, M.; Preckel, F. Crystallized Intelligence, Fluid Intelligence, and Need for Cognition: Their Longi-tudinal Relations in Adolescence. *J. Intell.* **2024**, *12*, 104. [CrossRef] [PubMed]
3. Wang, T.; Ren, X.; Altmeyer, M.; Schweizer, K. An account of the relationship between fluid intelligence and complex learning in considering storage capacity and executive attention. *Intelligence* **2013**, *41*, 537–545. [CrossRef]
4. Amin, H.U.; Malik, A.S.; Kamel, N.; Chooi, W.T.; Hussain, M. P300 correlates with learning & memory abilities and fluid intelligence. *J. Neuroeng. Rehabil.* **2015**, *12*, 1–14.
5. Neubauer, A.C.; Fink, A. Intelligence and neural efficiency. *Neurosci. Biobehav. Rev.* **2009**, *33*, 1004–1023. [CrossRef]
6. Liu, T.; Xiao, T.; Shi, J.; Zhao, D.; Liu, J. Conflict control of children with different intellectual levels: An ERP study. *Neurosci. Lett.* **2011**, *490*, 101–106. [CrossRef]
7. Raven, J.C. Raven standard progressive matrices. *J. Cogn. Dev.* **1936**. [CrossRef]
8. Raven, J. The Raven Progressive Matrices Tests: Their Theoretical Basis and Measurement Model. Uses and Abuses of Intelligence. Studies Advancing Spearman and Raven's Quest for Non-arbitrary Metrics (Part I). 2008. Available online: http://www.eyeonsociety.co.uk/resources/UAIChapter1.pdf (accessed on 17 February 2025).
9. Zurrin, R.; Wong, S.T.S.; Roes, M.M.; Percival, C.M.; Chinchani, A.; Arreaza, L.; Kusi, M.; Momeni, A.; Rasheed, M.; Mo, Z.; et al. Functional brain networks involved in the Raven's standard progressive matrices task and their relation to theories of fluid intelligence. *Intelligence* **2024**, *103*, 101807. [CrossRef]
10. Duncan, J.; Chylinski, D.; Mitchell, D.J.; Bhandari, A. Complexity and compositionality in fluid intelligence. *Proc. Natl. Acad. Sci. USA* **2017**, *114*, 5295–5299. [CrossRef]

11. Tanaka, F.; Kachi, T.; Yamada, T.; Sobue, G. Auditory and visual event-related potentials and flash visual evoked potentials in Alzheimer's disease: Correlations with Mini-Mental State Examination and Raven's Coloured Progressive Matrices. *J. Neurol. Sci.* **1998**, *156*, 83–88. [CrossRef]
12. Jung, R.E.; Haier, R.J. The Parieto-Frontal Integration Theory (P-FIT) of intelligence: Converging neuroimaging evidence. *Behav. Brain Sci.* **2007**, *30*, 135–154. [CrossRef] [PubMed]
13. Zacks, J.M. Neuroimaging studies of mental rotation: A meta-analysis and review. *J. Cogn. Neurosci.* **2008**, *20*, 1–19. [CrossRef] [PubMed]
14. Haier, R.J.; Siegel, B.; Tang, C.; Abel, L.; Buchsbaum, M.S. Intelligence and changes in regional cerebral glucose metabolic rate following learning. *Intelligence* **1992**, *16*, 415–426. [CrossRef]
15. Neubauer, A.C.; Grabner, R.H.; Fink, A.; Neuper, C. Intelligence and neural efficiency: Further evidence of the influence of task content and sex on the brain–IQ relationship. *Cogn. Brain Res.* **2005**, *25*, 217–225. [CrossRef]
16. Zygouris, N.C. Differences in children and adolescents with depression before and after a remediation program: An event-related potential study. *Brain Sci.* **2024**, *14*, 660. [CrossRef]
17. Karapetsas, A.V.; Zygouris, N.C. Event Related Potentials (ERPs) in prognosis, diagnosis and rehabilitation of children with dyslexia. *Encephalos* **2011**, *48*, 118–127.
18. Howe, A.S.; Bani-Fatemi, A.; De Luca, V. The clinical utility of the auditory P300 latency subcomponent event-related potential in preclinical diagnosis of patients with mild cognitive impairment and Alzheimer's disease. *Brain Cogn.* **2014**, *86*, 64–74. [CrossRef]
19. Demirayak, P.; Kıyı, İ.; İşbitiren, Y.Ö.; Yener, G. Cognitive load associates prolonged P300 latency during target stimulus processing in individuals with mild cognitive impairment. *Sci. Rep.* **2023**, *13*, 15956. [CrossRef]
20. Zhong, R.; Li, M.; Chen, Q.; Li, J.; Li, G.; Lin, W. The P300 event-related potential component and cognitive im-pairment in epilepsy: A systematic review and meta-analysis. *Front. Neurol.* **2019**, *10*, 943. [CrossRef]
21. Sternberg, R.J. (Ed.) *The Cambridge Handbook of Intelligence*; Cambridge University Press: Cambridge, UK, 2020.
22. Walhovd, K.B.; Nyberg, L.; Lindenberger, U.; Amlien, I.K.; Sørensen, Ø.; Wang, Y.; Mowinckel, A.M.; Kievit, R.A.; Ebmeier, K.P.; Bartrés-Faz, D.; et al. Brain aging differs with cognitive ability regardless of education. *Sci. Rep.* **2022**, *12*, 13886. [CrossRef]
23. Warchoł, Ł.; Zając-Lamparska, L. The Relationship of N200 and P300 Amplitudes with Intelligence, Working Memory, and Attentional Control Behavioral Measures In Young Healthy Individuals. *Adv. Cogn. Psychol.* **2023**, *19*, 63–75. [CrossRef]
24. Walhovd, K.B.; Fjell, A.M. One-year test–retest reliability of auditory ERPs in young and old adults. *Int. J. Psychophysiol.* **2002**, *46*, 29–40. [CrossRef] [PubMed]
25. Polich, J. Neuropsychology of P300. In *The Oxford Handbook of Event-Related Potential Components*; Kappenman, E.S., Luck, S.J., Eds.; Oxford University Press: Oxford, UK, 2011; pp. 160–188.
26. Sternberg, R.J. A theory of adaptive intelligence and its relation to general intelligence. *J. Intell.* **2019**, *7*, 23. [CrossRef] [PubMed]
27. Lozano-Blasco, R.; Quílez-Robres, A.; Usán, P.; Salavera, C.; Casanovas-López, R. Types of intelligence and academic performance: A systematic review and meta-analysis. *J. Intell.* **2022**, *10*, 123. [CrossRef]
28. Ren, X.; Schweizer, K.; Wang, T.; Chu, P.; Gong, Q. On the relationship between executive functions of working memory and components derived from fluid intelligence measures. *Acta Psychol.* **2017**, *180*, 79–87. [CrossRef]
29. Ren, X.; Wang, T.; Sun, S.; Deng, M.; Schweizer, K. Speeded testing in the assessment of intelligence gives rise to a speed factor. *Intelligence* **2018**, *66*, 64–71. [CrossRef]
30. Bazana, P.G.; Stelmack, R.M. Intelligence and information processing during an auditory discrimination task with backward masking: An event-related potential analysis. *J. Personal. Soc. Psychol.* **2002**, *83*, 998. [CrossRef]
31. De Pascalis, V.A.; Varriale, V.; Matteoli, A. Intelligence and P3 components of the event-related potential elicited during an auditory discrimination task with masking. *Intelligence* **2008**, *36*, 35–47. [CrossRef]
32. Troche, S.J.; Houlihan, M.E.; Stelmack, R.M.; Rammsayer, T.H. Mental ability, P300, and mismatch negativity: Analysis of frequency and duration discrimination. *Intelligence* **2009**, *37*, 365–373. [CrossRef]
33. Teixeira-Santos, A.C.; Pinal, D.; Pereira, D.R.; Leite, J.; Carvalho, S.; Sampaio, A. Probing the relationship between late endogenous ERP components with fluid intelligence in healthy older adults. *Sci. Rep.* **2020**, *10*, 11167. [CrossRef]
34. Polich, J. Updating P300: An integrative theory of P3a and P3b. *Clin. Neurophysiol.* **2007**, *118*, 2128–2148. [CrossRef] [PubMed]
35. Der, G.; Deary, I.J. The relationship between intelligence and reaction time varies with age: Results from three rep-resentative narrow-age age cohorts at 30, 50 and 69 years. *Intelligence* **2017**, *64*, 89–97. [CrossRef] [PubMed]
36. Kannen, K.; Aslan, B.; Boetzel, C.; Herrmann, C.S.; Lux, S.; Rosen, H.; Selaskowski, B.; Wiebe, A.; Philipsen, A.; Braun, N. P300 modulation via transcranial alternating current stimulation in adult attention-deficit/hyperactivity disorder: A crossover study. *Front. Psychiatry* **2022**, *13*, 928145. [CrossRef] [PubMed]
37. Jungeblut, H.M.; Hagemann, D.; Löffler, C.; Schubert, A.L. An investigation of the slope parameters of reaction times and P3 latencies in the Sternberg memory scanning task—A fixed-links model approach. *J. Cogn.* **2021**, *4*, 26. [CrossRef]
38. Schubert, A.L.; Löffler, C.; Hagemann, D.; Sadus, K. How robust is the relationship between neural processing speed and cognitive abilities? *Psychophysiology* **2023**, *60*, e14165. [CrossRef]

39. Schubert, A.L. A meta-analysis of the worst performance rule. *Intelligence* **2019**, *73*, 88–100. [CrossRef]
40. Regel, S.; Meyer, L.; Gunter, T.C. Distinguishing neurocognitive processes reflected by P600 effects: Evidence from ERPs and neural oscillations. *PLoS ONE* **2014**, *9*, e96840. [CrossRef]
41. Beldzik, E.; Ullsperger, M. A thin line between conflict and reaction time effects on EEG and fMRI brain signals. *Imaging Neurosci.* **2024**, *2*, 1–17. [CrossRef]
42. Raven, J. Raven progressive matrices. In *Handbook of Nonverbal Assessment*; Springer: Boston, MA, USA, 2003; pp. 223–237.
43. Jasper, H.H. Ten-twenty electrode system of the international federation. *Electroencephalogr. Clin. Neurophysiol.* **1958**, *10*, 371–375.
44. Sadus, K.; Schubert, A.L.; Löffler, C.; Hagemann, D. An explorative multiverse study for extracting differences in P3 latencies between young and old adults. *Psychophysiology* **2024**, *61*, e14459. [CrossRef]
45. Picton, T.W.; Bentin, S.; Berg, P.; Donchin, E.; Hillyard, S.A.; Johnson, R.; Miller, G.A.; Ritter, W.; Ruchkin, D.S.; Rugg, M.D.; et al. Guidelines for using human event-related potentials to study cognition: Recording standards and publication criteria. *Psychophysiology* **2000**, *37*, 127–152. [CrossRef] [PubMed]
46. Zygouris, N.C.; Avramidis, E.; Karapetsas, A.V.; Stamoulis, G.I. Differences in dyslexic students before and after a remediation program: A clinical neuropsychological and event related potential study. *Appl. Neuropsychol. Child* **2018**, *7*, 235–244. [CrossRef] [PubMed]
47. Zygouris, N.C.; Vlachos, F.; Stamoulis, G.I. ERPs in Children and Adolescents with Generalized Anxiety Disorder: Before and after an Intervention Program. *Brain Sci.* **2022**, *12*, 1174. [CrossRef] [PubMed]
48. OpenAI. ChatGPT [Large Language Model]. 2024. Available online: https://openai.com (accessed on 28 February 2025).
49. Cohen, J. Set correlation and contingency tables. *Appl. Psychol. Meas.* **1988**, *12*, 425–434. [CrossRef]
50. Beauchamp, C.M.; Stelmack, R.M. The chronometry of mental ability: An event-related potential analysis of an auditory oddball discrimination task. *Intelligence* **2006**, *34*, 571–586. [CrossRef]
51. Wongupparaj, P.; Sumich, A.; Wickens, M.; Kumari, V.; Morris, R.G. Individual differences in working memory and general intelligence indexed by P200 and P300: A latent variable model. *Biol. Psychol.* **2018**, *139*, 96–105. [CrossRef]
52. McGarry-Roberts, P.A.; Stelmack, R.M.; Campbell, K.B. Intelligence, reaction time, and event-related potentials. *Intelligence* **1992**, *16*, 289–313. [CrossRef]
53. Sur, S.; Sinha, V.K. Event-related potential: An overview. *Ind. Psychiatry J.* **2009**, *18*, 70–73. [CrossRef]
54. Gmaj, B.; Januszko, P.; Kamiński, J.; Drozdowicz, E.; Kopera, M.; Wołyńczyk-Gmaj, D.; Wojnar, M. EEG source activity during processing of neutral stimuli in subjects with anxiety disorders. *Acta Neurobiol. Exp.* **2016**, *76*, 75–85. [CrossRef]
55. Doebler, P.; Scheffler, B. The relationship of choice reaction time variability and intelligence: A meta-analysis. *Learn. Individ. Differ.* **2016**, *52*, 157–166. [CrossRef]
56. Tsai, Y.C.; Lu, H.J.; Chang, C.F.; Liang, W.K.; Muggleton, N.G.; Juan, C.H. Electrophysiological and behavioral evidence reveals the effects of trait anxiety on contingent attentional capture. *Cogn. Affect. Behav. Neurosci.* **2017**, *17*, 973–983. [CrossRef] [PubMed]
57. Jensen, A.R. *The Factor*; Prager: Westport, CT, USA, 1998.
58. Sanz, M.; Molina, V.; Martin-Loeches, M.; Calcedo, A.; Rubia, F.J. Auditory P300 event related potential and sero-tonin reuptake inhibitor treatment in obsessive-compulsive disorder patients. *Psychiatry Res.* **2001**, *101*, 75–81. [CrossRef] [PubMed]
59. Deary, I.J.; Cox, S.R.; Okely, J.A. Inspection time and intelligence: A five-wave longitudinal study from age 70 to age 82 in the Lothian Birth Cohort 1936. *Intelligence* **2024**, *105*, 101844. [CrossRef]
60. Coles, M.G.H.; Smid, H.G.O.M.; Scheffers, M.K.; Otten, L.J. Mental Chronometry and the study of human in-formation processing. In *Electrophysiology of Mind*; Oxford University Press: Oxford, UK, 1995; pp. 86–127.
61. Shaw, P.; Greenstein, D.; Lerch, J.; Clasen, L.; Lenroot, R.; Gogtay, N.E.E.A.; Evans, A.; Rapoport, J.; Giedd, J. Intellectual ability and cortical development in children and adolescents. *Nature* **2006**, *440*, 676–679. [CrossRef]
62. Kutas, M.; Federmeier, K.D. Thirty years and counting: Finding meaning in the N400 component of the event-related brain potential (ERP). *Annu. Rev. Psychol.* **2011**, *62*, 621–647. [CrossRef]
63. Wronka, E.; Kaiser, J.; Coenen, A.M. Psychometric intelligence and P3 of the event-related potentials studied with a 3-stimulus auditory oddball task. *Neurosci. Lett.* **2013**, *535*, 110–115. [CrossRef]
64. Luck, S.J. *An Introduction to the Event-Related Potential Technique*; MIT Press: Cambridge, MA, USA, 2014.
65. Zygouris, N.C.; Dermitzaki, I.; Karapetsas, A.V. Differences in brain activity of children with higher mental abilities. An Event Related Potentials study using the latency of P300 and N100 waveforms. *Int. J. Dev. Neurosci.* **2015**, *47*, 118–119. [CrossRef]
66. Gagné, F. From gifts to talents: The DMGT as a devalopmental model. In *Conceptions of Giftedness*; Sternberg, R.J., Davidson, J.E., Eds.; Cambridge University Press: New York, NY, USA, 2005; pp. 98–120.
67. Greer, K. Neural Assemblies as Precursors for Brain Function. *NeuroSci* **2022**, *3*, 645–655. [CrossRef]
68. Liu, X.; Yang, S.; Liu, Z. Predicting Fluid Intelligence via Naturalistic Functional Connectivity Using Weighted En-semble Model and Network Analysis. *NeuroSci* **2021**, *2*, 427–442. [CrossRef]
69. Karapetsas, A.; Zygouris, N. Charting the maturation of the prefrontal lobes at school aged children and adolescents, using Event Related Potentials. *Ann. Gen. Psychiatry* **2008**, *7* (Suppl. S1), S355. [CrossRef]

70. Merks, S. Elucidating Different Aspects of Speed of Information Processing: Comparison of Behavioral Response Latency and P300 Latency in a Modified Hick Reaction Time Task. Ph.D. Thesis, Universität Bern, Bern, Switzerland, 2016.
71. Lees, T.; Fry, C.M.; Terrell, S.; Jetha, M.K.; Segalowitz, S.J.; Gatzke-Kopp, L.M. Developmental changes in external and internal performance monitoring across middle childhood: An ERP study. *Int. J. Psychophysiol.* **2021**, *169*, 20–33. [CrossRef]
72. Raufi, B.; Longo, L. An evaluation of the EEG alpha-to-theta and theta-to-alpha band ratios as indexes of mental workload. *Front. Neuroinform.* **2022**, *16*, 861967. [CrossRef] [PubMed]
73. Rico-Picó, J.; Hoyo, Á.; Guerra, S.; Conejero, Á.; Rueda, M.R. Behavioral and brain dynamics of executive control in relation to children's fluid intelligence. *Intelligence* **2021**, *84*, 101513. [CrossRef]
74. De Zwarte, S.M.; Brouwer, R.M.; Agartz, I.; Alda, M.; Alonso-Lana, S.; Bearden, C.E.; Bertolino, A.; Bonvino, A.; Bramon, E.; Buimer, E.E.; et al. Intelligence, educational attainment, and brain structure in those at familial high-risk for schizophrenia or bipolar disorder. *Hum. Brain Mapp.* **2022**, *43*, 414–430. [CrossRef] [PubMed]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.