Supplementary Materials

Figure S1. Examples of digestion grades considered in the present study on bird ulnae. From left to right: not digested (0), slightly digested (1), moderately digested (2), and extremely digested (3). Pictures: V. Laroulandie.

Table S1. Detailed data on the proportions of the prey species found in the three studied pellet samples from the island of Dominica, by site and by observer. For Aves, observers 1 and 2 made counts and determinations together, and this explains why there is no difference between the two datasets.

	Grand Bay		Salisbury		Canefield		Total	
	NISP	MNI	NISP	MNI	NISP	MNI	NISP	MNI
Number of pellets	27		34		50		111	
Observer 1								
Aves	**186**	**23**	**206**	**24**	**295**	**34**	**687**	**81**
Size class 0	107	13	70	9	120	14	297	36
Size class 1	79	10	136	15	172	19	387	44
Size class 2	0	0	0	0	3	1	3	1
Squamata	**84**	**10**	**173**	**20**	**36**	**7**	**293**	**37**
Thecadactylus rapicauda	84	10	12	2	14	2	110	14
Anolis sp.	0	0	161	18	22	5	183	23
Rodentia	**27**	**5**	**39**	**6**	**207**	**26**	**273**	**37**
Mus musculus	3	1	35	5	61	7	99	13
Rattus rattus	24	4	4	1	146	19	174	24
Chiroptera	**103**	**17**	**116**	**16**	**21**	**4**	**240**	**37**
Artibeus jamaicensis	11	2	45	5	12	2	68	9
Artibeus schwartzi	0	0	0	0	2	1	2	1
Artibeus sp.	4	1	0	0	0	0	4	1
Brachyphylla cavernarum	23	6	18	3	7	1	48	10
Ardops nicholsii	44	5	53	8	0	0	97	13
Monophylus plethodon	14	2	0	0	0	0	14	2
Sturnira angelli	7	1	0	0	0	0	7	1
Chiroptera indet.	0	0	0	0	0	0	0	0
Total microvertebrates	**400**	**55**	**534**	**66**	**559**	**71**	**1493**	**192**
Observer 2								
Aves	**186**	**23**	**206**	**24**	**295**	**34**	**687**	**81**
Size class 0	107	13	70	9	120	14	297	36
Size class 1	79	10	136	15	172	19	387	44
Size class 2	0	0	0	0	3	1	3	1
Squamata	**67**	**9**	**145**	**19**	**37**	**8**	**249**	**36**
Thecadactylus rapicauda	67	9	9	2	13	2	89	13
Anolis sp.	0	0	136	17	24	6	160	23
Rodentia	**27**	**5**	**39**	**5**	**208**	**29**	**274**	**39**
Mus musculus	3	1	35	4	61	7	99	12
Rattus rattus	24	4	4	1	147	22	175	27
Chiroptera	**101**	**19**	**118**	**15**	**21**	**6**	**240**	**40**
Artibeus jamaicensis	12	2	35	4	0	0	47	6
Artibeus schwartzi	0	0	0	0	0	0	0	0
Artibeus sp.	0	0	6	1	6	3	12	4
Brachyphylla cavernarum	20	6	18	2	0	0	38	8
Ardops nicholsii	30	4	58	7	0	0	88	11
Monophylus plethodon	12	2	0	0	0	0	12	2
Sturnira angelli	7	1	0	0	0	0	7	1
Chiroptera indet.	20	4	1	1	15	3	36	8
Total microvertebrates	**381**	**56**	**508**	**63**	**561**	**77**	**1450**	**196**
Total (mean)								
Aves	**186.0**	**23.0**	**206.0**	**24.0**	**295.0**	**34.0**	**687.0**	**81.0**
Size class 0	107.0	13.0	70.0	9.0	120.0	14.0	297.0	36.0
Size class 1	79.0	10.0	136.0	15.0	172.0	19.0	387.0	44.0

Size class 2	0.0	0.0	0.0	0.0	3.0	1.0	3.0	1.0
Squamata	**75,5**	**9,5**	**159,0**	**20,0**	**36,5**	**7,3**	**271,0**	**36,0**
Thecadactylus rapicauda	75,5	9,5	10,5	1,8	13,5	2,0	99,5	13,3
Anolis sp.	0,0	0,0	148,5	17,5	23,0	5,3	171,5	22,8
Rodentia	**27,0**	**5,0**	**39,0**	**5,5**	**207,5**	**27,5**	**273,5**	**38,0**
Mus musculus	3,0	1,0	35,0	4,5	61,0	7,0	99,0	12,5
Rattus rattus	24,0	4,0	4,0	1,0	146,5	20,5	174,5	25,5
Chiroptera	**102,0**	**18,0**	**117,0**	**15,5**	**21,0**	**5,0**	**240,0**	**38,5**
Artibeus jamaicensis	11,5	2,0	40,0	4,5	6,0	1,0	57,5	7,5
Artibeus schwartzi	0,0	0,0	0,0	0,0	1,0	0,5	1,0	0,5
Artibeus sp.	2,0	0,5	3,0	0,5	3,0	1,5	8,0	2,5
Brachyphylla cavernarum	21,5	6,0	18,0	2,5	3,5	0,5	43,0	9,0
Ardops nicholsii	37,0	4,5	55,5	7,5	0,0	0,0	92,5	12,0
Monophylus plethodon	13,0	2,0	0,0	0,0	0,0	0,0	13,0	2,0
Sturnira angelli	7,0	1,0	0,0	0,0	0,0	0,0	7,0	1,0
Chiroptera indet.	10,0	2,0	0,5	0,5	7,5	1,5	18,0	4,0
Total microvertebrates	**390,5**	**55,5**	**521,0**	**65,0**	**560,0**	**73,8**	**1471,5**	**193,5**
+ Insect remains	**4**	**/**	**5**	**/**	**2**	**/**	**11**	**/**

Table S2. Detailed data on the size categories of the prey taxa found in the three studied pellet samples from the island of Dominica, by site and by observer. For Aves, observers 1 and 2 made counts and determinations together, and this explains why there is no difference between the two datasets.

	Grand Bay		Salisbury		Canefield		TOTAL	
	NISP	MNI	NISP	MNI	NISP	MNI	NISP	MNI
Observer 1								
Aves								
Size class 0	107	13	70	9	120	14	297	36
Size class 1	79	10	136	15	172	19	387	44
Size class 2	0	0	0	0	3	1	3	1
Squamata								
Size class 0	0	0	0	0	0	0	0	0
Size class 1	84	10	173	20	36	7	293	37
Size class 2	0	0	0	0	0	0	0	0
Rodentia								
Size class 0	0	0	0	0	0	0	0	0
Size class 1	25	4	39	6	207	26	271	36
Size class 2	2	1	0	0	0	0	2	1
Chiroptera								
Size class 0	0	0	0	0	0	0	0	0
Size class 1	103	17	116	16	21	4	240	37
Size class 2	0	0	0	0	0	0	0	0
Observer 2								
Aves								
Size class 0	107	13	70	9	120	14	297	36
Size class 1	79	10	136	15	172	19	387	44
Size class 2	0	0	0	0	3	1	3	1
Squamata								
Size class 0	0	0	0	0	0	0	0	0
Size class 1	67	9	145	19	37	8	249	36
Size class 2	0	0	0	0	0	0	0	0
Rodentia								
Size class 0	0	0	0	0	0	0	0	0
Size class 1	13	2	35	4	194	25	242	31
Size class 2	14	3	4	1	14	4	32	8
Chiroptera								
Size class 0	0	0	0	0	0	0	0	0
Size class 1	101	19	118	15	21	6	240	40
Size class 2	0	0	0	0	0	0	0	0
Total (mean)								
Aves								
Size class 0	107,0	13,0	70,0	9,0	120,0	14,0	297,0	36,0
Size class 1	79,0	10,0	136,0	15,0	172,0	19,0	387,0	44,0
Size class 2	0,0	0,0	0,0	0,0	3,0	1,0	3,0	1,0
Squamata								
Size class 0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0
Size class 1	75,5	9,5	159,0	19,5	36,5	7,5	271,0	36,5
Size class 2	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0
Rodentia								
Size class 0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0
Size class 1	19,0	3,0	37,0	5,0	200,5	25,5	256,5	33,5
Size class 2	8,0	2,0	2,0	0,5	7,0	2,0	17,0	4,5

Chiroptera								
Size class 0	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0
Size class 1	102,0	18,0	117,0	15,5	21,0	5,0	240,0	38,5
Size class 2	0,0	0,0	0,0	0,0	0,0	0,0	0,0	0,0

Table S3. Detailed data on the percentage of anatomical representation (PR) for each faunal group, by site and by observer. For Aves, observers 1 and 2 made counts and determinations together, and this explains why there is no difference between the two datasets.

	Grand Bay	Salisbury	Canefield	TOTAL
	PR	PR	PR	PR
Observer 1				
Aves				
Mandible	78.3	91.7	100.0	91.4
Humerus	100.0	97.9	98.5	98.8
Carpometacarpus	78.3	83.3	70.6	76.5
Femur	80.4	83.3	89.7	85.2
Tarsometatarsus	63.0	75.0	80.9	74.1
Squamata				
Dentary	55.0	95.0	92.9	83.8
Humerus	90.0	97.5	28.6	82.4
Ulna	100.0	67.5	28.6	68.9
Femur	90.0	87.5	64.3	83.8
Tibia	85.0	85.0	42.9	77.0
Rodentia				
Mandible	80.0	83.3	75.0	77.0
Humerus	60.0	66.7	73.1	70.3
Ulna	60.0	58.3	75.0	70.3
Femur	40.0	66.7	92.3	81.1
Tibia	40.0	66.7	88.5	78.4
Chiroptera				
Mandible	76.5	56.3	100.0	70.3
Humerus	55.9	81.3	75.0	68.9
Radius	50.0	81.3	25.0	60.8
Femur	61.8	75.0	50.0	66.2
Tibia	58.8	68.8	25.0	59.5
Observer 2				
Aves				
Mandible	78.3	91.7	100.0	91.4
Humerus	100.0	97.9	98.5	98.8
Carpometacarpus	78.3	83.3	70.6	76.5
Femur	80.4	83.3	89.7	85.2
Tarsometatarsus	63.0	75.0	80.9	74.1
Squamata				
Dentary	72.2	92.1	93.8	87.5
Humerus	100.0	97.4	31.3	83.3
Ulna	16.7	42.1	18.8	30.6
Femur	94.4	94.7	62.5	87.5
Tibia	88.9	55.3	25.0	56.9
Rodentia				
Mandible	80.0	100.0	78.0	81.4
Humerus	60.0	70.0	74.0	71.4
Ulna	60.0	70.0	80.0	75.7
Femur	40.0	70.0	94.0	82.9
Tibia	30.0	80.0	90.0	80.0
Chiroptera				
Mandible	68.4	66.7	66.7	67.5
Humerus	50.0	83.3	41.7	61.3

Radius	44.7	86.7	16.7	56.3
Femur	52.6	80.0	33.3	60.0
Tibia	50.0	76.7	16.7	55.0
Total (mean)				
Aves				
Mandible	78.3	91.7	100.0	91.4
Humerus	100.0	97.9	98.5	98.8
Carpometacarpus	78.3	83.3	70.6	76.5
Femur	80.4	83.3	89.7	85.2
Tarsometatarsus	63.0	75.0	80.9	74.1
Squamata				
Dentary	63.2	93.6	100.0	86.8
Humerus	94.7	97.4	32.1	84.0
Ulna	60.5	55.1	25.0	50.7
Femur	92.1	91.0	67.9	86.8
Tibia	86.8	70.5	35.7	68.1
Rodentia				
Mandible	80.0	90.9	76.5	79.2
Humerus	60.0	68.2	73.5	70.8
Ulna	60.0	63.6	77.5	72.9
Femur	40.0	68.2	93.1	81.9
Tibia	35.0	72.7	89.2	79.2
Chiroptera				
Mandible	72.2	61.3	80.0	67.9
Humerus	52.8	82.3	55.0	64.1
Radius	47.2	83.9	20.0	57.7
Femur	56.9	77.4	40.0	62.2
Tibia	54.2	72.6	20.0	56.4

Table S4. Ratios calculated with the considered anatomical elements for each faunal group and each site, all observers combined.

	Grand Bay	Salisbury	Canefield	TOTAL
Aves				
Cranial/Post-cranial	0.97	1.08	1.18	1.09
Anterior/Posterior	1.24	1.14	0.99	1.10
Stylopodia/Zeugopodia	1.28	1.14	1.24	1.22
Squamata				
Cranial/Post-cranial	0.76	1.19	2.49	1.20
Anterior/Posterior	0.87	0.94	0.55	0.87
Stylopodia/Zeugopodia	1.27	1.50	1.65	1.44
Rodentia				
Cranial/Post-cranial	1.64	1.33	0.92	1.04
Anterior/Posterior	1.05	0.94	0.83	0.89
Stylopodia/Zeugopodia	1.05	1.00	1.00	1.00
Chiroptera				
Cranial/Post-cranial	1.37	0.78	2.37	1.13
Anterior/Posterior	0.90	1.11	1.25	1.03
Stylopodia/Zeugopodia	1.08	1.04	2.38	1.11

Table S5. Detailed data on the Number of Identified Specimens (NISP) fragmented and Percentage of Fragmentation (PF) for each faunal group, by site and by observer.

	Humerus NISP	Carpo/Ul/Rad NISP	Femur NISP	Tarso/Tib NISP	TOTAL NISP
Observer 1					
Aves					
(0)	143	115	126	107	491
(1)	5	0	1	0	6
(2)	5	8	6	10	29
(3)	0	0	0	0	0
(4)	7	1	5	3	16
(5)	0	0	0	0	0
PF (%)	*10.6*	*7.3*	*8.7*	*10.8*	*9.4*
Squamata					
(0)	53	46	50	48	197
(1)	8	5	12	9	34
PF (%)	*13.1*	*9.8*	*19.4*	*15.8*	*14.7*
Rodentia					
(0)	46	50	57	53	206
(1)	1	0	0	0	1
(2)	1	1	0	1	3
(3)	0	0	0	0	0
(4)	1	0	0	0	1
(5)	0	0	0	0	0
PF (%)	*6.1*	*2.0*	*0.0*	*1.9*	*2.4*
Chiroptera					
(0)	40	36	43	41	160
(1)	2	1	3	1	7
(2)	4	2	0	0	6
(3)	0	4	0	0	4
(4)	3	1	3	0	7
(5)	0	1	0	0	1
PF (%)	*18.4*	*20.0*	*12.2*	*2.4*	*13.5*
Observer 2					
Aves					
(0)	143	118	125	105	491
(1)	6	0	0	0	6
(2)	4	5	7	10	26
(3)	0	0	0	1	1
(4)	7	1	6	3	17
(5)	0	0	0	1	1
PF (%)	*10.6*	*4.8*	*9.4*	*12.5*	*9.4*
Squamata					
(0)	57	21	60	39	177
(1)	3	1	3	2	9
PF (%)	*5.3*	*4.8*	*5.0*	*5.1*	*4.8*
Rodentia					
(0)	47	49	58	49	203
(1)	1	0	0	0	1
(2)	1	2	0	2	5
(3)	0	1	0	1	2
(4)	0	0	0	0	0

(5)	1	0	0	0	1
PF (%)	*6.0*	*5.8*	*0.0*	*5.8*	*4.2*
Chiroptera					
(0)	37	23	40	39	139
(1)	1	0	3	1	5
(2)	9	9	1	4	23
(3)	0	6	0	0	6
(4)	2	6	5	0	13
(5)	0	1	0	0	1
PF (%)	*24.5*	*48.9*	*18.4*	*11.4*	*25.7*
Total (mean)					
Aves					
(0)	143.0	116.5	125.5	106.0	491.0
(1)	5.5	0.0	0.5	0.0	6.0
(2)	4.5	6.5	6.5	10.0	27.5
(3)	0.0	0.0	0.0	0.5	0.5
(4)	7.0	1.0	5.5	3.0	16.5
(5)	0.0	0.0	0.0	0.5	0.5
PF (%)	*10.6*	*6.0*	*9.1*	*11.7*	*9.4*
Squamata					
(0)	55.0	33.5	55.0	43.5	187.0
(1)	5.5	3.0	7.5	5.5	21.5
PF (%)	*9.1*	*8.2*	*12.0*	*11.2*	*10.3*
Rodentia					
(0)	46.5	49.5	57.5	51.0	204.5
(1)	1.0	0.0	0.0	0.0	1.0
(2)	1.0	1.5	0.0	1.5	4.0
(3)	0.0	0.5	0.0	0.5	1.0
(4)	0.5	0.0	0.0	0.0	0.5
(5)	0.5	0.0	0.0	0.0	0.5
PF (%)	*6.1*	*3.9*	*0.0*	*3.8*	*3.3*
Chiroptera					
(0)	38.5	29.5	41.5	40.0	149.5
(1)	1.5	0.5	3.0	1.0	6.0
(2)	6.5	5.5	0.5	2.0	14.5
(3)	0.0	5.0	0.0	0.0	5.0
(4)	2.5	3.5	4.0	0.0	10.0
(5)	0.0	1.0	0.0	0.0	1.0
PF (%)	*21.4*	*34.4*	*15.3*	*7.0*	*19.6*

(0) complete bone, (1) only proximal part, (2) proximal part + shaft, (3) only shaft, (4) distal part + shaft, (5) only distal part, except for Squamates for which counts were only made as "not fragmented" (0) or "fragmented" (1).

Table S6. Detailed data on the Percentage of Digestion (PD) for each faunal group, by anatomical element and by observer. For Squamates, *Thecadactylus rapicauda* and *Anolis* sp. were separated, and total values are also given. For this group, proximal and distal parts were not considered separately. For Rodentia, distal unla was not considered for digestion.

	Md/ Dent	**Humerus**		**Carpo./Ulna/ Radius**		**Femur**		**Tarso./Tibia**		**TOTAL**
	NISP	**NISP prox.**	**NISP dist.**	**NISP prox.**	**NISP dist.**	**NISP prox.**	**NISP dist.**	**NISP prox.**	**NISP dist.**	**NISP**
Observer 1										
Aves										
0	25	101	91	52	76	96	69	49	72	631
1	45	39	31	44	23	24	27	47	21	301
2	3	5	22	19	14	6	22	8	11	110
3	0	4	5	6	3	4	11	8	3	44
PD (%)	*65.8*	*32.2*	*38.9*	*57.0*	*34.5*	*26.2*	*46.5*	*56.3*	*32.7*	*41.9*
Thecadactylus rapicauda										
0	8	6		4		5		5		28
1	9	3		13		6		9		40
2	2	6		5		4		5		22
3	0	7		1		9		3		20
PD (%)	*57.9*	*72.7*		*82.6*		*79.2*		*77.3*		*74.5*
***Anolis* sp.**										
0	19	23		18		17		21		98
1	9	9		6		10		7		41
2	12	4		4		8		5		33
3	3	3		0		3		2		11
PD (%)	*55.8*	*41.0*		*35.7*		*55.3*		*40.0*		*46.4*
Squamata										
0	27	29		22		22		26		126
1	18	12		19		16		16		81
2	14	10		9		12		10		55
3	3	10		1		12		5		31
PD (%)	*56.5*	*52.5*		*56.9*		*64.5*		*54.4*		*57.0*
Rodentia										
0	28	21	25	15	/	29	21	27	28	206
1	24	24	19	26	/	19	33	25	23	222
2	0	2	1	8	/	4	3	1	2	22
3	2	2	3	10	/	5	3	1	3	31
PD (%)	*48.1*	*57.1*	*47.9*	*74.6*	/	*49.1*	*65.0*	*50.0*	*50.0*	*57.2*
Chiroptera										
0	28	31	10	15	11	35	24	32	30	216
1	18	10	6	3	5	6	8	5	4	65
2	2	0	7	3	3	3	6	5	1	30
3	2	2	14	9	9	4	10	7	4	61
PD (%)	*44.0*	*27.9*	*73.0*	*50.0*	*60.7*	*27.1*	*50.0*	*34.7*	*23.1*	*41.9*
Observer 2										
Aves										
0	25	94	83	58	82	100	63	47	75	627
1	36	43	33	36	16	17	29	43	10	263
2	11	8	28	21	11	10	29	18	16	152
3	1	4	3	6	8	3	7	3	4	39
PD (%)	*65.8*	*36.9*	*43.5*	*52.1*	*29.9*	*23.1*	*50.8*	*57.7*	*28.6*	*42.0*

Thecadactylus rapicauda										
0	9	5		1		3		10		28
1	10	3		2		7		7		29
2	0	11		0		8		4		23
3	0	3		0		7		1		11
PD (%)	*52.6*	*77.3*		*66.7*		*88.0*		*54.5*		*69.2*
***Anolis* sp.**										
0	21	14		16		14		12		77
1	12	18		2		14		4		50
2	8	1		0		2		0		11
3	3	5		1		10		3		22
PD (%)	*52.3*	*63.2*		*15.8*		*65.0*		*36.8*		*51.9*
Squamata										
0	30	19		17		17		22		105
1	22	21		4		21		11		79
2	8	12		0		10		4		34
3	3	8		1		17		4		33
PD (%)	*52.4*	*68.3*		*22.7*		*73.8*		*46.3*		*58.2*
Rodentia										
0	43	14	27	15	/	9	13	17	19	157
1	13	9	16	30	/	2	23	14	3	110
2	0	4	1	7	/	0	4	2	0	18
3	0	1	0	0	/	0	0	0	1	2
PD (%)	*23.2*	*50.0*	*38.6*	*71.2*	*/*	*18.2*	*67.5*	*48.5*	*17.4*	*45.3*
Chiroptera										
0	33	20	4	15	9	31	19	26	24	181
1	20	16	18	8	10	8	15	14	11	120
2	6	2	11	5	5	3	7	3	3	45
3	0	5	5	4	6	0	1	0	0	21
PD (%)	*44.1*	*53.5*	*89.5*	*53.1*	*70.0*	*26.2*	*54.8*	*39.5*	*36.8*	*50.7*
Total (mean)										
Aves										
0	25.0	97.5	87.0	55.0	79.0	98.0	66.0	48.0	73.5	629.0
1	40.5	41.0	32.0	40.0	19.5	20.5	28.0	45.0	15.5	282.0
2	7.0	6.5	25.0	20.0	12.5	8.0	25.5	13.0	13.5	131.0
3	0.5	4.0	4.0	6.0	5.5	3.5	9.0	5.5	3.5	41.5
PD (%)	*65.8*	*34.6*	*41.2*	*54.5*	*32.2*	*24.6*	*48.6*	*57.0*	*30.7*	*41.9*
Thecadactylus rapicauda										
0	8.5	5.5		2.5		4.0		7.5		28.0
1	9.5	3.0		7.5		6.5		8.0		34.5
2	1.0	8.5		2.5		6.0		4.5		22.5
3	0.0	5.0		0.5		8.0		2.0		15.5
PD (%)	*55.3*	*75.0*		*80.8*		*83.7*		*65.9*		*72.1*
***Anolis* sp.**										
0	20.0	18.5		17.0		15.5		16.5		87.5
1	10.5	13.5		4.0		12.0		5.5		45.5
2	10.0	2.5		2.0		5.0		2.5		22.0
3	3.0	4.0		0.5		6.5		2.5		16.5
PD (%)	*54.0*	*51.9*		*27.7*		*60.3*		*38.9*		*49.0*
Squamata										
0	28.5	24.0		19.5		19.5		24.0		115.5
1	20.0	16.5		11.5		18.5		13.5		80.0
2	11.0	11.0		4.5		11.0		7.0		44.5

3	3.0	9.0	1.0	14.5	4.5	32.0				
PD (%)	*54.4*	*60.3*	*46.6*	*69.3*	*51.0*	*57.5*				
Rodentia										
0	35.5	17.5	26.0	15.0	/	19.0	17.0	22.0	23.5	175.5
1	18.5	16.5	17.5	28.0	/	10.5	28.0	19.5	13.0	151.5
2	0.0	3.0	1.0	7.5	/	2.0	3.5	1.5	1.0	19.5
3	1.0	1.5	1.5	5.0	/	2.5	1.5	0.5	2.0	15.5
PD (%)	*35.5*	*54.5*	*43.5*	*73.0*	/	*44.1*	*66.0*	*49.4*	*40.5*	*51.5*
Chiroptera										
0	30.5	25.5	7.0	15.0	10.0	33.0	21.5	29.0	27.0	198.5
1	19.0	13.0	12.0	5.5	7.5	7.0	11.5	9.5	7.5	92.5
2	4.0	1.0	9.0	4.0	4.0	3.0	6.5	4.0	2.0	37.5
3	1.0	3.5	9.5	6.5	7.5	2.0	5.5	3.5	2.0	41.0
PD (%)	*44.0*	*40.7*	*81.3*	*51.6*	*65.5*	*26.7*	*52.2*	*37.0*	*29.9*	*46.3*