

Article



Assessment of Fruit and Vegetable Residues Suitable for Renewable Energy Production: GIS-Based Model for Developing New Frontiers within the Context of Circular Economy

Roberta Selvaggi and Francesca Valenti *

Department of Agriculture, Food and Environment, University of Catania, Via Santa Sofia, 95123 Catania, Italy; roberta.selvaggi@unict.it

* Correspondence: francesca.valenti@unict.it

Abstract: Due to the necessity of developing renewable energy sources, the anaerobic digestion for producing biomethane has developed significantly in the last years, since it allows to both reduce disposal treatment and produce green energy. In this field, fruit and vegetable wastes have been recently put forward, since they could represent a suitable resource for producing biomethane as a new frontier within the context of a circular economy. This study aims at filling the gap in the knowledge of the production, quantities and biogas potential production of these residues. On this basis, a GIS-based model was developed and applied to the Sicily region by investigating the specific regulatory framework as well as by analysing descriptive statistics. The results of the GIS analyses enabled the localisation of the highest productive territorial areas and highlighted where fruit and vegetable wastes are abundantly located. In this regard, about 7 million Nm³ of biogas could be produced by reusing only the fruit and vegetable residues coming from the three most representative Sicilian wholesale markets among those considered. Finally, the regulatory framework is of crucial importance in inhibiting or supporting the use of the selected biomass in a specific sector, with regard to the case study considered.

Keywords: renewable energy; agricultural waste; recycling; biomethane; GIS; circular economy

1. Introduction

Nowadays, climate change is the most imminent environmental issue facing the world. Rising global temperatures will have major effects on human life, on food chains, ecosystems and wildlife. It is well known that global warming is due to the large-scale anthropogenic emission of greenhouse gases (GHG). The usage of renewable energy sources to produce electricity and heat could reduce the negative impact of fossil fuels on GHG released into the atmosphere [1,2]. So renewable resources could play a key role in current CO₂-mitigation policies [3]. In this context, biomasses are one of the most important renewable energy sources, also because they can guarantee a continuous power generation, unlike other discontinuous energy sources (i.e., wind or solar energy). To contribute to reaching the 2020 EU goals, among the different technologies and processes anaerobic digestion (AD) could be a suitable way to produce renewable energy [4,5]. In fact, in accordance with the environmental and sustainability policies adopted by European Member States, recently there has been an increase in the spread of this technological process.

AD is a biochemical degradation process that presents a double benefit, as it produces biogas and meanwhile treats biomass, especially agricultural residues, and agro-industrial by-products, reducing their disposal in landfills [6,7].

Citation: Selvaggi, R.; Valenti, F. Assessment of Fruit and Vegetable Residues Suitable for Renewable Energy Production: GIS-Based Model for Developing New Frontiers within the Context of Circular Economy. *Appl. Syst. Innov.* **2021**, *4*, 10. https://doi.org/10.3390/ asi4010010

Received: 7 January 2021 Accepted: 1 February 2021 Published: 3 February 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses /by/4.0/). Currently, AD plants are encouraged by the Indirect Land Use Change (ILUC) policy to produce advanced biofuels from feedstock that do not compete directly with food and feed crops, such as wastes and agricultural residues, non-food crops and algae. In detail, waste materials like sewage sludge, livestock manure, crop residues, the OFMSW (organic fraction of municipal solid waste) and fruit and vegetable waste are very important since they do not compete with food and feed crops as biomasses for the AD process [3,8]. In this regard, there are several types of waste creation areas in societies, for instance, farm-level waste [9], waste in construction sites [10], household-level solid waste [11], OFMSW [12] and vegetable and fruit waste [13]. Among these, fruit and vegetable wastes (FVWs) are a very important category of biomasses because they are produced in considerable quantities at the end of all agricultural, supermarket and wholesale market activites, and their landfill disposal is quite difficult due to their very high perishability [14,15].

Fruit and vegetable wastes contain a high percentage of organic matter that can be converted into biogas with a high yield. The presence of key micro, macro and trace elements allows the process to proceed without the addition of additives or other organic materials such as co-substrates [16,17]. It is important that fruits and vegetables are mixed in appropriate proportions to balance the excess or deficiency of nutrients and other inhibiting substances, or to limit the input of undesirable substances such as sulfur [18].

According to the estimation of the United Nations Food and Agriculture Organization (FAO), the waste generated by the fruit and vegetable sector is estimated to represent up to 60% of the food waste generated yearly in the world [19,20], and this estimation was reported to be much higher in Italy (about 87%) [21].

FVW generation increases the operating costs of markets due to the reduction of revenue and the increase of disposal costs [22]. In detail, FVWs represent a significant economic issue for companies and pose environmental problems due to their high moisture content and biodegradability [23]. According to the Goal 12.3 of the Sustainable Development Goals (SDGs) identified in 2015 by ONU, sustainable production and consumption is one advantage of reducing food wastes. Many governments and organisations around the world have committed to reducing food loss and waste. The valorisation of reusing them as renewable energy source meets the indication of Directive 2008/98/EC of the European Parliament, which establishes, following a hierarchy of residues, that the residues must be sent to disposal in sanitary landfills if they cannot be reused, recycled or recovered [24,25].

For these reasons, in recent years, large efforts have been dedicated to the development of actions and policies for FVW management [26]. So FVW valorisation processes concerning value-added derivatives have been deeply analysed in recent years. Despite the intense research activities, the current main destination of FVWs is still landfill disposal, although researchers agree that they can be use as feedstocks to produce renewable energy and fertilisers. Moreover, this could represent a suitable and more interesting alternative to both reduce environmental impact and produce clean energy within a green economy perspective [27]. In this regard, the valorisation of FVW is at an initial stage of development and more essential elements must still be assessed to verify its feasibility [28,29]. Firstly, data on the exact amount of both wastes obtained by agro-industrial companies, food processing and fruit and vegetable residues form markets is nowadays very limited. In fact, no official information is provided by the EU, given the commercial sensitivity of this data for all the producers.

This lack of official data related to the amount of waste, in terms of volume, and to the spatial localisation of the sites where these wastes are produced, is an important factor that has limited the reuse, exploitation, and valorisation of the wastes for sustainable processes.

Moreover, the demand for renewable and sustainable resources (e.g., green energy, water, investment for equipment and workforce) and for innovative valorisation strategies as compared to that of traditional waste management options should be evaluated. In fact, as reported by Valenti et al. [30] the implementation of new strategies is viable only if it

brings economic (e.g., reduction of disposal costs and increase of incomes from the valorisation of the by-products/waste) and environmental advantages (e.g., reduction of waste amount for the landfill and of emission) as compared to conventional disposal strategies.

Nonetheless, logistic-related problems make these residues difficult to valorise [31]. In this regard, geographical information system (GIS) tools have been intensively used to perform analyses of feedstock supply and logistics [32–35]. While fruit and vegetable processing side streams generated from agro-industrial chains have been evaluated for the production of enzymes, organic acids (e.g., succinic and lactic acid) and renewable energy [36,37], whole fruit and vegetables from open markets have not been considered as potential feedstocks by evaluating the high amounts disposed of in landfill as waste, as reported by Liakou et al. [38].

Given the uncertainty of data relating to biomass amounts, this research aims, by using GIS tools, to fill the gap in the knowledge of the production, localisation and yield of fruit and vegetable residues to support producers in waste valorisation actions, i.e., for energy uses [22,32,39,40]; to revisit regulations to facilitate their use is urgently needed.

This study investigates the suitability of using FVWs as a possible feedstock of the blend for the diet of the anaerobic digestion plants where the biogas sector is still developing and biomethane is a real and viable possibility. The production of energy is only one of the possible destinations of these wastes, but it represents the most reasonable. In fact, in the study area there are no farms or other possible outlets for these by-products, if not landfills.

Therefore, the collected data influenced the actual and real possibility of recycling fruit and vegetable residues. This explorative study is important for planning the valorisation of new biomasses that could be used as feedstock for the existing biogas plants or for developing new ones in a sustainable way. In fact, their use can have a positive impact with regard to environmental protection, and especially in terms of reduction of both GHG emissions from these biomasses logistic supplies and soil consumption for dedicated energy crops.

2. Materials and Methods

A methodology for assessing the suitability of using new biomasses suitable as a feedstock to produce biogas is described. In this regard, statistical tools were adopted, and a GIS analysis was carried out. In detail, the QGis software (ver. 3.10.11), an open-source GIS software, was used since it is a decision support tool appropriate for collecting, organising, analysing and localising geographical data.

Firstly, by adopting statistical tools, the agricultural sector and the production were analysed in order to find the trend of production and the cultivated area for the Sicilian horticultural sector from 2015 to 2019. Data were acquired from the national surveys database [41]. Then, by using spatial analysis GIS tools, the production areas were localised in the study area, and the most productive areas were considered in order to deeply investigate the wholesale market flows. A specific methodology was developed in order to quantify the processed fruit and vegetables, the obtained waste production and, therefore, the potential biogas production. In detail, FVW production was computed by using specific face-to-face surveys at the wholesale markets and data available from annual statistic tools. After obtaining specific authorisation to process the data, from the managers of the food markets, for each selected market (Catania, Ragusa and Palermo) eight pit owners were interviewed. They represent about 30% of the total pit owners for each considered wholesale market. This quantity was imposed on us by the wholesale market managers, due to the Covid-19 emergency that made further pit visits impossible. The questionnaire was very simple, with only one section. We noted the amount in transit through the pits in the various years (from 2015 to 2019) in terms of product input and biomass disposed of (net of plastics and other packaging). So we obtained an index related to the quantities of wastes and we used this index referring to the total amount of biomass disposed of by the wholesale market (data collected by interviewing the wholesale manager).

Finally, the amount of potential biogas producible was estimated according to the quantity of FVWs. In the following subsections are detailed the methodological approach and the materials utilised.

2.1. Case Study

Sicily is an autonomous region of Italy and the largest island in the Mediterranean Sea (Figure 1), covering approximately 26,000 km². It has a unique climate pattern characterised by hot/dry summers and mild/wet winters. These conditions are beneficial to renewable energy production using agro-ingustrial residues obtained from typical activities: Cultivation first of all of citrus, and grape, olive oil, wheat and other products, and sheep and cattle breeding. Therefore, it was chosen as the study area.

In Sicily, agriculture has a decisive position in the regional economy. Like in the other parts of southern Italy, employees in agriculture are plentiful but they have low wages.

Furthermore, in Sicily, about 3.9 million metric tons of matrix are yearly produced by agro-industrial activities and are available and potentially reusable in biogas plants. These biomasses include byproducts from the agro-industrial chains (i.e., whey, citrus pulps, olive pomace residues and others), wastes from livestock (i.e., cattle manure), agricultural crop residues (i.e., cereal straw and others field and process residues) and some dedicated feedstocks obtained from energy crops. Among them, agro-industrial biomasses and livestock manure represent more than 60% of the total biomasses produced [42].



Figure 1. Geographic position of the study area.

Sicily is the Italian region with the greatest extension of agricultural land [42]. Considering the area dedicated to agriculture, Sicily excels in cereal production and is the first Italian producer of oranges with half (52%) of the entire national output.

By elaborating on data from the Italian National Statistical Institute (Istat) database it was found that, in the last five years (from 2015 to 2019), with reference to the horticultural production area (both horticultural open field crops and greenhouses), in Italy an average of 356,089 ha/year were cultivated, while in Sicily about 58,357 ha/year (about 16%) were cultivated [41]. Table 1 shows the trend of the analysed period and the mean values, subdivided for the specific cultivation of horticultural open field crops and greenhouses ones, for the nine provinces of the study area.

		Cultivated Area (ha)				
	2015	2016	2017	2018	2019	Mean (ha)
Agrigento						
-Horticultural open field crops	17,519	17,449	17,358	17,898	18,557	17,756
	100.0	99.6	99.1	102.2	105.9	
-Horticultural greenhouses	346	342	329	329	329	335
	100.0	98.9	95.1	95.0	95.1	
Caltanissetta	10 102	10 221	0079	0107	0010	0707
-Horticultural open field crops	10,102	10,331	9978	9107	9019	9707
-Horticultural greenhouses	830	102.5 830	90.0 879	90.2 829	870	879
	100.0	100.0	99.9	99.9	99.9	029
Catania	100.0	100.0	,,,,	,,,,	,,,,	
	3846	3847	3232	3847	3849	3724
-Horticultural open field crops	100.0	100.0	84.0	100.0	100.1	
	15	0.2	0.2	0.2	0.2	3
-Horticultural greenhouses	100.0	1.0	1.0	1.0	1.0	
Enna						
Harticultural open field grops	560	560	560	560	560	560
Horicultural open field crops	100.0	100.0	100.0	100.0	100.0	
-Horticultural greenhouses	11	9.0	9.0	9.0	9.0	9
	100.0	85.7	85.7	85.7	85.7	
Messina	1212	1 = 0 2	1504	4 24 2	4 804	1 (20
-Horticultural open field crops	1243	1703	1706	1717	1781	1630
	100.0	137.0	137.2	138.1	143.3	10
-Horticultural greenhouses	100.0	0.9	0.9	0.9	0.9	12
Palermo	100.0	1.0	1.0	1.0	1.0	
	7185	7143	7663	7794	8015	7560
-Horticultural open field crops	100.0	99.4	106.7	108.5	111.6	
	61	58.3	58.3	57.8	58.9	59
-Horticultural greenhouses	100.0	96.3	96.3	95.5	97.4	
Ragusa						
-Horticultural open field crops	1835	1985	2095	2225	2170	2062
	100.0	108.2	114.2	121.3	118.3	
-Horticultural greenhouses	4470	4380.0	3970.0	3960.0	3925.0	4141
	100.0	98.0	88.8	88.6	87.8	
Syracuse	40.42	E 40E	E 4 5 0	5440	5 00 0	
-Horticultural open field crops	4943	5425	5453	5662	5802	5457
	100.0	109.8	110.3 1109 E	114.5	117.4	1100
-Horticultural greenhouses	1135 100 0	99 N	97 7	1093.0 96 3	1083.1 95 /	1109
Tranani	100.0	<i></i>)1.1	70.5	7 J. 4	
1. 1. 1. p. 1. 1. t.	2690	2315	2305	2195	1965	2294
-Horticultural open field crops	100.0	86.1	85.7	81.6	73.0	/ 1
-Horticultural greenhouses	1104	1094.0	1094.0	1134.0	1114.0	1108
	100.0	99.1	99.1	102.7	100.9	

Table 1. Sicilian cultivated area for horticultural crops (ha) and trend of evolution (%).

Within the selected interval, overall, as shown in Table 1, the areas cultivated with horticultural crops registered a slight increase (+1.9%) by rising from 57,949 ha to 59,069 ha.

Instead of, by considering greenhouse horticulture, leaving out the data for the provinces of Catania and Messina, which are outliers, there is a common trend across the region: Greenhouse horticulture areas decreased from 2015 to 2019. Only the province of Trapani recorded a slight increase in greenhouse cultivated areas: From 1104 ha in 2015 to 1114 ha in 2019.

Regarding open field horticultural areas, however, there is no common generalised trend: Some provinces have recorded a contraction of areas while others have recorded a marked increase (first of all Messina).

Greenhouse cultivation is mainly concentrated in the province of Ragusa, in the coastal area. More than 56% of the Sicilian greenhouse surface is located in this territorial area.

As regards greenhouse cultivation, the most common species are, in this order: Tomato, zucchini, bell pepper, eggplant and watermelon; whereas, for open field production, the most cultivated species are, in this order: Artichoke, tomato, melon, cauliflower and lettuce.

In Table 2 the production obtained in these cultivated areas, detailed for each Sicilian province, is shown. The trend for the period from 2015 to 2019 and the average values, by providing the detail of greenhouse and field crops, were reported.

	Production (t)					
	2015	2016	2017	2018	2019	Mean (t)
Agrigento						
-Horticultural open field crops	322,937	302,870	300,227	307,831	316,093	309,991
	100.0	93.8	93.0	95.3	97.9	
-Horticultural greenhouses	21,932	14,449	12,826	13,017	13,093	15,064
2 14 14 14	100.0	65.9	58.5	59.4	59.7	
Caltanissetta	00.204	114.050			F1 202	07.000
-Horticultural open field crops	99,284	114,852	115,566	85,641	71,303	97,329
1 1	100.0	115.7	116.4	86.3	71.8	41 700
-Horticultural greenhouses	41,725	41,725	41,698	41,696	41,696	41,708
Catavia	100.0	100.0	99.9	99.9	99.9	
Cutuniu	59 667	59 685	47 401	59 846	60 039	57 328
-Horticultural open field crops	100.0	100.0	79.4	100 3	100.6	57,520
	750	750	750	750	750	750
-Horticultural greenhouses	100.0	100.0	100.0	100.0	100.0	700
Enna	10010	10010	10010	10010	10010	
11114	7265	7970	8550	7750	7535	7814
-Horticultural open field crops	100.0	109.7	117.7	106.7	103.7	-
	695	1265	1265	1265	1265	1151
-Horticultural greenhouses	100.0	182.1	182.1	182.1	182.1	
Messina						
	18,290	36,379	36,555	37,055	37,445	33,145
-Horticultural open field crops	100.0	198.9	199.9	202.6	204.7	
Hartigultural groophousoo	3181	1705	1818	1818	1818	7794
-Horticultural greenhouses	100.0	5.4	5.7	5.7	5.7	
Palermo						
-Horticultural open field crops	85,253	84,611	106,318	106,939	109,989	98,622
-Horneuntariai open neia erops	100.0	99.2	124.7	125.4	129.0	
-Horticultural greenhouses	20,173	19,664	19,666	19,736	20,051	19,858
	100.0	97.5	97.5	97.8	99.4	
Ragusa						
-Horticultural open field crops	52,057	52,640	71,350	82,100	83,300	68,289
	100.0	101.1	137.1	157.7	160.0	
-Horticultural greenhouses	239,300	232,300	209,250	220,300	243,000	228,830
	100.0	97.1	87.4	92.1	101.5	
Syracuse	111.005	100 (14	100.000	104 (1)	105 501	
-Horticultural open field crops	111,035	122,614	123,920	124,616	125,591	121,555
1 1	100.0	110.4	111.6	112.2	113.1	15 000
-Horticultural greenhouses	45,240	44,224	45,320	44,815	45,412	45,002
Tranani	100.0	97.8	100.2	99.1	100.4	
тирині	22 158	27 120	25 280	2/ 200	21 000	26 250
-Horticultural open field crops	32,438 100 0	27,120 82.6	20,000 70 0	24,390 7E 1	∠1,700 67 ⊑	20,200
	0704	03.0	70.Z	10 474	10 424	10.014
-Horticultural greenhouses	7704 100 0	7704 100 0	9704 100 5	10,4/4 107 0	10,434	10,014
	100.0	100.0	100.5	107.9	107.5	

Table 2. Sicilian production of horticultural crops (t) and trend of evolution (%).

Concerning open field horticultural areas, and also horticultural greenhouse production, Messina data are outliers and cannot be considered.

Regarding horticultural greenhouse production, there is a common trend across the region: Greenhouse horticulture productions are constant from 2015 to 2019. Only the province of Enna recorded a doubled production: From 695 t in 2015 to 1265 t in 2019, with the same cultivated area of about 10 ha (as shown in Table 1)

Regarding open field horticultural production, however, there is no common generalised trend: Trapani, Caltanissetta and Agrigento have recorded a contraction of areas, while others have recorded a marked increase (first of all Messina and Ragusa). In these areas there has been an intensification of cultivation processes and a birth of numerous new companies.

Furthermore, by analysing the available data from the ISTAT database, it was found that the most productive horticultural greenhouse species are, in order, tomato, zucchini, eggplant, bell pepper and watermelon, while the most productive horticultural open field crop species are, in order, tomato, melon, artichoke, cauliflower and lettuce.

As for the surface area, Ragusa is also the province in which there is the greatest production (about 60.9% of total Sicilian production).

2.2. Data Analysis

In this study, an extensive database was improved according to statistical sources, i.e., ISTAT (years 2015–2019), to quantify horticultural production by GIS analysis.

The base maps used in the GIS included the Regional Technical Map (RTM 2008) as the base map for producing thematic maps. RTM 2008 is an upgrade of previous versions of the RTM 2005 numerical edition (sites CDE), the 2001 edition (sites 7-8-9), the 2004 edition (site B) and the 2003 edition (site A). RTM 2008 was created by using the digital colour ortho-photos ATA0708 with a geometrical resolution of 25 cm × 25 cm. All administrative boundaries adopted for map elaboration were taken from RTM 2008 layer I.

By analysing available data on the agricultural sector, the provinces with the highest volumes produced were identified and localised on GIS software. Then, by using QGIS software v.3.10.11, the overlay of data contained in the base maps and the use of the Jenks tool in the QGIS software produced an arrangement of the selected horticultural producing areas based on the maximisation of variance among the classes. The next step of the methodology was the analysis of the quantity of FVWs produced from the selected wholesale market.

In detail the amount of the produced waste was analysed to identify and to survey local wholesale markets located in the study area.

To quantify the production and obtain the residual amount, which could be considered waste production, the biggest markets sited in the study area were sampled. A questionnaire was given to each pit manager involved in the research to find out the production targets, the quantity of the products entering the pit and the amount of residual products coming out of the pit, especially for those wastes usable for the energy process.

The collected data was anonymously elaborated to specifically estimate the amounts of residual products and where they are available (spatial location) in the study area.

The last step of the study was to assess the potential production of biogas obtainable by the amount of estimated residual products. The theoretical biogas potential (Btot) was computed by using the formula (Equation (1)):

$$B_{\rm tot} = A \cdot Y \tag{1}$$

where A is the amount of FVWs (expressed as tons of fresh matter) and Y is the FVW biogas potential yield, which was considered equal to $158.1 \pm 18.7 \text{ Nm}^3$ /t of fresh matter,

as reported by Adani et al. [43], who adopted the simple and fast tests to predict potential biogas developed by Schievano et al. [44].

3. Results

Horticultural producing areas acquired by the ISTAT database were deeply analyzed for the different provinces by providing a GIS map to show their territorial distribution. In detail, ISTAT data were used to localise, on a territorial level, the distribution of cultivated areas dedicated to horticultural production. By adopting the Jenks tool, available in the QGis software, the Sicilian provinces were grouped in three different classes in order to maximise the differences in the cultivated areas. As a result, the areas that showed the highest horticultural surface areas were the provinces of Palermo, Agrigento and Caltanissetta, as shown in Figure 2.



Figure 2. Distribution of horticultural cultivated area within the Sicily region (ISTAT, from 2015 to 2019).

Then, the horticultural cultivated areas were subdivided into horticultural open field crops and horticultural greenhouses (Table 1) by showing different distributions within the Sicilian provinces, as reported in Figure 3a,b.



Figure 3. (a) Distribution of horticultural open field crop cultivated areas. (b) Distribution of horticultural greenhouse cultivated areas (ISTAT from 2015 to 2019).

The distribution of the horticultural cultivated areas allows the identification of two different areas for horticultural production located in the south-west and in the south-east of the island, respectively, for open field crops and greenhouses.

The second step was the analysis of the horticultural production, since the most productive areas could not correspond to the most cultivated areas.

The next step was focussed on wholesale market localisation. After a direct interview with the regional coordinator of wholesales market activities, the Sicilian active wholesale markets were located within Sicily region by considering their GPS coordinates, and a GIS map was made as shown in Figure 4.



Figure 4. Wholesale market localisation within Sicily region (data obtained by direct interviews).

In order to quantify wholesale market residues and waste production, firstly all the wholesale market data (i.e., production flows) were analysed based on the available database, and by using the GIS tool (i.e., Heatmap plugin) a heat map of their production was made (Figure 5).



Figure 5. Wholesale markets heat map distribution based on their flow production.

As reported in Figure 5, firstly (in orange) all the wholesale markets data were elaborated for the production of a heat map; then, among them, the wholesale markets with the biggest flow production were selected (in pink), and their data were elaborated by adopting again the heat map tool. An overlay of the two produced heat maps was carried out in order to show the high difference of flow production (Figure 5).

In detail, the largest and most important wholesale markets, sited in the provinces of Catania, Ragusa, and Palermo, were selected as a sample for carrying out a face-to-face survey analysis with the aim of estimating the quantity of FVWs.

The surveyed data are summarised in Table 3 in order to define an average percentage of potential horticultural wastes in the period between 2015 and 2019.

Site of the Wholesale Market ——	Incoming Products (Vegetables and Fruit)	Was	Wastes		
	(t)	(t)	(%)		
Catania	308,579	21,376	6.9%		
Vittoria	261,901	13,357	5.1%		
Palermo	183,500	11,744	6.4%		

Table 3. Quantities of estimated fruit and vegetable wastes (t).

As shown in Figure 6, by considering all the horticultural products, the provinces of Syracuse, Ragusa, Agrigento, Catania, Palermo and Caltanissetta were highlighted as the most productive ones. Furthermore, the results of the analysis of the production of the three selected biggest wholesale markets confirmed the results obtained with the ISTAT database on the horticultural production by highlighting three different areas (Figure 6).



Figure 6. Overlay result of the selected biggest wholesale markets and the horticultural production areas.

The last step of the study was the computation of the potential biogas production by recycling FVWs. As reported in Table 3, the FVWs were computed, and by using Equation (1) the potential biogas was estimated to be about 7 million of Nm³ [44]. In detail, as reported in Figure 7, the heat map with potential biogas production obtainable by the three selected wholesale markets was produced. Moreover, the potential production from each considered wholesale market amounts to about 3.3 million Nm³, 2.1 million Nm³ and 1.8 million Nm³, respectively, for the wholesale markets located in the municipalities of Catania, Vittoria and Palermo. As shown in Figure 7 the wholesale market located in Catania province is the biggest one, with the highest flow production and potential biogas production.



Figure 7. Estimation of biogas availability.

4. Discussion

Sicily and other southern Italian regions are well behind the rest of the country in terms of anaerobic digestion plants and the agro-energy chain in general [45]. The production potential of bio-methane in many parts of southern Italy is undeveloped and even unknown despite being potentially very high [46].

Furthermore, there is a great number of agro-industrial processing plants which could provide large quantities of "integrated biomasses" like citrus pulp and olive pomace, the last being absolutely appropriate for producing advanced biofuels [32–34]. In this context of green energy management, the possibility to reduce the wastes obtained by the wholesale market of fruits and vegetables is also a concrete option.

FVWs are produced in huge quantities by wholesale markets and represent an important category of residues that are difficult to dispose of because of their high perishability. They are easily degraded by the microbial bacteria with a speed dependent on the presence of mechanical damage or excessive ripening. This leads to significant environmental complications and high costs for fruit and vegetable markets both for disposal and for economic losses due to the lack of product sales. Anaerobic digestion may be the most appropriate technology for the disposal and energy valorisation of these residues [47].

Therefore, it would be preferable that the future policies of development of the biomethane sector take into account the availability and distribution at a territorial level of other agro-industrial biomasses [48], in order to find the optimal location of new anaerobic digestion plants [42–49].

The amount of FVWs estimated to be sent to anaerobic digestion plants is just under 46 thousand tons of biomasses. This means that this biomass could make a net contribution to the production of biomethane. However, there is a limit imposed by the Italian law about disposal and reuse. In particular, the by-products produced by the wholesale market are classified as "waste" (in accordance with art. n° 10 of Legislative Decree 205/2010 [50]) because the wholesale market is the final step of the chain before the sale: If the products are not suitable for sale, they are waste. Instead, they can be classified as "by-products" (in accordance with art. n° 12 of Legislative Decree 205/2010 [50]) if they are produced by a farm that sells to the wholesale market. So, if the anaerobic digestion plant is authorised only for the utilisation of by-products and if its manager wants to use digestate as an organic soil improver for agricultural soil, wholesale market FVWs cannot be used as feedstocks.

This is nonsense because the product at the market does not undergo manipulation or alteration. It is the same product that is considered differently only because its holder (wholesale market or farmer) is different.

If these biomasses were treated as by-products and not as wastes, producers would avoid landfill disposal costs and could only pay for the delivery to biogas plants. Even more, biogas plant owners might be willing to pay for having these biomasses. Therefore, a virtuous economic circle could be generated: From waste to by-product.

5. Conclusions

The study carried out on collected data and GIS-based maps fulfilled the objectives of the research proposed by providing both the spatial localisation of horticultural production in Sicily and the estimation of the amount of FVW production by both investigating the flow production of all the wholesale markets located in Sicily and faceto-face interviewing the selected three biggest ones. Then, these results allowed the assessment of the potential biogas production from FVW reuse.

Therefore, since anaerobic digestion plants need a blend of different biomasses for the diet, new biogas plants could be located as near as possible to where biomasses are produced. This condition is relevant to reducing the transport costs of these materials. Therefore, a further study, which is in progress, is necessary in order to assess the effective biogas production by BMP and AD tests with the aim of finding a suitable mix for the digesters and promoting among local authorities the use of these wastes as new biomasses. So, it will be possible to reduce the significant environmental impact of the wholesale market residues and increase bioenergy production within the context of a green and circular economy.

Finally, the importance of the regulatory framework to support or to inhibit the use of theses biomasses was highlighted with regard to the considered case study.

Author Contributions: Conceptualization, R.S. and F.V.; methodology, F.V.; software, F.V.; formal analysis, R.S.; investigation, R.S. and F.V.; data curation, R.S.; writing—original draft preparation, R.S. and F.V.; writing—review and editing, R.S. and F.V.; visualization, R.S.; supervision, F.V. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Data Availability Statement: Data available in a publicly accessible repository.

Acknowledgments: This research was carried out within the project entitled "Sostenibilità dell'agricoltura e dell'industria agro-alimentare Mediterranea" —Programma Operativo Nazionale (PON) FSE–FESR "Research and Innovation 2014–2020, D.D. 407/2018" Attraction and Mobility (AIM)–Proposal: A1M1848200, Line 1, CUP: E64I19002440007 supported by Italian Ministry of Education, University and Research (MIUR). Furthermore, it is in line with the research project funded by the University of Catania: (PIAno di inCEntivi per la RIcerca di Ateneo 2020/2022 (D.R. N. 1208 dell'11 Maggio 2020)—"Engineering solutions for sustainable development of agricultural buildings and land"-WP3: Ottimizzazione e valorizzazione dell'uso delle risorse naturali, sottoprodotti e scarti.

Conflicts of Interest: The authors declare no conflict of interest.

References

- 1. Di Maria, F.; Sordi, A.; Cirulli, G.; Micale, C. Amount of energy recoverable from an existing sludge digester with the codigestion with fruit and vegetable waste at reduced retention time. *Appl. Energy* **2015**, *150*, 9–14.
- 2. Ingrao, C.; Selvaggi, R.; Valenti, F.; Matarazzo, A.; Pecorino, B.; Arcidiacono, C. Life cycle assessment of expanded clay granulate production using different fuels. *Resour. Conserv. Recycl.* **2019**, *141*, 398–409.
- 3. Appels, L.; Lauwers, J.; Degrève, J.; Helsen, L.; Lievens, B.; Willems, K.; Van Impe, J.; Dewil, R. Anaerobic digestion in global bio-energy production: Potential and research challenges. *Renew. Sust. Energy Rev.* **2011**, *15*, 4295–4301.
- Beurskens, L.W.M.; Hekkenberg, M. European Environment Agency—Renewable Energy Projection as Published in the National Renewable Energy Action Plans of the European Members States. 2011. Available online: http://www.ecn.nl/docs/library/report/2010/e10069.pdf (accessed on 15 October 2020).
- 5. Directive 2001/77/EC of the European Parliament and of the Council of 27 September 2001 on the promotion of electricity produced from renewable energy sources in the internal electricity market. *Off. J. Eur. Communities* **2001**, *L283*, 33–44.
- Bolzonella, D.; Battistoni, P.; Susini, C.; Cecchi, F. Anaerobic codigestion of waste activated sludge and OFMSW: The experience of Viareggio and Treviso plants (Italy). *Water Sci. Technol.* 2006, *53*, 203–211.
- Lin, J.; Zuo, J.; Gan, L.; Li, P.; Liu, F.; Wang, K.; Chen, L.; Gan, H. Effects of mixture ratio on anaerobic co-digestion with fruit and vegetable waste and food waste of China. J. Environ. Sci. 2011, 23, 1403–1408.
- 8. Selvaggi, R.; Pappalardo, G.; Chinnici, G.; Fabbri, C. Assessing land efficiency of biomethane industry: A case study of Sicily. *Energy Policy* **2018**, *119*, 689–695.
- Mostafavi, M.; Tajrishi, M.; Babaeinejad, S.M. Technical and Economic Study of Wastes in Agricultural Industry; Research Project at Sharif University of Technology Environmental Office: Tehran, Iran, 2003.
- 10. Li, J.; Zuo, J.; Cai, H.; Zillante, G. Construction waste reduction behavior of contractor employees: An extended theory of planned behavior model approach. *J. Clean. Prod.* **2018**, 172, 1399–1408.
- 11. Williams, H.; Wikstrom, F.; Otterbring, T.; Lofgren, M.; Gustafsson, A. Reasons for household food waste with special attention to packaging. *J. Clean. Prod.* **2012**, *24*, 141–148.
- 12. Pattnaik, S.; Reddy, M.V. Assessment of municipal solid waste management in Puducherry (Pondicherry), India. *Resour. Conserv. Recycl.* 2010, *54*, 512–520.
- Augustin, M.A.; Sanguansria, L.; Foxa, E.M.; Cobiac, L.; Cole, M.B. Recovery of wasted fruit and vegetables for improving sustainable diets. *Trends Food Sci. Technol.* 2020, 95, 75–85.
- 14. Viturtia, M.A.; Mata-Alvarez, J.; Cecchi, F. Two-phase continuous anaerobic digestion of fruit and vegetable wastes. *Resour. Conserv. Recycl.* **1995**, *13*, 257–267.
- 15. Abadi, B.; Mahdavian, S.; Fattahi, F. The waste management of fruit and vegetable in wholesale markets: Intention and behavior analysis using path analysis. *J. Clean. Prod.* **2021**, *279*, 123802.
- 16. Borowski, S. Co-digestion of the hydromechanically separated organic fraction of municipal solid waste with sewage sludge. *J. Environ. Manag.* **2015**, *147*, 87–94.
- 17. Masebinu, S.O.; Akinlabi, E.T.; Muzenda, E.; Aboyade, A.O.; Mbohwa, C. Experimental and feasibility assessment of biogas production by anaerobic digestion of fruit and vegetable waste from Joburg Market. *Waste Manag.* **2018**, *75*, 236–250.
- 18. Chen, Y.; Cheng, J.J.; Creamer, K.S. Inhibition of anaerobic digestion process: A review. Bioresour. Technol. 2008, 99, 4044–4064.
- 19. FAO-Food and Agriculture Organization. Developing sustainable Food Value Chains-Guiding Principles; FAO: Rome, Italy, 2014.
- 20. Gustavsson, J.; Cederberg, C.; Sonesson, U.; Van Otterdijk, R.; Meybeck, A. *Global Food Losses and Food Waste: Extent, Causes and Prevention*; Food and Agriculture Organization of the United Nations (FAO): Rome, Italy, 2011.
- 21. Segrè, A.; Falasconi, L. Il Libro Nero Dello Spreco in Italia: Il Cibo; Ambiente Edizioni: Milan, Italy, 2011.
- Scano, E.A.; Asquer, C.; Pistis, A.; Ortu, L.; Demontis, V.; Cocco, D. Biogas from anaerobic digestion of fruit and vegetable wastes: Experimental results on pilot-scale and preliminary performance evaluation of a full-scale power plant. *Energy Convers. Manag.* 2014, *77*, 22–30.
- 23. Esparza, I.; Jimenez-Moreno, N.; Bimbela, F.; Ancín-Azpilicueta, C.; Gandía, L.M. Fruit and vegetable waste management: Conventional and emerging approaches. *J. Environ. Manag.* **2020**, *265*, 110510.
- 24. Directive 2008/98/EC of the European Parliament and of the Council of 19 November 2008 on waste and repealing certain Directives. *Off. J. Eur. Union* **2008**, *L*312/3, 1–32.
- 25. Pavi, S.; Kramer, L.E.; Gomes, L.P.; Miranda, L.A.S. Biogas production from co-digestion of organic fraction of municipal solid waste and fruit and vegetable waste. *Bioresour. Technol.* **2017**, *228*, 362–367.
- Directive 2006/12/EC of the European Parliament and of the Council of 5 April 2006 on Waste. Off. J. Eur. Union 2006, L114/9, 1– 14.
- 27. Pappalardo, G.; Selvaggi, R.; Lusk, J. Procedural invariance as a result of commitment costs: Evidence from an economic experiment on farmers' willingness to pay for digestate. *Appl. Econ. Lett.* **2019**, *26*, 1243–1246.
- 28. Cristobal, J.; Caldeira, C.; Corrado, S.; Sala, S. Techno-economic and profitability analysis of food waste biorefineries at European level. *Bioresour. Technol.* **2018**, 259, 244–252.
- 29. Heck, S.; Rogers, M. Century Resource Revolution: How to Capture the Biggest Business Opportunity in a Century; McKinsey Company: Seattle, WA, USA, 2014.
- Valenti, F.; Liao, W.; Porto, S.M.C. Life cycle assessment of agro-industrial by-product reuse: A comparison between anaerobic digestion and conventional disposal treatments. *Green Chem.* 2020, 22, 7119–7139.

- 31. Valenti, F.; Porto, S.M.C.; Selvaggi, R.; Pecorino, B. Co-digestion of by-products and agricultural residues: A bioeconomy perspective for a Mediterranean feedstock mixture. *Sci. Total Environ.* **2020**, *700*, 134440.
- Valenti, F.; Porto, S.M.C.; Chinnici, G.; Cascone, G.; Arcidiacono, C. A GIS-based model to estimate citrus pulp availability for biogas production: An application to a region of the Mediterranean Basin. *Biofuels Bioprod. Biorefining* 2016, 10, 710–727.
- Valenti, F.; Arcidiacono, C.; Chinnici, G.; Cascone, G.; Porto, S.M.C. Quantification of olive pomace availability for biogas production by using a GIS-based model. *Biofuels Bioprod. Biorefining* 2017, 11, 784–797.
- Valenti, F.; Liao, W.; Porto, S.M.C. A GIS-based spatial index of feedstock-mixture availability for anaerobic co-digestion of Mediterranean by-products and agricultural residues. *Biofuels Bioprod. Biorefining* 2018, 12, 362–378.
- 35. Valenti, F.; Porto, S.M.C. Net electricity and heat generated by reusing Mediterranean agro-industrial by-products. *Energies* **2019**, *12*, 470.
- 36. Panda, S.K.; Mishra, S.S.; Kayitesi, E.; Ray, R.C. Microbial-processing of fruit and vegetable wastes for production of vital enzymes and organic acids: Biotechnology and scopes. *Environ. Res.* **2016**, *146*, 161–172.
- Diaz, A.I.; Laca, A.; Laca, A.; Diaz, M. Treatment of supermarket vegetable wastes to be used as alternative substrates in bioprocesses. Waste Manag. 2017, 67, 59–66.
- Liakou, V.; Paterakia, C.; Palaiogeorgoua, AM.; Kopsahelisa, N.; de Castroc, M.M.; Freired, D.M.G.; Nychasa, G.J.E.; Papanikolaoua, S.; Koutinasa, A. Valorisation of fruit and vegetable waste from open markets for the production of 2,3butanediol. *Food Bioprod. Process.* 2018, 108, 27–36.
- 39. Manetto, G.; Pecorino, B.; Selvaggi, R. Sustainability of a consortial anaerobic fermentation plant in Sicily. *Qual. Access Success* **2016**, *17* (Suppl. 1), 106–112.
- 40. Plazzotta, S.; Cottes, M.; Simeoni, P.; Manzocco, L. Evaluating the environmental and economic impact of fruit and vegetable waste valorisation: The lettuce waste study-case. *J. Clean. Prod.* **2020**, *262*, 121435.
- 41. ISTAT-Several Years. Available online: www.dati.istat.it (accessed on 7 November 2020).
- 42. Valenti, F.; Porto, S.M.C.; Dale, B.E.; Liao, W. Spatial analysis of feedstock supply and logistics to establish regional biogas power generation: A case study in the region of Sicily. *Renew. Sustain. Energy Rev.* **2018**, *97*, 50–63.
- Adani, F.; Schievano, A.; D'Imporzano, G. I Fattori Che Rendono Ottimale la Razione Per il Digestore. L'Informatore Agrario 2008, 40/2008:19. Available online: https://www.ilbiogas.it/biogas-ricerche-e-studi/fattori-razione-digestore.pdf (accessed on 24 November 2020).
- 44. Schievano, A.; Pognani, M.; D'Imporzano, G.; Adani, F. Predicting anaerobic biogasification potential of ingestates and digestates of a full-scale biogas plant using chemical and biological parameters. *Bioresour. Technol.* **2008**, *99*, 8112–8117.
- 45. Pappalardo, G.; Selvaggi, R.; Bracco, S.; Chinnici, G.; Pecorino, B. Factors affecting purchasing process of digestate: Evidence from an economic experiment on Sicilian farmers' willingness to pay. *Agric. Food Econ.* **2018**, *6*, 1–12.
- 46. Giuseppe, M.; Emanuele, C.; Rita, P.; Roberta, S.; Biagio, P. Performance evaluation of digestate spreading machines in vineyards and citrus orchards: Preliminary trials. *Helyon* **2020**, *6*, e04257.
- 47. Viswanath, P.; Devi, S.S.; Nand, S.K. Anaerobic digestion of fruit and vegetable processing wastes for biogas production. *Biores. Technol.* **1992**, 40, 43–48.
- Chinnici, G.; Selvaggi, R.; D'Amico, M.; Pecorino, B. Assessment of the potential energy supply and biomethane from the anaerobic digestion of agro-food feedstocks in Sicily. *Renew. Sustain. Energy Rev.* 2018, 82, 6–13.
- 49. Selvaggi, R.; Valenti, F.; Pappalardo, G.; Rossi, L.; Bozzetto, S.; Pecorino, B.; Dale, B.E. Sequential crops for food, energy, and economic development in rural areas: The case of Sicily. *Biofuels Bioprod. Biorefining* **2018**, *12*, 22–28.
- Legislative Decree 205/2010 about Provisions Implementing Directive 2008/98/EC of the European Parliament and of the Council of 19 November 2008 on Waste and Repealing Certain Directives; (10G0235) (OJ General Series n.288 of 10-12-2010—Ordinary Supplement n. 269); Off. J. Italian: Italy, 2010.