particles

MDPI

Article

Wave-like Behavior in the Source–Detector Resonance

Ioannis Contopoulos

Research Center for Astronomy and Applied Mathematics, Academy of Athens, GR 11527 Athens, Greece; icontop@academyofathens.gr

Abstract: We consider a particular model of a *Source* of independent particles and a macroscopic *Detector* that are both tuned to the same resonance frequency $\nu_0 \equiv 1/P$. Particles are emitted by the Source at exact multiples of the resonance period P, and the Detector absorbs them with a certain probability at any one of its points. The Detector may also announce the detection of the absorbed particle. Any particle that is not absorbed at a certain point passes through to a deeper layer in the interior of the Detector. Eventually, all particles will be absorbed, i.e., detected. We calculate the probability of detection for two particle time series generated by the same Source reaching the Detector with a time delay of δt and show that it manifests the illusion of collective (wave-like) interference with particle number conservation. We conclude that wave phenomena may reflect the nature of detectors rather than fundamental laws of physics.

Keywords: computer simulation; wave mechanics interpretations

1. Introduction

Quantum mechanics encompasses various interpretations, including the Copenhagen, de Broglie–Bohm, von Neumann–Wigner, stochastic mechanics, and many worlds theories [1–12]. Despite these interpretations, challenges remain with the ontological duality of microscopic particles. When detected individually, these particles appear as distinct entities, but their collective detection reveals wave-like distributions, such as interference patterns. Some researchers have suggested that individual particles carry phase information, which is gathered by specialized detectors [3,9,13,14]. These detectors only exhibit wave-like characteristics after processing a significant number of particles. Contopoulos et al. [14] concluded that interference phenomena may be explained classically, and wave-like behavior may just be an illusion of our detectors.

In this work, we diverge from the model of [14] because we understood that individual particles like photons cannot carry any phase information. Instead, we now propose that the phase information is carried collectively by the time series of these particles. As we will see in Sections 2–4 below, this information may be transferred to the detector if the latter is in resonance with the source that produces these time series. In Section 5, we describe a numerical double-slit experiment with two resonant streams of individual particles. In the end, we reach the same conclusion as [14], namely, that wave phenomena may reflect the nature of detectors rather than fundamental laws of physics.

2. The Source–Detector Resonance

We would like to once again ask the question: is it possible to manifest wave interference characteristics with independent non-interacting elementary particles?, where by elementary we mean classical particles with no internal degrees of freedom (i.e., no hidden

check for updates

Academic Editor: Hideaki Ohgaki

Received: 27 December 2024
Revised: 20 February 2025
Accepted: 25 February 2025
Published: 3 March 2025

Citation: Contopoulos, I. Wave-like Behavior in the Source–Detector Resonance. *Particles* **2025**, *8*, 24. https://doi.org/10.3390/particles8010024

Copyright: © 2025 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https://creativecommons.org/licenses/by/4.0/).

variables). Feynman famously quoted that wave-like characteristics are 'impossible, absolutely impossible to explain in any classical way' (Feynman Lectures on Physics [15]). We will challenge this statement with our model of the Source–Detector Resonance (hereafter SDR) that we will now present below. Let us begin by considering a mass M attached to a macroscopic oscillator which follows the equation

$$M\ddot{x} = -k(x - L_0). \tag{1}$$

Here, $(x - L_0)$ is the particle displacement around the equilibrium position $x = L_0$. We do not define the particular physical form of the oscillator. The resonance frequency of the system is

$$\nu_0 = \frac{1}{2\pi}\sqrt{\frac{k}{M}}. \tag{2}$$

Let us now assume that this oscillator is at the heart of a classical system that releases one elementary particle of mass $m \ll M$ every period

$$P \equiv \frac{1}{\nu_0}. \tag{3}$$

This continuous periodic release of particles does not have any effect on this large classical system. In particular, its eigenfrequency ν_0 remains unchanged, and its supply of elementary particles is unphased and practically unlimited. We call such a system a *Source* (see Figure 1).



Figure 1. A *Source* contains a macroscopic oscillator of resonance frequency ν_0 that releases elementary particles of mass m with the same velocity v every resonance period $P = 1/\nu_0$. M is the oscillating mass, and L_0 is the equilibrium position.

The time series of particles emitted by the Source may be represented as in the top left panel of Figure 2 if particles are emitted every period P, or as in the other panels of Figure 2 if they are emitted randomly 80%, 50%, 20% of the time of the full time series respectively, at random integer multiples of the period P. In both cases, the Fourier power spectrum density (hereafter PSD) of the time series has strong peaks around $\nu = 0$ and $\nu = \nu_0$, plus some extra noise at all frequencies. Notice that the peak at $\nu = 0$ accounts for all the particles, while the peak at $\nu = \nu_0$ is slightly lower because of the particles that contribute to the noise spectrum. If the signal becomes too weak (i.e., if too many particles are missing from the continuous time series of the top left panel of Figure 2), the peak at $\nu = \nu_0$ will disappear inside the noise, and there will remain no information of periodicity in the signal. The PSD shown is rescaled so that the value $\mathrm{PSD}(\nu = 0) = 1$ corresponds to the power of the full time series of the top left panel of Figure 2.



Figure 2. Top–bottom rows of panels: Time series and corresponding spectra of particles emitted continuously and intermittently $80\%, 50\%, 20\%$ with fixed period P by the Source (from top left to bottom right, respectively).

Let us now consider a *Resonant Oscillator* that may absorb these particles only if the particle time series it receives has the same eigenfrequency ν_0. In other words, the periodic sequence of particles can stimulate this Resonant Oscillator only at its resonance frequency. It is therefore natural to expect that the amplitude of the resonant oscillation A_{res} will be proportional to the number of particles per unit time that reach it at the resonant frequency. Clearly, this number is proportional to the ratio of the time series PSD at its resonance frequency ν_0 over the time series PSD at zero frequency. We thus expect that

$$A_{\text{res}} \propto \frac{\text{PSD}(\nu = \nu_0)}{\text{PSD}(\nu = 0)}. \tag{4}$$

We will now assume that the probability of detecting and absorbing one particle in a resonant time series by a Resonant Oscillator is proportional to its established amplitude of oscillation A_{res}. Mathematically, we express this as

$$\text{Prob}_{\text{abs one particle}} \propto A_{\text{res}} \propto \frac{\text{PSD}(\nu = \nu_0)}{\text{PSD}(\nu = 0)}. \tag{5}$$

In our approach, the macroscopic *Detector* consists of a large (infinite) number of Resonant Oscillators that receive a time series of individual particles, a fraction $\text{PSD}(\nu = \nu_0)/\text{PSD}(\nu = 0)$, which consists of a time series that is resonant with the De-

tector. This resonant time series excites the individual oscillators (an equivalent term may be 'pre-conditions') into resonant oscillations with amplitudes proportional to the exciting resonant time series, namely $\mathrm{PSD}(\nu = \nu_0)/\mathrm{PSD}(\nu = 0)$. In other words, a collective oscillating pattern is established over the Resonant Oscillators that comprise our Detector. If we further assume that the detection probability of one particle in the exciting resonant time series is proportional to its established amplitude of oscillation, then the resonant detection probability at every point of the Detector is pre-conditioned according to the ratio $\mathrm{PSD}(\nu = \nu_0)/\mathrm{PSD}(\nu = 0)$ of the time series that reach that point.

We have thus established that the Resonant Oscillator responds and announces the detection of (i.e., detects) a particle with a finite probability given by Equation (5) only if it consists part of a resonant time series. In particular if the stream of particles reaches a Resonant Oscillator with a frequency different from its resonance frequency, these particles will not be fully absorbed (i.e., detected). Some fraction of them will be absorbed, and some fraction of them will continue their motion unimpeded behind the Resonant Oscillator (see Figure 3). Obviously, the probability of detecting a particle in a general (not fully resonant) time series by a Resonant Oscillator is proportional to the probability of detecting one of them times the fraction of particles that are in a resonant time series. Mathematically, we express this as

$$\mathrm{Prob}_{\mathrm{abs}} \propto A_{\mathrm{res}} \times \left(\frac{\mathrm{PSD}(\nu = \nu_0)}{\mathrm{PSD}(\nu = 0)} \right) \propto \left(\frac{\mathrm{PSD}(\nu = \nu_0)}{\mathrm{PSD}(\nu = 0)} \right)^2. \tag{6}$$

We thus conclude that: The probability of detecting a particle in a general time series by a Resonant Oscillator is proportional to the square of the ratio of the time series PSD at its resonance frequency ν_0 over the time series PSD at zero frequency. When the Resonant Oscillator announces a particle absorption, this will result in a *Detection*. Particles that are not absorbed/detected by the Resonant Oscillator pass through it unimpeded.



Figure 3. The *Resonant Oscillator* is tuned to the same resonance frequency ν_0 as the Source. A stream of elementary particles reaches it, but the Resonant Oscillator absorbs them only if they reach it at its resonance frequency. The Resonant Oscillator absorbs particles with a certain probability given by Equation (6). If a particle is not absorbed, it continues its motion unimpeded downstream from the Resonant Oscillator.

3. Interference Between Two Particle Streams

Let us assume that the Source emits random particle sequences (i.e., with particles emitted at random integer multiples of the resonance period P) that travel either through path 1 or through path 2 (Figure 4). Both particle time sequences reach a Resonant Oscillator at a certain position in which the particles going through path 2 have a certain time delay δt with respect to those going through path 1. Let us start with no time delay, or a time delay equal to an integer multiple of the period P. In that case, the Resonant Oscillator receives the time sequence and corresponding spectrum of the top left panel of Figure 5. For a time delay equal to 10%, 20%, 30%, 40% of P plus an integer multiple of the period P, the Resonant Oscillator receives the time sequences and corresponding spectra of Figure 5. Finally, for a time delay equal to 50% of P plus an integer multiple of the period P, the Resonant Oscillator receives the time sequence and corresponding spectrum of the bottom right panel

of Figure 5. In that particular case, the power of the spectrum at the resonance frequency $\nu = \nu_0$ disappears! The reason is that, because of the superposition, the shortest periodic time interval between particles that reach the Resonant Oscillator is $P/2$, not P as in the original time series, thus the fundamental frequency of the combined series is $2\nu_0$, not ν_0. In Figure 6 we repeated the calculations of Figure 5 for the superposition of two unequal time series of particles emitted intermittently by the Source, one with only 20% of the particles of the continuous time series, and another with only 50%. We clearly see that when $\delta t = P/2$, the spectrum power at resonance does not vanish because the two time series are of unequal strength, and therefore, they cannot cancel each other. In Figure 7, we performed multiple numerical experiments for multiple values of the time delay δt and plotted the probability of absorption of a particle in the combined time series according Equation (6). This is the distribution of the square of the ratio of the combined stream's PSD at resonance over the PSD of the combined stream at zero frequency (the total power of the combined stream) as a function of the time delay δt. We also plot the function $\cos^2(\pi \delta t/P)$ obtained from Equation (7) below for $\alpha = \beta = 0.8$. The fit to the numerical experiments is perfect, except around $\delta \approx P/2$ where the PSD at resonance reaches the noise level.



Figure 4. Sketch of interference between two different streams of particles that originate at the same Source and have a time delay of $\delta t = (L_1 - L_2)/v$ between themselves. Such superposition manifests interference characteristics at the Detector.

We see that when the two streams of particles have the same power and are in phase (i.e., $\delta t = 0$ or some integer multiple of P), the Resonant Oscillator detects the same number of particles as the combined two particle time series, i.e., the same number of particles as those emitted by the Source. No more, no less. At the other limit, when the two streams of particles have the same power and are opposite in phase (i.e., $\delta t = P/2$ plus some integer multiple of P), the combined stream's spectrum power at resonance vanishes. In general, in the case of a superposition of two intermittently periodic time series with magnitudes α and β, the probability of detection at the Resonant Oscillator/Detector as defined by Equation (6) is found numerically to be proportional to

$$\text{Prob}_{\text{abs}} \propto \left(\frac{\text{PSD}(\nu_0)}{\text{PSD}(0)}\right)^2 = \frac{\alpha^2+\beta^2}{(\alpha+\beta)^2}\left(1+\frac{2\alpha\beta}{\alpha^2+\beta^2}\cos\left(2\pi\frac{\delta t}{P}\right)\right) \tag{7}$$

(See Figure 7). Once again, particles from both particle streams that are not absorbed/detected by the Resonant Oscillator pass through it unimpeded, and reach a deeper point inside the Detector.



Figure 5. Top–bottom rows of panels: The superposition of two equal-power time series of particles emitted intermittently by the Source with fixed period P and a time delay between them equal to $\delta t = 0, 0.1, 0.2, 0.3, 0.4, 0.5$ times the period P (plus an integer multiple of P) from top left to bottom right respectively. These time series of particles are collected by the Resonant Oscillator/Detector. The corresponding spectra are also shown. We see very clearly that when $\delta t = 0.5P$, the spectrum power at resonance vanishes.



Figure 6. Top-Bottom rows of panels: The superposition of two unequal time series of particles emitted intermittently by the Source with fixed period P and a time delay between them equal to $\delta t = 0, 0.1, 0.2, 0.3, 0.4, 0.5$ times the period P (plus an integer multiple of P) from top left to bottom right respectively. The first time series contains only 20% of the particles of the continuous time series of the tope left panel of Figure 2, and the second only 50%. These time series of particles are collected by the Resonant Oscillator/Detector. The corresponding spectra are also shown. We see very clearly that when $\delta t = 0.5P$, the spectrum power at resonance does not vanish. The ratio of the number of particles and resonance over the total number of particles emitted by the Source is given by Equation (7).



Figure 7. Distribution of the square of the ratio of the PSD at resonance over the PSD at zero frequency as a function of the time delay δt. We consider here the combination of two power time series with $(\alpha, \beta) = (0.8, 0.8)/(0.8, 0.6)/(0.8, 0.4)/(0.8, 0.2/(0.8, 0.1)/(0.8, 0.05))$ from top left to bottom right, respectively. The fits to the expression of Equation (7) for the correponding values of α and β are also shown. Except for the case of zero time delay δt, this distribution leaves a fraction $(1 - \text{Prob}_{\text{abs}})$ of particles undetected. Clearly, when either one of α or β is equal to zero (i.e., if we have only one and not two combined particle time series), the detection probability Prob_{abs} is equal to unity.

4. A Double-Slit Experiment with Particles

According to our previous definition, a macroscopic Detector consists of a large (infinite) collection of Resonant Oscillators. It is important that the Detector has sufficient thickness to absorb (and thus also detect) all particles from the combined time series that reach it. The interference setup that we propose is that from a typical double-slit experiment. Particle series reach points of the Detector coming from both slits, and the time difference δt between the two time series is proportional to the difference in the particle trajectory lengths over the particle speed. Particles are absorbed by the Resonant Oscillators/Detectors at each layer of the Detector according to the probability distribution of Equation (7). The particles that are not absorbed continue to travel along the same line that they followed after emerging from their respective slit and are combined with the stream of non-absorbed particles that emerged from the other slit. Some of them will be absorbed/detected in that second layer of the Detector and so on and so forth. Eventually, all particles emerging from both slits will be absorbed, and the interference pattern shown schematically in Figure 8 will emerge in the interior of the Detector. No particle will be left undetected. What is important though is that a certain finite thickness is needed in order for the Detector to absorb/detect all particles.



Figure 8. Sketch of a double-slit experiment. Two particle streams from the two slits that reach the surface of the *Detector* with $\delta t = 0$ (plus an integer multiple of the period P) are directly absorbed. Double particle streams with $\delta t = P/2$ plus an integer multiple of P are not immediately absorbed, and their individual particle streams continue unimpeded in the interior of the Detector. It is clear that all such particles will be gradually absorbed within a certain depth inside the Detector as they reach positions with $0 \leq \delta t < P/2$ (plus an integer multiple of the period P). The Detector thus announces the same number of particles as those that reached it through the two slits.

It is interesting that the same distribution as that of Equation (7) is obtained when we superpose two delayed periodic signals of amplitudes α and β, respectively, expressed mathematically as $\alpha \exp(i2\pi t/P) + \beta \exp(i2\pi(t+\delta t)/P)$. This sum may represent for example the addition of the electric field of two interfering electromagnetic waves. Classically, the power of the combined signal is proportional to the square of the amplitude of the electric field superposition, namely $(\alpha + \beta\cos(2\pi\delta t/P))^2 + \beta^2 \sin^2(2\pi\delta t/P) = (\alpha^2+\beta^2)(1 + 2\alpha\beta\cos(2\pi\delta t/P)/(\alpha^2+\beta^2))$. If we further divide this expression by $(\alpha+\beta)^2$, we obtain the result of Equation (7). In our picture of periodic sequences of elementary particles, the power is proportional to the number of detected particles per unit of time at the Detector. Figure 7 is telling us that when

two independent periodic signals from our Source reach a certain point in the resonant Detector with a time delay of δt between them, the Resonant Oscillator at that position collects the sum of the two particle streams, i.e., a power that is double the power of each particle stream, but absorbs (detects, announces) a number of particles proportional to Equation (7) times the power of the Source particle stream. This result applies on average even if the periodic sequences of particles are not complete and certain particles are randomly missing. As long as particles of each stream are emitted and collected at integer multiples of the resonance period P, the same illusion of interference is obtained.

It should be obviously observed that the number of particles that reach the Detector per unit of time is equal to the sum of the number of particles per unit of time in the two particle streams, irrespective of the time delay δt between the two streams. In the case of destructive interference at a certain position ($\delta t = P/2$), the Detector announces zero particles at that position. As a result of this non-detection, the two particle streams continue their travel unimpeded, and eventually all of them are detected deeper inside the Detector (see Figure 8). In the case of constructive interference ($\delta t = 0$) the Detector absorbs the two streams of particles, but if we also count the particles also absorbed behind that point of constructive interference, the Detector announces on average double the number of particles detected at the first surface point of constructive interference. Thus, the Detector conserves the number of particles, provided we are willing to account for all detections, even those in the interior of the Detector.

If we collect all detections along the various angles of the experiment, we obtain the result of a double-slit experiment. The new result of our present numerical experiment is that the same illusion of interference is obtained only if we are willing to consider a Detector with a finite thickness. Had we considered only the surface layers of the Detector, we would have obtained the same interference pattern with one-half the number of particles that reach the Detector.

5. Conclusions

We have shown that interference patterns may arise from the superposition of two particle streams generated by the same classical Resonant Oscillator that reach a second classical Resonant Oscillator tuned to the same resonance frequency. These two Resonant Oscillators are

- The Source that emits a series of particles at integer multiples of its resonance period,
- The Detector that receives these particle streams at each point and announces detections at that point with a probability proportional to the square of the PSD of the particle time series at the Source–Detector Resonance (SDR) frequency. The particles that are not detected at that point continue their motion unimpeded deeper inside the Detector.

Essentially, two such streams of independent non-interacting particles emitted with the same velocity at times that are integer multiples of P manifest on average the characteristics of a continuous wave, provided the detector operates only at the resonance frequency $\nu_0 = 1/P$. We have shown this result with a simple numerical experiment of interference when there is a time delay δt between two such streams of particles. Our numerical result is valid even when the two streams carry unequal numbers of particles per unit time (on average). The resulting interference pattern is the same as that of continuous sinusoidal waves with equal or unequal amplitudes.

When we applied the above result to reproduce a double-slit experiment with particles, only one half of the particles were detected when they first reached the surface layer of the Detector. The other one half of the particles continued their respective motions in deeper layers of the Detector and were eventually absorbed/detected by it. The interference pattern

integrated over the thickness of the Detector, creates the same effect as the interference pattern resulting from two interfering sinusoidal waves. This result is currently only qualitative, and we plan to provide a rigorous proof in the future.

Several researchers have tried in the past to generate similar interference patterns with particles that carry some type of internal hidden variables in the form of an internal clock (e.g., [3,9,13,14]). The detector collects and processes that information before it announces a detection, similarly to our present model. The problem with such approaches is that they do not work for photons which cannot carry any internal clock. In our present model, the necessary information of the time delay is carried collectively by the two streams of particles, not by each individual particle as in the previous approaches quoted above.

Our current model applies only to free particles. Until this point, we have said nothing about the particle velocity v which is important for the manifestation of the time delay δt between the two different paths of lengths L_1 and L_2 ($\delta t = (L_1 - L_2)/v$). In the Appendix, we offer a tentative physical model of how these particles may have obtained their velocity v and their kinetic energy.

In summary, if there exist macroscopic sources and detectors that operate as proposed, wave-like interference phenomena may be explained if the macroscopic resonant detector is excited by the resonant time series into a wave-like oscillation pattern which pre-conditions the detection probability of each one of its individual oscillators according to Equation (6). It is interesting that this pre-conditioning of the detector is also able to explain classical wave phenomena like light refraction with resonant time series of individual particles/photons. We plan to design a numerical experiment with a large number of resonant physical oscillators to test our claims above. If all macroscopic detectors in nature operate as proposed in this paper, wave-like phenomena may exist as an illusion of the detectors and not a fundamental property of nature.

Funding: This research received no external funding.

Data Availability Statement: The datasets generated during the current study are available from the author upon reasonable request.

Acknowledgments: We acknowledge fruitful discussions with George Contopoulos, Konstantinos Gontikakis, and Christos Efthymiopoulos.

Conflicts of Interest: The author declares no conflict of interest.

Appendix A. The Particle Kinetic Energy

We will now discuss how particles of mass m emitted by the Source may have obtained their velocity v. We will tentatively argue that particles are emitted with a particular kinetic energy

$$\epsilon_{\text{kinetic}} = \frac{1}{2} m v^2 = h\nu_0 \,, \tag{A1}$$

where h is a constant with dimensions of angular momentum. It is clear that what is physically fundamental in our model is not the particle energy but the Source resonance frequency. Let us now see how Equation (1) may be modified to account for the emission of particles with the energy of Equation (A1). It is easy to see that in order to introduce the parameter h and keep the dimensions of Equation (1), one may add an extra dissipation term as

$$M\ddot{x} = -k(x - L_0) - \frac{h}{2\pi^2} \frac{\dot{x}}{L_0^2}. \tag{A2}$$

One sees directly in Figure A1 below that, starting from an initial displacement $x(t=0) = 2L_0$, $x(t) \approx L_0(1+\cos(2\pi\nu_0 t))$ and $\dot{x}(t) \approx -2\pi\nu_0 L_0 \sin(2\pi\nu_0 t)$ for small h, thus every period the Source oscillator loses a small amount of energy equal to

$$\epsilon = \frac{h}{2\pi^2}\int_{t=0}^{P}\frac{\dot{x}^2}{L_0^2}\mathrm{d}t \approx \frac{h}{2\pi^2}\frac{(2\pi\nu_0 L_0)^2}{L_0^2}\int_{t=0}^{P}\sin^2(2\pi\nu_0 t)\mathrm{d}t = h\nu_0\,, \tag{A3}$$

provided the energy ϵ lost per period is much smaller than the energy stored in the Source oscillator, namely, $h\nu_0 \ll \frac{1}{2}M(2\pi\nu_0 L_0)^2$.



Figure A1. Upper plot: $x = x(t)$ for the Source oscillator described by Equation (A2). $x(0) = 2L_0$. $\dot{x}(0) = 0$. Initial oscillator energy $\gg h\nu_0$. Time in units of the oscillator period P. Lower plot: Evolution of the oscillator energy (blue/orange/green lines: kinetic/potential/total energy respectively).

We thus propose the following model for the emission of elementary particles of mass m by the Source. Every oscillation period, the Source loses energy ϵ according to Equation (A3). Let us assume that right next to the Source oscillator exists a reservoir of elementary particles that, every period P, absorbs this energy ϵ and leaves the system. Thus, at every period of oscillation, a particle is emitted with kinetic energy

$$\epsilon_{\text{kinetic}} = \frac{1}{2}mv^2 = \epsilon = h\nu_0. \tag{A4}$$

This approach offers a different view of how particles obtain their kinetic energy. It is the macroscopic Source oscillator that loses parcels of energy $h\nu_0$ every period P, and these parcels of energy are somehow transferred to the particles.

The model proposed in this short paper considers only streams of particles with the same kinetic energy ϵ_{kin} and velocity v. These particles do not change their kinetic energy as they travel from the Source to the Detector. If we were to expand our model to a theory of particle dynamics, we must assume that, in order for a particle to change (increase or decrease) its kinetic energy, it must be absorbed by a macroscopic system which resonates at the frequency that corresponds to its current kinetic energy, and it must then be re-emitted at a different energy that corresponds to a different resonance frequency of the macroscopic system. Such a tentative model would require a special type of interaction between elementary particles and force fields.

References

1. Bacciagaluppi, G.; Valentini, A. *Quantum Theory at the Crossroads: Reconsidering the 1927 Solvay Conference*; Cambridge University Press: Cambridge, UK, 2009.
2. Bassi, A.; Lochan, K.; Satin, S.; Singh, T.P.; Ulbricht, H. Models of wave-function collapse, underlying theories, and experimental tests. *Rev. Mod. Phys.* **2013**, *85*, 471. [CrossRef]
3. Bohm, D. A Suggested Interpretation of the Quantum Theory in terms of 'Hidden' Variables, I and II. *Phys. Rev.* **1952**, *85*, 166–193. [CrossRef]
4. Everett, H., III. "Relative state" formulation of quantum mechanics. *Rev. Mod. Phys.* **1957**, *29*, 454. [CrossRef]
5. Harrigan, N.; Spekkens, R.W. Einstein, incompleteness, and the epistemic view of quantum states. *Found. Phys.* **2010**, *40*, 125–157. [CrossRef]
6. Holland, P.R. *The Quantum Theory of Motion: An Account of the de Broglie-Bohm Causal Interpretation of Quantum Mechanics*; Cambridge University Press: Cambridge, UK, 1995.
7. Maudlin, T. Foundations of Quantum Mechanics: An Exploration of the Physical Meaning of Quantum Theory. *Am. J. Phys.* **2018**, *86*, 953–955. [CrossRef]
8. Merzbacher, E. *Quantum Mechanics*; John Wiley & Sons: New York, NY, USA, 1998.
9. Nelson, E. Derivation of the Schrödinger equation from Newtonian mechanics. *Phys. Rev.* **1966**, *150*, 1079. [CrossRef]
10. Nelson, E. *Quantum Fluctuations*; Princeton University Press: Princeton, NJ, USA, 1985.
11. Nelson, E. Review of stochastic mechanics. *J. Phys. Conf. Ser.* **2012**, *361*, 012011. [CrossRef]
12. Styer, D.F.; Balkin, M.S.; Becker, K.M.; Burns, M.R.; Dudley, C.E.; Forth, S.T.; Gaumer, J.S.; Kramer, M.A.; Oertel, D.C.; Park, L.H.; et al. Nine formulations of quantum mechanics. *Am. J. Phys.* **2002**, *70*, 288–297. [CrossRef]
13. Jin, F.; Yuan, S.; De Raedt, H.; Michielsen, K.; Miyashita, S. Corpuscular model of two-beam interference and double-slit experiments with single photons. *J. Phys. Soc. Jpn.* **2010**, *79*, 074401. [CrossRef]
14. Contopoulos, I.; Tzemos, A.C.; Zanias, F.; Contopoulos, G. Interference with Non-Interacting Free Particles and a Special Type of Detector. *Particles* **2023**, *6*, 121. [CrossRef]
15. Feynman, R.P. *The Feynman Lectures on Physics*; Addison-Wesley: Boston, MA, USA , 1965; Volume III.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.