

Article

BFSS Matrix Model Cosmology: Progress and Challenges

Suddhasattwa Brahma^{1,*}, Robert Brandenberger² and Samuel Laliberte²¹ Higgs Centre for Theoretical Physics, School of Physics & Astronomy, University of Edinburgh, Edinburgh EH9 3FD, UK² Department of Physics, McGill University, Montreal, QC H3A 2T8, Canada

* Correspondence: suddhasattwa.brahma@gmail.com

Abstract: We review a proposal to obtain an emergent metric space-time and an emergent early universe cosmology from the Banks–Fischler–Shenker–Susskind (BFSS) matrix model. Some challenges and directions for future research are outlined.

Keywords: early universe; superstring cosmology; emergent cosmology

1. Introduction

The current models of the very early universe are generally based on effective field theory (EFT), i.e., on considering matter fields minimally coupled to Einstein gravity. This is true both for the inflationary scenario [1–4] and for most alternatives to it (see, e.g., [5–7] for a comparative review of alternatives to inflation). However, there are conceptual challenges that effective-field-theory descriptions of cosmology face. For example, in the case of scalar field matter—ubiquitously used in early universe models—the range of such fields [8] and the shape of [9] of their potentials are constrained if we demand that the effective fields descend from some fundamental physical theory (the best candidate of which is superstring theory). Independent of superstring theory, the Trans-Planckian Censorship Conjecture (TCC) [10] severely constrains inflationary models that are based on an EFT analysis [11] (see, e.g., [12,13] for discussions of some arguments in support of the TCC; note that scenarios of inflation that are not based on EFT analyses (see [14–19] for examples) are not necessarily subject to these constraints).

An EFT analysis of matter also leads to the famous cosmological constant problem: considering matter fields coupled to Einstein gravity, and canonically quantizing the fields by expanding them in plane wave modes and quantizing each mode like a harmonic oscillator, then the vacuum energy of all of these modes yields a contribution to the cosmological constant that is many orders of magnitude (120 orders in a non-supersymmetric model) larger than what is compatible with current observations (to avoid the Planck catastrophe, an ultraviolet (UV) cutoff scale must be imposed, the value of which determines the predicted cosmological constant).

The abovementioned problems indicate that one needs to go beyond point-particle-based EFT to obtain a consistent picture of cosmology. Here, we review an approach [20,21] to obtain a model of the very early universe that is not based on EFT. Our starting point is the Banks–Fischler–Shenker–Susskind (BFSS) matrix model [22], a proposed non-perturbative definition of superstring theory (see also [23,24] for early paper on the quantum theory of membranes, and [25–33] for other matrix models related to superstring theory). We outline an avenue of how to obtain an emergent space-time, an emergent metric, and an emergent early universe cosmology. We consider the matrix model to be in a thermal state. This leads to density fluctuations and gravitational waves on the emergent space-time, and we indicate how the resulting power spectra of these fluctuations are scale-invariant and consistent with current observations.

In this context, we remind the reader that the inflationary scenario [1–4] is not the only early universe scenario that is consistent with current cosmological observations (see,



Citation: Brahma, S.; Brandenberger, R.; Laliberte, S. BFSS Matrix Model Cosmology: Progress and Challenges. *Physics* **2023**, *5*, 1–10. <https://doi.org/10.3390/physics5010001>

Received: 17 October 2022

Revised: 20 November 2022

Accepted: 23 November 2022

Published: 22 December 2022



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

e.g., [5–7] for comparative reviews of inflation and alternatives). An alternate scenario is the *emergent scenario*, which is based on the assumption that there is a primordial phase that can be modelled as quasi-static (from the point of view of the Einstein frame) and that then undergoes a phase transition to the Standard Big Bang phase of expansion.

A model for an emergent scenario is *string gas cosmology* (SGC) [34] (see also [35] for earlier study and [36] for a review), where it is assumed that the universe begins in a quasi-static *Hagedorn* phase, a phase in which a gas of closed superstrings is in thermal equilibrium at a high temperature close to the limiting Hagedorn value [37]. The initial size of all of the spatial dimensions is taken to be the same. However, a dimension of space can only expand if the winding modes in that direction disappear, and since strings are two-dimensional world sheets, the intersection of winding strings (which is necessary for winding modes to disappear) is not possible in more than four large space–time dimensions (in this argument, it is assumed that there are no long-range attractive forces between winding strings. Since the local curvature perpendicular to gauge strings vanishes [38], it is reasonable to make this assumption). This helps to clarify why only three of the nine spatial dimensions that superstring theory predicts can become large. It can then be shown [39,40] that thermal fluctuations of this string gas lead to scale-invariant spectra of cosmological perturbations and gravitational waves, with a slight red tilt for the curvature fluctuations and a slight blue tilt for the gravity wave spectrum.

What was missing in SGC is a consistent dynamical framework for analyzing the cosmological evolution. To obtain such a framework, one needs to go beyond an EFT analysis, and this is where our study comes in (see also [41] for a different approach to obtaining an emergent cosmology from superstring theory that matches well with the SGC model).

Note that, as we review below, the computation of cosmological perturbations and gravitational waves in our BFSS model is based on the same formalism used in SGC, with very similar results.

In the next Section, we present a brief review of the BFSS matrix model. In Section 3, we then present our proposal for emergent time, space, and metric from the matrix model. On this basis, in Section 4 we develop a new picture of an emergent universe and compute the curvature fluctuations and gravitational waves that are predicted if one starts with a high temperature state of the BFSS matrix model. We use natural units in which the speed of light and Planck’s constant are set to one.

2. BFSS Matrix Model

Starting point is the BFSS matrix model [22], a quantum theory involving ten bosonic time-dependent $N \times N$ Hermitean matrices, $A_0(\tau)$ and $X_i(\tau)$, $i = 1, \dots, 9$, and their sixteen Fermionic superpartners $\theta_{a,b}(\tau)$ (also $N \times N$ matrices), which transform as spinors under the $SO(9)$. The Lagrangian is

$$L = \frac{1}{2g^2} \left[\text{Tr} \left\{ \frac{1}{2} (D_\tau X_i)^2 - \frac{1}{4} [X_i, X_j]^2 \right\} - \theta^T D_\tau \theta - \theta^T \gamma_i [\theta, X^i] \right], \quad (1)$$

where g is a coupling constant and the covariant derivate involves the temporal bosonic matrix A_0 :

$$D_\tau := \partial_\tau - i[A_0(\tau), \cdot], \quad (2)$$

with $\partial_\tau \equiv \partial/\partial\tau$.

In the above, τ is the BFSS time, and γ_i are Clifford algebra matrices (see, e.g., [42,43] for reviews of the BFSS matrix model).

This Lagrangian has both a $U(N)$ symmetry and a $SO(9)$ symmetry under rotations of the spatial matrices (and the corresponding transformation of the spinors). The partition function of the theory is given by the functional integral,

$$Z = \int dA d\theta e^{iS_{\text{BFSS}}}, \quad (3)$$

where S_{BFSS} is the action obtained by integrating the Lagrangian over time, and the integration measures are the Haar measure for the bosonic variables and the spinorial counterpart.

This matrix model was introduced [22] as a candidate for a non-perturbative definition of superstring theory (more precisely of M-theory). String theory emerges in the limit $N \rightarrow \infty$ with $\lambda \equiv g^2 N$ held fixed.

Here, we consider this matrix model in a high-temperature state, with the temperature being denoted by T . By this assumption, there is no evolution of the system in BFSS time τ . However, below we extract an emergent time, t (which is different from the time τ) in terms of which there are non-trivial dynamics.

In the high-temperature state considered here, the spatial matrices, X_i , are periodic in Euclidean BFSS time τ_E , and one can expand them in Matsubara modes:

$$X_i(\tau_E) = \sum_n X_i^n e^{2\pi n T \tau_E}, \quad (4)$$

where n runs over the integers (note that this integer n is unrelated to the integers n_i used in later Sections to introduce comoving coordinates). At high temperatures, the BFSS action is dominated by the $n = 0$ modes,

$$S_{\text{BFSS}} = S_{\text{IKKT}} + \mathcal{O}(1/T), \quad (5)$$

where S_{IKKT} is the bosonic part of the action of the Ishibashi–Kawai–Kitazawa–Tsuchiya (IKKT) matrix model [44], which, in terms of rescaled matrices, A_i ,

$$A_i \equiv T^{-1/4} X_i^0, \quad (6)$$

is given by

$$S = -\frac{1}{g^2} \text{Tr} \left(\frac{1}{4} [A^a, A^b] [A_a, A_b] \right). \quad (7)$$

In what follows, the matrices A_0 and A_i (i.e., the zero modes of the BFSS matrices A_0 and X_i) are used to extract an emergent space-time. The $n \neq 0$ modes of the matrices play an important role when obtaining cosmological fluctuations and gravitational waves.

3. Emergent Metric Space-Time

Without loss of generality, one can work in a basis in which the temporal matrix A_0 is diagonal. The expectation values of the diagonal elements are identified with the *emergent time*. We order the diagonal elements in increasing value,

$$A_0 = \text{diag}(t_1, \dots, t_N), \quad (8)$$

with $t_i \leq t_j$ for $i < j$.

Numerical simulations of the IKKT model [45] indicate that

$$\frac{1}{N} \langle \text{Tr} A_0^2 \rangle \sim \kappa N, \quad (9)$$

where κ is a (small) constant. Assuming that the eigenvalues are separated equidistantly, it then follows that the maximal eigenvalue (the maximal value of the emergent time) scales as

$$t_{\max} \sim \sqrt{N}, \quad (10)$$

and that hence the difference between the discrete values of time scales as

$$\Delta t \sim \frac{1}{\sqrt{N}}. \quad (11)$$

Thus, in the $N \rightarrow \infty$ limit, one obtains emergent continuous and infinite time. Since the eigenvalues are symmetric about $t = 0$, the time that emerges is infinite both in the past and in the future. Hence, there is neither a Big Bang nor a Big Crunch singularity.

Turning to the spatial matrices, we follow the proposal of [46] and consider, for each integer m , $n_i \times n_i$ (where n_i is an integer sufficiently smaller than N) spatial submatrices $\bar{A}_i(t(m))$ centered a distance m down the diagonal (see Figure 1):

$$(\bar{A}_i)_{IJ}(t(m)) \equiv (A_i)_{m+I, m+J}, \quad (12)$$

where $t(m)$ is the m 'th temporal eigenvalue. Following [46], one can define the *extent of space* parameters $x_i(t)$ in the i 'th spatial direction at the time t via

$$x_i(t)^2 \equiv \left\langle \frac{1}{n} \text{Tr}(\bar{A}_i(t))^2 \right\rangle. \quad (13)$$

With this definition of *emergent space*, there is non-trivial time evolution in terms of the emergent time.




Figure 1. In the basis in which the A_0 matrix is diagonal with the diagonal elements defining emergent time, we can use spatial $n_i \times n_i$ submatrices $\bar{A}_i(t)$ located a distance t down the diagonal to define the i 'th spatial direction at time t . The figure is taken from [42] with permission.

In the IKKT model, it has been observed via numerical simulations that, in the presence of the fermionic terms in the action, the state that minimizes the free energy breaks the $SO(9)$ symmetry into $SO(6) \times SO(3)$. Of the nine extent of space parameters, six remain small and three increase [47–52]. This result can also be confirmed by means of Gaussian expansion calculations [53,54]. The emergence of three macroscopic spatial dimensions with the other six remaining microscopic dimensions reminds one of the same phenomenon that was discovered in the context of string gas cosmology [34], where string winding modes prevent spatial dimensions from expanding. In order to obtain large dimensions, the string winding modes near these dimensions must be annihilated, and this is not possible in

more than three large spatial dimensions since string winding modes will have vanishing intersection probability (see [55,56] for a more detailed analysis).

Very recently, we have been able to show [57] that in the presence of the fermionic terms, the $SO(9)$ symmetry will also be spontaneously broken in the BFSS matrix model. This result was derived using a Gaussian expansion calculation. With this tool, it is not yet possible to determine the symmetry of the state that minimizes the free energy, but the intuition gained from string gas cosmology leads us to expect that the symmetry breaking pattern will be the same as what was observed in the IKKT model.

Making use of the Riemann–Lebesgue Lemma, it can be shown from the bosonic IKKT action that the off-diagonal elements of the spatial matrices decay once one departs beyond a critical distance, n_c , from the diagonal:

$$\sum_i \langle |A_i|_{ab}^2 \rangle \rightarrow 0 \text{ for } |a - b| > n_c, \quad (14)$$

where

$$n_c \sim \sqrt{N} \quad (15)$$

This result can be confirmed by numerical simulations (see [58–61] for recent investigations of the IKKT matrix model), which also indicate that

$$\sum_i \langle |A_i|_{ab}^2 \rangle \sim \text{constant when } |a - b| < n_c, \quad (16)$$

a result, for which we at present do not have any analytical evidence.

Our proposal [21] is to view n_i as the i 'th comoving spatial coordinate and

$$l_{\text{phys},i}^2(n_i, t) \equiv \langle \text{Tr}(\bar{A}_i(t))^2 \rangle, \quad (17)$$

as the physical length square of the curve from $n_i = 0$ to n_i along the i 'th coordinate direction. It then follows from Equation (16) that

$$l_{\text{phys},i}(n_i) \sim n_i \text{ for } n_i < n_c. \quad (18)$$

Given the definition of length and comoving coordinates, we can extract the spatial metric component g_{ii} in the usual way:

$$g_{ii}(n_i)^{1/2} = \frac{d}{dn_i} l_{\text{phys},i}(n_i) \quad (19)$$

and we obtain

$$g_{ii}(n_i, t) = \mathcal{A}(t) \delta_{ii} \quad i = 1, 2, 3, \quad (20)$$

where $\mathcal{A}(t)$ is an increasing function of t at late times and δ_{ii} is the Kronecker delta. Making use of the remnant $SO(3)$ symmetry, one obtains:

$$g_{ij}(\mathbf{n}, t) = \mathcal{A}(t) \delta_{ij} \quad i, j = 1, 2, 3, \quad (21)$$

where \mathbf{n} stands for the comoving 3-vector corresponding to the three large dimensions.

We have thus indicated how one may obtain emergent time, space, and spatial metrics from the zero modes of the BFSS matrix model. The emergent space-time is infinite both in temporal and spatial directions, and from Equation (21) it follows that the metric that emerges on the emergent large three-dimensional space is spatially flat. Hence, in the scenario that we are proposing the famous horizon and flatness problems of Standard Big Bang cosmology are automatically solved without the need for any phase of inflation [1–4] or superslow contraction [62,63]. In the next Section we show that the thermal fluctuations in the BFSS state automatically lead to approximately scale-invariant spectra for cosmological perturbations and gravitational waves.

4. Emergent Early Universe Cosmology

The cosmology that emerges from the BFSS matrix model is an *emergent cosmology*. The classical notion of space-time only makes sense at late emergent time. Solving the classical bosonic matrix equations for the three large spatial dimensions at late times yields [64–66],

$$\mathcal{A}(t) \sim t^{1/2}, \quad (22)$$

i.e., expansion like in a radiation-dominated universe. We expect that the inclusion of the fermionic sector will change this scaling and produce a matter component that scales as pressureless dust. Thus, in our cosmology there is a direct transition from the emergent phase to that of Standard Big Bang cosmology, like in string gas cosmology. Note that there is no sign of a cosmological constant. Thus, it appears that the cosmology that emerges from the BFSS matrix model will be free of the infamous cosmological constant problem. On the other hand, it is at the present time unclear how dark energy will emerge. Based on the swampland criteria [8,9] and the TCC [10], it follows that dark energy cannot be a cosmological constant. It may be a quintessence field [67–69] that emerges from the matter sector (see, e.g., [70] for an attempt to obtain a unified dark sector (dark matter and dark energy) from string theory).

Since we are considering the BFSS matrix model in a high-temperature state, there will be thermal fluctuations, and these will lead to curvature fluctuations and gravitational waves like in SGC (see [39,40] for analyses of the generation of cosmological perturbations and gravitational waves in SGC).

At late times, the classical metric of the $3 + 1$ dimensional space-time can be expressed at the level of linearized fluctuations as (see, e.g., [71,72] for reviews of the theory of cosmological perturbations)

$$ds^2 = a^2(\eta) \left((1 + 2\Phi)d\eta^2 - [(1 - 2\Phi)\delta_{ij} + h_{ij}]dx^i dx^j \right), \quad (23)$$

where Φ (which depends on space and time) represents the curvature fluctuations, η denotes the conformal time, and the transverse traceless tensor, h_{ij} , stands for the gravitational waves (in writing, the perturbed metric in this form we have chosen to work in longitudinal gauge).

According to the linearized Einstein equations, the curvature fluctuations of the geometry are determined in terms of the energy density perturbations via

$$\langle |\Phi(k)|^2 \rangle = 16\pi^2 G^2 k^{-4} \langle \delta T^0_0(k) \delta T^0_0(k) \rangle, \quad (24)$$

and the gravitational waves are given by the off-diagonal pressure fluctuations,

$$\langle |h(k)|^2 \rangle = 16\pi^2 G^2 k^{-4} \langle \delta T^i_j(k) \delta T^j_i(k) \rangle \quad (i \neq j), \quad (25)$$

where G is the Newtonian constant of gravitation.

Since the energy density perturbations in a region of radius R are determined by the specific heat capacity, $C_V(R)$, on that scale, the power spectrum of curvature fluctuations is given by

$$P(k) = k^3 (\delta\Phi(k))^2 = 16\pi^2 G^2 k^2 T^2 C_V(R), \quad (26)$$

where k is the momentum scale associated with R .

For thermal fluctuations, the matter correlation functions are given by taking partial derivatives of the finite temperature partition function. Since the partition function of the BFSS matrix model is our starting point, we are able to compute the observables. Specifically, the specific heat capacity is related to the internal energy, $E(R)$, via

$$C_V(R) = \frac{\partial}{\partial T} E(R), \quad (27)$$

and the internal energy is given by the derivative of the partition function with respect to the inverse temperature, β :

$$E = -\frac{\partial}{\partial \beta} \ln Z(\beta). \quad (28)$$

The computation of the internal energy $E(R)$ was carried out in Ref. [20], based on the study [73] in the context of the IKKT matrix model. We find:

$$E^2 = N^2 \langle \mathcal{E} \rangle_{\text{BFSS}} \quad (29)$$

with

$$\mathcal{E} = -\frac{3}{4N\beta} \int_0^\beta dt \text{Tr}([X_i, X_j]^2). \quad (30)$$

Evaluating \mathcal{E} , we obtain:

$$E^2 = \frac{3}{4} N^2 \chi_2 T - \frac{3}{4} N^4 \alpha \chi_1 T^{-1/2}, \quad (31)$$

where χ_2 is a constant while χ_1 depends on R via

$$\chi_1 = \langle R^2 \rangle_{\text{BFSS}} T^{-1/2}. \quad (32)$$

The first term in Equation (31) yields a Poisson spectrum that dominates on ultraviolet scales, while the second term yields a scale-invariant contribution to the curvature power spectrum that dominates on the infrared scales relevant to current cosmological observations.

5. Challenges and Future Directions

To summarize, we have presented a proposal to obtain time, space, and a metric as emergent phenomena from the BFSS matrix model, taken in a high-temperature state. In the $N \rightarrow \infty$ limit, the time that emerges is infinite in both directions and continuous. Similarly, the emergent space is of infinite extent. Given our proposal, the background metric that emerges is spatially flat and isotropic. Since we are considering a thermal state of the BFSS model, there are thermal fluctuations, and we have reviewed here how these perturbations yield scale-invariant spectra of curvature fluctuations and gravitational waves. Hence, our cosmological model solves the famous horizon and flatness problems of standard cosmology and provides a causal mechanism for the origin of scale-invariant spectra of curvature fluctuations and gravitational waves, all without the need to postulate a phase of cosmological inflation. The cosmology is non-singular, and the cosmological constant problem is absent.

The $SO(9)$ symmetry of the BFSS action is spontaneously broken. In the case of the IKKT model, the state that minimizes the energy has $SO(6) \times SO(3)$ symmetry, with the extent of space increasing only in three spatial dimensions. Based on solving the classical matrix equations, it appears that the cosmological expansion of the three large spatial dimensions is as in a radiation-dominated universe. The presence of fermions plays a key role in obtaining this symmetry breaking. On the other hand, the fermionic sector has not been included in studying the late time dynamics of the three large spatial dimensions. An important open problem is to include the fermionic sector. A careful study of the full BFSS matrix model may also shed some light on the nature of dark energy.

An important open problem is to study the dynamics of this phase transition and to determine the key physical reason for the distinguished symmetry breaking pattern. We speculate that the reason may be related to the reason why three dimensions become large in SGC. Once the dynamics of the phase transition are understood, one may be able to make predictions for the tilts n_s and n_t of the spectra of cosmological perturbations and gravitational waves, and determine consistency relations between the tilts. One may be able to recover the same consistency relation $n_t = 1 - n_s$ as in SGC (and as in the recently proposed ekpyrotic scenario mediated by an S-brane [74]).

While our scenario yields emergent space and time, and an emergent metric, it is important to investigate what effective action describes the evolution of localized matrix excitations. At low energies, we need to be able to recover Einstein gravity coupled to regular matter and radiation.

In conclusion, although we have indicated a promising approach to the emergence of space and time, and of early universe cosmology, a lot of work needs to be done before one will be able to make successful contact with testable low-energy physics.

Author Contributions: Conceptualization, S.B., R.B. and S.L.; Methodology, S.B., R.B. and S.L.; Investigation, S.B., R.B. and S.L.; Writing—original draft, S.B., R.B. and S.L. The authors contributed equally to this work. All authors have read and agreed to the published version of the manuscript.

Funding: S.B. is supported in part by the Higgs Fellowship. S.L. is supported in part by the Fonds de recherche du Québec - Nature et technologies (FRQNT), Canada. The research at McGill is supported in part by funds from the Natural Sciences and Engineering Research Council of Canada (NSERC) and from the Canada Research Chair program.

Data Availability Statement: Not applicable.

Acknowledgments: We thank B. Ydri for permission to use Figure 1, which is taken from [42].

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Guth, A.H. The inflationary universe: A possible solution to the horizon and flatness problems. *Phys. Rev. D* **1981**, *23*, 347–356. Reprinted: *Adv. Ser. Astrophys. Cosmol.* **1987**, *3*, 139–148. [CrossRef]
2. Brout, R.; Englert, F.; Gunzig, E. The creation of the universe as a quantum phenomenon. *Ann. Phys.* **1978**, *115*, 78–106. [CrossRef]
3. Starobinsky, A.A. A New type of isotropic cosmological models without singularity. *Phys. Lett. B* **1980**, *91*, 99–102. [CrossRef]
4. Sato, K. First order phase transition of a vacuum and expansion of the universe. *Mon. Not. R. Astron. Soc.* **1981**, *195*, 467–479. [CrossRef]
5. Brandenberger, R.H. Alternatives to the inflationary paradigm of structure formation. *Int. J. Mod. Phys. Conf. Ser.* **2011**, *1*, 67–79. [CrossRef]
6. Brandenberger, R.H. Cosmology of the very early universe. *AIP Conf. Proc.* **2010**, *1268*, 3–70.
7. Brandenberger, R.; Peter, P. Bouncing cosmologies: Progress and problems. *Found. Phys.* **2017**, *47*, 797–850. [CrossRef]
8. Ooguri, H.; Vafa, C. On the geometry of the string landscape and the swampland. *Nucl. Phys. B* **2007**, *766*, 21–33. [CrossRef]
9. Obied, G.; Ooguri, H.; Spodyneiko, L.; Vafa, C. De Sitter space and the swampland. *arXiv* **2018**, arXiv:1806.08362.
10. Bedroya, A.; Vafa, C. Trans-Planckian censorship and the swampland. *J. High Energy Phys.* **2020**, *2009*, 123. [CrossRef]
11. Bedroya, A.; Brandenberger, R.; Loverde, M.; Vafa, C. Trans-Planckian censorship and inflationary cosmology. *Phys. Rev. D* **2020**, *101*, 103502. [CrossRef]
12. Brandenberger, R. String cosmology and the breakdown of local effective field theory. *arXiv* **2021**, arXiv:2112.04082.
13. Brandenberger, R.; Kamali, V. Unitarity problems for an effective field theory description of early universe cosmology. *Eur. Phys. J. C* **2022**, *82*, 818. [CrossRef] [PubMed]
14. Dvali, G.; Gomez, C.; Zell, S. Quantum break-time of de Sitter. *J. Cosmol. Astropart. Phys.* **2017**, *6*, 028. [CrossRef]
15. Dvali, G.; Gomez, C. On exclusion of positive cosmological constant. *Fortsch. Phys.* **2019**, *67*, 1800092. [CrossRef]
16. Dvali, G.; Gomez, C.; Zell, S. Quantum breaking bound on de Sitter and swampland. *Fortsch. Phys.* **2019**, *67*, 1800094. [CrossRef]
17. Brahma, S.; Dasgupta, K.; Tatar, R. Four-dimensional de Sitter space is a Glauber-Sudarshan state in string theory. *J. High Energy Phys.* **2021**, *2021*, 114. [CrossRef]
18. Brahma, S.; Dasgupta, K.; Tatar, R. de Sitter space as a Glauber-Sudarshan state. *J. High Energy Phys.* **2021**, *2021*, 104. [CrossRef]
19. Bernardo, H.; Brahma, S.; Dasgupta, K.; Tatar, R. Crisis on infinite earths: Short-lived de Sitter vacua in the string theory landscape. *J. High Energy Phys.* **2021**, *2021*, 37. [CrossRef]
20. Brahma, S.; Brandenberger, R.; Laliberte, S. Emergent cosmology from matrix theory. *J. High Energy Phys.* **2022**, *3*, 67. [CrossRef]
21. Brahma, S.; Brandenberger, R.; Laliberte, S. Emergent metric space-time from matrix theory. *J. High Energy Phys.* **2022**, *9*, 31. [CrossRef]
22. Banks, T.; Fischler, W.; Shenker, S.H.; Susskind, L. M theory as a matrix model: A Conjecture. *Phys. Rev. D* **1997**, *55*, 5112–5128. [CrossRef]
23. Hoppe, J. Quantum Theory of a Massless Relativistic Surface. Ph.D. Thesis, Massachusetts Institute of Technology (MIT), Cambridge, MA, USA, 1982. Available online: <http://dspace.mit.edu/handle/1721.1/15717> (accessed on 5 December 2022).
24. de Witt, B.; Hoppe, J.; Nicolai, H. On the quantum mechanics of supermembranes. *Nucl. Phys. B* **1988**, *305*, 545–581. [CrossRef]
25. Das, S.R.; Jevicki, A. String field theory and physical interpretation of $D = 1$ strings. *Mod. Phys. Lett. A* **1990**, *5*, 1639–1650. [CrossRef]

26. Dijkgraaf, R.; Verlinde, E.P.; Verlinde, H.L. Matrix string theory. *Nucl. Phys. B* **1997**, *500*, 43–61. [\[CrossRef\]](#)
27. Craps, B.; Sethi, S.; Verlinde, E.P. A matrix big bang. *J. High Energy Phys.* **2005**, *10*, 5. [\[CrossRef\]](#)
28. Blau, M.; O’Loughlin, M. DLCQ and plane wave matrix big bang models. *J. High Energy Phys.* **2008**, *9*, 97. [\[CrossRef\]](#)
29. Berenstein, D.E.; Maldacena, J.M.; Nastase, H.S. Strings in flat space and pp waves from $N = 4$ superYang-Mills. *J. High Energy Phys.* **2002**, *4*, 13. [\[CrossRef\]](#)
30. Maldacena, J.M.; Nunez, C. Towards the large N limit of pure $N = 1$ superYang-Mills. *Phys. Rev. Lett.* **2001**, *86*, 588–591. [\[CrossRef\]](#)
31. Craps, B.; Evnin, O. Light-like big bang singularities in string and matrix theories. *Class. Quant. Grav.* **2011**, *28*, 204006. [\[CrossRef\]](#)
32. Smolin, L. M theory as a matrix extension of Chern-Simons theory. *Nucl. Phys. B* **2000**, *591*, 227–242. [\[CrossRef\]](#)
33. Yargic, Y.; Lanier, J.; Smolin, L.; Wecker, D. A cubic matrix action for the Standard Model and beyond. *arXiv* **2022**, arXiv:2201.04183.
34. Brandenberger, R.H.; Vafa, C. Superstrings in the early universe. *Nucl. Phys. B* **1989**, *316*, 391–410. [\[CrossRef\]](#)
35. Kripfganz, J.; Perlt, H. Cosmological impact of winding strings. *Class. Quant. Grav.* **1988**, *5*, 453. [\[CrossRef\]](#)
36. Brandenberger, R.H. String gas cosmology; In *String Cosmology*; Erdmenger, J., Ed.; Wiley-VCH Verlag GmbH & Co. KGaA: Weinheim, Germany, 2009; pp. 193–230. Available online: <https://www.lehmanns.ch/shop/technik/24731252-9783527628070-string-cosmology> (accessed on 5 December 2022).
37. Hagedorn, R. Statistical thermodynamics of strong interactions at high-energies. *Nuovo Cim. Suppl.* **1965**, *3*, 147–186. Reprinted: *Quark–Gluon Plasma: Theoretical Foundations*; Kapusta, J.L., Müller, B., Rafelski, J., Eds.; North-Holland: Amsterdam, The Netherlands; 2003; pp. 24–63. Available online: <http://cds.cern.ch/record/346206> (accessed on 5 December 2022).
38. Vilenkin, A. Gravitational field of vacuum domain walls and strings. *Phys. Rev. D* **1981**, *23*, 852–857. [\[CrossRef\]](#)
39. Nayeri, A.; Brandenberger, R.H.; Vafa, C. Producing a scale-invariant spectrum of perturbations in a Hagedorn phase of string cosmology. *Phys. Rev. Lett.* **2006**, *97*, 021302. [\[CrossRef\]](#)
40. Brandenberger, R.H.; Nayeri, A.; Patil, S.P.; Vafa, C. Tensor modes from a primordial Hagedorn phase of string cosmology. *Phys. Rev. Lett.* **2007**, *98*, 231302. [\[CrossRef\]](#)
41. Agrawal, P.; Gukov, S.; Obied, G.; Vafa, C. Topological gravity as the early phase of our universe. *arXiv* **2020**, arXiv:2009.10077.
42. Ydri, B. Review of M(atrrix)-theory, type IIB matrix model and Matrix String Theory. *arXiv* **2017**, arXiv:1708.00734.
43. Taylor, W. M(atrrix) theory: Matrix quantum mechanics as a fundamental theory. *Rev. Mod. Phys.* **2001**, *73*, 419–462. [\[CrossRef\]](#)
44. Ishibashi, N.; Kawai, H.; Kitazawa, Y.; Tsuchiya, A. A large- N reduced model as superstring. *Nucl. Phys. B* **1997**, *498*, 467–491. [\[CrossRef\]](#)
45. Aoki, H.; Iso, S.; Kawai, H.; Kitazawa, Y.; Tada, T. Space-time structures from IIB matrix model. *Prog. Theor. Phys.* **1998**, *99*, 713–746. [\[CrossRef\]](#)
46. Kim, S.W.; Nishimura, J.; Tsuchiya, A. Expanding (3+1)-dimensional universe from a Lorentzian matrix model for superstring theory in (9+1)-dimensions. *Phys. Rev. Lett.* **2012**, *108*, 011601. [\[CrossRef\]](#) [\[PubMed\]](#)
47. Nishimura, J.; Vernizzi, G. Spontaneous breakdown of Lorentz invariance in IIB matrix model. *J. High Energy Phys.* **2000**, *4*, 15. [\[CrossRef\]](#)
48. Ambjorn, J.; Anagnostopoulos, K.N.; Bietenholz, W.; Hotta, T.; Nishimura, J. Monte Carlo studies of the IIB matrix model at large N . *J. High Energy Phys.* **2000**, *7*, 11. [\[CrossRef\]](#)
49. Nishimura, J.; Sugino, F. Dynamical generation of four-dimensional space-time in the IIB matrix model. *J. High Energy Phys.* **2002**, *5*, 1. [\[CrossRef\]](#)
50. Kawai, H.; Kawamoto, S.; Kuroki, T.; Matsuo, T.; Shinohara, S. Mean field approximation of IIB matrix model and emergence of four-dimensional space-time. *Nucl. Phys. B* **2002**, *647*, 153–189. [\[CrossRef\]](#)
51. Aoyama, T.; Nishimura, J.; Okubo, T. Spontaneous breaking of the rotational symmetry in dimensionally reduced super Yang-Mills models. *Prog. Theor. Phys.* **2011**, *125*, 537–563. [\[CrossRef\]](#)
52. Nishimura, J.; Okubo, T.; Sugino, F. Systematic study of the SO(10) symmetry breaking vacua in the matrix model for type IIB superstrings. *J. High Energy Phys.* **2011**, *10*, 135. [\[CrossRef\]](#)
53. Kabat, D.N.; Lifschytz, G.; Lowe, D.A. Black hole thermodynamics from calculations in strongly coupled gauge theory. *Int. J. Mod. Phys. A* **2001**, *16*, 856–865. [\[CrossRef\]](#)
54. Kabat, D.N.; Lifschytz, G.; Lowe, D.A. Black hole entropy from nonperturbative gauge theory. *Phys. Rev. D* **2001**, *64*, 124015. [\[CrossRef\]](#)
55. Watson, S.; Brandenberger, R. Stabilization of extra dimensions at tree level. *J. Cosmol. Astrpart. Phys.* **2003**, *11*, 8. [\[CrossRef\]](#)
56. Patil, S.P.; Brandenberger, R.H. The cosmology of massless string modes. *J. Cosmol. Astrpart. Phys.* **2006**, *1*, 5. [\[CrossRef\]](#)
57. Brahma, S.; Brandenberger, R.; Laliberte, S. Spontaneous symmetry breaking in the BFSS model: Analytical results using the Gaussian expansion method. *arXiv* **2022**, arXiv:2209.01255.
58. Nishimura, J. New perspectives on the emergence of (3+1)D expanding space-time in the Lorentzian type IIB matrix model. *PoS* **2020**, CORFU2019, 178.
59. Hirasawa, M.; Anagnostopoulos, K.; Azuma, T.; Hatakeyama, K.; Ito, Y.; Nishimura, J.; Papadoudis, S.; Tsuchiya, A. A new phase in the Lorentzian type IIB matrix model and the emergence of continuous space-time. *PoS* **2022**, LATTICE2021, 428.
60. Hatakeyama, K.; Anagnostopoulos, K.; Azuma, T.; Hirasawa, M.; Ito, Y.; Nishimura, J.; Papadoudis, S.; Tsuchiya, A. Relationship between the Euclidean and Lorentzian versions of the type IIB matrix model. *PoS* **2022**, LATTICE2021, 341.

61. Hatakeyama, K.; Anagnostopoulos, K.; Azuma, T.; Hirasawa, M.; Ito, Y.; Nishimura, J.; Papadoudis, S.; Tsuchiya, A. Complex Langevin studies of the emergent space-time in the type IIB matrix model. In *Proceedings of the East Asia Joint Symposium on Fields and Strings 2021*; Iso, S., Itoyama, H., Maruyoshi, K., Nishinaka, T., Oota, T., Sakai, K., Tsuchiya, A., Eds.; World Scientific Co. Ltd.: Singapore, 2022; pp. 9–18. [\[CrossRef\]](#)
62. Khoury, J.; Ovrut, B.A.; Steinhardt, P.J.; Turok, N. The ekpyrotic universe: Colliding branes and the origin of the hot big bang. *Phys. Rev. D* **2001**, *64*, 123522. [\[CrossRef\]](#)
63. Khoury, J.; Ovrut, B.A.; Seiberg, N.; Steinhardt, P.J.; Turok, N. From big crunch to big bang. *Phys. Rev. D* **2002**, *65*, 086007. [\[CrossRef\]](#)
64. Kim, S.W.; Nishimura, J.; Tsuchiya, A. Late time behaviors of the expanding universe in the IIB matrix model. *J. High Energy Phys.* **2012**, *10*, 147. [\[CrossRef\]](#)
65. Ito, Y.; Nishimura, J.; Tsuchiya, A. Power-law expansion of the Universe from the bosonic Lorentzian type IIB matrix model. *J. High Energy Phys.* **2015**, *11*, 70. [\[CrossRef\]](#)
66. Hatakeyama, K.; Matsumoto, A.; Nishimura, J.; Tsuchiya, A.; Yosprakob, A. The emergence of expanding space–time and intersecting D-branes from classical solutions in the Lorentzian type IIB matrix model. *Prog. Theor. Exp. Phys.* **2020**, *2020*, 43B10. [\[CrossRef\]](#)
67. Wetterich, C. Cosmology and the fate of dilatation symmetry. *Nucl. Phys. B* **1988**, *302*, 668–696. [\[CrossRef\]](#)
68. Peebles, P.J.E.; Ratra, B. Cosmology with a time variable cosmological constant. *Astrophys. J.* **1988**, *325*, L17–L20. Available online: <https://articles.adsabs.harvard.edu/full/1988ApJ...325L..17P> (accessed on 5 December 2022). [\[CrossRef\]](#)
69. Ratra, B.; Peebles, P.J.E. Cosmological consequences of a rolling homogeneous scalar field. *Phys. Rev. D* **1988**, *37*, 3406–3427. [\[CrossRef\]](#)
70. Bernardo, H.; Brandenberger, R.; Fröhlich, J. Towards a dark sector model from string theory. *J. Cosmol. Astrpart. Phys.* **2022**, *9*, 40. [\[CrossRef\]](#)
71. Mukhanov, V.F.; Feldman, H.A.; Brandenberger, R.H. Theory of cosmological perturbations. *Phys. Rep.* **1992**, *215*, 203–333. [\[CrossRef\]](#)
72. Brandenberger, R. Lectures on the theory of cosmological perturbations. In *The Early Universe and Observational Cosmology*; Bretón, N., Cervantes-Cota, J.L., Salgado, M., Eds.; Springer: Berlin/Heidelberg, Germany, 2004; pp. 127–167. [\[CrossRef\]](#)
73. Kawahara, N.; Nishimura, J.; Takeuchi, S. High temperature expansion in supersymmetric matrix quantum mechanics. *J. High Energy Phys.* **2007**, *12*, 103. [\[CrossRef\]](#)
74. Brandenberger, R.; Wang, Z. Nonsingular ekpyrotic cosmology with a nearly scale-invariant spectrum of cosmological perturbations and gravitational waves. *Phys. Rev. D* **2020**, *101*, 063522. [\[CrossRef\]](#)

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.