

## Supporting Information

# Activated Carbon from Sugarcane Bagasse: A Low-Cost Approach towards Cr(VI) Removal from Wastewater

Rana Ahmed<sup>1</sup>, Inga Block<sup>1</sup>, Fabian Otte<sup>1</sup>, Christina Günter<sup>2</sup>, Alysson Duarte-Rodrigues<sup>3</sup>, Peter Hesemann<sup>3</sup>, Amitabh Banerji<sup>1</sup> and Andreas Taubert<sup>1,\*</sup>

<sup>1</sup> Institute of Chemistry, University of Potsdam, D-14476 Potsdam, Germany

<sup>2</sup> Institute of Geosciences, University of Potsdam, D-14476 Potsdam, Germany

<sup>3</sup> ICGM, Université de Montpellier-CNRS-ENSCM, 34095 Montpellier, France


\* Correspondence: ataubert@uni-potsdam.de; Tel.: +49-(0)331-977-5773.

### Contents:


#### IR-Spectra of SB1 to SB7

S2	raw sugar cane bagasse (raw SCB)
S3	SB1
S4	SB2
S5	SB3
S6	SB4
S7	SB5
S8	SB6
S9	SB7


Raw sugar cane bagasse (raw SCB)




SB1




SB2




SB3




SB4



SB5



SB6





SB7

