

Supplementary material

Table S1. Reported diagnoses (multiple diagnoses selections were allowed-not mutually exclusive).

Disorder	N	%
Depression	282	34.06
Generalized anxiety disorder	277	33.45
Skin Picking	90	10.87
Tinnitus	80	9.66
Social anxiety disorder	77	9.30
Obsessive-Compulsive Disorder (OCD)	67	8.09
Panic Disorder	37	4.47
Do you have a diagnosis of a substance abuse disorder?	34	4.11
Attention deficit disorder (ADD)	30	3.62
Attention deficit hyperactivity disorder (ADHD)	28	3.38
Autism spectrum disorder (ASD)	25	3.02
Binge eating disorder	25	3.02
Bipolar disorder	22	2.66
Persistent depressive disorder (dysthymia)	22	2.66
Dyslexia	21	2.54
Hyperacusis	20	2.42
Borderline personality disorder	17	2.05
Anorexia nervosa	16	1.93
Bulimia	15	1.81
Trichotillomania (hair pulling)	15	1.81
Obsessive-compulsive personality disorder	13	1.57
Auditory processing disorder	10	1.21
Dyspraxia/developmental coordination disorder	6	0.72
Antisocial personality disorder	5	0.60
Narcissistic personality disorder	1	0.12

Table S2. Descriptive indices of measures-complete sample.

Scale	N	Mean (SD)	Mode (min-max)	Median (Q1-Q3)
A-MISO-S total score	737	9.1 (6.3)	0 (0 - 26)	8 (4 - 14)
GAD7 total score	1582	6.9 (5.8)	0 (0 - 21)	6 (2 - 11)
MQ: Misophonia Symptom Scale	576	14.9 (6.9)	13 (0 - 32)	15 (10 - 20)
MQ: Misophonia Emotions and Behaviours	567	17.1 (9.7)	19 (0 - 41)	18 (9 - 25)
MQ: Misophonia Severity	696	5.1 (3.5)	4a (0 - 15)	5 (2 - 7)
MQ total	576	31.7 (15.4)	34 (0 - 65)	33 (19 - 45)
PHQ9 total score	1561	8.1 (6.4)	0 (0 - 27)	7 (3 - 12)
WSAS total score	831	15.7 (9)	18 (0 - 40)	16 (9 - 22)