

Article

# Einstein's $E = mc^2$ Derivable from Heisenberg's Uncertainty Relations

Sibel Başkal <sup>1</sup>, Young S. Kim <sup>2,\*</sup>, and Marilyn E. Noz <sup>3</sup><sup>1</sup> Department of Physics, Middle East Technical University, 06800 Ankara, Turkey; sibelbaskal@gmail.com<sup>2</sup> Center for Fundamental Physics, University of Maryland, College Park, MD 20742, USA<sup>3</sup> Department of Radiology, New York University, New York, NY 10016, USA; marilyne.noz@gmail.com

\* Correspondence: yskim@umd.edu; Tel.: +1-301-937-1306

Received: 12 September 2019; Accepted: 7 November 2019; Published: 9 November 2019



**Abstract:** Heisenberg's uncertainty relation can be written in terms of the step-up and step-down operators in the harmonic oscillator representation. It is noted that the single-variable Heisenberg commutation relation contains the symmetry of the  $Sp(2)$  group which is isomorphic to the Lorentz group applicable to one time-like dimension and two space-like dimensions, known as the  $O(2, 1)$  group. This group has three independent generators. The one-dimensional step-up and step-down operators can be combined into one two-by-two Hermitian matrix which contains three independent operators. If we use a two-variable Heisenberg commutation relation, the two pairs of independent step-up, step-down operators can be combined into a four-by-four block-diagonal Hermitian matrix with six independent parameters. It is then possible to add one off-diagonal two-by-two matrix and its Hermitian conjugate to complete the four-by-four Hermitian matrix. This off-diagonal matrix has four independent generators. There are thus ten independent generators. It is then shown that these ten generators can be linearly combined to the ten generators for Dirac's two oscillator system leading to the group isomorphic to the de Sitter group  $O(3, 2)$ , which can then be contracted to the inhomogeneous Lorentz group with four translation generators corresponding to the four-momentum in the Lorentz-covariant world. This Lorentz-covariant four-momentum is known as Einstein's  $E = mc^2$ .

**Keywords:**  $E = mc^2$  from Heisenberg's uncertainty relations; one symmetry for quantum mechanics and special relativity

## 1. Introduction

Let us start with Heisenberg's commutation relations

$$[x_i, P_j] = i \delta_{ij}, \quad (1)$$

with

$$P_i = -i \frac{\partial}{\partial x_i}, \quad (2)$$

where  $i = 1, 2, 3$ , corresponds to the  $x, y, z$  coordinates respectively.

With these  $x_i$  and  $P_i$ , we can construct the following three operators,

$$J_i = \epsilon_{ijk} x_j P_k. \quad (3)$$

These three operators satisfy the closed set of commutation relations:

$$[J_i, J_j] = i \epsilon_{ijk} J_k. \quad (4)$$

These  $J_i$  operators generate rotations in the three-dimensional space. In mathematics, this set is called the Lie algebra of the rotation group. This is a direct consequence of Heisenberg's commutation relations.

In quantum mechanics, each  $J_i$  corresponds to the angular momentum along the  $i$  direction. A remarkable fact is that it is also possible to construct the same Lie algebra with two-by-two matrices. These matrices are of course the Pauli spin matrices, leading to the observable angular momentum not seen in classical mechanics.

As the expression shows in Equation (2), each  $P_i$  generates a translation along the  $i^{\text{th}}$  direction. Thus, the three translation generators, together with the three rotation generators constitute the Lie algebra of the Galilei group, with the additional commutation relations:

$$[J_i, P_j] = i\epsilon_{ijk}P_k. \quad (5)$$

This set of commutation relations together with those of Equation (4) constitute a closed set for both  $P_i$  and  $J_i$ . This set is called the Lie algebra of the Galilei group. This group is the basic symmetry group for the Schrödinger or non-relativistic quantum mechanics.

In the Schrödinger picture, the generator  $P_i$  corresponds to the particle momentum along the  $i$  direction. In addition, the time translation operator is

$$P_0 = i\frac{\partial}{\partial t}. \quad (6)$$

This corresponds to the energy variable.

Let us go to the Lorentzian world. Here we have to take into account the generators of the boosts. The generators thus include the time variable, and the generator of boosts along the  $i$  direction is

$$K_i = i\left(x_i\frac{\partial}{\partial t} + t\frac{\partial}{\partial x_i}\right). \quad (7)$$

These generators satisfy the commutation relations

$$[K_i, K_j] = -i\epsilon_{ijk}J_k. \quad (8)$$

Thus, these three boost generators alone cannot constitute a closed set of commutation relations (Lie algebra).

With  $J_i$ , these boost generators satisfy

$$[J_i, K_j] = i\epsilon_{ijk}K_k. \quad (9)$$

With  $P_i$ , they satisfy the relation

$$[P_i, K_i] = i\delta_{0i}P_0. \quad (10)$$


Thus, the commutation relations of Equations (4),(5),(8–10) constitute a closed set of the ten generators. This closed set is commonly called the Lie algebra of the Poincaré symmetry.

The three rotation and three translation generators are contained in, or are derivable from, Heisenberg's commutation relations, and the time translation operator is seen in the Schrödinger equation. They are all Hermitian operators corresponding to dynamical variables. On the other hand, the three boost generators of Equation (7) are not derivable from the Heisenberg relations. Furthermore, they do not appear to correspond to observable quantities [1].

The purpose of this paper is to show that the Lie algebra of the Poincaré symmetry is derivable from the Heisenberg commutation relations. For this purpose, we first examine the symmetry of the Heisenberg commutation relation using the Wigner function in the phase space. It is noted that the

single-variable relation contains the symmetry of the Lorentz group applicable to two space-like and one time-like dimensions.

As Dirac noted in 1963 [2], two coupled oscillators lead to the symmetry of the  $O(3,2)$  or the Lorentz group applicable to the three space-like directions and two time-like directions. As is illustrated in Figure 1, it is possible to contract one of those two time variables of this  $O(3,2)$  group into the inhomogeneous Lorentz group, consisting of the Lorentz group applicable to the three space-like dimensions and one time-like direction, plus four translation generators corresponding to the energy-momentum four-vector. This of course leads to Einstein’s energy–momentum relation of  $E = mc^2$ .



**Figure 1.** The Inönü–Wigner contraction procedure interpreted as squeeze transformations. In (a), the square becomes a narrow rectangle during the squeeze process. When the rectangle becomes narrow enough, the point A can be moved to the horizontal axis. Then, the inverse squeeze brings back the rectangle to the original shape. The point A remains on the horizontal axis. In (b), both the hyperbola and the circle become flattened to the horizontal axis, during the initial squeeze. The point on the curve moves to the horizontal axis. This point moves back to its finite position during the inverse squeeze.

In Section 2, it is noted that the best way to study the symmetry of the Heisenberg commutation relation is to use the Wigner function for the Gaussian function for the oscillator state. In the Wigner phase space, this function contains the symmetry for the Lorentz group applicable to two space-like dimensions and one time-like dimension. This group has three generators. This operation is equivalent to constructing a two-by-two block-diagonal Hermitian matrix with quadratic forms of the step-up and step-down operators.


In Section 3, we consider two oscillators. If these oscillators are independent, it is possible to construct a four-by-four block diagonal matrix, where each block consists of the two-by-two matrix for each operator defined in Section 2. Since the oscillators are uncoupled, this four-by-four block-diagonal Hermitian matrix contains six independent generators.

If the oscillators are coupled, then to keep the overall four-by-four block-diagonal matrix Hermitian, we need one off-diagonal block matrix, with four independent quadratic forms. Thus, the overall four-by-four matrix contains ten independent quadratic forms of the creation and annihilation operators.

It is shown that these ten independent generators can be linearly combined into the ten generators constructed by Dirac for the the Lorentz group applicable to three space-like dimensions and two time-like dimensions, commonly called the  $O(3,2)$  group.

In Section 4, using the boosts belonging to one of its time-like dimensions, we contract  $O(3,2)$  to produce the Lorentz group applicable to one time dimension and four translations leading to the four-momentum. This Lorentz-covariant four-momentum is commonly known as Einstein’s  $E = mc^2$ .

This paper is essentially based on Dirac’s paper published in 1949 and 1963 [1,2]. As is illustrated in Figure 2, we show here that the space-time symmetry of quantum mechanics mentioned in his 1949 paper is derivable from his two-oscillator system discussed in 1963. The route is the group contraction procedure of Inönü and Wigner [3].



**Figure 2.** According to Dirac’s 1949 paper, the task of constructing quantum mechanics is essentially constructing a representation of the inhomogeneous Lorentz group. In his 1963 paper, Dirac constructed the Lie algebra of the  $O(3,2)$  de Sitter group from the algebra of two harmonic oscillators, which is a direct consequence of Heisenberg’s uncertainty commutation relations. It is possible to derive the Lie algebra of the inhomogeneous Lorentz group from that of  $O(3,2)$  using the group-contraction procedure of Inönü and Wigner [3].

Indeed, from 1927 Dirac made lifelong efforts to synthesize quantum mechanics and special relativity [4]. In 1949 and before, he treated quantum mechanics and special relativity as two separate scientific disciplines, and then in 1949 he attempted to synthesize them. Thus, it is of interest to see how Dirac’s ideas evolved during the period 1929–1949. We shall give a brief review of Dirac’s efforts during the period in Appendix A.

## 2. Symmetries of the Single-Mode States

Heisenberg’s uncertainty relation for a single Cartesian variable takes the form

$$[x, p] = i \tag{11}$$

with

$$p = -i \frac{\partial}{\partial x}.$$

Very often, it is more convenient to use the operators

$$a = \frac{1}{\sqrt{2}}(x + ip), \quad a^\dagger = \frac{1}{\sqrt{2}}(x - ip) \tag{12}$$

with

$$[a, a^\dagger] = 1. \tag{13}$$

This aspect is well known.

The representation based on  $a$  and  $a^\dagger$  is known as the harmonic oscillator representation of the uncertainty relation and is the basic language for the Fock space for particle numbers. This representation is therefore the basic language for quantum optics.

Let us next consider the quadratic forms:  $aa, a^\dagger a^\dagger, aa^\dagger$ , and  $a^\dagger a$ . Then the linear combination

$$aa^\dagger - a^\dagger a = 1, \tag{14}$$

according to the uncertainty relation. Thus, there are three independent quadratic forms, and we are led to the following two-by-two matrix:

$$\begin{pmatrix} (aa^\dagger + a^\dagger a) / 2 & aa \\ a^\dagger a^\dagger & (aa^\dagger + a^\dagger a) / 2 \end{pmatrix}. \quad (15)$$

This matrix leads to the following three independent operators:

$$J_2 = \frac{1}{2} (aa^\dagger + a^\dagger a), \quad K_1 = \frac{1}{2} (a^\dagger a^\dagger + aa), \quad K_3 = \frac{i}{2} (a^\dagger a^\dagger - aa). \quad (16)$$

They produce the following set of closed commutation relations:

$$[J_2, K_1] = -iK_3, \quad [J_2, K_3] = iK_1, \quad [K_1, K_3] = iJ_2. \quad (17)$$

This set is commonly called the Lie algebra of the  $Sp(2)$  group, locally isomorphic to the Lorentz group applicable to one time and two space coordinates.

The best way to study the symmetry property of these operators is to use the Wigner function for the ground-state oscillator which takes the form [5–8]

$$W(x, p) = \frac{1}{\pi} \exp \left[ - (x^2 + p^2) \right]. \quad (18)$$

This distribution is concentrated in the circular region around the origin. Let us define the circle as

$$x^2 + p^2 = 1. \quad (19)$$

We can use the area of this circle in the phase space of  $x$  and  $p$  as the minimum uncertainty. This uncertainty is preserved under rotations in the phase space and also under squeezing. These transformations can be written as

$$\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} x \\ p \end{pmatrix}, \quad \begin{pmatrix} e^\eta & 0 \\ 0 & e^{-\eta} \end{pmatrix} \begin{pmatrix} x \\ p \end{pmatrix}, \quad (20)$$

respectively. The rotation and the squeeze are generated by

$$J_2 = -i \left( x \frac{\partial}{\partial p} - p \frac{\partial}{\partial x} \right), \quad K_1 = -i \left( x \frac{\partial}{\partial x} - p \frac{\partial}{\partial p} \right). \quad (21)$$


If we take the commutation relation with these two operators, the result is

$$[J_2, K_1] = -iK_3, \quad (22)$$

with

$$K_3 = -i \left( x \frac{\partial}{\partial p} + p \frac{\partial}{\partial x} \right). \quad (23)$$

Indeed, as before, these three generators form the closed set of commutation which form the Lie algebra of the  $Sp(2)$  group, isomorphic to the Lorentz group applicable to two space and one time dimensions. This isomorphic correspondence is illustrated in Figure 3.



**Figure 3.** Rotations and squeezes in the phase space produced by the  $Sp(2)$  transformations. The squeeze along the  $x$  direction corresponds to the Lorentz boost along the  $z$  direction, while the squeeze along the  $45^\circ$  angle corresponds to the boost along the  $x$  direction. The rotation by  $45^\circ$  corresponds to the rotation by  $90^\circ$  around the  $y$  axis.

Let us consider the Minkowski space of  $(x, y, z, t)$ . It is possible to write three four-by-four matrices satisfying the Lie algebra of Equation (17). The three four-by-four matrices satisfying this set of commutation relations are:

$$J_2 = \begin{pmatrix} 0 & 0 & i & 0 \\ 0 & 0 & 0 & 0 \\ i & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad K_1 = \begin{pmatrix} 0 & 0 & 0 & i \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ i & 0 & 0 & 0 \end{pmatrix}, \quad K_3 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & i \\ 0 & 0 & i & 0 \end{pmatrix}. \quad (24)$$

However, these matrices have null second rows and null second columns. Thus, they can generate Lorentz transformations applicable only to the three-dimensional space of  $(x, z, t)$ , while the  $y$  variable remains invariant. Thus, this single-oscillator system cannot describe what happens in the full four-dimensional Minkowski space.

Yet, it is interesting that the oscillator system can produce three different representations sharing the same Lie algebra with the  $(2 + 1)$ -dimensional Lorentz group, as shown in Table 1.

**Table 1.** Transformation for the Gaussian function, in terms of harmonic oscillators, two-dimensional phase space, and the four-dimensional Minkowski space.

Generators	Oscillator	Phase Space	Lorentz
$J_2$	$\frac{1}{2} (aa^\dagger + a^\dagger a)$	$\frac{1}{2}\sigma_2$	$\begin{pmatrix} 0 & 0 & i & 0 \\ 0 & 0 & 0 & 0 \\ i & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$
$K_1$	$\frac{1}{2i} (a^\dagger a^\dagger + aa)$	$\frac{i}{2}\sigma_1$	$\begin{pmatrix} 0 & 0 & 0 & i \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ i & 0 & 0 & 0 \end{pmatrix}$
$K_3$	$\frac{1}{2} (a^\dagger a^\dagger - aa)$ ,	$\frac{i}{2}\sigma_3$	$\begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & i \\ 0 & 0 & i & 0 \end{pmatrix}$

### 3. Symmetries from Two Oscillators

In order to generate Lorentz transformations applicable to the full Minkowskian space, we may need two Heisenberg commutation relations. Indeed, Paul A. M. Dirac started this program in 1963 [2]. It is possible to write the two uncertainty relations using two harmonic oscillators as

$$[a_i, a_j^\dagger] = \delta_{ij}, \tag{25}$$

with

$$a_i = \frac{1}{\sqrt{2}} (x_i + ip_i), \quad a_i^\dagger = \frac{1}{\sqrt{2}} (x_i - ip_i), \tag{26}$$

and

$$x_i = \frac{1}{\sqrt{2}} (a_i + a_i^\dagger), \quad p_i = \frac{i}{\sqrt{2}} (a_i^\dagger - a_i), \tag{27}$$

where  $i$  and  $j$  could be 1 or 2.

As in the case of the two-by-two matrix given in Equation (15), we can consider the following four-by-four block-diagonal matrix if the oscillators are not coupled:

$$\begin{pmatrix} (a_1 a_1^\dagger + a_1^\dagger a_1) / 2 & a_1 a_1 & 0 & 0 \\ a_1^\dagger a_1^\dagger & (a_1 a_1^\dagger + a_1^\dagger a_1) / 2 & 0 & 0 \\ 0 & 0 & (a_2 a_2^\dagger + a_2^\dagger a_2) / 2 & a_2 a_2 \\ 0 & 0 & a_2^\dagger a_2^\dagger & (a_2 a_2^\dagger + a_2^\dagger a_2) / 2 \end{pmatrix}. \tag{28}$$

There are six generators in this matrix.

We are now interested in coupling them by filling in the off-diagonal blocks. The most general forms for this block are the following two-by-two matrix and its Hermitian conjugate:

$$\begin{pmatrix} a_1^\dagger a_2 & a_1 a_2 \\ a_1^\dagger a_2^\dagger & a_1 a_2^\dagger \end{pmatrix} \tag{29}$$

with four independent generators. This leads to the following four-by-four matrix with ten (6 + 4) generators:

$$\begin{pmatrix} (a_1 a_1^\dagger + a_1^\dagger a_1) / 2 & a_1 a_1 & a_1^\dagger a_2 & a_1 a_2 \\ a_1^\dagger a_1^\dagger & (a_1 a_1^\dagger + a_1^\dagger a_1) / 2 & a_1^\dagger a_2^\dagger & a_1 a_2^\dagger \\ a_1 a_2^\dagger & a_1 a_2 & (a_2 a_2^\dagger + a_2^\dagger a_2) / 2 & a_2 a_2 \\ a_1^\dagger a_2^\dagger & a_1^\dagger a_2 & a_2^\dagger a_2^\dagger & (a_2 a_2^\dagger + a_2^\dagger a_2) / 2 \end{pmatrix}. \tag{30}$$

With these ten elements, we can now construct the following four rotation-like generators:

$$\begin{aligned} J_1 &= \frac{1}{2} (a_1^\dagger a_2 + a_2^\dagger a_1), & J_2 &= \frac{1}{2i} (a_1^\dagger a_2 - a_2^\dagger a_1), \\ J_3 &= \frac{1}{2} (a_1^\dagger a_1 - a_2^\dagger a_2), & S_0 &= \frac{1}{2} (a_1^\dagger a_1 + a_2 a_2^\dagger), \end{aligned} \tag{31}$$

and six squeeze-like generators:

$$\begin{aligned}
 K_1 &= -\frac{1}{4} \left( a_1^\dagger a_1^\dagger + a_1 a_1 - a_2^\dagger a_2^\dagger - a_2 a_2 \right), \\
 K_2 &= +\frac{i}{4} \left( a_1^\dagger a_1^\dagger - a_1 a_1 + a_2^\dagger a_2^\dagger - a_2 a_2 \right), \\
 K_3 &= +\frac{1}{2} \left( a_1^\dagger a_2^\dagger + a_1 a_2 \right),
 \end{aligned} \tag{32}$$

and

$$\begin{aligned}
 Q_1 &= -\frac{i}{4} \left( a_1^\dagger a_1^\dagger - a_1 a_1 - a_2^\dagger a_2^\dagger + a_2 a_2 \right), \\
 Q_2 &= -\frac{1}{4} \left( a_1^\dagger a_1^\dagger + a_1 a_1 + a_2^\dagger a_2^\dagger + a_2 a_2 \right), \\
 Q_3 &= +\frac{i}{2} \left( a_1^\dagger a_2^\dagger - a_1 a_2 \right).
 \end{aligned} \tag{33}$$

There are now ten operators from Equations (31)–(33), and they satisfy the following Lie algebra as was noted by Dirac in 1963 [2]:

$$\begin{aligned}
 [J_i, J_j] &= i\epsilon_{ijk} J_k, & [J_i, K_j] &= i\epsilon_{ijk} K_k, \\
 [J_i, Q_j] &= i\epsilon_{ijk} Q_k, & [K_i, K_j] &= [Q_i, Q_j] = -i\epsilon_{ijk} J_k, \\
 [K_i, Q_j] &= -i\delta_{ij} S_0, & [J_i, S_0] &= 0, & [K_i, S_0] &= -iQ_i, & [Q_i, S_0] &= iK_i.
 \end{aligned} \tag{34}$$

Dirac noted that this set is the same as the Lie algebra for the  $O(3, 2)$  de Sitter group, with ten generators. This is the Lorentz group applicable to the three-dimensional space with two time variables. This group plays a very important role in space-time symmetries.

In the same paper, Dirac pointed out that this set of commutation relations serves as the Lie algebra for the four-dimensional symplectic group commonly called  $Sp(4)$ . For a dynamical system consisting of two pairs of canonical variables  $x_1, p_1$  and  $x_2, p_2$ , we can use the four-dimensional phase space with the coordinate variables defined as [9]

$$(x_1, p_1, x_2, p_2). \tag{35}$$

Then the four-by-four transformation matrix  $M$  applicable to this four-component vector is canonical if [10,11]

$$MJ\tilde{M} = J, \tag{36}$$

where  $\tilde{M}$  is the transpose of the  $M$  matrix, with

$$J = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix}, \tag{37}$$

which we can write in the block-diagonal form as

$$J = i \begin{pmatrix} I & 0 \\ 0 & I \end{pmatrix} \sigma_2, \tag{38}$$

where  $I$  is the unit two-by-two matrix.

According to this form of the  $J$  matrix, the area of the phase space for the  $x_1$  and  $p_1$  variables remains invariant, and the story is the same for the phase space of  $x_2$  and  $p_2$ .

We can then write the generators of the  $Sp(4)$  group as [12]

$$J_1 = -\frac{1}{2} \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix} \sigma_2, \quad J_2 = \frac{i}{2} \begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix} I, \quad J_3 = \frac{1}{2} \begin{pmatrix} -I & 0 \\ 0 & I \end{pmatrix} \sigma_2, \quad S_0 = \frac{1}{2} \begin{pmatrix} I & 0 \\ 0 & I \end{pmatrix} \sigma_2, \quad (39)$$

and

$$K_1 = \frac{i}{2} \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix} \sigma_1, \quad K_2 = \frac{i}{2} \begin{pmatrix} I & 0 \\ 0 & I \end{pmatrix} \sigma_3, \quad K_3 = -\frac{i}{2} \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix} \sigma_1, \\ Q_1 = -\frac{i}{2} \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix} \sigma_3, \quad Q_2 = \frac{i}{2} \begin{pmatrix} I & 0 \\ 0 & I \end{pmatrix} \sigma_1, \quad Q_3 = \frac{i}{2} \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix} \sigma_3. \quad (40)$$

Among these ten matrices, six of them are in block-diagonal form. They are  $S_0, J_3, K_1, K_2, Q_1,$  and  $Q_2$ . In the language of two harmonic oscillators, these generators do not mix up the first and second oscillators. There are six of them because each operator has three generators for its own  $Sp(2)$  symmetry. These generators, together with those in the oscillator representation, are tabulated in Table 2.

**Table 2.** Transformation generators for the two-oscillator system.

Generators	Two Oscillators	Phase Space
$J_1$	$\frac{1}{2} (a_1^\dagger a_2 + a_2^\dagger a_1)$	$-\frac{1}{2} \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix} \sigma_2$
$J_2$	$\frac{1}{2i} (a_1^\dagger a_2 - a_2^\dagger a_1)$	$\frac{i}{2} \begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix} I$
$J_3$	$\frac{1}{2} (a_1^\dagger a_1 - a_2^\dagger a_2),$	$\frac{1}{2} \begin{pmatrix} -I & 0 \\ 0 & I \end{pmatrix} \sigma_2$
$S_0$	$\frac{1}{2} (a_1^\dagger a_1 + a_2^\dagger a_2),$	$\frac{1}{2} \begin{pmatrix} I & 0 \\ 0 & I \end{pmatrix} \sigma_2$
$K_1$	$-\frac{1}{4} (a_1^\dagger a_1^\dagger + a_1 a_1 - a_2^\dagger a_2^\dagger - a_2 a_2)$	$\frac{i}{2} \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix} \sigma_1$
$K_2$	$+\frac{i}{4} (a_1^\dagger a_1^\dagger - a_1 a_1 + a_2^\dagger a_2^\dagger - a_2 a_2)$	$\frac{i}{2} \begin{pmatrix} I & 0 \\ 0 & I \end{pmatrix} \sigma_3$
$K_3$	$\frac{1}{2} (a_1^\dagger a_2^\dagger + a_1 a_2)$	$-\frac{i}{2} \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix} \sigma_1$
$Q_1$	$-\frac{i}{4} (a_1^\dagger a_1^\dagger - a_1 a_1 - a_2^\dagger a_2^\dagger + a_2 a_2)$	$-\frac{i}{2} \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix} \sigma_3$
$Q_2$	$-\frac{1}{4} (a_1^\dagger a_1^\dagger + a_1 a_1 + a_2^\dagger a_2^\dagger + a_2 a_2)$	$\frac{i}{2} \begin{pmatrix} I & 0 \\ 0 & I \end{pmatrix} \sigma_1$
$Q_3$	$\frac{i}{2} (a_1^\dagger a_2^\dagger - a_1 a_2)$	$\frac{1}{2} \begin{pmatrix} I & 0 \\ 0 & I \end{pmatrix} \sigma_2$

The off-diagonal matrix  $J_2$  couples the first and second oscillators without changing the overall volume of the four-dimensional phase space. However, in order to construct the closed set of commutation relations, we need the three additional generators:  $J_1, K_3$ , and  $Q_3$ . The commutation relations given in Equations (34) are clearly consequences of Heisenberg’s uncertainty relations.

As for the  $O(3, 2)$  group, the generators are five-by-five matrices, applicable to  $(x, y, z, t, s)$ , where  $t$  and  $s$  are time-like variables. These matrices can be written as

$$\begin{aligned}
 J_1 &= \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -i & 0 & 0 \\ 0 & i & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}, & J_2 &= \begin{pmatrix} 0 & 0 & i & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ -i & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}, & J_3 &= \begin{pmatrix} 0 & -i & 0 & 0 & 0 \\ i & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \\
 K_1 &= \begin{pmatrix} 0 & 0 & 0 & i & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ i & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}, & K_2 &= \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & i & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & i & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}, & K_3 &= \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & i & 0 \\ 0 & 0 & i & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \\
 Q_1 &= \begin{pmatrix} 0 & 0 & 0 & 0 & i \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ i & 0 & 0 & 0 & 0 \end{pmatrix}, & Q_2 &= \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & i \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & i & 0 & 0 & 0 \end{pmatrix}, & Q_3 &= \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & i \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & i & 0 & 0 \end{pmatrix}, \\
 S_0 &= \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -i \\ 0 & 0 & 0 & i & 0 \end{pmatrix}.
 \end{aligned} \tag{41}$$

Next, we are interested in eliminating all the elements in the fifth row. The six generators  $J_i$  and  $K_i$  are not affected by this operation, but  $Q_1, Q_2, Q_3$ , and  $S_0$  become

$$\begin{aligned}
 P_1 &= \begin{pmatrix} 0 & 0 & 0 & 0 & i \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}, & P_2 &= \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & i \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}, & P_3 &= \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & i \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \\
 P_0 &= \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -i \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix},
 \end{aligned} \tag{42}$$

respectively. While  $J_i$  and  $K_i$  generate Lorentz transformations on the four dimensional Minkowski space, these  $Q_i$  and  $S_0$  in the form of the  $P_i, P_0$  matrices generate translations along the  $x, y, z$ , and  $t$  directions respectively. We shall study this aspect in detail in Section 4.

#### 4. Contraction of $O(3,2)$ to the Inhomogeneous Lorentz Group

We can contract  $O(3,2)$  according to the procedure introduced by Inönü and Wigner [3]. They introduced the procedure for transforming the four-dimensional Lorentz group into the three-dimensional Galilei group. Here, we shall contract the boost generators belonging to the time-like  $s$  variable,  $Q_i$ , along with the rotation generator between the two time-like variables,  $S_0$ .

Here, we illustrate the Inönü-Wigner procedure using the concept of squeeze transformations. For this purpose, let us introduce the squeeze matrix

$$C(\epsilon) = \begin{pmatrix} 1/\epsilon & 0 & 0 & 0 & 0 \\ 0 & 1/\epsilon & 0 & 0 & 0 \\ 0 & 0 & 1/\epsilon & 0 & 0 \\ 0 & 0 & 0 & 1/\epsilon & 0 \\ 0 & 0 & 0 & 0 & \epsilon \end{pmatrix}. \tag{43}$$

This matrix commutes with  $J_i$  and  $K_i$ . The story is different for  $Q_i$  and  $S_0$ .

For  $Q_1$ ,

$$C Q_1 C^{-1} = \begin{pmatrix} 0 & 0 & 0 & 0 & i/\epsilon^2 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ i\epsilon^2 & 0 & 0 & 0 & 0 \end{pmatrix}, \tag{44}$$

which, in the limit of small  $\epsilon$ , becomes

$$Q'_1 = \begin{pmatrix} 0 & 0 & 0 & 0 & i/\epsilon^2 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}. \tag{45}$$

We then make the inverse squeeze transformation:

$$C^{-1} Q'_1 C = \begin{pmatrix} 0 & 0 & 0 & 0 & i \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}. \tag{46}$$

Thus, we can write this contraction procedure as

$$P_1 = \lim_{\epsilon \rightarrow 0} \left( \epsilon^2 C Q_1 C^{-1} \right), \tag{47}$$

where the explicit five-by-five matrix is given in Equation (42). Likewise

$$P_2 = \lim_{\epsilon \rightarrow 0} \left( \epsilon^2 C Q_2 C^{-1} \right), \quad P_3 = \lim_{\epsilon \rightarrow 0} \left( \epsilon^2 C Q_3 C^{-1} \right), \quad P_0 = \lim_{\epsilon \rightarrow 0} \left( \epsilon^2 C S_0 C^{-1} \right). \tag{48}$$

These four contracted generators lead to the five-by-five transformation matrix, as can be seen from

$$\exp \{ -i (aP_1 + bP_2 + cP_3 + dP_0) \} \tag{49}$$

performing translations in the four-dimensional Minkowski space:

$$\begin{pmatrix} 1 & 0 & 0 & 0 & a \\ 0 & 1 & 0 & 0 & b \\ 0 & 0 & 1 & 0 & c \\ 0 & 0 & 0 & 1 & -d \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \\ t \\ 1 \end{pmatrix} = \begin{pmatrix} x + a \\ y + b \\ z + c \\ t - d \\ 1 \end{pmatrix}. \quad (50)$$

In this way, the space-like directions are translated and the time-like  $t$  component is shortened by an amount  $d$ . This means the group  $O(3,2)$  derivable from the Heisenberg's uncertainty relations becomes the inhomogeneous Lorentz group governing the Poincaré symmetry for quantum mechanics and quantum field theory. These matrices correspond to the differential operators

$$P_x = -i\frac{\partial}{\partial x}, \quad P_y = -i\frac{\partial}{\partial y}, \quad P_z = -i\frac{\partial}{\partial z}, \quad P_0 = i\frac{\partial}{\partial t}, \quad (51)$$

respectively. These translation generators correspond to the Lorentz-covariant four-momentum variable with

$$p_1^2 + p_2^2 + p_3^2 - p_0^2 = \text{constant}. \quad (52)$$

This energy-momentum relation is widely known as Einstein's  $E = mc^2$ .

## 5. Concluding Remarks

According to Dirac [1], the problem of finding a Lorentz-covariant quantum mechanics reduces to the problem of finding a representation of the inhomogeneous Lorentz group. Again, according to Dirac [2], it is possible to construct the Lie algebra of the group  $O(3,2)$  starting from two oscillators. We have shown in our earlier paper [12] that this  $O(3,2)$  group can be contracted to the inhomogeneous Lorentz group according to the group contraction procedure introduced by Inönü and Wigner [3].

In this paper, we noted first that the symmetry of a single oscillator is generated by three generators. Two independent oscillators thus have six generators. We have shown that there are four additional generators needed for the coupling of the two oscillators. Thus there are ten generators. These ten generators can then be linearly combined to produce ten generators which were spelled out in Dirac's 1963 paper.

For the two-oscillator system, there are four step-up and step-down operators. There are therefore sixteen quadratic forms [9]. Among those, only ten of them are in Dirac's 1963 paper [2]. Why ten? Dirac needed those ten to construct the Lie algebra for the  $O(3,2)$  group. At the end of the same paper, he stated that this Lie algebra is the same as that for the  $Sp(4)$  group, which preserves the minimum uncertainty for each oscillator.

In this paper, we started with the block-diagonal matrix given in Equation (28) for two totally independent oscillators with six independent generators. We then added one two-by-two Hermitian matrix of Equation (29) with four generators for the off-diagonal blocks. The result is the four-by-four Hermitian matrix given in Equation (30). This four-by-four Hermitian matrix has ten independent operators which can be linearly combined to the ten operators chosen by Dirac. Thus, in this paper, we have shown how the two-oscillators are coupled, and how this coupling introduces additional symmetries.

Paul A. M. Dirac made life-long efforts to make quantum mechanics consistent with special relativity, starting from 1927 [4]. While we exploited the contents of his paper published in 1963 [2], it is of interest to review his earlier efforts. In his earlier papers, Dirac started with quantum mechanics and special relativity as two different branches of science based on two different mathematical bases.

In this paper, based on Dirac's two papers [1,2], we concluded that both quantum mechanics and special relativity can be derived from the same mathematical base. A brief review of Dirac's earlier efforts is given in Appendix A.

**Author Contributions:** Each author contributed equally to this article.

**Funding:** This research received no external funding.

**Conflicts of Interest:** The authors declare no conflict of interest.


### Appendix A

As we all know, quantum mechanics and special relativity were developed along two separate routes. As early as 1927, Dirac was interested in understanding whether these two scientific disciplines are compatible with each other. In his paper of 1927 [4], Dirac noted the the existence of the time-energy uncertainty relation without excitations. He called this the “c-number” time–energy uncertainty relation. Dirac pointed out that the space-time asymmetry makes it difficult to construct the uncertainty relation in the Lorentz-covariant world.

In 1945, Dirac considered the four-dimensional harmonic oscillator wave functions applicable to the four-dimensional space and time. In so doing, Dirac was considering localized bound states. The space and time variables in his case are the separations between two constituents, like the proton and electron in the hydrogen atom.

It was shown later that Dirac’s concern about the c-number time–energy uncertainty is not necessary in view of the fact that a massive particle at rest has only three space-like dimensions [13]. According to Wigner [14], the little group for the massive particle is isomorphic to  $O(3)$  [14]. With this understanding, we can use a circle in the  $z t$  plane as shown in Figure A1, where  $z$  and  $t$  are longitudinal and time separations respectively.

In his 1949 paper [15], Dirac introduced the light-cone coordinate system which tells us that the Lorentz boost is a squeeze transformation. This aspect is also illustrated in Figure A1. It is then not difficult to see how the circle looks to a moving observer.



**Figure A1.** Dirac’s three papers. His 1927 and 1945 papers can be described by a circle in the longitudinal space-like and time-like coordinate. Dirac introduced the light-cone coordinate system in 1949. In this system, the Lorentz boost is a squeeze transformation. It is then natural to synthesize these two figures to a squeezed circle or an ellipse. Figure A2 will illustrate how this elliptic squeeze manifests itself in the real world.


The next question is whether this elliptic squeeze has anything to do with the real world. One hundred years ago, Niels Bohr and Albert Einstein met occasionally to discuss physics. Their interests were different. Bohr was worrying about the electron orbit in the hydrogen atom. Einstein was interested in how things look to moving observers. Then the question arises: How would the hydrogen atom look to a moving observer? This was a metaphysical issue during the period of Bohr and Einstein, as there were no hydrogen atoms moving fast enough to exhibit this Einstein effect.

Fifty years later, the physics world was able to produce many protons from particle accelerators. In 1964 [16], Gell-Mann observed that the proton is a bound state of the more fundamental particles called “quarks” according to the quantum mechanics applicable also to the hydrogen atom.

However, according to Feynman [17,18], when the proton moves very fast, it appears as a collection of a large number of free-moving light-like partons with a wide-spread momentum distribution, as described in Figure A2. Feynman’s parton picture was entirely based on what we observe in laboratories.

### Lorentz-squeezed Hadron

Quarks → Partons



**Figure A2.** In the harmonic-oscillator regime, the momentum–energy wave function takes the same mathematical form as that of the space-time wave functions. This figure shows that the quark model and the parton model are two different aspects of one Lorentz-covariant entity. In 1969 [17], Feynman observed that the fast-moving proton appears as a collection of a large number of light-like partons with a wide-spread momentum distribution, and short interaction time with the external signal. This figure is a graphical illustration of the 1977 paper by Kim and Noz [19]. This figure is from a recent book by the present authors [20].

Unlike the hydrogen atom, the proton can become accelerated, and its speed could be very close to that of light. Thus the Bohr–Einstein issue became the Gell-Mann–Feynman issue, as illustrated in Figure A3. The question is whether Gell-Mann’s quark model and Feynman’s parton picture are two different aspects of one Lorentz-covariant entity. This question was addressed by Kim and Noz 1977 [19] and was explained in detail by the present authors with a graphical illustration given in Figure A2.



100 years ago, Bohr was worrying about the orbit of the hydrogen atom.

Einstein was interested in how things look to moving observers. Then how the hydrogen atom would look to moving observers? This was a metaphysical question for them.



50 years ago, the proton became a bound state of the quarks sharing the same quantum mechanics as that for the hydrogen atom, according to Gell-Mann. If it moves with a speed close to that of light, the proton appears as a collection of partons, according to Feynman.

**Question. Does the proton appear like a collection of Feynman's partons to a moving observer?**



Photo of Gell-Mann by Y.S.Kim (2010), all others photos are from the public domain.

**Figure A3.** The Bohr–Einstein issue is 100 years old. Fifty years later, it became the quark–parton puzzle, based on observations made in high-energy laboratories.

## References

1. Dirac, P.A.M. Forms of Relativistic Dynamics. *Rev. Mod. Phys.* **1949**, *21*, 392–399. [[CrossRef](#)]
2. Dirac, P.A.M. A Remarkable Representation of the  $3 + 2$  de Sitter Group. *J. Math. Phys.* **1963**, *4*, 901–909. [[CrossRef](#)]
3. İnönü, E.; Wigner, E.P. On the Contraction of Groups and their Representations. *Proc. Natl. Acad. Sci. USA* **1953**, *39*, 510–524. [[CrossRef](#)] [[PubMed](#)]
4. Dirac, P.A.M. The Quantum Theory of the Emission and Absorption of Radiation. *Proc. R. Soc. (London)* **1927**, *A114*, 243–265. [[CrossRef](#)]
5. Han, D.; Kim, Y.S.; Noz, M.E. Linear canonical transformations of coherent and squeezed states in the Wigner phase space. *Phys. Rev. A* **1988**, *37*, 807–814. [[CrossRef](#)] [[PubMed](#)]
6. Kim, Y.S.; Wigner, E.P. Canonical transformation in quantum mechanics. *Am. J. Phys.* **1990**, *58*, 439–447. [[CrossRef](#)]
7. Kim, Y.S.; Noz, M.E. *Phase Space Picture of Quantum Mechanics*; World Scientific Publishing Company: Singapore, 1991.
8. Dodonov, V.V.; Man'ko, V.I. *Theory of Nonclassical States of Light*; Taylor & Francis: London, UK; New York, NY, USA, 2003.
9. Han, D.; Kim, Y.S.; Noz, M.E.  $O(3,3)$ -like Symmetries of Coupled Harmonic Oscillators. *J. Math. Phys.* **1995**, *36*, 3940–3954. [[CrossRef](#)]
10. Abraham, R.; Marsden, J.E. *Foundations of Mechanics*, 2nd ed.; Benjamin Cummings: Reading, MA, USA, 1978.
11. Goldstein, H. *Classical Mechanics*, 2nd ed.; Addison-Wesley: Reading, MA, USA, 1980.
12. Başkal, S.; Kim, Y.S.; Noz, M.E. Poincaré Symmetry from Heisenberg's Uncertainty Relations. *Symmetry* **2019**, *11*, 409. [[CrossRef](#)]
13. Kim, Y.S.; Noz, M.E.; Oh, S.H. Representations of the Poincaré group for relativistic extended hadrons. *J. Math. Phys.* **1979**, *20*, 1341–1344. [[CrossRef](#)]
14. Wigner, E. On unitary representations of the inhomogeneous Lorentz group. *Ann. Math.* **1939**, *40*, 149–204. [[CrossRef](#)]
15. Dirac, P.A.M. Unitary Representations of the Lorentz Group. *Proc. R. Soc. (London)* **1945**, *A183*, 284–295.
16. Gell-Mann, M. A Schematic Model of Baryons and Mesons. *Phys. Lett.* **1964**, *8*, 214–215. [[CrossRef](#)]
17. Feynman, R.P. Very High-Energy Collisions of Hadrons. *Phys. Rev. Lett.* **1969**, *23*, 1415–1417. [[CrossRef](#)]

18. Bjorken, J.D.; Paschos, E.A. Electron-Proton and  $\gamma$ -Proton Scattering and the Structure of the Nucleon. *Phys. Rev.* **1969**, *185*, 1975–1982. [[CrossRef](#)]
19. Kim, Y.S.; Noz, M.E. Covariant harmonic oscillators and the parton picture. *Phys. Rev. D* **1977**, *15*, 335–338. [[CrossRef](#)]
20. Başkal, S; Kim, Y.S.; Noz, M.E. *Physics of the Lorentz Group, IOP Concise Physics*; Morgan & Claypool Publisher: San Rafael, CA, USA; IOP Publishing: Bristol, UK, 2015.



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).