

Article

Error Distribution Model to Standardize LPUE, CPUE and Survey-Derived Catch Rates of Target and Non-Target Species

Régis Santos^{1,2*}, Osman Crespo^{1,2}, Wendell Medeiros-Leal^{1,2}, Ana Novoa-Pabon¹ and Mário Pinho^{1,2}

¹Okeanos-UAc Instituto de Investigação em Ciências do Mar, Universidade dos Açores. Rua Prof. Dr. Frederico Machado, 4, 9900-138 – Horta, Faial, Portugal.

²IMAR Instituto do Mar, Departamento de Oceanografia e Pescas, Universidade dos Açores. Rua Prof. Dr. Frederico Machado, 4, 9901-862 – Horta, Faial, Portugal.

* Correspondence: regisvinicius@gmail.com, Tel. (+351) 292 200 469

Figure S1. Diagnostic plot to detect the adequacy of the assumed variance function in generalized linear models: Pearson residuals plotted versus fitted values for thornback ray *Raja clavata* (RJC), blackbelly rosefish *Helicolenus dactylopterus* (BRF), and common mora *Mora moro* (RIB). The null pattern is a no trend in the residuals. The red line is the *loess* smoother through the plotted values. A: LPUE (kg landing⁻¹ vessel⁻¹); B: CPUE (kg days at sea⁻¹ vessel⁻¹); C: RPN (ind. 10⁻³ hooks).





