

Supplemental Table S2. Laboratory Analysis of Bloodwork and Joint Aspirate and Pathology of Intraoperative Specimens

Descriptive	All Outcomes				Prosthesis Survival			Aseptic Loosening			Recurrent PJI		
	All Patients	Male	Female	p Value	Male	Female	p Value	Male	Female	p Value	Male	Female	p Value
<i>Joint Aspirate</i>													
Absolute Cell Number [/nL]	20,294±32,438	26,868±41,471	13,719±16,399	0.111	23,776±38,645	14,528±16,263	0.377	18,818±36,209	8,911±15,583	0.178	33,337±46,363	26,499±16,283	0.830
PMN [/nL]	18,398±29,955	24,074±38,130	12,721±15,734	0.166	21,708±35,163	13,212±16,095	0.419	3,488±33,519	8,507±14,990	0.312	32,551±42,466	24,316±14,940	0.787
PMN [%]	64.7±26.9%	63.1±27.7%	66.4±25.4%	0.662	64.7±27.9%	65.7±26.4%	0.925	52.5±27.9%	62.8±25.8%	0.858	57.8±26.4%	81.6±12.6%	0.235
<i>CRP [mg/L]</i>													
Preoperative Prosthesis Removal	39.8±59.2	50.7±77.1	31.2±36.4	0.092	47.8±77.6	31.0±33.5	0.264	41.2±73.8	21.2±33.5	0.128	56.2±79.2	46.7±43.4	0.761
Postoperative Prosthesis Removal	9.9±13.4	8.4±7.6	11.0±16.5	0.316	9.8±8.5	10.5±12.1	0.789	3.9±8.2	12.5±17.5	0.445	6.7±4.6	10.7±8.9	0.246
Preoperative Prosthesis Reimplantation	49.3±33.2	54.7±37.0	44.8±28.6	0.149	54.9±36.5	45.4±28.3	0.288	36.7±34.2	38.7±29.1	0.278	60.9±41.4	52.3±23.3	0.619
<i>Pathology</i>													
Type I [%] (No. of Patients)	19.3% (21)	20.4% (10)	18.3% (11)	0.962	15.6% (5)	17.1% (6)	0.673	42.9% (3)	57.1% (4)	0.531	25.0% (3)	10.0% (1)	0.411
Type II [%] (No. of Patients)	37.6% (41)	34.7% (17)	40.0% (24)	0.653	37.5% (12)	40.0% (14)	0.870	14.3% (1)	71.4% (5)	0.426	33.3% (4)	40.0% (4)	0.543
Type III [%] (No. of Patients)	25.7% (28)	28.6% (14)	23.3% (14)	0.591	31.3% (10)	22.9% (8)	0.508	14.3% (1)	42.9% (3)	0.770	25.0% (3)	30.0% (3)	0.823
Type IV [%] (No. of Patients)	15.6% (17)	16.3% (8)	15.0% (9)	0.862	15.6% (5)	17.1% (6)	0.878	28.6% (2)	42.9% (3)	0.694	16.7% (2)	0.0% (0)	0.197
N/A [%] (No. of Patients)	19.3% (21)	20.4% (10)	18.3% (11)	0.366	0.0% (0)	2.9% (1)	-	0.0% (0)	0.0% (0)	-	0.0% (0)	10.0% (1)	0.273