

Article

A Regional Perspective of Socio-Ecological Predictors for Fruit and Nut Tree Varietal Diversity Maintained by Farmer Communities in Central Asia

Muhabbat Turdieva ¹, Agnès Bernis-Fonteneau ^{2,3,*} , Maira Esenalieva ⁴, Abdihalil Kayimov ⁵, Ashirmuhammed Saparmyradov ⁶, Khursandi Safaraliev ⁷, Kairkul Shalpykov ⁸, Paolo Colangelo ⁹ , and Devra I. Jarvis ^{2,10,11} 

- ¹ Office for Central Asia, Bioersivity International, c/o ICARDA, Osiyo Str., 6, Tashkent 100000, Uzbekistan; m.turdieva@cgiar.org
 - ² The Raffaella Foundation Platform for Agrobiodiversity Research (PAR), 80 Myer Creek Rd, Twisp, WA 98856, USA; d.jarvis@raffaellafoundation.org
 - ³ Department of Environmental Biology, Sapienza University of Rome, Piazzale Aldo Moro 5, 00185 Rome, Italy
 - ⁴ Horticulture Department, Kazakh National Agricultural Research University, 137, Valihanov Str., Almaty 050000, Kazakhstan; maira81@mail.ru
 - ⁵ Forestry Department, Tashkent State Agrarian University, 2, Universitetskaya Str., Tashkent 100140, Uzbekistan; a.kayimov@mail.ru
 - ⁶ Agricultural Sciences Department, Academy of Sciences, Turkmenistan, 15, Bitarap Turkmenistan Str., Ashgabat 744000, Turkmenistan; keremli@mail.ru
 - ⁷ Non Commercial Cooperative “Sarob”, 52/46 Ayni Str., Dushanbe 734003, Tajikistan; khursandi@mail.ru
 - ⁸ Institute of Chemistry and Phytotechnologies, 267, Chuy Avenue, Bishkek 720071, Kyrgyzstan; alhor6464@mail.ru
 - ⁹ National Research Council, Research Institute on Terrestrial Ecosystems, Via Salaria km 29,300, 00015 Rome, Italy; paolo.colangelo@cnr.it
 - ¹⁰ Bioersivity International, Via di San Domenico, 1, 00153 Rome, Italy
 - ¹¹ Department of Crop and Soil Sciences, Washington State University, Pullman, WA 99164, USA
- * Correspondence: agnes.bernisfonteneau@uniroma1.it



Citation: Turdieva, M.; Bernis-Fonteneau, A.; Esenalieva, M.; Kayimov, A.; Saparmyradov, A.; Safaraliev, K.; Shalpykov, K.; Colangelo, P.; Jarvis, D.I. A Regional Perspective of Socio-Ecological Predictors for Fruit and Nut Tree Varietal Diversity Maintained by Farmer Communities in Central Asia. *World* **2024**, *5*, 22–35. <https://doi.org/10.3390/world5010002>

Academic Editor: Manfred Max Bergman

Received: 9 November 2023

Revised: 5 January 2024

Accepted: 9 January 2024

Published: 11 January 2024



Copyright: © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: The five independent countries of Central Asia, namely Kazakhstan, Kyrgyzstan, Tajikistan, Turkmenistan, and Uzbekistan, contain one of the richest areas in the world for the specific and intraspecific diversity of temperate fruit and nut tree species. Research was carried out via the collaboration of national research and education institutes with local community-based agencies and farmer communities. Raw data (2014 observations) for almond, apple, apricot, cherry plum, currant, grapevine, pear, pomegranate, and walnut were collected at the household (HH) level across the five countries: Uzbekistan, Kazakhstan, Tajikistan, Turkmenistan, and Kyrgyzstan. A set of models was used, including household variety richness as the dependent variable, to understand the influence of socio-ecological variables on the amount and distribution of crop varietal diversity in the farmers' production systems. Four variables were included as explanatory variables of variety richness (fixed factors): ecoregion, ethno-linguistic group, management, and abiotic stress. The results show clear evidence that abiotic stress determines a higher richness of intra-specific diversity in the form of local varieties grown by farmers living in climatically unfavorable areas. The results for the studied ecoregions follow the same trend, with ecoregions with harsher conditions displaying a higher positive correlation with diversity. Mild environments such as the Central Asian riparian woodlands show an unexpectedly lower diversity than other harsher ecoregions. Ethno-linguistic groups also have an effect on the level of varietal diversity used, related to both historic nomadic practices and a culture of harvesting wild fruit and nuts in mountainous areas. The home garden management system hosts a higher diversity compared to larger production systems such as orchards. In Central Asia, encouraging the cultivation of local varieties of fruit and nut trees provides a key productive and resilient livelihood strategy for farmers living under the harsh environmental conditions of the region while providing a unique opportunity to conserve a genetic heritage of global importance.

Keywords: intraspecific diversity; resilience; environmental stress; horticultural crops; risk management; climate adaptation; ethno-linguistic diversity; Central Asia

1. Introduction

The five independent countries of Central Asia, namely Kazakhstan, Kyrgyzstan, Tajikistan, Turkmenistan, and Uzbekistan, contain one of the richest areas in the world for specific and intraspecific diversity of temperate fruit and nut tree species [1]. Important horticultural crops commonly used in temperate farmland worldwide, including Apple (*Malus domestica*), apricot (*Prunus armeniaca*), pear (*Pyrus communis*), cherry plum (*Prunus cerasifera*), currant (*Ribes nigrum*), pomegranate (*Punica granatum*), grape (*Vitis vinifera*), almond (*Amygdalus communis*), and walnut (*Juglans regia*), have their primary centers of diversity in Central Asia. These perennial crops, domesticated in the region, spread to temperate areas globally, following commercial trade routes [2–15]. Diverse, extreme natural and climatic conditions, coupled with human influence over the past three to four millennia, marked by the selection and development of varieties of these perennial fruit and nut trees, have provided productivity and resilience to the production systems of smallholder farmers in the region [9,16–18]. This dense reservoir of varieties adapted to specific environments of Central Asia is of high value for today's farmers and future breeding and development [18–21].

Today, Central Asia continues to host a remarkable diversity of many of these species [9,18,22,23]. This genetic base of Central Asia's cultivated fruit and nut trees continues to be expanded today by the constant interaction of cultivated varieties with wild materials collected by farmers, such as rootstock and graft wood adapted to local harsh abiotic conditions and biotic stress [9]. The genetic pool is maintained and shaped by traditional knowledge transmitted from one generation to the other in the farming population [18]. Local grafting practices in orchards and home gardens are passed down through generations to incorporate locally adapted varieties into the production systems of these harsh environments [24].

The development of arboriculture in Central Asia represents an investment for longer-term returns compared to those gained from annual crops, which has, in turn, shaped both the agricultural landscapes and local livelihood strategies toward durable benefits and resilience [25–27]. Horticulture in Central Asia has been increasingly studied in the past decade with the aim of improving the conservation of fruit and nut trees in the area [28]. Despite the collectivization of agriculture during the Soviet era, the production of fruit and nut trees remained important at the family level in Central Asian countries. Perennial crop production has increased since independence, supporting the need for households to be resilient during the economic transition [10] as new independent countries in Central Asia moved out of mainly cooperative farms into individual farming households. Market demand for local fruit and nut crop varieties has increased during the past ten years, combined with farmers' interest in local fruit crop varieties, especially apple, pear, and grape [18].

From 2006 to 2013, a coordinated global partnership of researchers working with farming communities in five Central Asia countries measured the amount and distribution of fruit and nut tree genetic diversity in the form of traditional, improved (improved varieties from regional materials) and introduced (exotic materials from outside the region) varieties in farmers' fields of nine fruit and nut tree species (Figure 1). Via this partnership, countries worked together, using globally applicable diversity indices, developed for annual crops [29], to compare across farmer households and socioecological zones the amount and distribution of perennial fruit and nut tree varietal diversity. This paper (i) synthesizes the total body of fruit and nut tree diversity data gathered on-farm in the study; (ii) demonstrates that considerable fruit and nut tree genetic diversity continues to be maintained in family farms; (iii) examines socio-ecological variables, abiotic stress,

management space, ecoregion, and ethno-linguistic group, to better understand how these factors affect and explain the observed varietal richness in fruit and nut trees of Central Asia; and (iv) provides an analytical framework for determining the role of on-farm fruit and nut tree genetic diversity in the production and resilience of family farms. The readaptation of fruit and nut tree cultivation once out of the Soviet era and the long lifespan of these perennial crops make the dataset collected extremely valuable for analysis for policy recommendations in the region today.




Figure 1. Sampled households across country sites (red dots).

2. Materials and Methods

2.1. Study Sites

Research was carried out in Kazakhstan, Kyrgyzstan, Tajikistan, Turkmenistan, and Uzbekistan via the collaboration of national research and education institutes with local community-based agencies and farmer communities. Economic development ranged from less-developed (lower middle income) economies in Kyrgyzstan, Tajikistan, and Uzbekistan, to the more-developed (upper middle income) economies such as Kazakhstan and Turkmenistan, which are oil-based (World Bank Country and Lending Groups webpage (<https://datahelpdesk.worldbank.org/knowledgebase/articles/906519>, accessed on 11 September 2023)). Seventeen social economic zones were selected to encompass environmental, cultural, technological, and economic differences (see Tables S1 and S2 in Supplementary Materials for details). Figure 1 presents the locations of the sampled households across the five studied countries.

The survey spanned socio-economic zones, covering 11 ecoregions as defined by Olson et al. [30]: the Alai-Western Tian Shan steppe; the Badkhyz and Karabil semi-desert; the Central Asian northern desert; the Central Asian riparian woodlands; the Central Asian southern desert; the Gissaro-Alai open woodlands; the Kopet Dag semi-desert; the Kopet Dag woodlands and forest steppe; the Pamir alpine desert and tundra; the Tian Shan foothill arid steppe; and the Tian Shan montane conifer forests. Site elevation ranged from 100 m a.s.l. to 1800 m a.s.l. Precipitation in the sites ranges from 80 mm to 1000 mm, falling predominantly from October to May. In the south, temperatures reach over 45 °C in the summer months and can drop to as low as −25 °C in the coldest months throughout Central Asia in the plains. Eight major ethno-linguistic groups, Kazakh, Kyrgyz, Russian, Tajik, Turkish, Turkmen, Uygur, and Uzbek, were located in the communities surveyed. The average size of home gardens was 0.026 hectares. The average orchard size per crop was approximately one hectare but ranged from 0.002 ha in Turkmenistan for cherry plums to 95 ha for grapes in Kazakhstan.

2.2. Sampling on Farms and in Communities

The farms were sampled to represent a broad picture of the fruit and nut tree varietal diversity in the five Central Asian countries. Raw data (2014 observations) for almond, apple, apricot, cherry plum, currant, grapevine, pear, pomegranate, and walnut were collected at the household (HH) level across the five countries: Uzbekistan, Kazakhstan, Tajikistan, Turkmenistan, and Kyrgyzstan. At each HH, data were collected on all varieties and genotypes grown by the farmer for the targeted fruit and nut trees of the study, using a standardized participatory diagnostic survey adapted from Jarvis and Campilan [31] and the Diversity Assessment Tool for Agrobiodiversity and Resilience (DATAR) [32]. Information was collected on variety names, number of trees, ethno-linguistic group of the farmer (inferred from the language used for the interview), ecoregion, and management type (orchard or home garden). Fruit and nut tree varieties are characterized by Central Asians into three groups: traditional, modern, and introduced. Traditional varieties (synonym for landrace) are defined as farmer recognized crop varieties possessing a certain genetic integrity that evolved in cultivation over long periods, adapted to a specific local environment or purpose. Modern varieties are defined as varieties improved, registered, and released in the national research breeding systems in Central Asia Institutes. Introduced varieties are defined as certified varieties developed outside Central Asia. The nomenclature, identification, and consistency of variety names follow the methods described in Jarvis et al. (2008) [29]. Subsequently, data were aggregated for the analysis in order to obtain variety richness for each fruit tree species at the HH level, also considering the type of management arrangement.

2.3. Data Analysis

A set of generalized linear mixed models (GLMM) was used, including household variety richness as the dependent variable. The GLMM was used to evaluate the effect of the socio-ecological variables on the distribution of fruit and nut tree diversity in Central Asia for data collected from 388 households, with ecoregion, ethno-linguistic group, management, and abiotic stress as fixed effects. Data were analyzed at a regional level, pooling together data from different fruit tree crops and countries. This type of approach presents some difficulties due to the nature of the data: (1) data are unevenly distributed in each country (i.e., different sampling efforts and different interests in specific fruit tree crops); and (2) different fruit tree species are characterized by their own variety richness due to biological reason and selection effort by farmers. Richness data (count data) do not follow a normal distribution. In order to control all these factors, we used a GLMM that can handle non-normal data (richness) and control for bias due to the structure of the data [33].

Ethno-linguistic groups were defined on the basis of the language used for the interview of the farmers; management represents the arrangement of fruit trees in small groups for domestic or local use (*home garden*) or in larger lands and devoted mainly to market (*orchard*). Ecoregions included semi-desert, northern desert, riparian woodlands, southern desert, open woodlands, woodlands and forest steppe, alpine desert and tundra, montane foothill arid steppe, and montane conifer forests across Central Asia. Ecoregions were obtained from the map of terrestrial ecoregions provided by Olson et al. [30]. Ethno-linguistic groups were based on the main language identified by the household: Kyrgyz, Russian, Tajik, Turkish, Turkmen, Uygur, and Uzbek. Finally, climatic stress represents a combined variable derived from the worldclim database (<https://worldclim.org/>, accessed on 11 September 2023), from which we extracted BIO1 (annual mean temperature); BIO5 (maximum temperature of the warmest month); BIO6 (minimum temperature of the coldest month); BIO12 (annual precipitation); B016 (precipitation in the wettest quarter); and B017 (precipitation of driest quarter), as they are known to influence fruit and nut tree productivity [34]. Given that these variables are known to be correlated with each other and in order to avoid an inflation of variables in the models, we performed a Principal Component Analysis (PCA) of the five variables and took the first PC score (representing 75% of the total variance) as a proxy for abiotic stress.

A full GLMM was obtained, including the four fixed effects and two random effects: Country and Crop. Successively, we compared the full model with all the possible reduced models (with a combination of 3 or fewer variables) and a null model. The best-fitting model was chosen based on the Akaike Information Criterion corrected for finite sample size (AICc).

All the analyses were conducted using the R statistical environment [35]. The GLMM was performed using the package lme4 [36]. Comparisons of AICc scores between full and reduced models were carried out using the package MuMIn [37].

3. Results

3.1. Overall Diversity Estimates

Intra-specific varietal diversity for each studied fruit or nut tree at the household level, median value, and confidence intervals are presented in Figure 2. The richness of intra-specific diversity at the household level is high and variable. All the studied fruit and nut trees have an average variety richness at a household level above 2.2; Apple, Apricot, Cherry plum, Grapevine, and Pear are even above 3.7, with the highest average richness at the household level for Apricot at 4.8. Four of these crop trees show a very wide range of variety richness at the household level, ranging from 1 to 62 for Apple, 1 to 24 for Apricot, 1 to 58 for Grapevine, and 1 to 27 for Pear. The median is at 2 for Almond, Apple, Currant, Grapevine, Pear, and Pomegranate; at 3 and 4, respectively, for Apricot and Cherry plum; and 1.5 for Walnut. Detailed results of varietal diversity at the household level are available in Table S3 in the Supplementary Materials. They are also presented in Figure 2 with box plots summarizing these results.




Figure 2. Intra-specific varietal richness at household level by crop (crop names on the x-axis and number of varieties on the y-axis). Bold line = median value; box = 50% of the confidence interval; whiskers = 95% of the confidence interval.

Across the five Central Asian countries, the variety richness shows very high values for apple, 188; Apricot, 79; and Grapevine, 100. These results are summarized in Table 1. The table also presents the percentages of variety types: traditional, modern, and introduced. In Table S4 of the Supplementary Materials, the full list of variety names (in Russian) by crop and their type (traditional, modern, or introduced) is also made available.

Table 1. Variety richness and percentages by type per crop at project level.

Crop	Total Number of Varieties Per Crop	% Traditional Varieties (Landraces)	% Modern Varieties Bred in Central Asia Institutes	% Introduced Improved Varieties from Outside Central Asia
Almond	11	91%	9%	0%
Apple	188	72%	6%	22%
Apricot	79	89%	1%	10%
Cherry plum	11	36%	0%	64%
Currant	7	86%	14%	0%
Grapevine	100	72%	1%	27%
Pear	47	74%	4%	21%
Pomegranate	29	86%	0%	14%
Walnut	24	83%	0%	17%
Total number/mean %	496	76%	3%	20%

3.2. Social Ecological Predictors

According to the Akaike Information Criterion corrected for finite sample size (AICc), we found that the full model is the best fitting model to explain the fruit tree diversity in Central Asia with an AIC delta of 8.06 with the next best fitting model (model that excludes ethno-linguistic groups). The model selection table is available in Table S5 in Supplementary Materials. This allowed us to robustly reject reduced models in favor of a multifactorial full model.

We then estimated the effects and their significance for the full model. Coefficient estimates, standard error, and significance are calculated for the full model with reference effects categories where management is set to the home garden, ecoregion to Alai-Western Tian Shan steppe, and Kazakh for the ethno-linguistic group. The results are presented in Table 2.

According to the estimated values for fixed factors (Table 2, Figure 3), we found that diversity increases significantly with the increase in abiotic stress; the trend is clear, and confidence intervals (CI) are narrow around the coefficient estimate. Figure 3A shows this trend clearly. Among management practices, home gardens harbor a significantly higher diversity with respect to orchards. The coefficient estimate for the management factor, despite a wide CI, shows that varieties under orchard management have clearly less varietal diversity compared to home gardens; Figure 3B gives a good visualization of the differences between the two management types. Among the ecoregions, as represented in Figure 3C, we found large fluctuations in the estimated diversity in many cases, such as the Central Asian northern desert (C), Pamir alpine desert and tundra (I), Tian Shan foothill arid steppe (J), and the Tian Shan montane conifer forests (K), which showed higher estimated values of richness whereas Badghyz and Karabil semi-desert (B), Central Asian riparian woodlands (D), Central Asian southern desert (E), and Kopet Dag semi-desert (G) showed a low diversity. Among the different ethno-linguistic groups (Figure 3D), we found significant differences, especially in the Kyrgyz, Russian, Turkmen, and Uzbek groups, which are related to a higher tree diversity. However, it is difficult to evaluate the effect of this variable due to the general wide confidence interval. Graphs in Figure 3 provide a good visualization of correlations and confidence intervals.




Figure 3. Relationships among variety richness and the four socio-ecological variables estimated by the full GLMM. Plots contain a confidence band, prediction line, and partial residuals (points). Ecoregion names were abbreviated as follows: Alai-Western Tian Shan steppe (A); Badghyz and Karabil semi-desert (B), Central Asian northern desert (C), Central Asian riparian woodlands (D), Central Asian southern desert (E), Gissaro-Alai open woodlands (F), Kopet Dag semi-desert (G), Kopet Dag woodlands and forest steppe (H), Pamir alpine desert and tundra (I), Tian Shan foothill arid steppe (J), and Tian Shan montane conifer forests (K). Ethno-linguistic groups were abbreviated as follows: Kazakh (Kaz), Kyrgyz (Kyr), Russian (Rus), Tajik (Tjk), Turkish (Tur), Turkmen (Trk), Uygur (Uyr), and Uzbek (Uzb).

Table 2. Estimated values for fixed factors. For each fixed, effects coefficient (coef.) estimate, standard (std.) error, z value, and significance (p) are reported. For multilevel variables (management, ecoregion, and ethno-linguistic group), the significance is relative to the first level (in alphabetical order) of the belonging variable group. For the random effects crop and country, the model estimated a variance and standard deviation of 0.2075 ± 0.4556 and 0.0865 ± 0.2941 , respectively.

	Coef. Estimate	Std. Error	z Value	p ($> z $)
(Intercept)	1.211071	0.263782	4.591	4.41×10^{-6}
Management: Orchard	-0.346055	0.061785	-5.601	2.13×10^{-8}
Ecoregion: Badghyz and Karabil semi-desert	-0.484916	0.138486	-3.502	0.000463
Ecoregion: Central Asian northern desert	0.798726	0.146242	5.462	4.72×10^{-8}
Ecoregion: Central Asian riparian woodlands	-0.624811	0.133364	-4.685	2.80×10^{-6}
Ecoregion: Central Asian southern desert	-0.454274	0.112785	-4.028	5.63×10^{-5}
Ecoregion: Gissaro-Alai open woodlands	-0.155917	0.096973	-1.608	0.107868
Ecoregion: Kopet Dag semi-desert	-0.729989	0.193984	-3.763	0.000168
Ecoregion: Kopet Dag woodlands and forest steppe	-0.341358	0.178067	-1.917	0.055235
Ecoregion: Pamir alpine desert and tundra	0.159150	0.223831	0.711	0.477066
Ecoregion: Tian Shan foothill arid steppe	0.272411	0.130293	1.351	0.036550

Table 2. Cont.

	Coef. Estimate	Std. Error	z Value	p (> z)
Ecoregion: Tian Shan montane conifer forests	0.234397	0.173525	1.526	0.176760
Abiotic stress	0.082108	0.023349	3.517	0.000437
Ethno-linguistic group: Kyrgyz	0.439634	0.161478	2.723	0.006478
Ethno-linguistic group: Russian	0.261656	0.113879	2.298	0.021580
Ethno-linguistic group: Tajik	−0.009328	0.182743	−0.051	0.959290
Ethno-linguistic group: Turkish	−0.077775	0.235427	−0.330	0.741131
Ethno-linguistic group: Turkmen	0.349178	0.459837	0.759	0.447642
Ethno-linguistic group: Uygur	−0.331758	0.169417	−1.958	0.050203
Ethno-linguistic group: Uzbek	0.192472	0.157862	1.219	0.222754

4. Discussion

The aim of this research was to understand what socio-ecological variables can explain the observed varietal richness in fruit trees of Central Asia. Our analysis provides an overall picture of the main drivers of fruit tree diversity in agroecosystems of Central Asia. As expected, diversity is the product of multifactorial drivers and an interplay between environmental, abiotic, and cultural factors.

The variety richness shows impressively high results for apple, grapevine, and apricot. Traditional varieties represent over 70% of all varieties of all crops except cherry plum, showing these traditional varieties are key to agricultural production and cultural choices and are still predominantly used by farmers rather than modern or introduced varieties.

According to the estimated values and graphic representations obtained with the model (Table 1 and Figure 3), we found that all four variables tested are good predictors of fruit tree diversity in Central Asia. Our results show not only that different environmental conditions found in different ecoregions can determine different levels of varietal diversity but, unexpectedly, that more stressful conditions in an agroecosystem can trigger the planting and management of a larger number of varieties per crop than in more benign environments. We found that more extreme temperatures and low precipitation conditions are linked to higher levels of intra-specific (varietal) diversity (Figure 3A). Results for the studied ecoregions follow the same trend, with ecoregions having harsher conditions displaying a higher positive correlation with diversity, examples being the Central Asian northern desert (C), Pamir alpine desert, and tundra (I), Tian Shan foothill arid steppe (J), and Tian Shan montane conifer forests (K).

The more extreme environments in Central Asia have created conditions for higher levels of managed fruit and nut tree diversity compared to more benign environments. Numerous studies have shown that farmers living in extreme environments use higher levels of intra-specific diversity both to adapt to niche environments [38] and as a risk management strategy [39,40] to meet their production needs under changing stressful conditions [41]. Our analysis clearly shows an increase in fruit and nut tree intra-specific diversity with an increase in abiotic stress (high and low temperature and low rainfall). Most temperate fruit trees are obligatory outcrossers, requiring pollination from a compatible donor tree of a different variety before they can set fruit. Levels of pollinators are also affected by climatic conditions [42–44]. The more extreme environments in Central Asia contain a greater risk of unpredictable late frosts during flowering. Frosts just after pollination can also damage the first stages of fruit formation [45]. Farmers may use varietal diversity as a risk management strategy for both main varieties and pollinator donor trees in regions of increased unpredictability of frost onset and the length of the frost period. This strategy has also recently been recommended for apples in the Appalachian Mountains for changing frost patterns in the eastern United States [46].

Many of the ecoregions of the studied sites are subject to a dry climate. Although some have been labeled “deserts” under international classifications, the terminology can be misleading. These dry areas also contain numerous oases along their main rivers where fruit and nut trees thrive well. Ecoregions are correlated with climates. Temperatures and rainfalls vary according to the latitude and altitude. There is a wide range of ecoregions in this study; they nonetheless display some common features, such as the large fluctuation of temperatures between night and day and by seasons with a very cold winter and hot summer, and the occurrence of the limited precipitations mainly during the winter season. Overall, the results show more crop and varietal diversity in mountain areas. These results may be explained by the fact that these mountain areas host a patchwork of small econiches with various microclimates, which require the use of a diversity of adapted crops and varieties. Perennial fruit and nut tree crops display great potential for the mitigation and adaptation to climate change. Trees regulate water stress and excess on farms and are key to enhanced soil health [26,47,48], particularly important in mountain areas.

Different levels of soil salinity are also found across the ecoregions studied. Soil salinity is not correlated with ecoregions but with geographical characteristics, as is the case of saline soils in the irrigated areas of the valleys where groundwater tables are close to the surface. Farmers, faced with more or less salinity in their fields and orchards, experiment with different varieties with tolerance traits to soil salinity [49,50]. More stressful climate and soil conditions also affect levels of resistance for perennial crop species to pests and diseases. Farmers use varietal diversity to constantly adapt to evolving pests and diseases, looking for tolerance and resistance traits but also by using mixtures of different varieties within a field or orchard as part of their risk-management strategy [51–53]. Several studies have demonstrated that apple varieties and their wild relatives display various tolerance and resistance traits to diseases [54–58]. The use of varietal mixture can also play such a role [59]. By limiting the spread of epidemics, apple varietal diversity participates in disease regulation. Demestihis and colleagues (2017) [60] demonstrated the high potential for fruit trees in orchards to provide multiple ecosystem services, including pest and disease regulation and pollination, while Montanaro et al. (2017) [61] point to similar results for peach, apricot, olive groves, and vineyards in the Mediterranean area.

Among the different ethno-linguistic groups studied, we found significant differences; Kyrgyz, Russian, Turkmen, and Uzbek groups are related to a higher crop tree diversity, while Tajik and Kazakhs had significantly less diversity at the household level. These differences can be related to several cultural differences. An earlier econometric analysis in Central Asia revealed a linkage between Uzbek participation in community groups and the levels of fruit and nut tree diversity managed by households [62]. In contrast, Kazakhs are historically nomadic people and have only recently started to carry out crop farming [63], which could possibly explain the low presence of varietal diversity in this ethno-linguistic group. Tajik people commonly live in mountainous areas that host a rich wild nut and fruit tree diversity; rather than planting a wide set of varieties for each species, they have the conditions and a culture to collect a large part of their fruits and nuts from the wild [18,64]. Farmers in countries of Central Asia where populations of wild relatives of perennial crops can rely on genetic materials extracted from wild populations present in potential surrounding natural environments. These natural environments, when they are present, provide natural laboratories for farmers to select from. It is often the case in mountainous areas, where farmers select genotypes that display stronger tolerance and resistance to the biotic and abiotic local conditions. Farmers establish their own trials and selection. They are continuing the domestication process, bringing wild materials into their cultivated environments [3,16,65].

Home gardens in the study sites harbor significantly higher varietal diversity with respect to orchards. This result is consistent with studies of home gardens globally [66]. Home gardens support the rural population’s livelihood, improve food security, nutrition, and income, and provide an efficient risk management option against economic and environmental fluctuations [67]. Their contribution to overcoming the economic crisis

when transitioning from state to more private-led agriculture in Central Asia has also been documented [68]. These characteristics support the use of home gardens for sustainable agriculture, respect for the environment, supporting food security, and stimulating economic growth [67]. Important diversity, including crop intraspecific diversity in the form of varieties still commonly found in home gardens that are not present to that extent in larger fields [9,29,65,67,69]. Home gardens can also be a crucial management strategy to diversify household incomes and dietary diversity in the agricultural systems of former soviet republics of Central Asia. These gardens, of small size, about one hectare on average, have been spaces where farmers could exert their own choice even during the era of collective farming [9]. They usually grow vegetables, fruits, and nuts for their own consumption and for the provision of extra income to the household. Although home gardens primarily sustain household consumption, farmers are favoring plant varieties that are easily saleable on the market to be able to liquidate their surplus production [9]. In addition, home gardens are also places where farmers experiment with new materials from the wild and test management practices for perennial crops, including fruit and nut species [16,22]. Home gardens in Central Asia are also important hotspots of fruit tree varietal diversity because they maintain rare and differentiated varieties that can provide genetic material for future agronomic practices while at the same time meeting changing market demands. Traditional knowledge, built using farmers' experience passed from one generation to the other, is favored in home gardens and embeds the importance of crop diversity and varietal diversity.

Not only do Kazakhstan, Kyrgyzstan, Tajikistan, Turkmenistan, and Uzbekistan share a common wealth of genetic reserves of cultivated fruit and nut species, but they also share similar traditional farming systems and the same recent historical, economic and political legacy. The five countries have similar legislative frameworks regarding land reforms and cropping patterns, with implications for national and regional governments to provide future support to the use of traditional fruit and nut tree varieties in the farmers' production systems [22]. The dramatic changes since the breakup of the Soviet Union in 1991 have greatly influenced individual farmers' freedom and capacity to cultivate different types of crops and varieties, as seen from the results of our survey. The work presented here provides a scientific basis for policy development to include the use of fruit and nut tree varietal diversity in national development planning actions.

5. Conclusions

In Central Asia, encouraging the cultivation of local varieties of fruit and nut trees provides a key productive and resilient livelihood strategy for farmers living under the harsh environmental conditions of the region while at the same time providing a unique opportunity to conserve a genetic heritage of global importance. Both ecological and social-economic conditions can drive the choice of farmers, especially of smallholders, in maintaining a large portfolio of crop varieties that can ensure a good level of productivity under unfavorable and changing conditions. There is clear evidence that abiotic stress determines a higher richness of intra-specific diversity in the form of local varieties grown by farmers living in climatically unfavorable areas. The results for the studied ecoregions follow the same trend, with ecoregions with harsher conditions displaying a higher positive correlation with diversity. Mild environments such as the Central Asian riparian woodlands show an unexpectedly lower diversity than other harsher ecoregions. Ethno-linguistic groups also have an effect on the level of varietal diversity used, related both to historic nomadic practices and a culture of harvesting wild fruit and nuts in mountainous areas. Home garden management systems host a higher diversity compared to larger production systems, such as orchards. The integration of the varietal diversity in home gardens into national research and extension systems could facilitate the national conservation and development of fruit and nut trees for the benefit of farmers living in the diverse environmental habitats of Central Asia.

Supplementary Materials: The following supporting information can be downloaded at <https://www.mdpi.com/article/10.3390/world5010002/s1>. Table S1. Description of the selected agroecosystems and project sites in the region; Table S2. Characteristics of Households (HH): crop, management space, country, agroecological zone, ethnic groups, and ecoregion; Table S3. Varietal diversity at household level; Table S4. Variety names and types per crop at project level; Table S5. Model selection table.

Author Contributions: M.T., M.E., A.K., A.S., K.S. (Khursandi Safaraliev), K.S. (Kairkul Shalpykov), and D.I.J. designed the research; M.T., M.E., A.K., A.S., K.S. (Khursandi Safaraliev), K.S. (Kairkul Shalpykov) and D.I.J. performed research; P.C. and D.I.J. developed the analytic framework; P.C., D.I.J., M.T. and A.B.-F. analyzed data; M.T., D.I.J., P.C. and A.B.-F. wrote the paper; M.T., A.B.-F., M.E., A.K., A.S., K.S. (Khursandi Safaraliev), K.S. (Kairkul Shalpykov), P.C. and D.I.J. reviewed the manuscripts and provided meaning full intellectual contributions. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the United Nations Environment Programme (UNEP), Global Environment Facility (GEF) In Situ/On-Farm Conservation and Use of Agricultural Biodiversity (Horticultural Crops and Wild Fruit Species) in Central Asia: Grant number: GEF Project 1025; IMIS number GFL-2328-2715-4893.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: Data are available from the senior author upon reasonable request.

Acknowledgments: Our sincere and profound thanks go to the local communities and local authorities in Kazakhstan, Kyrgyzstan, Tajikistan, Turkmenistan, and Uzbekistan, who made this study possible. We also wish to thank the local community researchers and government officials in Kazakhstan, Kyrgyzstan, Tajikistan, Turkmenistan, and Uzbekistan for their support in field operations.

Conflicts of Interest: The authors declare no conflicts of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

References

- Vavilov, N.I. *Origin and Geographical Distribution of Cultivated Plants*; Nauka: Leningrad, Russia, 1987.
- Purugganan, M.D.; Fuller, D.Q. The Nature of Selection during Plant Domestication. *Nature* **2009**, *457*, 843–848. [[CrossRef](#)] [[PubMed](#)]
- Miller, A.J.; Gross, B.L. From Forest to Field: Perennial Fruit Crop Domestication. *Am. J. Bot.* **2011**, *98*, 1389–1414. [[CrossRef](#)] [[PubMed](#)]
- Pollegioni, P.; Woeste, K.E.; Chiocchini, F.; Del Lungo, S.; Olimpieri, I.; Tortolano, V.; Clark, J.; Hemery, G.E.; Mapelli, S.; Malvolti, M.E. Ancient Humans Influenced the Current Spatial Genetic Structure of Common Walnut Populations in Asia. *PLoS ONE* **2015**, *10*, e0135980. [[CrossRef](#)]
- Janick, J. The Origins of Fruits, Fruit Growing, and Fruit Breeding. In *Plant Breeding Reviews*; John Wiley & Sons, Ltd.: Hoboken, NJ, USA, 2005; Volume 25, pp. 255–321, ISBN 9780470650301.
- Groppi, A.; Liu, S.; Cornille, A.; Decroocq, S.; Bui, Q.T.; Tricon, D.; Cruaud, C.; Arribat, S.; Belser, C.; Marande, W.; et al. Population Genomics of Apricots Unravels Domestication History and Adaptive Events. *Nat. Commun.* **2021**, *12*, 3956. [[CrossRef](#)]
- Liu, S.; Cornille, A.; Decroocq, S.; Tricon, D.; Chague, A.; Eyquard, J.P.; Liu, W.S.; Giraud, T.; Decroocq, V. The Complex Evolutionary History of Apricots: Species Divergence, Gene Flow and Multiple Domestication Events. *Mol. Ecol.* **2019**, *28*, 5299–5314. [[CrossRef](#)]
- Cornille, A.; Antolín, F.; Garcia, E.; Vernesi, C.; Fietta, A.; Brinkkemper, O.; Kirleis, W.; Schlumbaum, A.; Roldán-Ruiz, I. A Multifaceted Overview of Apple Tree Domestication. *Trends Plant Sci.* **2019**, *24*, 770–782. [[CrossRef](#)]
- Vinceti, B.; Elias, M.; Azimov, R.; Turdieva, M.; Aaliev, S.; Bobokalonov, F.; Butkov, E.; Kaparova, E.; Mukhsimov, N.; Shamuradova, S.; et al. Home Gardens of Central Asia: Reservoirs of Diversity of Fruit and Nut Tree Species. *PLoS ONE* **2022**, *17*, e0271398. [[CrossRef](#)]
- Van Dusen, E. Agricultural Biodiversity in Transition Agriculture: Fruit Tree Genetic Resources in Rural Uzbekistan. Available online: https://www.bioecon-network.org/pages/7th_2005/VanDusen.pdf (accessed on 11 September 2023).
- Zhebentyayeva, T.; Ledbetter, C.; Burgos, L.; Llácer, G. Apricots. In *Fruit Breeding*; Springer: New York, NY, USA, 2012; pp. 415–458, ISBN 9781441907639.
- Azarov, A.; Polesny, Z.; Darr, D.; Kulikov, M.; Verner, V.; Sidle, R.C. Classification of Mountain Silvopastoral Farming Systems in Walnut Forests of Kyrgyzstan: Determining Opportunities for Sustainable Livelihoods. *Agriculture* **2022**, *12*, 2004. [[CrossRef](#)]

13. Bourguiba, H.; Scotti, I.; Sauvage, C.; Zhebentyayeva, T.; Ledbetter, C.; Krška, B.; Remay, A.; D'Onofrio, C.; Iketani, H.; Christen, D.; et al. Genetic Structure of a Worldwide Germplasm Collection of *Prunus Armeniaca* L. Reveals Three Major Diffusion Routes for Varieties Coming from the Species' Center of Origin. *Front. Plant Sci.* **2020**, *11*, 638. [[CrossRef](#)]
14. Riaz, S.; de Lorenzis, G.; Velasco, D.; Koehmstedt, A.; Maghradze, D.; Bobokashvili, Z.; Musayev, M.; Zdunic, G.; Laucou, V.; Andrew Walker, M.; et al. Genetic diversity analysis of cultivated and wild grapevine (*Vitis vinifera* L.) accessions around the Mediterranean basin and Central Asia. *BMC Plant Biol.* **2018**, *18*, 137. [[CrossRef](#)]
15. Mir-Makhamad, B.; Bjørn, R.; Stark, S.; Spengler, R.N. Pistachio (*Pistachio vera*) Domestication and Dispersal Out of Central Asia. *Agronomy* **2022**, *12*, 1758. [[CrossRef](#)]
16. Jarvis, D.I.; Hodgkin, T.; Brown, A.H.D.; Tuxill, J.; Noriega, I.; Smale, M.; Sthapit, B. *Crop Genetic Diversity in the Field and on the Farm: Principles and Applications in Research Practices*; Yale Agrarian Studies Series; Yale University Press: New Haven, CT, USA; London, UK, 2016; ISBN 0300161123.
17. *Wild Apple and Fruit Trees of Central Asia*; Janick, J. (Ed.) John Wiley & Sons: New York, NY, USA, 2003; Volume 29, ISBN 0471219681.
18. Turdieva, M.K.; Kayimov, A.K.; Baymetov, K.I.; Mustafina, F.U.; Butkov, E.A. Conservation and Sustainable Use of Biodiversity of Fruit Crops and Wild Fruit Species. In Proceedings of the International Scientific and Practical Conference, Tashkent, Uzbekistan, 23–26 August 2011.
19. Ryabov, I.N. Mobilization of Traditional Breeding Varieties and Their Use. In *Proceedings of the Scientific Conference*; Aystan Publishing House: Yerevan, Armenia, 1970; pp. 199–202. (In Russian)
20. Oleichenko, S.; Yegizbayeva, T.K.; Apushev, A.K.; Nusipzhanov, N.S. *Assessment of Promising Local Walnut Forms for the South and South-East of Kazakhstan*; National Academy of Sciences of the Republic of Kazakhstan: Almaty, Kazakhstan, 2020; Volume 5.
21. Martínez-García, P.J.; Hartung, J.; Pérez, L.; Cobos, F. Temporal Response to Drought Stress in Several *Prunus* Rootstocks and Wild Species. *Agronomy* **2020**, *10*, 1383. [[CrossRef](#)]
22. *Conservation of Fruit Tree Diversity in Central Asia: Policy Options and Challenges*; Lapenña, I.; Turdieva, M.; Noriega, I.L.; Ayad, W.G. (Eds.) Bioversity International: Rome, Italy, 2014; ISBN 9789290439202.
23. Gross, B.L.; Henk, A.D.; Richards, C.M.; Fazio, G.; Volk, G.M. Genetic Diversity in *Malus × domestica* (Rosaceae) through Time in Response to Domestication. *Am. J. Bot.* **2014**, *101*, 1770–1779. [[CrossRef](#)] [[PubMed](#)]
24. Vahdati, K.; Sarikhani, S.; Arab, M.M.; Leslie, C.A.; Dandekar, A.M.; Aletà, N.; Bielsa, B.; Gradziel, T.M.; Montesinos, Á.; Rubio-Cabetas, M.J.; et al. Advances in Rootstock Breeding of Nut Trees: Objectives and Strategies. *Plants* **2021**, *10*, 2234. [[CrossRef](#)]
25. Fuller, D.Q.; Stevens, C.J. Between Domestication and Civilization: The Role of Agriculture and Arboriculture in the Emergence of the First Urban Societies. *Veg. Hist. Archaeobotany* **2019**, *28*, 263–282. [[CrossRef](#)]
26. Djanibekov, U.; Villamor, G.B.; Dzhakypbekova, K.; Chamberlain, J.; Xu, J. Adoption of Sustainable Land Uses in Post-Soviet Central Asia: The Case for Agroforestry. *Sustainability* **2016**, *8*, 1030. [[CrossRef](#)]
27. Sidle, R.C.; Khan, A.A.; Caiserman, A.; Qadamov, A.; Khojazoda, Z. Food security in high mountains of Central Asia: A broader perspective. *BioScience* **2023**, *73*, 347–363. [[CrossRef](#)]
28. Rajametov, S.; Abdullaev, S. Horticulture Research In Central Asia: A review of Papers from Scopus Database Published for The Period of 2000–2020. Available online: <https://www.preprints.org/manuscript/202202.0325/v1> (accessed on 11 September 2023).
29. Jarvis, D.I.; Brown, A.H.D.; Cuong, P.H.; Collado-Panduro, L.; Latournerie-Moreno, L.; Gyawali, S.; Tanto, T.; Sawadogo, M.; Mar, I.; Sadiki, M.; et al. A Global Perspective of the Richness and Evenness of Traditional Crop-Variety Diversity Maintained by Farming Communities. *Proc. Natl. Acad. Sci. USA* **2008**, *105*, 5326–5331. [[CrossRef](#)]
30. Olson, D.M.; Dinerstein, E.; Wikramanayake, E.D.; Burgess, N.D.; Powell, G.V.N.; Underwood, E.C.; D'Amico, J.A.; Itoua, I.; Strand, H.E.; Morrison, J.C.; et al. Terrestrial Ecoregions of the World: A New Map of Life on Earth. *Bioscience* **2001**, *51*, 933–938. [[CrossRef](#)]
31. Jarvis, D.I.; Turdieva, M.K.; Campilian, D.M. *Crop Genetic Diversity to Reduce Pest and Disease Pressures on On-Farm: Participatory Diagnosis Guidelines. Version I*; Bioversity International: Rome, Italy, 2006; 101p.
32. Bernis-Fonteneau, A.; Alcadi, R.; Frangella, M.; Jarvis, D.I. Scaling Up Pro-Poor Agrobiodiversity Interventions as a Development Option. *Sustainability* **2023**, *15*, 10526. [[CrossRef](#)]
33. Bolker, B.M.; Brooks, M.E.; Clark, C.J.; Geange, S.W.; Poulsen, J.R.; Stevens, M.H.H.; White, J.S.S. Generalized linear mixed models: A practical guide for ecology and evolution. *Trends Ecol. Evol.* **2009**, *24*, 127–135. [[CrossRef](#)]
34. Fick, S.E.; Hijmans, R.J. WorldClim 2: New 1-km Spatial Resolution Climate Surfaces for Global Land Areas. *Int. J. Climatol.* **2017**, *37*, 4302–4315. [[CrossRef](#)]
35. R Core Team. *R: A Language and Environment for Statistical Computing*; R Foundation for Statistical Computing: Vienna, Austria, 2023; Available online: <https://www.R-project.org/> (accessed on 11 September 2023).
36. Bates, D.; Mächler, M.; Bolker, B.; Walker, S. Fitting Linear Mixed-Effects Models Using lme4. *J. Stat. Softw.* **2015**, *67*, 1–48. [[CrossRef](#)]
37. Bartoń, K. MuMIn: Multi-Model Inference. R Package Version 1.47.5. 2023. Available online: <https://CRAN.R-project.org/package=MuMIn> (accessed on 11 September 2023).
38. Bisht, I.S.; Mehta, P.S.; Bhandari, D.C. Traditional crop diversity and its conservation on-farm for sustainable agricultural production in Kumaon Himalaya of Uttaranchal state: A case study. *Genet. Resour. Crop Evol.* **2007**, *54*, 345–357. [[CrossRef](#)]

39. Barry, M.B.; Pham, J.L.; Noyer, J.L.; Courtois, B.; Billot, C.; Ahmadi, N. Implications for in situ genetic resource conservation from the ecogeographical distribution of rice genetic diversity in Maritime Guinea. *Plant Genet. Resour.* **2007**, *5*, 45–54. [[CrossRef](#)]
40. Bhandari, B. Summer Rainfall Variability and the Use of Rice (*Oryza Sativa* L.) Varietal Diversity for Adaptation: Farmers' Perceptions and Responses in Nepal. Master's Thesis, CBM Swedish Biodiversity Centre, Uppsala, Sweden, 2009.
41. Bellon, M.R. The dynamics of crop infraspecific diversity: A conceptual framework at the farmer level. *Econ. Bot.* **1996**, *50*, 26–39. [[CrossRef](#)]
42. Klein, A.-M.; Vaissière, B.E.; Cane, J.H.; Steffan-Dewenter, I.; Cunningham, S.A.; Kremen, C.; Tscharntke, T. Importance of pollinators in changing landscapes for world crops. *Proc. R. Soc. B Biol. Sci.* **2007**, *274*, 303–313. [[CrossRef](#)]
43. Rai, R.; Joshi, S.; Roy, S.; Singh, O.; Samir, M.; Chandra, A. Implications of Changing Climate on Productivity of Temperate Fruit Crops with Special Reference to Apple. *J. Hortic.* **2015**, *2*, 135. [[CrossRef](#)]
44. Richards, A.J. Does Low Biodiversity Resulting from Modern Agricultural Practice Affect Crop Pollination and Yield? *Ann. Bot.* **2001**, *88*, 165–172. [[CrossRef](#)]
45. Rodrigo, J. Spring frosts in deciduous fruit trees—Morphological damage and flower hardiness. *Sci. Hortic.* **2000**, *85*, 155–173. [[CrossRef](#)]
46. Veteto, J.R.; Carlson, S.B. Climate change and apple diversity: Local perceptions from Appalachian North Carolina. *J. Ethnobiol.* **2014**, *34*, 359–382. [[CrossRef](#)]
47. Koskela, J.; Buck, A.; Cros, T. Proceedings of the conference EUFORGEN Climate Change and Forest Genetic Diversity: Implications for Sustainable Forest Management in Europe, Paris, France, 15–16 March 2006.
48. Orlandi, F.; Marrapodi, S.; Proietti, C.; Ruga, L.; Fornaciari, M. Ecosystem Functions of Fruit Woody Species in an Urban Environment. *Environ. Monit. Assess.* **2023**, *195*, 118. [[CrossRef](#)] [[PubMed](#)]
49. Thant, A.A.; Teutscherova, N.; Vazquez, E.; Kalousova, M.; Phyo, A.; Singh, R.K.; Lojka, B. On-farm rice diversity and farmers' preferences for varietal attributes in Ayeyarwady Delta, Myanmar. *J. Crop Improv.* **2020**, *34*, 549–570. [[CrossRef](#)]
50. Witcombe, J.R.; Hollington, P.A.; Howarth, C.J.; Reader, S.; Steele, K.A. Breeding for abiotic stresses for sustainable agriculture. *Philos. Trans. R. Soc. B Biol. Sci.* **2008**, *363*, 703–716. [[CrossRef](#)]
51. Finckh, M.R.; Wolfe, M.S. Diversification strategies. In *The Epidemiology of Plant Disease*; Cooke, B.M., Jones, D.G., Kaye, B., Eds.; Springer: New York, NY, USA, 2006; pp. 269–308.
52. Mulumba, J.V.; Nankya, R.; Adokorach, J.; Kiwuka, C.; Fadda, C.; De Santis, P.; Jarvis, D.I. A risk-minimizing argument for traditional crop varietal diversity use to reduce pest and disease damage in agricultural ecosystems of Uganda. *Agric. Ecosyst. Environ.* **2012**, *157*, 70–86. [[CrossRef](#)]
53. Garrett, K.A.; Mundt, C.C. Epidemiology in mixed host populations. *Phytopathology* **1999**, *89*, 984–990. [[CrossRef](#)]
54. Forsline, P.L.; Aldwinckle, H.S. Evaluation of *Malus Sieversii* Seedling Populations for Disease Resistance and Horticultural Traits. *Acta Hortic.* **2004**, *663*, 529–534. [[CrossRef](#)]
55. Miller, D.D. Evaluation of *Malus Sieversii* Seedlings from Kazakhstan for Disease Resistance and Time of Leafing. *Acta Hortic.* **2004**, *663*, 535–538. [[CrossRef](#)]
56. Fazio, G.; Aldwinckle, H.S.; Volk, G.M.; Richards, C.M.; Janisiewicz, W.J.; Forsline, P.L. Progress in Evaluating *Malus Sieversii* for Disease Resistance and Horticultural Traits. *Acta Hortic.* **2009**, *814*, 59–66. [[CrossRef](#)]
57. Jurick, W.M.; Janisiewicz, W.J.; Saftner, R.A.; Vico, I.; Gaskins, V.L.; Park, E.; Forsline, P.L.; Fazio, G.; Conway, W.S. Identification of Wild Apple Germplasm (*Malus* Spp.) Accessions with Resistance to the Postharvest Decay Pathogens *Penicillium expansum* and *Colletotrichum acutatum*. *Plant Breed.* **2011**, *130*, 481–486. [[CrossRef](#)]
58. Sestras, A.F.; Pamfil, D.; Dan, C.; Bolboaca, S.D.; Jäntschi, L.; Sestras, R.E. Possibilities to Improve Apple Scab (*Venturia inaequalis* (Cke.) Wint.) and Powdery Mildew [*Podosphaera leucotricha* (Ell. et Everh.) Salm.] Resistance on Apple by Increasing Genetic Diversity Using Potentials of Wild Species. *Australian J. Crop Sci.* **2011**, *5*, 748–755.
59. Didelot, F.; Brun, L.; Parisi, L. Effects of Cultivar Mixtures on Scab Control in Apple Orchards. *Plant Pathol.* **2007**, *56*, 1014–1022. [[CrossRef](#)]
60. Demestihias, C.; Plénet, D.; Génard, M.; Raynal, C.; Lescourret, F. Ecosystem Services in Orchards. *A Review. Agron. Sustain. Dev.* **2017**, *37*, 12. [[CrossRef](#)]
61. Montanaro, G.; Xiloyannis, C.; Nuzzo, V.; Dichio, B. Orchard Management, Soil Organic Carbon and Ecosystem Services in Mediterranean Fruit Tree Crops. *Sci. Hortic.* **2017**, *217*, 92–101. [[CrossRef](#)]
62. Van Dusen, M.E.; Dennis, E.; Ilyasov, J.; Lee, M.; Treshkin, S.; Smale, M. Social Institutions and Seed Systems: The Diversity of Fruits and Nuts in Uzbekistan. In *Valuing Crop Biodiversity: On-Farm Genetic Resources and Economic Change*; Smale, M., Ed.; IPGRI: Maccaresse, Italy, 2006; p. 192, ISBN 9780851990835.
63. Kerven, C.; Robinson, S.; Behnke, R. Pastoralism at Scale on the Kazakh Rangelands: From Clans to Workers to Ranchers. *Front. Sustain. Food Syst.* **2021**, *4*, 590401. [[CrossRef](#)]
64. Haider, L.J.; Boonstra, W.J.; Akobirshoeva, A.; Schlüter, M. Effects of development interventions on biocultural diversity: A case study from the Pamir Mountains. *Agric. Hum. Values* **2020**, *37*, 683–697. [[CrossRef](#)]
65. Galluzzi, G.; Eyzaguirre, P.; Negri, V. Home Gardens: Neglected Hotspots of Agro-Biodiversity and Cultural Diversity. *Biodivers. Conserv.* **2010**, *19*, 3635–3654. [[CrossRef](#)]
66. Eyzaguirre, P.B.; Linares, O.F. *Home Gardens and Agrobiodiversity*; Smithsonian Book: Washington, DC, USA, 2004.

67. Galhena, D.H.; Freed, R.; Maredia, K.M. Home Gardens: A Promising Approach to Enhance Household Food Security and Wellbeing. *Agric. Food Secur.* **2013**, *2*, 8. [[CrossRef](#)]
68. Lerman, Z.; Sedik, D. *Sources of Agricultural Productivity Growth in Central Asia: The Case of Tajikistan and Uzbekistan*; FAO Regional Office for Europe and Central Asia, Policy Studies on Rural Transition, No. 2009-5; FAO: Rome, Italy, 2009.
69. Pushpakumara, G.; Sokolow, J.; Sthapit, B.; Sujarwo, W.; Hunter, D. Keeping It Close to Home. In *Home Gardens for Improved Food Security and Livelihoods*; Routledge: London, UK, 2020; pp. 46–77, ISBN 9781315471778.

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.