

**Supplementary Table S1:** Prevalence of *Toxoplasma gondii* infection in TBI according to sociodemographic characteristics.

Characteristic	Overall N = 160	Negative N = 108	Positive N = 52	P-value
<b>Age</b>	32 (26, 39)	32 (27, 38)	30 (23, 39)	0.6
<b>Gender</b>				0.072
Female	16 (10%)	14 (13%)	2 (3.8%)	
Male	144 (90%)	94 (87%)	50 (96%)	
<b>Profession</b>				0.5
Bike riders	43 (27%)	28 (26%)	15 (29%)	
Employee in service	30 (19%)	21 (19%)	9 (17%)	
Manual workers	24 (15%)	16 (15%)	8 (15%)	
Others	26 (16%)	15 (14%)	11 (21%)	
Student	16 (10%)	10 (9.3%)	6 (12%)	
Unemployed	21 (13%)	18 (17%)	3 (5.8%)	
<b>Classification of TBI</b>				
Mild		42 (39%)	24 (46%)	0.7
Moderate		38 (35%)	17 (33%)	
Severe		28 (26%)	11 (21%)	

Median (IQR); n (%) *Wilcoxon rank sum test*; *Pearson's Chi-squared test*

TBI: Traumatic brain injury