

Proceeding Paper

Impact of Water Deficit on Primary Metabolism at the Whole Plant Level in Bread Wheat Grown under Elevated CO₂ and High Temperature at Different Developmental Stages †

Nara Bueno-Ramos ^{1,‡}, Ana I. González-Hernández ^{1,‡}, Emilio L. Marcos-Barbero ¹,
Jon Miranda-Apodaca ^{1,2}, Ouardia Bendou ¹, Ismael Gutiérrez-Fernández ¹, Juan B. Arellano ¹
and Rosa Morcuende ^{1,*}

- ¹ Institute of Natural Resources and Agrobiological Sciences (IRNASA), Consejo Superior de Investigaciones Científicas (CSIC), 37008 Salamanca, Spain; nara.bueno@irnasa.csic.es (N.B.-R.); ana.gonzalez@irnasa.csic.es (A.I.G.-H.); emiliol.marcos@irnasa.csic.es (E.L.M.-B.); jon.miranda@ehu.es (J.M.-A.); ouardia.bendou@irnasa.csic.es (O.B.); ismael.gutierrez@irnasa.csic.es (I.G.-F.); juan.arellano@irnasa.csic.es (J.B.A.)
- ² Department of Plant Biology and Ecology, Faculty of Science and Technology, University of the Basque Country (UPV/EHU), Apdo. 644, 48080 Bilbao, Spain
- * Correspondence: rosa.morcuende@irnasa.csic.es
- † Presented at the 1st International Online Conference on Agriculture—Advances in Agricultural Science and Technology (IOCAG2022), 10–25 February 2022; Available online: <https://iocag2022.sciforum.net/>.
- ‡ These authors contributed equally to this work.



Citation: Bueno-Ramos, N.; González-Hernández, A.I.; Marcos-Barbero, E.L.; Miranda-Apodaca, J.; Bendou, O.; Gutiérrez-Fernández, I.; Arellano, J.B.; Morcuende, R. Impact of Water Deficit on Primary Metabolism at the Whole Plant Level in Bread Wheat Grown under Elevated CO₂ and High Temperature at Different Developmental Stages. *Chem. Proc.* **2022**, *10*, 6. <https://doi.org/10.3390/IOCAG2022-12232>

Academic Editor: Bin Gao

Published: 10 February 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Predicted increases in the atmospheric CO₂ concentration and the earth's mean surface temperature will be accompanied by a higher incidence of drought events. These environmental changes are likely to adversely affect crop productivity and quality, including wheat, an essential food in the human diet. We investigated the primary C-N metabolism response to drought stress at the whole-plant level and its dependence on plant development in bread wheat grown under combined elevated CO₂ and temperature. With this aim, the content of carbohydrates, nitrate, proteins and amino acids, together with the biomass were assessed in flag leaves and roots of wheat grown in controlled environment chambers at both ear emergence and anthesis stages. Multifactorial analysis revealed that the organ was the main factor explaining data variation. The physiological and biochemical traits in the flag leaves were more affected by drought than growth stage, leading to an accumulation of soluble carbohydrates, nitrate and amino acids. By contrast, roots were affected by the developmental stages but not by the treatment. The root content of fructose, glucose, starch and amino acids was higher at ear emergence than anthesis, whereas the accumulation of sucrose, fructans, proteins and nitrate increased at the latest growth stage. This study provides new insights into the reprogramming of primary metabolism at whole plant level throughout the development in response to the future climate scenario, which could help to select traits ensuring sustainable food production systems that strengthen capacity for adaptation to climate change following the Sustainable Development Goals of 2030 Agenda.

Keywords: wheat; climate change; drought; carbohydrates; N compounds; leaves; roots

1. Introduction

Bread wheat (*Triticum aestivum* L.) is the third largest crop in the world, feeding more than 4.5 billion people worldwide [1]. The necessity for increasing food supply by about 70% by 2050 [2], together with the detrimental effects of climate change on global crop production and quality, reveals that more efforts to improve wheat performance under sustainable conditions are required to ensure food security. The atmospheric CO₂ concentrations are expected to rise to 1000 ppm by the end of this century and the Earth's mean surface temperature is also expected to rise by 2.6 and 4.8 °C. These environmental

changes may lead to an increase in the frequency of other climate events such as irregular rainfalls and drought, which may spell serious threats to crop yield, including wheat [3–7].

Drought stress has an important impact on wheat performance throughout its different developmental stages, being the reproductive growth stage the most sensitive [8–10]. Morpho–physiological and biochemical changes in leaves and roots provide protection against drought stress, decreasing water losses [11–13]. One of the physiological responses include the stomata closure to avoid water loss through transpiration, which seems to be related to either hydraulic or chemical signals mediated by the abscisic acid, but may also lead to a limitation in carbon fixation and nutrient absorption by the roots [14–17]. At biochemical level, the osmotic adjustment counters the effect of water deficit by accumulating different osmolytes such as carbohydrates or amino acids [18,19]. Nevertheless, elevated CO₂ may mitigate the inhibition of photosynthesis under drought stress improving water use efficiency because a synergistic effect of both factors on stomatal closure [20]. These responses reveal that the regulation of primary metabolism is crucial for drought stress tolerance in higher plants [21]. Thus, the aim of this work was to assess the physiological and biochemical changes in the flag leaves and roots of wheat plants subjected to drought stress under simultaneous increases in atmospheric CO₂ and temperature in interaction with its developmental stages, including ear emergence and anthesis. The results of our study will help to select traits ensuring sustainable food production systems that strengthen the capacity for the adaptation to future climate change.

2. Methods

2.1. Plant Material and Growing Conditions

Seeds of bread wheat (*Triticum aestivum* L. var. Gazul), a high-yielding variety in Castilla y León, were sown in 2-L pots containing a mixture of perlite:vermiculite (3:1, *v/v*) and grown in controlled environmental growth chambers (3.6 m length × 4.8 m width × 2.4 m height) under combined elevated CO₂ (900 ppm) and temperature (4 °C above current temperatures of Salamanca region, 26 °C). The daily cycle was 16/8 h light/dark, with a light intensity of 400 μmol m^{−2} s^{−1} and relative humidity of 40%/60% day/night. At the beginning of the experiment, the maximum soil volumetric water content of each pot was evaluated as the difference between pot weight after watering with the excess water drained (field capacity) and the pot dry weight. Pots with two seeds per pot were irrigated from the seed germination until the tillering stage. After that, the water deficit was imposed and one half of the plants was maintained under well-watered conditions until the end of the experiment, whereas the other half was subjected to water deficit conditions (65% of field capacity). Plants were watered with a nutrient solution twice per week and with water once per week to avoid excessive salt accumulation. The nutrient solution contained NO₃[−] and it was supplied to a concentration of 10 mM provided by 6 mM KNO₃ and 2 mM Ca(NO₃)₂, while the other macro- and micronutrients were supplied as described by Vicente et al. [22].

2.2. Sampling Procedure

Plant material was harvested at ear emergence (Zadoks stage 59) and anthesis (Zadoks stage 65), respectively [23]. Flag leaves of the main stems and roots were harvested 3–4 h after the onset of the photoperiod. Each leaf was cut and immediately plunged into liquid N₂ under illumination. The roots were separated from the aerial part of the plant and they were collected after washing with water to remove substrate residues and nutrient solution, blotted dry on tissue paper, and quickly immersed in liquid N₂. The samples were stored at −80 °C until use for analyses. The experiments included a total of four replicates for each treatment and developmental growth stage.

2.3. C-N Metabolites Quantification

Frozen leaf and root plant material was used for ethanol/water extraction for carbohydrate determination according to Morcuende et al. [24]. Glucose, fructose, sucrose and

fructans were measured in the ethanol/water extracts with a spectrophotometric assay coupled to NADP⁺ reduction as described by Morcuende et al. [24], while the starch content was measured in the insoluble residue. Total amino acids content was also determined in the ethanol/water extracts by the ninhydrin colorimetric method, as well as NO₃[−] and protein contents following the methods described by Vicente et al. [25].

2.4. Determination of Biomass Parameters

At ear emergence, plants were harvested to determine the fresh (FW) and dry matter (DW) accumulation. The aerial part of the plants was separated from the roots. The dry weight of each of the parts separated from the plants was recorded after drying in an oven at 60 °C for 48 h. Humidity (H) and leaf dry matter content (LDMC) were estimated as $H = (FW - DW)/FW$ and $LDMC = DW/FW$, respectively.

2.5. Statistical Analyses

Descriptive analyses and graphs of variables and individuals from multifactorial analyses were carried out as described by Marcos-Barbero et al. [26].

3. Results and Discussion

Bread wheat (*Triticum aestivum* L.) is one of the most widely grown staple crop in the world threatened by the future climate change conditions. The impact of these conditions on wheat varies across the world, so specific adaptation strategies need to be studied in order to increase crop adaptation to the global climate change [27]. Furthermore, wheat growth and productivity under drought conditions needs an in-depth study of the physiological and biochemical mechanisms involved in the response to the conditions anticipated by global climate change by the end of this century and its dependence on development at the whole plant level. In this work, wheat plants were subjected to drought stress under combined elevated atmospheric CO₂ and temperature and flag leaves and roots were harvested at ear emergence and anthesis. Multifactorial analysis revealed that the organ was the main factor explaining data variation, whereas the developmental stage and drought treatment induced lower effects on the studied parameters (Figure 1A). The correlation circle showed that the flag leaves were associated with higher C and N metabolites, while the roots were mostly associated with fresh and dry weights and humidity percentage (Figure 1B,C). In accordance with these results, Kumar et al. [21] reported that sugars and amino acids increased significantly in the aerial organs of different crops under drought stress. In addition, greater growth reduction was observed in shoots than in roots of barley plants subjected to drought [28].

When the flag leaves and root organs were studied separately, the results reflected that the physiological and biochemical traits in the flag leaves were more affected by drought than by the developmental growth stage, leading to an accumulation of soluble carbohydrates, nitrate and amino acids (Figure 2A–C). These findings are in accordance with those reported by Guo et al. [29], suggesting that sugars, organic acids and amino acids played a pivotal role to enhance drought tolerance of the aerial part of the plant. Furthermore, enhanced accumulation of water-soluble carbohydrates (glucose, fructose, sucrose and fructans) was found in drought tolerant wheat genotypes [30,31], whereas a reduction in sucrose, fructose, mannose and tagatose seems to be associated with sensitive genotypes under water deficit stress [32]. This fact suggests that sugars may function as compatible solutes contributing to cell osmoregulation and the maintenance of foliar water content, providing a solvation shell around proteins, stabilizing subcellular structures and protecting membrane integrity [32–34]. In addition, amino acids content was also increased in wheat plants grown under water deficit conditions, in which proline, tryptophan, leucine, isoleucine and valine were notably accumulated in leaf tissues [32], protecting cellular structures during dehydration and contributing to redox balance [35]. The increase in NO₃[−] in leaves could be explained by the inability of plants to reduce nitrate, being used as an osmolyte receiving a better response to drought stress [36]. By contrast, roots were

mainly affected by the developmental stages, but not by the treatment (Figure 2D). The root content of fructose, glucose, starch and amino acids was higher at ear emergence than anthesis, whereas the accumulation of sucrose, fructans, proteins and nitrate increased at the latest growth stage (Figure 2E,F). The increase in glucose and amino acids enhanced the capacity to reduce and transport nitrate to other organs [37]. At anthesis, the increase in nitrate levels could be related to a limitation of nitrate reduction with the evolution of plant development [38]. Moreover, the increase in sucrose levels has been previously reported by Shao et al. [39].




Figure 1. Multifactorial analyses for the C and N metabolites (C-met and N-met) and biomass parameters evaluated in the flag leaves and roots (L; R) of wheat plants subjected to different water regimes (d: drought; w: water) under elevated CO₂ and high temperature at different developmental stages (E: ear emergence; A: anthesis). The considered factors were organ, treatment and stage. The statistical analyses included groups representation (A), individual factor map (B) and correlation circle (C).




Figure 2. Multifactorial analyses for the primary C and N metabolites (C-met: glucose, fructose, sucrose and starch; N-met: nitrate, amino acids and proteins) and biomass parameters (DW, FW, LDMC and H) evaluated in the flag leaves and roots of wheat plants subjected to different water regimes (D: drought; W: water) under elevated CO₂ and high temperature at different developmental stages (EEm: ear emergence; Ant: anthesis). The statistical analyses for each wheat organ included groups representation (A,D), correlation circle (B,E) and MFA (C,F).

4. Conclusions

The comparative analyses of the physiological and biochemical traits in the flag leaves and roots of wheat plants under concurrent drought stress, elevated CO₂ and high temperature showed that leaves were more affected by the drought treatment than by the growth stage, leading to an accumulation of soluble carbohydrates, nitrate and amino acids, whereas roots were more affected by the developmental stages accumulating more fructose, glucose, starch and amino acids at ear emergence rather than anthesis, whereas the accumulation of sucrose, fructans, proteins and nitrate increased at the latest growth stage. Thus, this study provides new evidence of the shift of primary C-N metabolism at whole plant level at ear emergence and anthesis developmental stages in response to the predicted elevated CO₂ and temperatures, and more lasting drought events. This could help us to select specific traits that are useful for wheat breeding programs to ensure food production systems with more resilient wheat varieties to climate change conditions following the Sustainable Development Goals of 2030 Agenda.

Author Contributions: J.B.A. and R.M. conceived and designed the experiment. N.B.-R., A.I.G.-H., E.L.M.-B., J.M.-A., J.B.A. and R.M. carried out the experiments and performed the measurements of the different assays. N.B.-R., A.I.G.-H. and E.L.M.-B. carried out the statistical data analyses. O.B. and I.G.-F. contributed to the experimental work. N.B.-R. and A.I.G.-H. wrote the proceeding paper. J.B.A. and R.M. revised the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: This research was supported by the Spanish National R&D&I Plan (Project PID2019-107154RB-I00; AGL2016-79589-R) and the Regional Government of Castilla y León (CSI260P20). The Project “CLU—2019—05—IRNASA/CSIC Unit of Excellence” funded by Junta de Castilla y León and co-financed by the European Union (ERDF “Europe drives our growth”) is also acknowledged. Ms. N. Bueno-Ramos and Mr. E. L. Marcos-Barbero are the recipients of a pre-doctoral contract from the Junta de Castilla y León (E-37-2020-0042432 and E-37-2017-0066125, respectively); Dr. A.I.

González-Hernández holds a JCyL postdoctoral contract funded by the project CSI260P20, Ms. O. Bendou holds an Algerian Government fellowship; Mr. I. Gutiérrez-Fernández holds a contract funded by the project CLU-2019-05.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Data are available upon request from the corresponding authors.

Acknowledgments: We want to thank A. Verdejo and M.A. Boyero for plant growth and sample preparation.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Asseng, S.; Foster, I.; Turner, N.C. The impact of temperature variability on wheat yields. *Glob. Chang. Biol.* **2011**, *17*, 997–1012. [CrossRef]
- FAO. How to Feed the World in 2050. High-Level Experts Forum 2009. Available online: https://www.fao.org/fileadmin/templates/wsfs/docs/expert_paper/How_to_Feed_the_World_in_2050.pdf (accessed on 30 December 2021).
- Porter, J.R.; Semenov, M.A. Crop responses to climatic variation. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **2005**, *360*, 2021–2035. [CrossRef] [PubMed]
- Weisheimer, A.; Palmer, T.N. Changing frequency of occurrence of extreme seasonal temperatures under global warming. *Geophys. Res. Lett.* **2005**, *32*, L20721. [CrossRef]
- Semenov, M.A.; Shewry, P.R. Modelling predicts that heat stress, not drought, will increase vulnerability of wheat in Europe. *Sci. Rep.* **2011**, *1*, 66. [CrossRef]
- Lobell, B.D.; Sibley, A.; Ortiz-Monasterio, J.I. Extreme heat effects on wheat senescence in India. *Nat. Clim. Chang.* **2012**, *2*, 186–189. [CrossRef]
- Asseng, S.; Ewert, F.; Martre, P.; Rötter, R.P.; Lobell, D.B.; Cammarano, D.; Kimball, B.A.; Ottman, M.J.; Wall, G.W.; White, J.W.; et al. Rising temperatures reduce global wheat production. *Nat. Clim. Chang.* **2015**, *5*, 143–147. [CrossRef]
- Barnabas, B.; Jäger, K.; Fehér, A. The effect of drought and heat stress on reproductive processes in cereals. *Plant Cell Environ.* **2008**, *31*, 11–38. [CrossRef]
- Farooq, M.; Hussain, M.; Siddique, K.H.M. Drought stress in wheat during flowering and grain-filling periods. *CRC Crit. Rev. Plant Sci.* **2014**, *33*, 331–349. [CrossRef]
- Senapati, N.; Stratonovitch, P.; Paul, M.J.; Semenov, M.A. Drought tolerance during reproductive development is important for increasing wheat yield potential under climate change in Europe. *J. Exp. Bot.* **2019**, *70*, 2549–2560. [CrossRef]
- Izanloo, A.; Condon, A.G.; Langridge, P.; Tester, M.; Schnurbusch, T. Different mechanisms of adaptation to cyclic water stress in two South Australian bread wheat cultivars. *J. Exp. Bot.* **2008**, *59*, 3327–3346. [CrossRef]
- Comas, L.H.; Becker, S.R.; Cruz, V.M.; Byrne, P.F.; Dierig, D.A. Root traits contributing to plant productivity under drought. *Front. Plant Sci.* **2013**, *4*, 442. [CrossRef] [PubMed]
- Fahad, S.; Bajwa, A.A.; Nazir, U.; Anjum, S.A.; Farooq, A.; Zohaib, A.; Sabia, S.; Nasim, W.; Adkins, S.; Saud, S.; et al. Crop production under drought and heat stress: Plant responses and management options. *Front. Plant Sci.* **2017**, *8*, 1147. [CrossRef] [PubMed]
- Steudle, E. Water uptake by roots: Effects of water deficit. *J. Exp. Bot.* **2000**, *51*, 1531–1542. [CrossRef] [PubMed]
- Rizhsky, L.; Liang, H.; Mittler, R. The combined effect of drought stress and heat shock on gene expression in tobacco. *Plant Physiol.* **2002**, *130*, 1143–1151. [CrossRef]
- Mahajan, S.; Tuteja, N. Cold, salinity and drought stresses: An overview. *Arch. Biochem. Biophys.* **2005**, *444*, 139–158. [CrossRef]
- Monneveux, P.; Rekika, D.; Acevedo, E.; Merah, O. Effect of drought on leaf gas exchange, carbon isotope discrimination, transpiration efficiency and productivity in field grown durum wheat genotypes. *Plant Sci.* **2006**, *170*, 867–872. [CrossRef]
- Slama, I.; Abdelly, C.; Bouchereau, A.; Flower, T.; Savouré, A. Diversity, distribution and roles of osmoprotective compounds accumulated in halophytes under abiotic stress. *Ann. Bot.* **2015**, *115*, 433–447. [CrossRef]
- Sharma, A.; Shahzad, B.; Rehman, A.; Bhardwaj, R.; Landi, M.; Zheng, B. Response of phenylpropanoid pathway and the role of polyphenols in plants under abiotic stress. *Molecules* **2019**, *24*, 2452. [CrossRef]
- Medina, S.; Vicente, R.; Amador, A.; Araus, J.L. Interactive effects of elevated [CO₂] and water stress on physiological traits and gene expression during vegetative growth in four durum wheat genotypes. *Front. Plant Sci.* **2007**, *7*, 1738.
- Kumar, M.; Kumar-Patel, M.; Kumar, N.; Bajpai, A.B.; Siddique, K.H.M. Metabolomics and molecular approaches reveal drought stress tolerance in plants. *Int. J. Mol. Sci.* **2021**, *22*, 9108. [CrossRef]
- Vicente, R.; Pérez, P.; Martínez-Carrasco, R.; Usadel, B.; Kostadinova, S.; Morcuende, R. Quantitative RT-PCR platform to measure transcript levels of C and N metabolism-related genes in durum wheat: Transcript profiles in elevated [CO₂] and high temperature at different levels of N supply. *Plant Cell Physiol.* **2015**, *56*, 1556–1573. [CrossRef] [PubMed]
- Zadoks, J.C.; Chang, T.; Konzak, C.F. A decimal code for the growth stages of cereals. *Weed Res.* **1974**, *14*, 415–421. [CrossRef]

24. Morcuende, R.; Kostadinova, S.; Pérez, P.; Martín del Molino, I.M.; Martínez-Carrasco, R. Nitrate is a negative signal for fructan synthesis, and the fructosyltransferase-inducing trehalose inhibits nitrogen and carbon assimilation in excised barley leaves. *New Phytol.* **2004**, *161*, 749–759. [[CrossRef](#)] [[PubMed](#)]
25. Vicente, R.; Pérez, P.; Martínez-Carrasco, R.; Feil, R.; Lunn, J.E.; Watanabe, M.; Arrivault, S.; Stitt, M.; Hoefgen, R.; Morcuende, R. Metabolic and transcriptional analysis of durum wheat responses to elevated CO₂ at low and high nitrate supply. *Plant Cell Physiol.* **2016**, *57*, 2133–2146. [[CrossRef](#)] [[PubMed](#)]
26. Marcos-Barbero, E.L.; Pérez, P.; Martínez-Carrasco, R.; Arellano, J.B.; Morcuende, R. Screening for higher grain yield and biomass among sixty bread wheat genotypes grown under elevated CO₂ and high-temperature conditions. *Plants* **2021**, *10*, 1596. [[CrossRef](#)] [[PubMed](#)]
27. Noleto Luz Pequeno, D.; Hernandez-Ochoa, I.M.; Reynolds, M.; Sonder, K.; Molero-Milan, A.; Robertson, R.; Lopes, M.S.; Xiong, W.; Kropff, M.; Asseng, S.; et al. Climate impact and adaptation to heat and drought stress of regional and global wheat production. *Environ. Res. Lett.* **2021**, *16*, 054070. [[CrossRef](#)]
28. Boudiar, R.; Casas, A.M.; Gioia, T.; Fiorani, F.; Nagel, K.A.; Igartua, E. Effects of low water availability on root placement and shoot development in landraces and modern barley cultivars. *Agronomy* **2020**, *10*, 134. [[CrossRef](#)]
29. Guo, R.; Shi, L.; Jiao, Y.; Li, M.; Zhong, X.; Gu, F.; Liu, Q.; Xia, X.; Li, H. Metabolic responses to drought stress in the tissues of drought-tolerant and drought-sensitive wheat genotype seedlings. *AoB Plants* **2018**, *10*, ply016. [[CrossRef](#)]
30. Goggin, D.E.; Setter, T.L. Fructosyltransferase activity and fructan accumulation during development in wheat exposed to terminal drought. *Funct. Plant Biol.* **2004**, *31*, 11–21. [[CrossRef](#)]
31. Wardlaw, I.F.; Willenbrink, J. Mobilization of fructan reserves and changes in enzyme activities in wheat stems correlate with water stress during kernel filling. *New Phytol.* **2000**, *148*, 413–422. [[CrossRef](#)]
32. Bowne, J.B.; Erwin, T.A.; Juttner, J.; Schnurbusch, T.; Langridge, P.; Bacic, A.; Roessner, U. Drought responses of leaf tissues from wheat cultivars of differing drought tolerance at the metabolite level. *Mol. Plant* **2012**, *5*, 418–429. [[CrossRef](#)] [[PubMed](#)]
33. Hoekstra, F.A.; Golovina, E.A.; Buitink, J. Mechanisms of plant desiccation tolerance. *Trends Plant Sci.* **2001**, *6*, 431–438. [[CrossRef](#)]
34. Anjum, S.A.; Xie, X.Y.; Wang, L.C.; Saleem, M.F.; Man, C.; Lei, W. Morphological, physiological and biochemical responses of plants to drought stress. *Afr. J. Agric. Res.* **2011**, *6*, 2026–2032.
35. Michaletti, A.; Naghavi, M.R.; Toorchi, M.; Zolla, L.; Rinalducci, S. Metabolomics and proteomics reveal drought-stress responses of leaf tissues from spring-wheat. *Sci. Rep.* **2018**, *8*, 5710. [[CrossRef](#)] [[PubMed](#)]
36. Miranda-Apodaca, J.; Agirresarobe, A.; Martínez-Goñi, X.S.; Yoldi-Achalandabaso, A.; Pérez-López, U. N metabolism performance in *Chenopodium quinoa* subjected to drought or salt stress conditions. *Plant Physiol. Biochem.* **2020**, *155*, 725–734. [[CrossRef](#)]
37. Oaks, A.; Hirel, B. Nitrogen metabolism in roots. *Annu. Rev. Plant Physiol.* **1985**, *36*, 345–365. [[CrossRef](#)]
38. Barillot, R.; Chambon, C.; Andrieu, B. CN-Wheat, a functional–structural model of carbon and nitrogen metabolism in wheat culms after anthesis. II. Model evaluation. *Ann. Bot.* **2016**, *118*, 1015–1031. [[CrossRef](#)]
39. Shao, K.; Bai, Z.; Li, M.; Yu, C.; Shao, J.; Sun, Y.; Li, G.; Zhang, S.; Wang, R. Sucrose metabolism enzymes affect sucrose content rather than root weight in sugar beet (*Beta vulgaris*) at different growth stages. *Sugar Tech.* **2020**, *22*, 504–517. [[CrossRef](#)]